Published in final edited form as:

Curr Opin Chem Biol. 2012 April ; 16(0): 45–53. doi:10.1016/j.cbpa.2012.03.004.

Iron sulfur cluster proteins and microbial regulation: implications for understanding tuberculosis

Vikram Saini¹, Aisha Farhana¹, Joel N Glasgow¹, and Adrie JC Steyn^{1,2,3,4,5}

¹ Department of Microbiology, University of Alabama at Birmingham, AL 35294, USA

² Centers for AIDS Research, University of Alabama at Birmingham, AL 35294, USA

³ Free Radical Biology, University of Alabama at Birmingham, AL 35294, USA

⁴ KwaZulu-Natal Research Institute for Tuberculosis and HIV, Durban 4001, South Africa

⁵ Department of Pathology, Nelson Mandela School of Medicine, University of KwaZulu-Natal, Durban 4001, South Africa

Abstract

All pathogenic and nonpathogenic microbes are continuously exposed to environmental or endogenous reactive oxygen and nitrogen species, which can critically effect survival and disease. Iron–sulfur [Fe–S] cluster containing prosthetic groups provide the microbial cell with a unique capacity to sense and transcriptionally respond to diatomic gases (e.g. NO and O₂) and redox-cycling agents. Recent advances in our understanding of the mechanisms for how the FNR and SoxR [Fe–S] cluster proteins respond to NO and O₂ have provided new insights into the biochemical mechanism of action of the *Mycobacterium tuberculosis* (*Mtb*) family of WhiB [Fe–S] cluster proteins. These insights have provided the basis for establishing a unifying paradigm for the *Mtb* WhiB family of proteins. *Mtb* is the etiologic agent for tuberculosis (TB), a disease that affects nearly one-third of the world's population.

Introduction

Iron sulfur clusters (Fe–S) represent one of nature's most ubiquitous, dynamic and likely most ancient prosthetic groups necessary to perform distinct cellular functions. In most Fe–S proteins, the clusters function as electron-transfer groups, but alternative functions have been described over the years including maintenance of protein structure, enzyme catalysis, metabolic regulation, regulation of gene expression and protein folding [1,2]. In this review, we will focus specifically on the *Mycobacterium tuberculosis* (*Mtb*) WhiB family of Fe–S cluster proteins.

Fe–S cluster assembly in microbes and sources of iron and sulfur

Fe–S cluster assembly in microbes is achieved *via* complex protein systems that construct nascent clusters on scaffold proteins and then transfer the cluster into recipient apo-proteins. Three genetic Fe–S cluster assembly systems are conserved among bacteria: the *nif* operon (nitrogen fixation), the *isc* operon (iron–sulfur clusters) and the *sufABCDSE* operon

(mobilization of sulfur) [3,4]. Many organisms possess more than one system, while the *suf* system is the only system in certain bacteria and cyanobacteria (reviewed in [2–4]).

The sulfur used by Fe–S systems is usually procured from L-cysteine with the help of cysteine desulfurylases such as NifS, IscS and SufSE. However, some archaea like methanococcus may use sulfide as a proximal sulfur donor [5*]. The source of iron for the Fe–S cluster, however, remains controversial [6]. Recent work suggests that Fe–S clusters are part of a cellular chelatable iron pool (CIP) [7,8**].

Prototype Fe–S cluster proteins: FNR and SoxR

Fumarate nitrate reductase (FNR)

FNR is involved primarily in the transition from the aerobic to anaerobic phase and regulates over 100 genes involved in various pathways. Under anaerobic conditions, FNR exists as a homodimer wherein each monomer contains one $[4\text{Fe–4S}]^{2+}$ cluster bound to four Cys residues. In the presence of O_2 , the FNR $[4\text{Fe–4S}]^{2+}$ cluster is converted to an $[2\text{Fe–2S}]^{2+}$ cluster, leading to the reversible dissociation of the dimer and loss of DNA binding activity (Figure 1) [9**]. FNR is a cytoplasmic enzyme and diffusion of O_2 into the cytoplasm takes place even at very low O_2 tension ($1\ \mu\text{M}$) leading to inactivation of $[4\text{Fe–4S}]^{2+}$ cluster and hence FNR [10]. However, the cell continues to produce FNR protein and reactivate it. The intracellular redox state plays an important role here as the presence of the major redox buffer glutathione ($E_{hc} = -250\ \text{mV}$ [10 mM]) acts as a reductant to reduce sulfur from persulfide (in a NifS-mediated cluster transfer reaction, S/HS^- ; $E^\circ = -260\ \text{mV}$), as well as Cys present in FNR [11]. Under oxic conditions, the S–S linked Cys ligands of apo-FNR are reduced to the thiol state for cluster incorporation. Thus, there is constant cycling of FNR between its three forms: apo-FNR, 4Fe-FNR and 2Fe-FNR [12]. The onset of anaerobiosis leads to an accumulation of 4Fe-FNR monomers which dimerize to produce the active transcription factor. The FNR $[4\text{Fe–4S}]^{2+}$ cluster is also responsive to NO to form protein-bound DNIC–FNR and serves as a functional switch in NO-responsive control of gene expression [13].

FNR regulates a large number of genes and operons, and the core set of target genes in *Escherichia coli* is fairly conserved across different bacterial species. The variation in so-called ‘extended regulons’ could be the outcome of niche-specific or host-specific adaptation of the organisms [14*].

SoxR

SoxR (named for superoxide response) is a constitutively expressed homodimeric protein of the MerR family of transcriptional regulators. Dimeric SoxR acts as a sensor-transmitter with its 2Fe–2S centers being reduced during normal growth and becoming oxidized during oxidative stress or by changes in intracellular NADPH levels (Figure 1) [15,16**]. The Fe–S center of SoxR is not required to maintain protein structure or DNA binding, as apo-SoxR attains a similar dimeric structure and binds to *soxS* DNA with the same affinity as that of holoenzyme [17]. Rather, redox changes induce structural alterations in the SoxR $[2\text{Fe–2S}]$ cluster that are communicated to the DNA-binding domain leading to distortion of *soxS* promoter DNA of oxidized SoxR [18].

Two prevailing models are emerging wherein either SoxR is directly activated by redox cycling compounds [19**,20] and controls the transcriptional activity of a few select genes, or the classical model wherein SoxR binds to *soxS*, and then SoxS in turn binds to more than 100 genes to modulate their expression [17]. However, despite the functional diversity of SoxR in different species, the biochemical activation mechanism appears to be conserved

across species. SoxR from *E. coli* and *Pseudomonas aeruginosa* have similar midpoint potentials (approximately -290 mV) [17] and might be kept in their reduced inactive state under physiological conditions by $\text{NADP}^+/\text{NADPH}$ ($E^\circ = -340$ mV) [15,16^{**}]. Binding to DNA shifts the midpoint potential of SoxR to $+200$ mV leading to a ~ 500 mV shift between the free and DNA-bound states of SoxR [21]. This implies that cellular thiols like glutathione could perhaps act as reductants for DNA-bound SoxR keeping it in its inactive form *in vivo*. Only in the presence of strong oxidants does DNA-bound SoxR actively regulate the genes associated with mitigation of oxidative stress and protection of the organism [21].

The *Mtb* WhiB protein family

Mtb WhiB proteins are the members of WhiB-like (Wbl) family that contain conserved Cys residues [Cys(x_n)Cys(x_2)Cys(x_5)Cys], which coordinate an Fe–S cluster. Members of the Wbl protein family are mostly restricted to *Actinobacteria* and were initially discovered in *Streptomyces* as various *wbl* mutants developed white colored aerial mycelium instead of the normal gray, and exhibited defects in sporulation of aerial hyphae [22]. However, mycobacterial Wbl proteins are evolutionarily closer to *Corneybacterium* and *Nocardia* and phylogenetic evidence suggest that they have been laterally acquired from plasmids and phages [23^{**}]. *Mtb* has seven members of WhiB family, namely, WhiB1 (Rv3219), WhiB2 (Rv3260c), WhiB3 (Rv3416), WhiB4 (Rv3681c), WhiB5 (Rv0022c), WhiB6 (Rv3862c) and WhiB7 (Rv3197A). The low sequence homology among these proteins and their restricted distribution within different mycobacterial lineages (e.g. the absence of WhiB5 in non-pathogenic species) implies that each WhiB member may perform diverse functions despite each containing a common Fe–S cluster [23^{**}]. Indeed, WhiB proteins in mycobacteria have been implicated in a plethora of functions including cell division (WhiB2), fatty acid metabolism and pathogenesis (WhiB3), oxidative stress (WhiB6) and antibiotic resistance (WhiB7) [24–27]. Most mycobacterial WhiB proteins harbor an AT hook-like DNA binding motif, consistent with the proposed role of these proteins as transcription factors. Garg *et al.* proposed these proteins to be general disulfide reductases since WhiB1 is able to reduce disulfide bonds in the (1,4)-glucan branching enzyme GlgB [28]. However, no such activity was observed in *Streptomyces* WhiD, a WhiB3 homolog, or in WhiB1 [29]. More recently, studies have shown that *Mtb* apo-WhiB1 functions as a transcriptional repressor [30].

As expected, Fe–S clusters of *Mtb* [30,31,32^{**},33^{*}] and *Streptomyces* [34] Wbl proteins show differential sensitivity and stability toward oxidative stress, including O_2 and NO exposure. *Streptomyces* WhiD and *Mtb* WhiB1 react with eight NO molecules per [4Fe–4S] cluster in a four-step reaction that is 10^4 -fold faster than their reaction with O_2 to form DNIC complexes [34]. These differential responses and susceptibilities of Wbl proteins to various environmental stimuli can be partially attributed to different oxidation states of the Fe in the cluster or the amino acid residues adjacent to the Fe–S cluster that modulate the redox potential of the cluster.

Mtb WhiB3 is the most studied member of WhiB family and represents the current paradigm (for a review see [23^{**}]) (Figure 1). Originally, *Mtb* WhiB3 was shown to physically interact with the principle sigma factor SigA, which suggested that Wbl proteins have DNA-binding activity [35]. Singh *et al.* [33^{*}] demonstrated that WhiB3 harbors an [4Fe–4S] cluster that reacts with NO and slowly with O_2 , and this contrasts with the rapid reaction of O_2 with FNR. From these studies, it was established that WhiB3 acts as a physiological sensor of NO and O_2 in *Mtb* and controls expression of genes involved in intermediary metabolism [32^{**}, 33^{*}]. More recently, the WhiB3 paradigm was further expanded by demonstrating that WhiB3 acts as a sensor of reductive stress in *Mtb*, and modulates lipid biosynthesis by binding to the upstream sequence of the virulence polyketide synthase genes *pks2* and *pks3*

[32^{••},36^{••}]. This is particularly important as a role for reductive stress in bacterial pathogenesis is unexplored [36^{••}]. While both apo-WhiB3 and holo-WhiB3 bind DNA, oxidized apo-WhiB3 demonstrates the strongest DNA-binding, illustrating the redox dependence of WhiB3. The conserved Cys residues in WhiB3 not only coordinate the Fe–S cluster, but also form intramolecular disulfide bonds following cluster loss. It is the formation of these bonds that enables apo-WhiB3 to form a strong protein–DNA complex [32^{••}]. This mechanism stands in contrast to that of *E. coli* FNR where cluster loss due to O₂ exposure leads to the formation of apo-FNR, which lacks DNA-binding activity [9^{••},37]. Another critical factor contributing toward the flexibility of the WhiB3 Fe–S cluster is the presence of multiple Arg residues (Cys₂₃-Arg₂₄-Cys₅₃-Arg₅₄-Arg₅₅-Cys₅₆-Cys₆₂-Arg₆₃) adjacent to the Fe–S cluster-coordinating Cys residues. At neutral pH, the Arg residues located close to Cys residues can significantly lower the *pK_a* of Cys, resulting in the formation of Cys thiolates, which are highly susceptible to oxidation, and loss of the Fe–S cluster. Not surprisingly, *Mtb whiB3* was highly upregulated when exposed to acidic stress [38].

Another member of the *Mtb* WhiB family, WhiB1, was also shown to possess a NO sensitive [4Fe–4S]²⁺ cluster, which may act as a NO sensor, at least *in vitro*. Exposure to NO converts holo-WhiB1 from a non-DNA-binding form into a DNA-binding form capable of regulating transcription. Only apo-WhiB1 and NO-treated holo-WhiB1 (but not holo-WhiB1) bind to the *whiB1* promoter region to repress its own transcription [30]. Similarly, *Mtb* apo-WhiB2 and its mycobacteriophage homologue apo-WhiBTM4 also exhibit strong DNA binding, as opposed to the holo-form of these proteins, which bind DNA weakly or not at all [27]. *Mtb* WhiB2 and WhiBTM4 bind to a conserved promoter sequence upstream of *whiB2* to regulate its transcription [27].

Bioinformatic analyses performed by us have revealed that *Mtb* contains at least 50 proteins that harbor different structural types of Fe–S clusters. The number and distribution of Fe–S cluster proteins are variable across different mycobacterial species and may be associated with intra-cellular adaptations (see Box 1). An important question is: How are these clusters shielded during aerobic growth? One possibility is that mycothiol, the major low-molecular-mass thiol present in mycobacteria, and thioredoxin play a protective role against cluster loss *in vitro* [29]. The redox machinery of mycobacteria is distinct from other intracellular pathogens as it lacks homologues of FNR, SoxR, ArcAB and OxyR (a pseudogene in *Mtb*), which could either protect the bacilli against host-generated oxidative stress or facilitate the transition from an aerobic to anaerobic environments. Furthermore, mycobacteria lack the glutathione system and use mycothiol as a redox buffer as well as ergothioneine. Examining the role of the WhiB proteins in the redox biology of *Mtb* may be critical to the understanding of *Mtb* virulence and persistence. Lastly, it is tempting to speculate that *Mtb* Wbbs play a role in virulence or persistence by controlling gene expression in response to host-generated NO or hypoxia, which are two host factors implicated in disease [39].

Emerging insights into Fe–S cluster containing proteins

Advances in protein sciences, computational biology and spectroscopy have led to the identification of new functions for Fe–S clusters. For example, the membrane-bound respiratory [NiFe]-hydrogenase of *Hydrogenovibrio marinus* catalyzes the oxidation of dihydrogen and harbors a unique [4Fe–3S] cluster coordinated by six Cys residues [40^{••}]. The presence of additional Cys provides oxygen tolerance to the organism through release of two electrons and a proton that reduces O₂ and prevents enzymatic inactivation [40^{••}]. Recently, Fe–S clusters have been shown to play a critical role in substrate specificity of assimilatory sulfonucleotide reductases such as APR (adenosine 5'-phosphosulfate reductase) and PAPR (3'-phosphoadenosine 5'-phosphosulfate reductase), which bind substrates that

differ by a single 3'-phosphate group despite similar sequences, structure, and thiol reaction chemistry [41*].

The *S. aureus* two component system AirSR (anaerobic iron sulfur cluster containing redox sensor regulator) represents an interesting case as it responds to O₂, H₂O₂ and NO *via* a redox active 2Fe–2S cluster in the sensor kinase, AirS [42]. AirS has a Cys arrangement (Cys-X₇-CysXCys-X₁₇-Cys) similar to that of SoxR and ferredoxins. Surprisingly, however, it harbors an N-terminal GAF domain, which is typically associated with heme that can bind NO, CO and O₂. Excess oxidation or reduction of AirS both lead to the loss of the Fe–S cluster with resultant loss of kinase activity [42].

Conclusions

During the past few years, the combination of biochemistry, biophysics and structural biology has significantly extended our knowledge of Fe–S cluster characterization, assembly and trafficking in model microorganisms. However, a substantial gap in the Fe–S cluster field is the absence of phylogenomic studies that focus on the genomes of non-model organisms such as *Mtb* (and many others), which is likely due to the pathogenic nature and slow growth of the organisms. Nonetheless, these non-model microbes represent unexplored avenues of research. For example, a fundamental challenge in the redox biology of *Mtb* is to understand how oxidative and nitrosative stress modulate *Mtb* disease. The role of the diatomic gases oxygen and NO, which are highly reactive toward Fe–S cluster proteins and affect redox homeostasis, are highly relevant to the disease caused by this global pathogen. For instance, *Mtb* persists in a dormant, drug-resistant state inside hypoxic granulomas (pO₂ < 1.99 mmHg), reactivation occurs in the oxygen rich upper lobe of the human lung, and a clear role for NO in TB has been demonstrated (for a review see [39]).

Intriguingly, although *Mtb* is an obligate aerobe, it appears that the number of Fe–S cluster motifs correlates with anaerobic microbes, rather than with aerobic microbes (see Box 1). The relative expression profiles of *Mtb* Fe–S cluster and Fe–S cluster assembly proteins under oxidative and nitrosative stress conditions, including upon macrophage infection, suggest that Fe–S cluster proteins are targeted by host reactive oxygen and nitrogen species to affect biogenesis and repair (Figure 2). An important finding was demonstrating for the first time that a Wbl member, *Mtb* WhiB3, binds DNA in a redox-dependent manner. *In vivo* DNA-binding studies under distinct environmental conditions should reveal important insights into the effect of the cytoplasmic redox state on WhiB3-mediated gene expression.

Important questions relevant to all microbes (especially the *Mtb* Wbl proteins) include: What is the *in vivo* redox state, and cellular reductants, of Fe–S cluster proteins? This is important as elegant *in vitro* studies have implicated mycothiol and thioredoxin as potential reductants of Wbl proteins [29]. Also, since many Fe–S cluster proteins (e.g. SoxR, FNR, WhiB3) are asserted to be intracellular redox sensors, the precise redox state of the microbial cell becomes important. Unfortunately, this is an understudied area of research, probably because of the lack of tools to accurately measure the ambient redox state of the microbial cytoplasm.

In conclusion, Fe–S cluster proteins play a central role in virtually all organisms, and *Mtb* WhiB [Fe–S] cluster proteins, which modulate redox homeostasis, oxidative stress, cell division, drug resistance and virulence, are ideal candidates for further study [23**]. It is strongly anticipated that the examination of non-model genomes (e.g. *Mtb*) for new genes and strategies for assembling Fe–S cluster proteins will open up new, exciting areas of investigation.

Acknowledgments

Research in the laboratory is supported in whole or in part by the National Institutes of Health Grants AI058131, AI076389 (to A.J.C.S.). This work is also supported by the University of Alabama at Birmingham (UAB) Center for AIDS Research, and UAB Center for Free Radical Biology (A.J.C.S.). A.J.C.S. is a Burroughs Wellcome Investigator in the Pathogenesis of Infectious Diseases.

References and recommended reading/References

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Py B, Moreau PL, Barras F. Fe–S clusters, fragile sentinels of the cell. *Curr Opin Microbiol.* 2011; 14:218–223. [PubMed: 21288764]
 2. Xu XM, Moller SG. Iron sulfur clusters: biogenesis, molecular mechanisms, and their functional significance. *Antioxid Redox Signal.* 2011; 15:271–307. [PubMed: 20812788]
 3. Wollers S, Layer G, Garcia-Serres R, Signor L, Clemancey M, Latour JM, Fontecave M, Ollagnier deChoudens S. Iron–Sulfur (Fe–S) cluster assembly: the SufBCD complex is a new type of Fe–S scaffold with a flavin redox cofactor. *J Biol Chem.* 2010; 285:23331–23341. [PubMed: 20460376]
 4. Ayala-Castro C, Saini A, Outten FW. Fe–S cluster assembly pathways in bacteria. *Microbiol Mol Biol Rev.* 2008; 72:110–125. [PubMed: 18322036]
 - 5•. Liu Y, Sieprawaska-Lupa M, Whitman WB, White RH. Cysteine is not the sulfur source for iron–sulfur cluster and methionine biosynthesis in the methanogenic archaeon *Methanococcus maripaludis*. *J Biol Chem.* 2010; 285:31923–31929. [PubMed: 20709756] [This is the first demonstration that sulfide can be used as a source of sulfur during Fe–S cluster biosynthesis. Usually organisms that stay in sulfide-rich habitats, mostly those belonging to archaea, use sulfide rather than cysteine to make Fe–S clusters.]
 6. Yasmin S, Andrews SC, Moore GR, Le Brun NE. A new role for heme, facilitating release of iron from the bacterioferritin iron biomineral. *J Biol Chem.* 2011; 286:3473–3483. [PubMed: 21106523]
 7. Hickok JR, Sahni S, Shen H, Arvind A, Antoniou C, Fung LW, Thomas DD. Dinitrosyliron complexes are the most abundant nitric oxide-derived cellular adduct: biological parameters of assembly and disappearance. *Free Radic Biol Med.* 2011; 51:1558–1566. [PubMed: 21787861]
 - 8••. Landry AP, Duan X, Huang H, Ding H. Iron–sulfur proteins are the major source of protein-bound dinitrosyl iron complexes formed in *Escherichia coli* cells under nitric oxide stress. *Free Radic Biol Med.* 2011; 50:1582–1590. [PubMed: 21420489] [There has been a longstanding debate about the source of iron in Fe–S clusters. In this study, using different *E. coli* mutants, it was conclusively demonstrated that Fe–S clusters are part of an intracellular chelatable iron pool.]
 - 9••. Crack JC, Green J, Cheesman MR, Le Brun NE, Thomson AJ. Superoxide-mediated amplification of the oxygen-induced switch from [4Fe–4S] to [2Fe–2S] clusters in the transcriptional regulator FNR. *Proc Natl Acad Sci U S A.* 2007; 104:2092–2097. [PubMed: 17267605] [This study elegantly demonstrates the steps involved in cluster conversion in FNR and provides mechanistic insights into the role of FNR as an oxygen sensor.]
 10. Uden G, Achebach S, Holighaus G, Tran HQ, Wackwitz B, Zeuner Y. Control of FNR function of *Escherichia coli* by O₂ and reducing conditions. *J Mol Microbiol Biotechnol.* 2002; 4:263–268. [PubMed: 11931557]
 11. Tran QH, Arras T, Becker S, Holighaus G, Ohlberger G, Uden G. Role of glutathione in the formation of the active form of the oxygen sensor FNR ([4Fe–4S] FNR) and in the control of FNR function. *Eur J Biochem.* 2000; 267:4817–4824. [PubMed: 10903516]
 12. Grainger DC, Aiba H, Hurd D, Browning DF, Busby SJW. Transcription factor distribution in *Escherichia coli*: studies with FNR protein. *Nucleic Acids Res.* 2007; 35:269–278. [PubMed: 17164287]

13. Poole RK, Hughes MN. New functions for the ancient globin family: bacterial responses to nitric oxide and nitrosative stress. *Mol Microbiol.* 2000; 36:775–783. [PubMed: 10844666]
- 14•. Dufour YS, Kiley PJ, Donohue TJ. Reconstruction of the core and extended regulons of global transcription factors. *PLoS Genet.* 2010; 6:e1001027. [PubMed: 20661434] [This study revealed that a core set of target genes is conserved across many species for FNR-type regulators. Using microarrays and Chip-chip studies, it was shown that the observed variation in the extended regulons for FNR-type regulators among closely related species denotes species-specific adaptation within a defined niche.]
15. Ding H, Hidalgo E, Demple B. The redox state of the [2Fe–2S] clusters in soxR protein regulates its activity as a transcription factor. *J Biol Chem.* 1996; 271:33173–33175. [PubMed: 8969171]
- 16••. Krapp AR, Humbert MV, Carrillo N. The soxRS response of *Escherichia coli* can be induced in the absence of oxidative stress and oxygen by modulation of NADPH content. *Microbiology.* 2011; 157:957–965. [PubMed: 21178165] [This study demonstrates that the *E. coli* soxRS regulon can be activated by depletion of the NADPH stock, rather than by accumulation of superoxide per se, following exposure to superoxide-generating compounds.]
17. Gaudu P, Moon N, Weiss B. Regulation of the soxRS oxidative stress regulon. Reversible oxidation of the Fe–S centers of soxR in vivo. *J Biol Chem.* 1997; 272:5082–5086. [PubMed: 9030573]
18. Kobayashi K, Mizuno M, Fujikawa M, Mizutani Y. Protein conformational changes of the oxidative stress sensor, soxR, upon redox changes of the [2Fe–2S] cluster probed with ultraviolet resonance raman spectroscopy. *Biochemistry.* 2011; 50:9468–9474. [PubMed: 21988663]
- 19••. Gu M, Imlay JA. The soxRS response of *Escherichia coli* is directly activated by redox cycling drugs rather than by superoxide. *Mol Microbiol.* 2011; 79:1136–1150. [PubMed: 21226770] [In this seminal study the authors demonstrated that redox cycling compounds can directly activate SoxR by oxidation of its [2Fe–2S] cluster, and that superoxide is not the signal sensed by SoxR.]
20. Dietrich LEP, Kiley PJ. A shared mechanism of soxR activation by redox cycling compounds. *Mol Microbiol.* 2011; 79:1119–1122. [PubMed: 21338412]
21. Gorodetsky AA, Dietrich LEP, Lee PE, Demple B, Newman DK, Barton JK. DNA binding shifts the redox potential of the transcription factor soxR. *Proc Natl Acad Sci U S A.* 2008; 105:3684–3689. [PubMed: 18316718]
22. Chater KF. A morphological and genetic mapping study of white colony mutants of *Streptomyces coelicolor*. *J Gen Microbiol.* 1972; 72:9–28. [PubMed: 4561048]
- 23••. Saini V, Farhana A, Steyn AJC. *Mycobacterium tuberculosis* WhiB3: a novel iron-sulfur cluster protein that regulates redox homeostasis and virulence. *Antioxid Redox Signal.* 2011 doi: 10.1089/ars.2011.4341. [This article represents the first comprehensive review of WhiB proteins with a focus on the regulation of *Mtb* WhiB3. The authors also performed bioinformatic and phylogenetic analyses, which provide new insight into the distribution and domain architecture of WhiB proteins in prokaryotes.]
24. Agarwal N, Raghunand TR, Bishai WR. Regulation of the expression of whiB1 in *Mycobacterium tuberculosis*: role of cAMP receptor protein. *Microbiology.* 2006; 152:2749–2756. [PubMed: 16946269]
25. Gomez JE, Bishai WR. whmD is an essential mycobacterial gene required for proper septation and cell division. *Proc Natl Acad Sci U S A.* 2000; 97:8554–8559. [PubMed: 10880571]
26. Morris RP, Nguyen L, Gatfield J, Visconti K, Nguyen K, Schnappinger D, Ehrt S, Liu Y, Heifets L, Pieters J, et al. Ancestral antibiotic resistance in *Mycobacterium tuberculosis*. *Proc Natl Acad Sci U S A.* 2005; 102:12200–12205. [PubMed: 16103351]
27. Rybníček J, Nowag A, Van Gumpel E, Nissen N, Robinson N, Plum G, Hartmann P. Insights into the function of the WhiB-like protein of mycobacteriophage TM4 — a transcriptional inhibitor of WhiB2. *Mol Microbiol.* 2010; 77:642–657. [PubMed: 20545868]
28. Garg S, Alam MS, Bajpai R, Kishan KVR, Agrawal P. Redox biology of *Mycobacterium tuberculosis* H37Rv: protein–protein interaction between GlgB and WhiB1 involves exchange of thiol-disulfide. *BMC Biochem.* 2009; 10:1. [PubMed: 19121228]

29. Crack JC, den Hengst CD, Jakimowicz P, Subramanian S, Johnson MK, Buttner MJ, Thomson AJ, Le Brun NE. Characterization of [4Fe–4S]-containing and cluster-free forms of *Streptomyces* WhiD. *Biochemistry*. 2009; 48:12252–12264. [PubMed: 19954209]
30. Smith LJ, Stapleton MR, Fullstone GJM, Crack JC, Thomson AJ, Le Brun NE, Hunt DM, Harvey E, Adinolfi S, Buxton RS, et al. *Mycobacterium tuberculosis* WhiB1 is an essential DNA-binding protein with a nitric oxide sensitive iron–sulphur cluster. *Biochem J*. 2010; 432:417–427. [PubMed: 20929442]
31. Alam MS, Garg SK, Agrawal P. Studies on structural and functional divergence among seven WhiB proteins of *Mycobacterium tuberculosis* H37Rv. *FEBS J*. 2009; 276:76–93. [PubMed: 19016840]
- 32•. Singh A, Crossman DK, Mai D, Guidry L, Voskuil MI, Renfrow MB, Steyn AJC. *Mycobacterium tuberculosis* WhiB3 maintains redox homeostasis by regulating virulence lipid anabolism to modulate macrophage response. *PLoS Pathog*. 2009; 5:e1000545. [PubMed: 19680450] [This article describes the role of the *Mtb* intracellular redox sensor WhiB3 in the redox-mediated regulation of complex virulence lipids. The concept of reductive stress emerged from these findings.]
- 33•. Singh A, Guidry L, Narasimhulu KV, Mai D, Trombley J, Redding KE, Giles GI, Lancaster JR Jr, Steyn AJ. *Mycobacterium tuberculosis* WhiB3 responds to O₂ and nitric oxide via its [4Fe–4S] cluster and is essential for nutrient starvation survival. *Proc Natl Acad Sci U S A*. 2007; 104:11562–11567. [PubMed: 17609386] [This article links mycobacterial metabolism with the redox signaling molecules NO and O₂ through the *Mtb* WhiB3 [4Fe–4S] cluster. Importantly, WhiB3 was shown to function as an intracellular redox sensor involved in the metabolic switchover to the preferred *in vivo* carbon source, fatty acids.]
34. Crack JC, Smith LJ, Stapleton MR, Peck J, Watmough NJ, Buttner MJ, Buxton RS, Green J, Oganessian VS, Thomson AJ, et al. Mechanistic insight into the nitrosylation of the [4Fe–4S] cluster of WhiB-like proteins. *J Am Chem Soc*. 2011; 133:1112–1121. [PubMed: 21182249]
35. Steyn AJ, Collins DM, Hondalus MK, Jacobs WR Jr, Kawakami RP, Bloom BR. *Mycobacterium tuberculosis* WhiB3 interacts with RpoV to affect host survival but is dispensable for *in vivo* growth. *Proc Natl Acad Sci U S A*. 2002; 99:3147–3152. [PubMed: 11880648]
- 36•. Farhana A, Guidry L, Srivastava A, Singh A, Hondalus MK, Steyn AJC. Reductive stress in microbes: implications for understanding *Mycobacterium tuberculosis* disease and persistence. *Adv Microb Physiol*. 2010; 57:43–117. [PubMed: 21078441] [This is a comprehensive review paper that describes the role of reductive stress in mycobacterial physiology and virulence.]
37. Khoroshilova N, Popescu C, Munck E, Beinert H, Kiley PJ. Iron–sulfur cluster disassembly in the FNR protein of *Escherichia coli* by O₂: [4Fe–4S] to [2Fe–2S] conversion with loss of biological activity. *Proc Natl Acad Sci U S A*. 1997; 94:6087–6092. [PubMed: 9177174]
38. Geiman DE, Raghunand TR, Agarwal N, Bishai WR. Differential gene expression in response to exposure to antimycobacterial agents and other stress conditions among seven *Mycobacterium tuberculosis* whiB-like genes. *Antimicrob Agents Chemother*. 2006; 50:2836–2841. [PubMed: 16870781]
39. Kumar A, Farhana A, Guidry L, Saini V, Hondalus M, Steyn AJ. Redox homeostasis in mycobacteria: the key to tuberculosis control? *Expert Rev Mol Med*. 2011; 13:e39. [PubMed: 22172201]
- 40•. Shomura Y, Yoon KS, Nishihara H, Higuchi Y. Structural basis for a [4Fe–3S] cluster in the oxygen-tolerant membrane-bound [NiFe]-hydrogenase. *Nature*. 2011; 479:253–256. [PubMed: 22002607] [This study reveals the structure of oxygen-tolerant membrane-bound respiratory [NiFe]-hydrogenase (MBH) and show that the proximal iron-sulfur (Fe–S) cluster of MBH has a [4Fe–3S] structure coordinated by six cysteine residues. The authors discuss different features of [4Fe–3S] cluster and suggest that in addition to being a part of electron relay machinery, the cluster also donates two electrons and one proton crucial for the reduction of O₂ to prevent the formation of the inactive state of the enzyme.]
- 41•. Carroll KS, Bhawe DP, Hong JA, Keller RL, Krebs C. Iron–sulfur cluster engineering provides insight into the evolution of substrate specificity among sulfonucleotide reductases. *ACS Chem Biol*. 2011 doi: 10.1021/cb200261n. [Using metalloprotein engineering, spectroscopy and

enzyme kinetics, this study shows for the first time that the Fe–S cluster in the sulfonucleotide reductase APS imparts substrate specificity and carries out catalysis.]

42. Sun F, Ji Q, Jones MB, Deng X, Liang H, Frank B, Telser J, Peterson SN, Bae T, He C. AirSR, a [2Fe–2S] cluster-containing two-component system, mediates global oxygen sensing and redox signaling in *Staphylococcus aureus*. *J Am Chem Soc*. 2012; 134:305–314. [PubMed: 22122613]
43. Boutet E, Lieberherr D, Tognolli M, Schneider M, Bairoch A. UniProtKB/Swiss-Prot. *Methods Mol Biol*. 2007; 406:89–112. [PubMed: 18287689]
44. Major TA, Burd H, Whitman WB. Abundance of 4Fe–4S motifs in the genomes of methanogens and other prokaryotes. *FEMS Microb Lett*. 2004; 239:117–123. [The authors studied the distribution of 4Fe–4S motifs in different prokaryotic species and demonstrated that the distribution of these motifs is closely associated with the lifestyle of the organisms.]
45. Betts JC, Lukey PT, Robb LC, McAdam RA, Duncan K. Evaluation of a nutrient starvation model of *Mycobacterium tuberculosis* persistence by gene and protein expression profiling. *Mol Microbiol*. 2002; 43:717–731. [PubMed: 11929527]
46. Schnappinger D, Ehrt S, Voskuil MI, Liu Y, Mangan JA, Monahan IM, Dolganov G, Efron B, Butcher PD, Nathan C, et al. Transcriptional adaptation of *Mycobacterium tuberculosis* within macrophages. *J Exp Med*. 2003; 198:693–704. [PubMed: 12953091]
47. Voskuil MI, Schnappinger D, Visconti KC, Harrell MI, Dolganov GM, Sherman DR, Schoolnik GK. Inhibition of respiration by nitric oxide induces a *Mycobacterium tuberculosis* dormancy program. *J Exp Med*. 2003; 198:705–713. [PubMed: 12953092]
48. Voskuil MI, Bartek IL, Visconti K, Schoolnik GK. The response of *Mycobacterium tuberculosis* to reactive oxygen and nitrogen species. *Front Microbiol*. 2011; 2:105. [PubMed: 21734908]
49. Banerjee S, Nandyala AK, Raviprasad P, Ahmed N, Hasnain SE. Iron-dependent RNA-binding activity of *Mycobacterium tuberculosis* aconitase. *J Bacteriol*. 2007; 189:4046–4052. [PubMed: 17384188]

Box 1 Repertoire of *Mtb* Fe–S cluster proteins

Analysis of the UniProtKB protein database [43] revealed more than 121 220 entries for Fe–S cluster-containing proteins. The majority of these proteins (~68.5%) are likely of the 4Fe–4S cluster type with ~21% of the 2Fe–2S type and only 3% of the 3Fe–4S cluster type. Considering the diversity in protein domain architecture, it is assumed that the distribution of Fe–S cluster-containing proteins within a species correlates with the dynamic lifestyle and associated features of the organism. Despite the availability of large databases and related bioinformatics tools, information regarding the full repertoire of Fe–S cluster proteins and their functions in mycobacteria is still lacking. To determine the number of Fe–S cluster-coordinating proteins in mycobacteria, protein sequences of *Mtb*, *M. avium paratuberculosis* and *M. leprae* from the NCBI genome database were subjected to PRATT and PROSITE-based analysis of amino acid pattern/signatures known to coordinate diverse types of Fe–S clusters. We observed that the *Mtb* genome contains 50 genes likely to encode Fe–S cluster proteins, which is roughly half of the known Fe–S proteins in *E. coli*. Expression profiles of these 50 genes and genes encoding the Fe–S cluster assembly machinery are indicated in Figure 2. Almost 50% of the *Mtb* Fe–S proteins, including those of the WhiB family, coordinate a 4Fe–4S type cluster. Using a dataset of over 100 prokaryotic species, it has been previously shown that there is an apparent skew in the distribution of 4Fe–4S-cluster coordinating motifs associated with the lifestyle of the organism [44*]. In this regard, anaerobic bacteria have significantly more motifs (7.4 motifs/1000 ORFs) compared to that of aerobic bacteria (2.8 motifs/1000 ORFs) [44*]. We determined that *Mtb* harbors ~6.5 motifs/1000 ORFs, which is significantly higher than aerobic bacterial species. It is possible that the increased prevalence of 4Fe–4S-clusters may contribute to persistent *Mtb* cells within the hypoxic granulomas. We also observed that some *Mtb* proteins, including other mycobacterial species contain structural features known to coordinate different cluster types such as 2Fe–2S, GATASE Type II, nitrogenase 2, rubredoxins and others (Figure 2). However, the distribution of Fe–S cluster coordinating proteins varies even within mycobacterial species. For example, *M. avium paratuberculosis* and *M. leprae* likely encode only 45 and 28 Fe–S cluster harboring proteins, respectively. *M. leprae* encodes very few proteins coordinating 2Fe–2S ferredoxin, 4Fe–4S ferredoxin and rubredoxin type motifs and could be because of excessive genome decay.

To classify the predicted functions of the 50 putative *Mtb* Fe–S proteins, we used Tuberculist and COG (Clusters of Orthologous Groups of Proteins databases) (Figure 3). Using the Tuberculist functional classification guidelines, we observed that ~70% (34/50) of *Mtb* Fe–S proteins are involved in intermediary metabolism and respiration functions, which is consistent with the reported functions of Fe–S proteins in other microbial species. Using COG, a more detailed analysis revealed that over 40% (21/50) of *Mtb* Fe–S proteins participate in energy production and conversion function (Figure 3). This is followed by transcriptional regulation (primarily the WhiB family) and amino acid transport and metabolism for 15% and 10% of the Fe–S proteins, respectively. In sum, we observed that Fe–S proteins perform a variety of functions in *Mtb* ranging from metabolism to defense mechanisms and inorganic ion transport (Figure 3) and hence, could be critical to *Mtb* virulence and persistence.

Current Opinion in Chemical Biology

Figure 1.

Schematic representation of the mechanism of action of *E. coli* FNR, *E. coli* SoxR and *Mtb* WhiB3. (a) FNR exists in its apo form under aerobic conditions, wherein prolonged exposure to O₂ inactivates FNR. With decreased O₂ concentration, IscS transfers an [2Fe-2S]²⁺ cluster to the monomeric apo-FNR, leading to the formation of the [4Fe-4S]²⁺ FNR dimer in a stepwise process. The dimer is capable of binding DNA to activate or repress target gene expression. Conversion of the Fe-S cluster and monomerization can lead to the loss of binding, wherein [4Fe-4S]²⁺ is converted to [4Fe-4S]³⁺ upon oxidation, with the release of superoxide (O₂^{•-}) and hydrogen peroxide (H₂O₂). O₂^{•-} can cause loss of the FNR Fe-S cluster and provides an additional mechanism of regulation of FNR DNA binding. Besides O₂, the active dimeric FNR [4Fe-4S]²⁺ is also responsive to NO to form a DNIC-FNR complex, which leads to its monomerization and dissociation from DNA. (b) SoxR exists as a homodimer in a [2Fe-2S]¹⁺ state in its inactive form, which is capable of sensing and responding to oxidative stress, intracellular NADPH, redox-active and superoxide-generating compounds and NO. Upon exposure to such stresses, oxidation of SoxR [2Fe-2S]¹⁺ leads to the formation of [2Fe-2S]²⁺, which induces DNA binding and transcription of target genes as well as its downstream signaling protein, SoxS. Besides this emerging model, the classical SoxR regulation demonstrates SoxR binding to SoxS in a complex leading to structural changes in SoxS that allow its binding to target DNA. This modulates the transcription of genes of the SoxRS regulon such as superoxide dismutase, xenobiotic efflux pumps and carbon metabolism enzymes. In addition to external stimuli, the intracellular redox environment also affects SoxR binding to DNA. (c) *Mtb* WhiB3 is a Fe-S

cluster protein present exclusively in members of *Actinobacteria* and harbors a [4Fe–4S] cluster, which upon exposure to O₂ generate [3Fe–4S], [2Fe–2S], and eventually apo, oxidized WhiB3. The WhiB3 [4Fe–4S] cluster can also react with NO to generate a stable DNIC complex. The DNA binding properties of WhiB3 are distinct from that of FNR and SoxR as oxidized apo-WhiB3 binds DNA exceptionally strongly compared to WhiB3 [4Fe–4S]²⁺ or [4Fe–4S]¹⁺, and FNR and SoxR. Whether WhiB3 exists as a monomer or dimer in its active or inactive form is the subject of further investigation. Vertical color triangles represent gradients of O₂ or NO.

Figure 2.

Microarray-based expression profiles of *Mtb* Fe-S cluster and Fe-S cluster assembly genes. Using bioinformatics-based amino acid pattern searches and manual curation of the literature, we identified 50 genes encoding putative *Mtb* Fe-S cluster proteins. Expression profiles of these 50 genes, and genes encoding the Fe-S cluster assembly machinery (Rv1460-Rv1466, Rv3025c) were extracted from independent *Mtb* expression studies [45–48] and a representative expression heat-map was generated. The heat-map represents differential expression profiles of these genes under diverse stress conditions such as aerated and non-aerated starvation, hypoxia, macrophage infection and exposure to NO and H₂O₂. The data suggest that most genes exposed to H₂O₂ and NO are weakly to highly upregulated

compared to starvation and hypoxic conditions. The expression profiles under hypoxic conditions, macrophage infection and starvation conditions are similar to some degree. Intriguingly, the Fe-S cluster assembly genes are downregulated during hypoxia, but are highly upregulated within macrophages, and upon exposure to H₂O₂ and NO. The dramatic upregulation of Fe-S cluster assembly genes is probably to compensate for the loss of Fe-S clusters due to oxidative stress. Note that aconitase, a bifunctional enzyme involved in iron homeostasis during oxidative stress [49], is the most highly upregulated gene, followed by several of the Fe-S cluster assembly genes. Color scale represents fold changes in mRNA levels in wt *Mtb* compared to wt *Mtb* grown under the 'control' conditions. Gray boxes; value not available, Rrf — ribosome recycling factor.

Figure 3.

Functional classification of *Mtb* Fe-S cluster proteins. Using Tuberculist and the COG databases, 50 putative Fe-S cluster containing proteins were analyzed. Approximately 70% of these proteins belong to intermediary metabolism and respiration followed by regulatory functions for ~15% of these proteins. Over 40% of Fe-S proteins in *Mtb* participate in energy production and conversion. This is followed by transcription regulation (mainly the WhiB family) and amino acid transport and metabolism for ~15% and 10% of Fe-S proteins, respectively.