Published in final edited form as:

Biochem Biophys Res Commun. 2014 February 21; 444(4): 634–637. doi:10.1016/j.bbrc.2014.01.143.

PITX2 associates with PTIP-containing histone H3 lysine 4 methyltransferase complex

Yan Liu^{a,1}, Yue Huang^{b,1}, Jun Fan^c, and Guo-Zhang Zhu^{b,*}

^aDepartment of Ophthalmology, Huashan Hospital Affiliated to Fudan University, Shanghai, 200040, People's Republic of China

^bDepartment of Biological Sciences, Marshall University, Huntington, WV, USA

^cDepartment of Biochemistry and Microbiology, Genomic Core Facility, School of Medicine, Marshall University, Huntington, WV, USA

Abstract

Pituitary homeobox 2 (PITX2), a Paired-like homeodomain transcription factor and a downstream effector of Wnt/ β -catenin signaling, plays substantial roles in embryonic development and human disorders. The mechanism of its functions, however, is not fully understood. In this study, we demonstrated that PITX2 associated with histone H3 lysine 4 (H3K4) methyltransferase (HKMT) mixed-lineage leukemia 4 (MLL4/KMT2D), Pax transactivation domain-interacting protein (PTIP), and other H3K4 HKMT core subunits. This association of PITX2 with H3K4 HKMT complex was dependent on PITX2's homeodomain. Consistently, the PITX2 protein complex was shown to possess H3K4 HKMT activity. Furthermore, the chromatin immunoprecipitation result revealed co-occupancy of PITX2 and PTIP on the promoter of the PITX2's transcriptional target. Taken together, our data provide new mechanistic perspectives on PITX2's functions and its related biological processes.

Keywords

PITX2; MLL4/KMT2D; PTIP; histone H3 lysine 4 (H3K4); histone lysine methyltransferase (HKMT)

1. Introduction

Pituitary homeobox 2 (PITX2) is a transcription factor with a *Paired*-like homeodomain. It is evolutionarily conserved in many animal species including zebrafish and human. Previous studies have demonstrated that PITX2 plays significant roles in embryonic development and human disorders. *Pitx2* knockout mice die by embryonic day 15 due to severe developmental defects [1–3]. It is critical for the establishment of the left-right body axis, the asymmetrical development of the heart, lungs, liver, and spleen, twisting of the gut and stomach [1–3]. It is also responsible for the development of the eye [4–5], tooth [6] and

© 2013 Elsevier Inc. All rights reserved.

*Corresponding author. Department of Biological Sciences, Marshall University, One John Marshall Drive, Huntington, WV 25705, USA. Fax: +1 304 696 7136. Tel: +1 304 696 2429. zhu4@marshall.edu (G.-Z. Zhu).

¹These authors contributed equally to this work.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

cardiac outflow tract [3, 7]. Mutations in the *PITX2* gene have been linked to several human disorders, including Axenfeld-Rieger syndrome, iridogoniodysgenesis syndrome, sporadic cases of Peters anomaly, and atrial fibrillation [4, 8]. We and others recently reported an oncogenic role of *PITX2* in human thyroid cancer [9], ovarian cancer [10], colon cancer [11], renal cancer [12] and esophageal squamous cell carcinoma [13].

Mechanistically, *PITX2* has been shown to serve as a downstream effector of Nodal and Wnt signaling [3, 14]. It can transcriptionally activate expression of genes required for cell proliferation (e.g., c-Myc, Cyclin Ds) [9, 15] or cell differentiation (e.g., MyoD) [16]. Several *PITX2*-interacting transcriptional regulators, including β -catenin, LEF/TCF and YB-1, have been reported previously [17–18].

In our previous study, we demonstrated that transient overexpression of *PITX2* in HEK293 cells resulted in 868 upregulated genes and 191 downregulated genes [18]. Among these regulated genes, 15 were imprinted genes including *H19*, *IGF2* and *CDKN1C*. This remarkable change of global gene expression prompted us to hypothesize that *PITX2* might execute its effect through interacting with chromatin-remodeling molecules. In this study, we tested this hypothesis and demonstrated that *PITX2* formed complex with histone lysine methyltransferase. This intriguing finding provides new insights on *PITX2*'s functions and its related biological processes.

2. Materials and methods

2.1. Cell Culture

HEK293 cells (CRL-1573TM, ATCC, VA) were cultured in Dulbecco's modified Eagle's medium with 4.5 g/l glucose, 10% fetal bovine serum, and penicillin-streptomycin (100 IU/ml). Cell culture media and supplements were purchased from ATCC. Cells were incubated at 37°C in a humidified atmosphere with 5% CO₂.

2.2. Plasmid Construction

Construction of the pEGFP-NFLAG-PITX2c plasmid (expressing N-terminal FLAG tagged full-length human PITX2C, amino acids 1–324) and the pEGFP-NFLAG-PITX2c- Δ HHD plasmid (expressing N-terminal FLAG tagged, homeodomain-deleted PITX2c, amino acids 1–131) has been reported in our previous works [9, 18]. The other truncated constructs of PITX2c were cloned from pEGFP-NFLAG-PITX2c and inserted into either the NheI/KpnI sites (for PITX2c amino acids 152–324) or the HindIII/KpnI sites (for others) of the pEGFP vector (Clontech Laboratories). Enhanced GFP (green fluorescent protein) was not fused with PITX2 in the abovementioned plasmids and its expression was solely served as an internal marker to monitor transfection efficiency.

2.3. Co-immunoprecipitation and Immunoblotting

The plasmids with FLAG-tagged, full-length or truncated PITX2 were transfected into HEK293 cells that had been cultured overnight in 6-well plates by Fugene HD transfection reagent (Roche, IN). Transfection efficiency was monitored by green fluorescence 48 hours later to ensure the consistency of experiments. Transfected cells were then lysed by the cell lysis buffer (50 mM Tris-HCl pH 7.4, 150 mM NaCl, 5 mM EDTA, 1% Triton X-100, protease inhibitors) and incubated with primary antibodies at 4°C overnight before being precipitated by protein A beads (Pierce). Immunoprecipitated samples were boiled in the SDS loading buffer and subjected to immunoblotting analysis as described previously [9]. Antibodies used for immunoprecipitation were: FLAG M2 (Sigma, MO); PTIP, RBBP5 and ASH2L (Bethyl laboratory); MLL4/KMT2D (a gift from Dr. Kai Ge, NIDDK/NIH).

Antibodies used for immunoblotting were: PITX2 (Abnova); PTIP, RBBP5 and ASH2L (Bethyl laboratory); hnRNP U (Abcam).

2.4. Histone Methyltransferase Assay

The nuclear extract (NE) of HEK293 cells was prepared by a NE-PER kit (Thermo Scientific Pierce) according to the manufacture's instruction. The PITX2, PTIP, or MLL4/KMT2D immunoprecipitated complex was prepared as described above. The NE or the immunoprecipitated complex was incubated with human histone H3.1 (New England Biolabs) and S-adenosylmethionine (SAM) (New England Biolabs) at room temperature for 1 hour. After incubation, the reaction was subjected to immunoblotting by the antibody against H3K4 mono-methylation (H3K4me1) (Cell Signaling Technology).

2.5. Chromatin Immunoprecipitation (ChIP) Assay

HEK293 cells were transiently transfected with pEGFP-NFLAG-PITX2c or pEGFP-NFLAG-PITX2c-ΔHD. After 48 hours of initial transfection, cells were cross-linked by 1% formaldehyde for 15 minutes. Cell genomic DNA was then sheared into fragments by sonication. Cell lysates were pre-cleared by protein A beads (Pierce, IL) for 2 h at 4°C. Anti-FLAG M2 conjugated agarose beads (Sigma), anti-PTIP (Bethyl laboratory), anti-H3K4me2 (Cell Signaling Technology), anti-HDAC1 (Cell Signaling Technology) or IgG antibody was then incubated with pre-cleared cell lysates at 4°C overnight. Beads were washed by the high salt and LiCl washing solution for eight times. Samples were reverse cross-linked in the high salt solution at 65°C overnight. DNA was purified by phenol-chloroform extraction and precipitated by isopropanol. PCR was then used to analyze precipitated DNA samples. The PCR primers (ATGGAAACGCTCCCGCTAGGT and AGGACCAAGTGTCTGAGGGATT) covered the promoter (−133 to −7) of the human Cyclin A1. PCR parameters are: 94°C for 2 min; 94°C for 20 sec, 58°C for 20 sec, 72°C for 50 sec, 36 cycles; 72°C for 6 min.

3. Results

3.1. PITX2 forms complex with MLL4, ASH2L, RBBP5 and PTIP

We previously reported that transiently overexpressed PITX2 significantly altered global gene expression patterns [18]. This finding prompted us to hypothesize that PITX2 might synergistically interact with chromatin-remodeling molecules. As a first attempt to test this hypothesis, we looked at histone H3 lysine 4 (H3K4) methyltransferases, which have been linked to transcriptional activation [19–20]. As shown in Figure 1, PITX2 co-immunoprecipitated with H3K4 methyltransferase mixed-lineage leukemia 4 (MLL4/KMT2D). ASH2L and RBBP5, two core subunits of the MLL family of H3K4 methyltransferases, were also found to associate with PITX2. Consistently, we observed the association of PITX2 with PTIP, which is a unique subunit for the MLL3 and MLL4 methyltransferase complex [21]. On the other hand, hnRNP U, a PITX2 interacting protein identified in our previous study [18], did not form complex with MLL4 or other subunits. Collectively, our data demonstrate that PITX2 associates with MLL4, ASH2L, RBBP5 and PTIP in one complex.

3.2. The homeodomain of PITX2 is required for its association with the H3K4 methyltransferase complex

To determine which domain of PITX2 is required for its association with the H3K4 methyltransferase complex, we performed co-immunoprecipitation on various truncated forms of PITX2. RBBP5, a core subunit of the MLL family, was selected as a read-out for this experiment. The result revealed that PITX2 (FL, aa1-324) (full-length PITX2), PITX2 (aa1-180) (lacking C-terminal half) and PITX2 (aa91-324) associated with RBBP5, but PITX2 (aa1-131) (lacking C-terminal half and part of homeodomain) and PITX2

(aa152-324) (lacking N-terminal and homeodomain) did not (Fig. 2), indicating that the homeodomain of PITX2 is critical for this association.

3.3. PITX2 complex methylates histone H3 on K4

The above results showed that PITX2 formed complex with MLL4/KMT2D. Thus, we asked whether the PITX2 complex possessed H3K4 histone lysine methyltransferase (HKMT) activity. As shown in Figure 3, the immunoprecipitates of PITX2, PTIP and MLL4 all efficiently methylated histone H3 on K4. In contrast, the immunoprecipitates of IgG control and homeodomain-deleted PITX2 failed to do so. These data indicate that PITX2 *bona fide* associates with H3K4 methyltransferase complex. This previously uncharacterized finding suggests that PITX2 may transcriptionally activate expression of its target genes by linking H3K4 HKMT to the target's promoter, a point to be investigated next.

3.4. PITX2 and PTIP co-occupy the promoter of the PITX2's transcriptional target

To determine whether PITX2 and H3K4 HKMT co-occupy the promoter of PITX2's target genes, we performed chromatin immunoprecipitation (ChIP) assays. Unfortunately, the MLL4 antibody did not work in ChIP. Thus, we alternatively chose PTIP, a unique subunit for the MLL4 methyltransferase complex [21]. The gene tested in ChIP was Cyclin A1, which was newly identified as PITX2's transcriptional target in our recent works [18, 22]. Consistent with our previous observation that Cyclin A1 transcription was turned on by full-length PITX2 but not by homeodomain-deleted PITX2 (PITX2 Δ HHD), our ChIP result revealed co-occupancy of PITX2 and PTIP on the Cyclin A1 promoter around the transcription start site (TSS) (Fig. 4). Similarly, we observed increased enrichment of H3K4 di-methylation (H3K4me₂) on the Cyclin A1 promoter (Fig. 4). Conversely, the ChIP signal of histone deacetylase 1 (HDAC1) was overtly decreased (Fig. 4). Taken together, these data suggest that PITX2 can recruit H3K4 HKMT to the promoter of its transcriptional target and, consequently, activate gene transcription.

4. Discussion

Homeodomain transcription factor PITX2 plays substantial roles in embryonic development and human disorders. Thus, elucidating the mechanisms of its action is of great interest to many researchers. Herein, we report that PITX2 associates with PTIP-containing H3K4 methyltransferase complex. This finding is significant since it uncovers a novel, epigenetic mechanism for PITX2's functions and its related biological processes.

Although our data demonstrate the association of PITX2 and MLL4/KMT2D, by no means we can discount the possibility that PITX2 might form complex with other H3K4 methyltransferases or other chromatin-remodeling molecules in general. In fact, this study just opens a whole new avenue to investigate the physical and functional interaction between PITX2 and epigenetic regulators.

Our results indicate that the homeodomain of PITX2 is necessary for its association with H3K4 methyltransferase complex and associated HKMT activity. This is consistent with a previous report that demonstrated the requirement of homeodomain in the interaction between mouse Pitx2 and WDR5 (another core subunit of the MLL family of H3K4 methyltransferases) [23]. However, how PITX2 interacts with H3K4 HKMT complex, e.g., which HKMT subunit is the direct binding partner, is still unclear and should be addressed next.

The finding of PITX2-PTIP complex in this study deserves further attention. PTIP has been assigned three distinct functions, one for regulating gene transcription [24], one for promoting immunoglobulin class switch recombination [25–26], and the other for

facilitating DNA damage repair (DDR) [27–30]. Although it is straightforward to line up PITX2-PTIP in gene transcription as previously reported PAX2-PTIP [31], it would be interesting to further ask whether PITX2 has a role in immunoglobulin class switch recombination and DDR through interacting with PTIP.

While we were preparing this manuscript, three recent reports have demonstrated that MLL4 is a major HKMT responsible for H3K4 mono- and di-methylation in mammalian cells and a majority of MLL4 binding sites are located in regions of potential enhancer elements [32–34]. With this regard, our finding of PITX2-MLL4 association might link the involvement of MLL4 in PITX2-dependent cell differentiation and development.

Acknowledgments

This work was partially supported by the U.S. National Science Foundation under Cooperative Agreement No. EPS-1003907 and the West Virginia state Research Challenge Fund. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

References

1. Lu MF, Pressman C, Dyer R, et al. Function of Rieger syndrome gene in left-right asymmetry and craniofacial development. *Nature*. 1999; 401:276–278. [PubMed: 10499585]
2. Gage PJ, Suh H, Camper SA. Dosage requirement of Pitx2 for development of multiple organs. *Development*. 1999; 126:4643–4651. [PubMed: 10498698]
3. Kioussi C, Briata P, Baek SH, et al. Identification of a Wnt/Dvl/[beta]-catenin --> Pitx2 pathway mediating cell-type-specific proliferation during development. *Cell*. 2002; 111:673–685. [PubMed: 12464179]
4. Lines MA, Kozlowski K, Walter MA. Molecular genetics of Axenfeld-Rieger malformations. *Hum Mol Genet*. 2002; 11:1177–1187. [PubMed: 12015277]
5. Gage PJ, Camper SA. Pituitary homeobox 2, a novel member of the bicoid-related family of homeobox genes, is a potential regulator of anterior structure formation. *Hum Mol Genet*. 1997; 6:457–464. [PubMed: 9147650]
6. Li X, Florez S, Wang J, et al. Dact2 represses PITX2 transcriptional activation and cell proliferation through Wnt/beta-catenin signaling during odontogenesis. *PLoS One*. 2013; 8:e54868. [PubMed: 23349981]
7. Liu C, Liu W, Palie J, et al. Pitx2c patterns anterior myocardium and aortic arch vessels and is required for local cell movement into atrioventricular cushions. *Development*. 2002; 129:5081–5091. [PubMed: 12397115]
8. Gudbjartsson DF, Arnar DO, Helgadóttir A, et al. Variants conferring risk of atrial fibrillation on chromosome 4q25. *Nature*. 2007; 448:353–357. [PubMed: 17603472]
9. Huang Y, Guigon CJ, Fan J, et al. Pituitary homeobox 2 (PITX2) promotes thyroid carcinogenesis by activation of cyclin D2. *Cell Cycle*. 2010; 9:1333–1341. [PubMed: 20372070]
10. Fung FK, Chan DW, Liu VW, et al. Increased expression of PITX2 transcription factor contributes to ovarian cancer progression. *PLoS One*. 2012; 7:e37076. [PubMed: 22615897]
11. Hirose H, Ishii H, Mimori K, et al. The significance of PITX2 overexpression in human colorectal cancer. *Ann Surg Oncol*. 2011; 18:3005–3012. [PubMed: 21479692]
12. Lee W, Chakraborty PK, Thevenod F. Pituitary homeobox 2 (PITX2) protects renal cancer cell lines against doxorubicin toxicity by transcriptional activation of the multidrug transporter ABCB1. *Int J Cancer*. 2013; 133:1–12. [PubMed: 23280579]
13. Zhang JX, Tong ZT, Yang L, et al. PITX2: A promising predictive biomarker of patients' prognosis and chemoradioresistance in esophageal squamous cell carcinoma. *Int J Cancer*. 2013; 132:2567–2577. [PubMed: 23132660]

14. Bamforth SDBJ, Farthing CR, Schneider JE, et al. Cited2 control left-right patterning and heart development through a Nodal-Pitx2c pathway. *Nat Genet.* 2004; 36:1189–1196. [PubMed: 15475956]
15. Baek SH, Kioussi C, Briata P, et al. Regulated subset of G1 growth-control genes in response to derepression by the Wnt pathway. *Proc Natl Acad Sci USA.* 2003; 100:3245–3250. [PubMed: 12629224]
16. L'honoré A, Ouimette JF, Lavertu-Jolin M, et al. Pitx2 defines alternate pathways acting through MyoD during limb and somitic myogenesis. *Development.* 2010; 137:3847–3856. [PubMed: 20978076]
17. Ai D, Wang J, Amen M, et al. Nuclear Factor 1 and T-Cell Factor/LEF Recognition Elements Regulate Pitx2 Transcription in Pituitary Development. *Mol Cell Biol.* 2007; 27:5765–5775. [PubMed: 17562863]
18. Huang Y, Huang K, Boskovic G, et al. Proteomic and genomic analysis of PITX2 interacting and regulating networks. *FEBS Lett.* 2009; 583:638–642. [PubMed: 19174163]
19. Eissenberg JC, Shilatifard A. Histone H3 lysine 4 (H3K4) methylation in development and differentiation. *Dev Biol.* 2010; 339:240–249. [PubMed: 19703438]
20. Ruthenburg AJ, Allis CD, Wysocka J. Methylation of lysine 4 on histone H3: intricacy of writing and reading a single epigenetic mark. *Mol Cell.* 2007; 25:15–30. [PubMed: 17218268]
21. Cho YW, Hong T, Hong S, et al. PTIP associates with MLL3- and MLL4-containing histone H3 lysine 4 methyltransferase complex. *J Biol Chem.* 2007; 282:20395–20406. [PubMed: 17500065]
22. Liu Y, Huang Y, Zhu GZ. Cyclin A1 is a transcriptional target of PITX2 and overexpressed in papillary thyroid carcinoma. *Mol Cell Biochem.* 2013; 384:221–227. [PubMed: 24002705]
23. Gan Q, Thiébaud P, Thézé N, et al. WD repeat-containing protein 5, a ubiquitously expressed histone methyltransferase adaptor protein, regulates smooth muscle cell-selective gene activation through interaction with pituitary homeobox 2. *J Biol Chem.* 2011; 286:21853–21864. [PubMed: 21531708]
24. Cho YW, Hong S, Jin Q, et al. Histone methylation regulator PTIP is required for PPARgamma and C/EBPalpha expression and adipogenesis. *Cell Metab.* 2009; 10:27–39. [PubMed: 19583951]
25. Daniel JA, Santos MA, Wang Z, et al. PTIP promotes chromatin changes critical for immunoglobulin class switch recombination. *Science.* 2010; 329:917–923. [PubMed: 20671152]
26. Schwab KR, Patel SR, Dressler DR. Role of PTIP in class switch recombination and long-range chromatin interactions at the immunoglobulin heavy chain locus. *Mol Cell Biol.* 2011; 31:1503–1511. [PubMed: 21282469]
27. Gong Z, Cho YW, Kim JE, et al. Accumulation of Pax2 transactivation domain interaction protein (PTIP) at sites of DNA breaks via RNF8-dependent pathway is required for cell survival after DNA damage. *J Biol Chem.* 2009; 284:7284–7293. [PubMed: 19124460]
28. Wu J, Prindle MJ, Dressler GR, et al. PTIP regulates 53BP1 and SMC1 at the DNA damage sites. *J Biol Chem.* 2009; 284:18078–18084. [PubMed: 19414588]
29. Wang X, Takenaka K, Takeda S. PTIP promotes DNA double-strand break repair through homologous recombination. *Genes Cells.* 2010; 15:243–254.
30. Munoz IM, Jowsey PA, Toth R, et al. Phospho-epitope binding by the BRCT domains of hPTIP controls multiple aspects of the cellular response to DNA damage. *Nucleic Acids Res.* 2007; 35:5312–5322. [PubMed: 17690115]
31. Patel SR, Kim D, Levitan I, et al. The BRCT-domain containing protein PTIP links PAX2 to a histone H3, lysine 4 methyltransferase complex. *Dev Cell.* 2007; 13:580–592. [PubMed: 17925232]
32. Hu D, Gao X, Morgan MA, et al. The MLL3/MLL4 branches of the COMPASS family function as major histone H3K4 monomethylases at enhancers. *Mol Cell Biol.* 2013; 33:4745–4754. [PubMed: 24081332]
33. Guo C, Chen LH, Huang Y, et al. KMT2D maintains neoplastic cell proliferation and global histone H3 lysine 4 monomethylation. *Oncotarget.* 2013; 4:2144–2153. [PubMed: 24240169]
34. Lee JE, Wang C, Xu S, et al. H3K4 mono- and di-methyltransferase MLL4 is required for enhancer activation during cell differentiation. *Elife.* 2013; 2:e01503. [PubMed: 24368734]

1. PITX2 forms complex with histone H3 lysine 4 (H3K4) methyltransferase
2. PITX2 complex methylates H3K4
3. PITX2 and PTIP co-occupy the promoter of the PITX2's transcriptional target

Fig. 1.

PITX2 associates with MLL4 histone methyltransferase complex. The antibodies for FLAG (PITX2), PTIP, MLL4, ASH2L and RBBP5 were used to precipitate respective antigens and their associated proteins from HEK293 cells expressing FLAG-PITX2. Immunoblotting was used to examine the components of respective protein complex.

Fig. 2.

The association of PITX2 with the H3K4 methyltransferase complex is dependent on its homeodomain. The plasmids containing full-length or truncated PITX2 were expressed in HEK293 cells. Anti-FLAG was used to co-immunoprecipitate PITX2 associated protein complex. RBBP5, a core subunit of the MLL family of H3K4 methyltransferases, was used as a read-out of the H3K4 methyltransferase complex. HD, homeodomain; OAR, Otp and Aristaless domain.

Fig. 3.

PITX2 complex methylates histone H3 on K4. The nuclear extract (NE) of HEK293 cells, the IgG immunoprecipitate complex (prepared from HEK293 cells), the PITX2 Δ HD immunoprecipitate complex (prepared from HEK293 cells expressing homeodomain-deleted PITX2), the PITX2 immunoprecipitate complex (prepared from HEK293 cells expressing full-length PITX2), the PTIP immunoprecipitate complex (prepared from HEK293 cells) or the MLL4 immunoprecipitate complex (prepared from HEK293 cells) was used as a potential source of H3K4 methyltransferases. H3K4 methylation assays were carried out as described in Materials and methods. Immunoblotting of H3 served as loading control.

Fig. 4. PITX2 recruits PTIP-containing H3K4 methyltransferase to the Cyclin A1 promoter. Chromatin immunoprecipitation (ChIP) assays were performed on HEK293 cells transiently expressing homeodomain-deleted PITX2 (PITX2ΔHD) or full-length PITX2 (PITX2). The PCR primers covered the Cyclin A1 promoter (–133 to –7, with the transcription start site as position 1).