Published in final edited form as:

Differentiation. 2013 October ; 86(3): 77–86. doi:10.1016/j.diff.2013.05.001.

Mechanobiology of bone marrow stem cells: from myosin-II forces to compliance of matrix and nucleus

Jae-Won Shin, Joe Swift, Irena Ivanovska, Kyle R. Spinler, Amnon Buxboim, and Dennis E. Discher

Biophysical Engineering Laboratory, University of Pennsylvania, Philadelphia, PA 19104

Abstract

Adult stem cells and progenitors are of great interest for their clinical application as well as their potential to reveal deep sensitivities to microenvironmental factors. The bone marrow is a niche for at least two types of stem cells, and the prototype is the hematopoietic stem cell/progenitors (HSC/Ps), which have saved many thousands of patients for several decades now. In bone marrow, HSC/Ps interact functionally with marrow stromal cells that are often referred to as mesenchymal stem cells (MSCs) or derivatives thereof. Myosin and matrix elasticity greatly effect MSC function, and these mechanobiological factors are now being explored with HSC/Ps both *in vitro* and *in vivo*. Also emerging is a role for the nucleus as a mechanically sensitive organelle that is semi-permeable to transcription factors which are modified for nuclear entry by cytoplasmic mechanobiological pathways. Since therapies envisioned with induced pluripotent stem cells and embryonic stem cells generally involve *in vitro* commitment to an adult stem cell or progenitor, a very deep understanding of stem cell mechanobiology is essential to progress with these multi-potent cells.

Keywords

Hematopoiesis; myosin; matrix; nucleus; mechanobiology

1. Introduction

Since the first human bone marrow transplantation (BMT) (Thomas et al., 1957) and the revelation of hematopoietic stem cell/progenitors (HSC/Ps) by clonal assay (Becker et al., 1963), adult stem cells of many tissue types have become an intense focus of research. These cells can either integrate in a target tissue or act as vehicles to deliver important signals to a tissue without integration (Daley and Scadden, 2008). More than 50,000 BMT procedures were performed in the year 2006 alone, saving a large percentage of treated patients but not all (Gratwohl et al., 2010). Such clinical advances and limitations have motivated investigators to elucidate mechanisms by which a few or perhaps just a single stem cell can generate an entire, regenerating tissue. Indeed, the demands are great in human hematopoiesis with 10^5 to 10^6 nucleated blood cells, anucleated red blood cells (RBCs), and platelets produced every second in marrow to enter blood. New approaches for HSC and HSC-derived blood cell transplantation are still needed to generate potent donor cells that

engraft efficiently. Factors that regulate proliferation, differentiation, and dispersion of adult stem cells continue to be elaborated, and include physical factors (eg. Figure 1) that are the focus of this Review. Some of the latest findings and methods for blood cells are likely to prove applicable to other types of adult stem cells and progenitors such as bone-marrow derived mesenchymal stem/stromal cells (MSCs) (Friedenstein et al., 1976). MSCs are also rapidly emerging as clinically-relevant (Hare et al., 2012) and have proven to be highly mechanosensitive – as will be reviewed. Such findings have stimulated new mechanical approaches to other stem cells including HSC/Ps.

The hierarchical nature of blood cell development has been elucidated through advances in the identification and isolation of HSC/Ps *versus* derived lineages. Particularly critical has been the prospective isolation of HSC/Ps by flow cytometry with a specific set of cell surface antigens (Weissman and Shizuru, 2008) combined with limiting dilution transplantations that quantify HSC frequency (Szilvassy et al., 1990). Aided by these methods, a number of soluble growth factors have been identified that support the differentiation of various blood lineages. However, much less is known about physical factors *in vivo* that might govern the self-renewal and function of HSCs. A stem cell “niche” or “microenvironment” was proposed to be a requirement for long-term hematopoiesis from HSCs (Schofield, 1978). Deeper understanding of the niche is likely to foster strategies to direct HSC/Ps *ex vivo* in order to enhance *in vivo* engraftment in patients. Such an effort will also benefit sustained *ex vivo* production of more mature blood cell types, especially RBCs and platelets that are transfused from donors in massive quantities today. While human erythropoietin (Miyake et al., 1977) and thrombopoietin (Kaushansky et al., 1994) have emerged as major factors to respectively facilitate erythroid and megakaryocyte (MK) differentiation from HSC/Ps, sustained generation *in vitro* of terminally differentiated cells has remained a major challenge. Indeed, none of the growth factors, cytokines or chemokines has yet proven sufficient to reproduce artificial marrow-like environments conducive to generation of all blood cell-types from HSC/Ps.

Mechanotransduction refers to the conversion of extracellular mechanical inputs to intracellular signals, both biochemical and biophysical (Wang et al., 2009). It is mediated in part by intracellular tension that is sustained by adhesion to matrix or other cells and generated by the actin-myosin-based cytoskeleton (Discher et al., 2005) which ultimately couples to the nucleus (e.g. Pajerowski et al., 2007). Given that HSC/Ps, MSCs and derived lineages interact in many ways with their microenvironments, mechanical aspects of microenvironment can in principle regulate stem and progenitor cell functionality. As reviewed here, stem cells generate forces in processes ranging from cell division to migration, while external stresses in stem cell microenvironments that include fluid flows and pressures will impact adhesion and associated signaling as do mechanical factors such as matrix elasticity for MSCs (Discher et al., 2005) and nuclear elasticity of HSC/Ps (e.g. Pajerowski et al., 2007; Shin et al., 2011). All of these structures with mechanical functions can in principle impact stem cell maintenance, lineage specification, and trafficking. We will discuss recent progress in the mechanobiology of adult stem cells and progenitors, with particular emphasis on bone marrow derived HSC/Ps and MSCs. Functional roles of actomyosin forces in stem cells are first introduced, followed by both cell-intrinsic and extrinsic biophysical properties. We then discuss the applicability of mechanobiology across different topics as well as the engineering of artificial marrow models. Throughout we try to highlight how insights gained from studies with MSCs can be relevant to the underlying mechanobiology of HSC/Ps and vice versa.

2. Actomyosin force regulation in stem cell and progenitor functions

In response to extracellular mechanical processes that range from matrix adhesion to fluid shear, cells generate intracellular forces with actin and myosin. This force generation process contributes to diverse biological functions that are relevant to differentiation, including cytokinesis, cortical tension, and migration. Here, we discuss the molecular basics of mammalian non-muscle myosin II (NMM-II), emphasizing hematopoietic and mesenchymal tissues and stem cells/progenitor contexts.

Actin is a principal filamentous constituent of the cytoskeleton that also translocates (as monomer) into the nucleus. Myosin motor proteins pull on actin filaments by dynamically crosslinking and translating along the filaments driven by the hydrolysis energy of ATP (Pollard and Korn, 1973). Such activities can impact actin monomer pools (Wilson et al., 2010) that might even regulate the serum response factor (SRF) pathway. Prototypical muscle myosins are type-II myosins that self-assemble into microns-long bipolar filaments, which register in striations (Brown et al., 2009). Non-muscle cells express isoforms of NMM-II that assemble into smaller mini-filaments that appear far less ordered than in muscle. The three isoforms of the NMM-II heavy chain, A, B and C, are encoded by the genes *MYH9*, *MYH10* and *MYH14*. *MYH9* expression is regulated in part by SRF which is an actin regulated transcription factor (Olson and Nordheim, 2010). NMM-IIA is widely expressed across different tissues, while NMM-IIB is expressed in cardiac and neural tissue, and NMM-IIC appears largely restricted to epithelial cells (Ma et al., 2010). At the molecular level, NMM-IIA has the highest rate of ATP hydrolysis compared to other isoforms (Kim et al., 2005), while NMM-IIB has a higher affinity for ADP, leading to a longer period of binding to actin filaments during force generation (higher duty ratio) than NMM-IIA (Wang et al., 2003). Hematopoietic cells and MSCs express A and B isoforms but not C, and these cells show dynamic expression patterns during differentiation (Engler et al., 2006).

Genetic methods show that NMM-II is essential to diverse biological functions, including embryo survival and cytokinesis, cortical tension and stiffness generation, and also sensing of matrix and shear stress. NMM-IIA knockout is embryonic lethal to mice at stage E6.5 with defects in visceral endoderm morphogenesis and overall tension of the embryo, which appears flaccid (Conti et al., 2004): essentially, no differentiation occurs even though the cells adhere and divide. NMM-IIB knockout mice survive longer but die at E14.5 due to abnormalities in heart and brain that include more binucleated cells (Takeda et al., 2003; Tullio et al., 1997). NMM-IIC knockout mice remain viable to adulthood with no obvious difference in phenotypes compared to wild-type mice, although the double B, C knockout show impaired cytokinesis compared to single NMM-IIB knockout mice (Ma et al., 2010). Deletion of myosin-II in the amoeba *Dictyostelium* leads to increased distortion of the cell membrane and multi-nucleation in suspension cultures where cells cannot crawl apart to facilitate division (Merkel et al., 2000). Pharmacological inhibition of NMM-II in adherent cells by blebbistatin (Bleb) likewise tends to increase the number of binucleate cells, indicating that NMM-II contributes to efficient cytokinesis (Straight et al., 2003). Importantly, NMM-II also plays a critical role in matrix sensing: Bleb eliminates the effect of matrix elasticity on lineage specification of MSCs (Engler et al., 2006) and also on maintenance of HSC/P numbers on highly flexible tropoelastin surfaces (Holst et al., 2010), as elaborated upon below.

NMM-II levels vary greatly between cell types and during differentiation, but NMM-II also responds rapidly under physiological settings to intracellular and extracellular forces through changes in its phosphorylation states. NMM-II consists of two heavy chains (230 kDa), two essential light chains (17 kDa) and two regulatory light chains. Myosin heavy

chain consists of an ATP-hydrolyzing N-terminal motor or head domain, which directly interacts with F-actin, and a ~150 nm long C-terminal helical tail which forms coiled-coil dimers and can further assemble into myosin filaments (Adelstein et al., 1972). Essential light chains are required for stabilizing the heavy chain (Hernandez et al., 2007). A number of studies (see (Matsumura, 2005)) have shown that regulatory light chains (RLC) modulate NMM-II activity via phosphorylation by numerous kinases, including myosin light chain kinase (MLCK) and Rho-associated protein kinase (ROCK), both of which can phosphorylate the Thr18 and Ser19 RLC residues. In heavy chains such as NMM-IIA, phosphorylation sites exist in the tails (e.g. Ser1943) and heads (e.g. Tyr277). The tail site has been found in MSCs to be more phosphorylated on soft matrix, where NMM-IIA is down-regulated in terms of expression levels, assembly state, and contributions to traction forces (Raab et al., 2012). Protein kinase C and casein kinase-II phosphorylate the heavy chain but additional kinases are likely relevant (Dulyaninova et al., 2005; Even-Faitelson and Ravid, 2006). Since heavy chains bind F-actin directly, regulation of the heavy chain seems likely to have the most direct impact on NMM-II activity compared to RLC phosphorylation. Indeed, recent mutagenesis studies indicate that NMM-IIA phosphorylation regulates the polyploidization of MKs more dramatically than, for example, inhibitors of MLCK or ROCK (Shin et al., 2011).

Although myosin-II's functions in early HSC/Ps remain unclear, it plays critical roles in differentiated hematopoietic lineages. In lymphoid cells, NMM-IIA is required for cytolytic granule exocytosis in human natural killer cells (Andzelm et al., 2007), endocytosis of stromal derived factor-1 (SDF-1) receptor (Rey et al., 2007), and polarization of endocytic clathrin at the uropod of T-cells (Samaniego et al., 2007). NMM-IIA is also required for immunological synapse stabilization and T-cell receptor microcluster assembly and trafficking (Ilani et al., 2009) as well as T-cell motility in three-dimensions (Jacobelli et al., 2010). NMM-II is required for B-cell receptor-driven antigen presentation (Vascotto et al., 2007). In myeloid cells, dendritic cell migration upon cytokine stimulation (Fernandez et al., 2011) or via CD74 (Faure-Andre et al., 2008) requires NMM-II as does CD47-mediated deactivation of macrophage phagocytosis (Tsai and Discher, 2008) and traction force generation by neutrophils in chemotaxis (Shin et al., 2010). In erythroid lineages, actin-mediated forces contribute to enucleation (Koury et al., 1989), and both colony forming unit-erythroid cytokinesis and erythroid enucleation are blocked by NMM-II inhibition, with NMM-IIB but not IIA required for enucleation (Ubukawa et al., 2011). In MK lineages, *MYH9* mutations associate with May-Hegglin anomaly, which is characterized by low platelet count (thrombocytopenia) and giant platelets (Kelley et al., 2000; Seri et al., 2000). Such genetic changes have been recapitulated recently in mice that harbor mutations of *MYH9* and show longer bleeding times and defects in clot retraction (Zhang et al., 2011). While *MYH9* mutations appear to impair platelet function, paradoxically, other previous studies using *in vitro* MKs show increased proplatelet formation upon NMM-II knockout or pharmacological inhibition (Chen et al., 2007; Eckly et al., 2009). Indeed, Bleb promotes platelet-like fragmentation by the fluid stresses that mimic shear stresses in capillaries of the marrow (Shin et al., 2011). Despite these enigmatic issues, actomyosin forces are clearly an attractive target to mechanically modulate hematopoiesis by synergizing with soluble factors in lineage specification and/or function.

3. Trafficking out of the Niche: Mechanical control *via* cortical and nuclear deformability

For stem cells and the lineages they generate, adhesion, retention, migration, and escape from a niche are all programmatic processes predicated in part on cell mechanics. While some mature blood cells such as RBC must circulate throughout the body for their function,

at least some of the immature HSC/Ps must be retained in marrow to sustain hematopoiesis. It has long been appreciated that the deformability of hematopoietic cells changes during differentiation, and so deformability is a likely factor in regulating cell traffickability under mechanically-constrained conditions that involve migration across the marrow-blood endothelial barrier and into 2–3 μm diameter capillaries. An early micropipette study showed that a 20-fold increase in aspiration pressure was required to deform myeloid progenitors (immature myeloblasts) compared to highly deformable mature granulocytes (Lichtman, 1970). Myeloblasts also cannot transit through 1 – 8 μm pores under a chemotatic gradient, whereas mature granulocytes can (Giordano and Lichtman, 1973). While studies of other soft tissue cell types, such as glioma cells have established a critical role for NMM-IIA in migration through capillary-size pores of 3 μm diameter (Beadle et al., 2008), how hematopoietic differentiation leads to changes in the physical deformability of cells and hence affects trafficking has not been deeply investigated. The same challenge applies to MSCs traffickability, although recent studies of MSC migration through soft, 3D collagen gels have revealed a key role for NMM-IIA (via heavy chain phosphorylation) in elasticity sensing and NMM-IIB in directional persistence (Raab et al., 2012). More generally, both cell cortex and nucleus likely contribute to overall cell deformability and thereby impact cell motility, niche retention, and trafficking.

3.1. Mechanical properties of membrane cortex and myosin-II contributions

In light of the flow stresses on blood cells, membrane deformability has been characterized for platelets (White et al., 1984), megakaryocytes (Smith et al., 1989), granulocyte and lymphocytes (Dong et al., 1991), and red blood cells at a detailed molecular level (Discher et al., 1994). It is clear from all of these studies that the specialized cytoskeletons are particularly key and part of each cell types program. In contrast to granulocyte differentiation, immature erythroid progenitors are stiffer than mature RBCs (Waugh et al., 2001), which could be coupled with nuclear condensation during erythropoiesis. Nucleus stiffening could be a driving factor for enucleation to allow egress of RBCs (Lichtman et al., 1989). In contrast to differentiated cells, membrane cortex elasticity remains to be characterized further for early HSC/Ps, but studies with other systems illustrate a common theme that may be relevant to HSC/Ps. In general, primitive stem cells tend to be soft, perhaps due to the immature cytoskeletal architecture; and thus allow cell plasticity to be particularly susceptible to physical changes in external forces in their surroundings. Undifferentiated ESCs are, for example, 10-fold softer than differentiated progeny and possess higher sensitivity to a local cyclic stress that triggers cell spreading and differentiation (Chowdhury et al., 2010). Likewise, primitive MSCs possess a lower cortical elastic modulus compared to differentiated fibroblasts (Titushkin and Cho, 2007).

Since myosin-II regulates cortical stiffness, whether HSC/Ps are softer than differentiated cells is likely to depend on the absolute abundance of myosin-II isoforms during hematopoiesis. Conversely, myosin-II itself is sensitive to external force; for example, myosin-II becomes polarized and assembles in regions of high stress (Reichl et al., 2008). In addition, both NMM-IIA and NMM-IIB are upregulated in MSCs upon adhesion to stiff matrix compared to soft matrix (Engler et al., 2006). In migration of MSCs from soft to stiff matrix, NMM-IIB becomes polarized rearward, while NMM-IIA remains unpolarized (Raab et al., 2012), indicating that different isoforms have distinct functional roles in cortical deformability, migration and trafficking. In the epithelial-to-mesenchymal transition, cells undergo myosin-II isoform switching from C to B, promoting cell motility and invasion (Beach et al., 2011). This insight has now been extended to the hematopoietic system where isoform switching is seen to regulate differentiation and motility (Shin et al., in revision).

3.2. Mechanical properties of the nucleus

In contrast to MSCs, hematopoietic cells have a small cytoplasmic volume, and so in addition to cell cortex, the nucleus is likely to contribute to overall cell deformability and trafficking. Nuclear deformability is largely understudied across the hematopoietic system. We have begun to approach this issue with two micropipette approaches involving either aspiration of isolated nuclei or aspiration of nuclei in live cells treated with actin polymerization inhibitors (Ivanovska et al., 2010). Our previous studies have indicated that nuclei in human hematopoietic CD34⁺ cells from marrow were slightly stiffer than nuclei in human ESCs but considerably softer than other differentiated cell nuclei, suggesting that nuclear deformability is developmentally regulated (Pajerowski et al., 2007). The same study further demonstrated that nuclear stiffness is dependent on lamins, which are intermediate filaments that assemble as coiled-coil dimers at the nuclear envelope and associate at least indirectly with chromatin (Shimi et al., 2010). While our initial micropipette study suggested the involvement of lamins and nuclear stiffness in 3D entry into pores, and a recent review speculated that this concept also applies to cancer cell migration and metastasis (Friedl et al., 2011), direct evidence to support the relevance of nuclear deformability in the hematopoietic system is just emerging (Shin et al., in review). One notion, for example, is that the polyploid nucleus of an MK is too large and rigid with a complex lamina to allow transmigration into blood, but the consequent marrow anchorage still allows these cells to extend anuclear proplatelet projections into blood that then fragment under shear flow to make platelets as seen recently (Junt et al., 2007). As another example, lamin-A,C is strongly upregulated upon T-cell activation (Andrade et al., 2003) and this could similarly favor retention of a T-cell at a site of inflammation. Similar to the hypothesis that NMM-II actively regulates deformability of the cell cortex, it appears reasonable to hypothesize that differential lamin expression patterns also regulate nuclear stiffness throughout hematopoiesis, influencing the ability of cells to migrate through and within the dense marrow or tissue and across an endothelium into blood. MSCs also express lamins and their levels can at least affect differentiation (Akter et al., 2009), but any relationship to nuclear mechanics, retention in a niche, or broader trafficking is thus far unclear.

4. Mechanical control of differentiation by forces during cell division

Since cytoskeleton proteins contribute cell division forces through the mitotic spindle and also in the cytokinetic ring of actin and NMM-II (Scholey et al., 2003), it is likely that these forces contribute to lineage specification. In particular, for hematopoietic progenitors and T-cells, proteins segregate asymmetrically during asymmetric division, providing an important basis for self-renewal and differentiation (Chang et al., 2007; Giebel et al., 2006; Ting et al., 2012). However, involvement of cytoskeletal machinery in the asymmetry has not been addressed and clearly needs to be. For example, polarized adhesion of one of the dividing daughter cells to an antigen-presenting cell can asymmetrically segregate a number of proteins such as atypical protein kinase C (Chang et al., 2007). A couple of cytoskeletal components, including Rac GTPase activating protein 1 and Tropomodulin 1, segregate asymmetrically during division, and overexpression of these proteins enhance HSC engraftment in vivo (Ting et al., 2012). Pharmacological inhibition of NMM-II has also been found to combine with soft matrix to maximize MK polyploidization (Shin et al., 2011), consistent with the more recent finding that the NMM-IIIB isoform specifically regulates MK maturation (Lordier et al., 2012). Distinct roles of NMM-II isoforms in HSC/P differentiation are just beginning to be elucidated (Shin et al., in revision), and more attention is generally needed to asymmetric division of MSCs in differentiation processes.

5. Niches and Microenvironment mechanical properties

MSCs were originally isolated as the plastic adherent cells from marrow and certainly have characteristics typical of cells that adhere within solid tissues (Engler et al., 2006). HSC/Ps and derived lineages do not adhere to plastic but do thrive in a range of solid-like tissues, including microenvironments in bone marrow (BM: HSC/Ps and different lineages), spleen (B-cells, MK/erythroid lineages, and HSC/Ps) and thymus (T-cells). However, unlike solid tissue cells such as MSCs, many blood cell types are also viable for days to months or longer while suspended in blood, which is fluid of course. Only a few studies have been conducted recently to elucidate roles of cell-extrinsic mechanical factors on hematopoietic functions (e.g. fluid suspensions *versus* adhesion to soft or stiff solids), but studies over the last 20 years have helped to identify cell types in niches as well as molecular factors that influence hematopoietic differentiation and trafficking.

The endosteum in BM has been viewed traditionally as a major contributor to the HSC niche. Osteoblasts (OBs) – which MSCs can differentiate to (Engler et al., 2006) – are located at the endosteum and provide necessary cytokines to facilitate myeloid differentiation and B lymphopoiesis from HSC/Ps (Taichman and Emerson, 1994; Zhu et al., 2007). In addition, genetic manipulations show that increased OB numbers lead to increased HSC numbers (Calvi et al., 2003; Zhang et al., 2003). Specific molecular signals emanating from OBs are known to regulate HSC functions and include Angiopoietin-1 and Tie-2 which maintain HSC quiescence (Arai et al., 2004) as well as Osteopontin, which suppresses HSC numbers (Nilsson et al., 2005; Stier et al., 2005). The notion that the endosteum is the main HSC niche component has been challenged recently by the discovery that long-term HSCs (LT-HSCs) are frequently located near BM vascular cells (Kiel et al., 2005); the BM vasculature consists of sinusoids, which are thinwalled blood vessels lined by endothelial cells. While it appears difficult to distinguish vascular niches from nearby endosteal niches (Kiel et al., 2005), the ability of HSCs to self-renew and differentiate during fetal development in spleen (Morita et al., 2011) and liver (Taniguchi et al., 1996) indicates that HSC functions can be maintained without the involvement of bone, although the species specific nature of such findings (mouse *versus* man) remains unresolved.

Sinusoidal endothelial cells are required for long-term hematopoietic reconstitution by donor HSCs, since deletion of vascular endothelial growth factor receptor-2 leads to failure in regeneration of the endothelium after sub-lethal irradiation, thereby preventing restoration of hematopoiesis (Hooper et al., 2009). Furthermore, endothelial cells facilitate functional expansion of HSC/Ps upon protein kinase B activation, while mitogen-activated protein kinase activation leads to inhibition of HSC/P expansion and shifts cells towards maintenance and differentiation (Kobayashi et al., 2010). In addition, endothelial cells support myeloid and MK differentiation in culture (Rafii et al., 1995). Given that HSCs are now further divided into dormant and active populations, a recent hypothesis is that dormant HSCs tend to be located at the endosteum, which is devoid of endothelial cells, while active HSCs may be located at the vascular niche, and that the HSCs can be stimulated or stressed to traffic between the niches (Trumpp et al., 2010) – even if these niches are not so anatomically distinct as mentioned. The cell and molecular nature of HSC/P niches has become even more complicated because other regulators have been demonstrated, including SDF-1 abundant reticular stromal niches (Sugiyama et al., 2006), osteoprogenitors nearby BM sinusoids (Sacchetti et al., 2007), BM adipocytes (Naveiras et al., 2009), and nestin-positive pericyte-like cells (Mendez-Ferrer et al., 2010), all of which are either MSCs or derivable from MSCs.

Mechanics of marrow environments at the sub-cellular level can in principle be quantified by methods such as atomic force microscopy (AFM), but studies are thusfar limited.

Macroscopic measurements of extracted bone marrow suggest an elasticity of ~0.3 kPa (Winer et al., 2009) although removal of marrow can be highly disruptive to structure and mechanics. *In situ* micro-scale measurements with AFM of marrow show the central marrow to be moderately heterogeneous but generally soft (< 0.3 kPa) while bone is invariably rigid (> 1,000 kPa) (Shin et al., in revision). Osteoblasts in culture secrete a thin, endosteal-like extracellular matrix (ECM) measured by AFM to be 20–40 kPa. Remarkably, when MSCs are grown on collagen-I gels in this range of matrix elasticity, osteogenesis predominates, whereas softer gels that mimic muscle (~10 kPa) prime MSCs for a myogenic lineage (Engler et al., 2006). Myosin-II activity has been found to be required for such lineage specification *in vitro*, but controlled *in vivo* studies are needed to thoroughly address molecular mechanisms. As illustration of the important molecular questions that arise, calcifications in the heart have been seen to result from mouse-MSCs that are injected into a myocardial infarct (Breitbach et al., 2007), which remodels to a stiff fibrotic scar ($E \sim 35\text{--}70$ kPa) that is several-fold stiffer than normal myocardium and mechanically more like endosteal-like ECM (Berry et al., 2006).

Consideration of how MSCs differentiate across different elasticities has important functional implications for understanding MSCs as well as HSC/P differentiation. While some of the known cell types that are key to HSC niches are of mesenchymal origin, relative contributions of immature MSCs and mature lineages to HSC functions remains to be quantified in parallel. This could be challenging using gene or promoter specific approaches. For instance, while nestin⁺ MSCs appear more pericyte-like and distinct from osteochondrial lineages, parathyroid hormone may activate both of them and accelerate differentiation of MSCs into osteoblasts to increase HSC number (Mendez-Ferrer et al., 2010). The col2.3 promoter transgene bearing mice have been widely used to deplete osteoblasts (Visnjic et al., 2001) to study HSC functions (Calvi et al., 2003). However, advances in MSC biology have led to speculation that other mesenchymal lineages – which express and respond to type I collagen (Engler et al., 2006) – could be affected as well. Therefore, controlled approaches to modulate physical properties of ECM *in vitro* and *in vivo* (by direct implantation, see below) could be useful to further characterize MSC and HSC niches.

Studies in other systems have since confirmed that stem cell maintenance and/or differentiation is indeed regulated by tissue microelasticity. Neural stem cells develop into neurons on soft matrices (< 1 kPa) whereas glial cells dominate on stiffer matrices (Saha et al., 2008). Skeletal muscle stem cells cultured on compliant muscle-mimetic hydrogels (~10 kPa) in contrast to rigid plastic dishes are able to maintain stemness and contribute to muscle regeneration *in vivo* (Gilbert et al., 2010). Particularly relevant to the present focus, HSC/P number is maintained (and perhaps increased) on highly flexible tropoelastin surfaces compared to stiff, heavily cross-linked tropoelastin (Holst et al., 2010), and myosin-II activity once again seems critical. Whether a soft, elastic environment generically facilitates stem cell expansion *versus* maintenance or differentiation needs to be studied in greater mechanistic depth, especially in light of a recent finding that functional tumor-initiating cells are enriched, rather than expanded, in soft 3D fibrin gels (Liu et al., 2012). Regardless, the *in vitro* studies reviewed above provide compelling evidence that the mechanics of a niche can have a regulatory role in stem cell fates.

As implied by extramedullary hematopoiesis (eg. liver and spleen), microenvironment factors can also regulate hematopoietic trafficking. A number of soluble factors, including SDF-1 and granulocyte stimulating factor (G-CSF) impact the trafficking of HSC/Ps and mature cells (Mazo et al., 2011), since they influence direct interaction between HSC/Ps and stromal or endothelial cell niches (Dar et al., 2005; Sugiyama et al., 2006), and are exploited clinically for HSC/P mobilization (DiPersio et al., 2009). In addition, receptors for ECM-

engaging factors, such as integrins $\alpha 2$, $\alpha 4/\alpha 5/\beta 1$, $\alpha 6$ that respectively interact with collagen, fibronectin, and laminin, have been implicated in HSC trafficking and self-renewal (Benveniste et al., 2010; Gu et al., 2003; Notta et al., 2011; Papayannopoulou et al., 1998). Consistent with findings for marrow-derived, $CD34^-$ MSCs (Engler et al., 2006), adhesion and motility of human $CD34^+$ cells are enhanced on stiff compared to soft fibronectin matrix (Lee-Thedieck et al., 2012), and cell spreading and morphology depend on substrate elasticity and dimensionality of collagen hydrogels (Choi and Harley, 2012). While both of these *in vitro* studies reveal matrix elasticity as a potential factor in regulation of hematopoietic trafficking, the *in vivo* relevance of these results as well as roles of other cell extrinsic mechanical factors require further investigation.

Some insight is perhaps provided by studies of focal adhesion kinase (FAK), which has an important role in mechanotransduction of signals from stiff matrix to most solid tissue cell types (Wang et al., 2001); while the formation of focal adhesions is unexpected for hematopoietic cells due to a lack of stress fibers in cells on stiff matrix, deletion of the FAK gene nonetheless increases MK ploidy and platelet number (Hitchcock et al., 2008). This seems consistent with the phenotype observed with myosin-II inhibition and soft matrix (Shin et al., 2011), although platelet adhesion and spreading are predictably impaired with ablation of FAK. Activation of FAK also associates with B-cell differentiation and trafficking (Glodek et al., 2003). FAK deletion does not intrinsically impair long-term HSC functions and steady-state hematopoiesis (Lu et al., 2012), but FAK signaling does maintain *primitive* acute leukemia cells (Despeaux et al., 2012). Such genetic approaches to key mechanosensing molecules (e.g. myosin-II isoforms) within stem cells in health and disease are likely to provide even more insight in the future.

6. Shear flow physiologically modulates lineage differentiation

Another important extrinsic biophysical factor to consider in differentiation is fluid shear forces, which have been investigated most extensively with endothelial cells (due to direct exposure to blood flow) but also to some extent with osteocytes which experience distinct fluid flows as bone is stressed in daily activity (Weinbaum et al., 1994). Endothelial cells can be differentiated from ESCs upon shear flow *in vitro* via the downstream activation of histone deacetylase 3 (Zeng et al., 2006), and osteoblast differentiation from BM-derived MSCs is facilitated by continuous shear *in vitro* (Stiehler et al., 2009). In addition, controlled shear flow *in vitro* induces differentiation from embryoid-body-derived cells into hematopoietic precursors as indicated by a strong upregulation of the Runx1 transcription factor (Adamo et al., 2009). Zebrafish models were likewise used to show that both chemical and genetic ablation of blood flow severely reduces HSC number (North et al., 2009).

An interplay between intrinsic and extrinsic mechanical factors likely contributes to cell fates. Fluid shear produces forces that are balanced by cell adhesions and the actomyosin cytoskeleton, or else fluid shear can be so strong that it dominates to convect and disperse cells or even fragment them such as shown in platelet biogenesis (Junt et al., 2007). Moreover, an absence of adhesion to external surfaces for many cell types limits cytokinesis, with cells tending to become multi-nucleated in suspension (if they survive) whereas adhesion-based traction forces help cells pull apart (Ben-Ze'ev and Raz, 1981). MK polyploidization is thus inhibited on stiff matrix where increased adhesion maximizes the traction forces for cell division even when myosin-II is inhibited (Shin et al., 2011). On the other hand, cell cycle progression can also be inhibited by soft matrix for at least some adherent cells (Klein et al., 2009). Adhesion, matrix elasticity, and external forces thus couple to myosin in processes central to stem cell fates such as cell division – whether it is symmetric, asymmetric, or suppressed.

7. Engineering stem cell niches: *In vivo* material approaches that control mechanics

Ectopic implants of materials with bone marrow cells or fragments have long been known to support hematopoiesis. A cellulose membrane placed in the peritoneum of mice will predictably produce the usual foreign body response with macrophages and other adherent cells recruited to its surface within days, but when such mice are irradiated and donor marrow cells are infused intraperitoneally (not intravenously), hematopoietic colonies form on the cell layer (Seki, 1973). Macrophages contribute to HSC niches (Winkler et al., 2010) and certainly have a key role in erythroid development (Chasis and Mohandas, 2008), but the colonies on cellulose were mostly granulocytic with limited erythroid and no MK or lymphoid lineages. A tubular cellulose membrane pre-coated with stromal cells derived either from the spleen or marrow (and probably including MSCs) is capable of full trilineage reconstitution upon intraperitoneal transplantation of donor marrow cells (Knospe, 1989).

With a better understanding of biomaterials and stem cells, a number of new ectopic transplant models have been proposed recently as useful tools to examine roles of microenvironment in stem cell biology. One study performed subcutaneous transplantation of mouse marrow stromal cells (MSCs) encapsulated in gelatin sponges to form an ossicle, but microvessel density, lineage number, and HSC/P number in the ossicle proves to be far lower than in the femur (Song et al., 2010). Interestingly, adipocyte number is ~30-fold higher in the ossicle than the femur, consistent with a previous observation that marrow adipocytes negatively regulate HSC number and engraftment (Naveiras et al., 2009). In another study, a polyacrylamide-based hydrogel scaffold was seeded with *human* marrow stromal cells, and subcutaneous transplantation of this cell-laden scaffold was found to first induce angiogenesis by recruiting *mouse* endothelial cells, followed by recruitment of endogenous Lin⁻Sca-1⁻c-Kit⁺ (LSK) progenitor cells (Lee et al., 2012). While factors released from human cells may thus be able to recruit mouse niche cells, the LSK frequency in this model is ~5-fold lower than that in femur and the functionality of the recruited HSC/Ps still needs to be confirmed *in vivo*. The hydrogel scaffold in the cited study has a stiffness about 50-fold higher than that of central marrow (0.3 kPa, see above), and the % engraftment of human marrow blood cells in the scaffold is far lower than that achieved in marrow by intravenous transplantation. Recent efforts with softer ECM-based scaffolds appear to show the *in vivo* formation of a bone organ with better capabilities to recruit and/or maintain HSCs. One study used matrigel that has an elastic modulus of 0.5kPa (Soofi et al., 2009), and showed that the % engraftment of human blood cells in the extramedullary bone is comparable to that in femur (Chen et al., 2012). Most recently, human MSCs were seeded in collagen-based scaffolds (1~2kPa elastic modulus) and supplemented with IL-1 β to induce endochondral bone formation followed by implantation *in vivo*, demonstrating recruitment of endogenous functional mouse HSCs comparable to femoral HSCs (Scotti et al., 2013). These implanted material models are thus increasingly promising mimics of bone marrow.

While the recent studies highlight useful strategies to engineer stem cell niches, our understanding of how mechanical factors regulate stem cells remains to be leveraged systematically to improve such strategies. Further biomaterial strategies to better mimic hematopoietic microenvironments *in vivo* could be adapted from bone regeneration studies where growth factors are gradually delivered by subcutaneous implantation of scaffolds to generate marrow-containing ectopic bones in a cell-free manner (Alsberg et al., 2002). This type of approach could therefore be revealing when combined with insights into the role of matrix mechanics as well as niche geometries, and/or porosity.

8. Mechanotransduction lessons from bone marrow derived MSCs

Although little work has been done to study mechanotransduction in hematopoietic systems, the mechanistic insights gained from studies with MSCs – or variations thereof – are likely applicable. Cultures of MSCs on collagen-coated polyacrylamide hydrogels of tuned elasticity have provided evidence of lineage specification in terms of cell morphology, protein expression, and mRNA profiles consistent with neurogenesis on soft brain-like matrix (0.1 – 1 kPa), muscle on stiffer matrix (8 – 17 kPa), and bone on the stiffest matrix (25 – 40 kPa) (Engler et al., 2006). Cell fates to fat and bone can also be affected by constraining cell size and geometry (Fu et al., 2010; McBeath et al., 2004). The cascade of events that lead from mechanical stimulation exterior to the cell to modulation of a presumed transcriptional response within the nucleus is not understood. A direct pathway of mechanotransduction from the microenvironment seems possible with mechanical linking of ECM through focal adhesions to the cytoskeleton and across the nuclear membrane via the linking nucleus to cytoskeleton (LINC) complex to the nuclear lamina and chromatin. This could bring about changes in physical conformation at any step, particularly those that distinguish active euchromatin from tightly packed, inactive heterochromatin that tends to concentrate near the lamina. An application of force to the surface of HeLa cells *via* magnetic beads can alter chromatin organization and deform the nucleus (Iyer et al., 2012b), which could be important given the spatial location of particular chromosomal territories within the nucleus that broadly regulate transcriptional programs (Iyer et al., 2012a).

Force-induced changes in protein conformation, complex formation and/or post-translational modification can influence cellular or nuclear localization of key molecules (perhaps including translocation from cytoplasm to nucleus) or direct activation or inactivation of transcription factors. Force sensitive mechanisms at focal adhesions are clearly able to regulate actin assembly and the force-generating machinery in cells (Geiger et al., 2009). In human keratinocytes, the assembly state of actin – which couples to matrix stiffness that drives cell spreading – reportedly dictates the cellular location of megakaryocytic acute leukemia protein (MAL or MKL) and its interaction with SRF, thus regulating differentiation (Connelly et al., 2010; Iyer et al., 2012b). The cellular location of transcription factor yes-associated protein (YAP) in MSCs and endothelial cells is also influenced by matrix softness as well as constraints on cell area, both of which favor cytoplasmic YAP whereas stiff matrix and unconstrained cell area promotes translocation to the nucleus (Dupont et al., 2011). The impact on differentiation requires tension in the actomyosin cytoskeleton (inhibited by Bleb) and the activity of Rho GTPase but is not governed by the Hippo/ Large tumor suppressor (LATS) cascade that is normally associated with YAP in pathways controlling organ size (Zhao et al., 2011). Co-factors and/or post-translational modifications that might control this translocation are an active topic of research, with hints of regulation by the nuclear lamina (Swift et al., in revision).

In the hematopoietic system, deletion of SRF increases HSC/P numbers by 3-fold without long-term modification of cell cycle dynamics, but it severely impairs cell retention in BM partly due to downregulation of integrins and hence of adhesion (Ragu et al., 2010). MAL deletion inhibits MK maturation and proplatelet formation (Gilles et al., 2009), while MAL overexpression increases MK differentiation as long as SRF is expressed normally (Cheng et al., 2009). YAP can be detected in a broad range of tissues, spanning from the very soft (brain) to the very stiff (bone), although protein levels do not appear to be a monotonic function of tissue stiffness (Swift et al., in revision). YAP is detectable at low levels in long-term HSCs but is barely detectable in other HSC/P or lineages, consistent with the marrow being very soft (see above). Overexpression of YAP does not alter normal hematopoiesis (Jansson and Larsson, 2012), although effects of YAP deletion have not yet been reported. As described earlier, the composition of the nuclear lamina varies dramatically across the

hematopoietic lineages, and so one might predict a corresponding modulation of cellular tension and therefore a means of directing activity of transcriptional co-factors such as MAL, SRF, and/or YAP (Shin et al., in review).

9. Conclusions

The prototypical hematopoietic stem cell/progenitors has saved many thousands of patients for several decades now. Hematopoietic cells show not only intrinsic differences in mechanical properties but also extrinsic sensitivity to mechanical cues from their niches, both of which could result in a variety of signaling and behavioral responses. Previous studies have revealed the importance of mechanotransduction pathways in stem cells, such as matrix elasticity effects on various transcription co-factors, some of which are being explored with HSC/Ps both *in vitro* and in well-established *in vivo* analyses. Novel therapies envisioned with induced pluripotent stem cells and ESCs generally involve *in vitro* commitment to adult stem cells or progenitors, and so a deep understanding of pathways, such as those reviewed here, in adult stem cell mechanobiology seem broadly important for the many other sub-fields of stem cells. Therefore, adult stem cells and progenitors remain of great interest for their clinical application, and this interest motivates further studies to reveal mechanistic insights into the complex interactions between cells and their physical surroundings.

Acknowledgments

We thank the National Institutes of Health (P01DK032094; R01HL062352; NCATS-8UL1TR000003) and the Human Frontier Science Program (I.I., D.E.D.).

Abbreviations

3D	Three-dimensional
AFM	Atomic force microscopy
Bleb	Blebbistatin
BM	Bone marrow
BMT	Bone marrow transplantation
ECM	Extracellular matrix
ESC	Embryonic stem cell
G-CSF	Granulocyte colony-stimulating factor
HSC/P	Hematopoietic stem cell/progenitor
kPa	kilopascal, units of Elasticity
MAL/MKL	Megakaryocytic acute leukemia protein
MK	Megakaryocyte
MLCK	Myosin light chain kinase
MSC	Mesenchymal stem cell
NMM-II	Non-muscle myosin-II
OB	Osteoblast
RBC	Red blood cell

RLC	regulatory light chains
ROCK	Rho-associated protein kinase
SDF-1	Stromal derived factor-1
SRF	Serum response factor
YAP	Yes-associated protein

References

- Adamo L, Naveiras O, Wenzel PL, McKinney-Freeman S, Mack PJ, Gracia-Sancho J, Suchy-Dicey A, Yoshimoto M, Lensch MW, Yoder MC, Garcia-Cardena G, Daley GQ. Biomechanical forces promote embryonic haematopoiesis. *Nature*. 2009; 459:1131–1135. [PubMed: 19440194]
- Adelstein RS, Conti MA, Johnson GS, Pastan I, Pollard TD. Isolation and characterization of myosin from cloned mouse fibroblasts. *Proc Natl Acad Sci U S A*. 1972; 69:3693–3697. [PubMed: 4264772]
- Akter R, Rivas D, Geneau G, Drissi H, Duque G. Effect of lamin A/C knockdown on osteoblast differentiation and function. *J Bone Miner Res*. 2009; 24:283–293. [PubMed: 18847334]
- Alsberg E, Anderson KW, Albeiruti A, Rowley JA, Mooney DJ. Engineering growing tissues. *Proc Natl Acad Sci U S A*. 2002; 99:12025–12030. [PubMed: 12218178]
- Andrade R, Alonso R, Pena R, Arlucea J, Arechaga J. Localization of importin alpha (Rch1) at the plasma membrane and subcellular redistribution during lymphocyte activation. *Chromosoma*. 2003; 112:87–95. [PubMed: 12883947]
- Andzelm MM, Chen X, Krzewski K, Orange JS, Strominger JL. Myosin IIA is required for cytolytic granule exocytosis in human NK cells. *J Exp Med*. 2007; 204:2285–2291. [PubMed: 17875677]
- Arai F, Hirao A, Ohmura M, Sato H, Matsuoka S, Takubo K, Ito K, Koh GY, Suda T. Tie2/angiopoietin-1 signaling regulates hematopoietic stem cell quiescence in the bone marrow niche. *Cell*. 2004; 118:149–161. [PubMed: 15260986]
- Beach JR, Hussey GS, Miller TE, Chaudhury A, Patel P, Monslow J, Zheng Q, Keri RA, Reizes O, Bresnick AR, Howe PH, Egelhoff TT. Myosin II isoform switching mediates invasiveness after TGF-beta-induced epithelial-mesenchymal transition. *Proc Natl Acad Sci U S A*. 2011; 108:17991–17996. [PubMed: 22025714]
- Beadle C, Assanah MC, Monzo P, Vallee R, Rosenfeld SS, Canoll P. The role of myosin II in glioma invasion of the brain. *Mol Biol Cell*. 2008; 19:3357–3368. [PubMed: 18495866]
- Becker AJ, Mc CE, Till JE. Cytological demonstration of the clonal nature of spleen colonies derived from transplanted mouse marrow cells. *Nature*. 1963; 197:452–454. [PubMed: 13970094]
- Ben-Ze'ev A, Raz A. Multinucleation and inhibition of cytokinesis in suspended cells: reversal upon reattachment to a substrate. *Cell*. 1981; 26:107–115. [PubMed: 6799205]
- Benveniste P, Frelin C, Janmohamed S, Barbara M, Herrington R, Hyam D, Iscove NN. Intermediate-term hematopoietic stem cells with extended but time-limited reconstitution potential. *Cell Stem Cell*. 2010; 6:48–58. [PubMed: 20074534]
- Berry MF, Engler AJ, Woo YJ, Pirolli TJ, Bish LT, Jayasankar V, Morine KJ, Gardner TJ, Discher DE, Sweeney HL. Mesenchymal stem cell injection after myocardial infarction improves myocardial compliance. *Am J Physiol Heart Circ Physiol*. 2006; 290:H2196–H2203. [PubMed: 16473959]
- Breitbach M, Bostani T, Roell W, Xia Y, Dewald O, Nygren JM, Fries JW, Tiemann K, Bohlen H, Hescheler J, Welz A, Bloch W, Jacobsen SE, Fleischmann BK. Potential risks of bone marrow cell transplantation into infarcted hearts. *Blood*. 2007; 110:1362–1369. [PubMed: 17483296]
- Brown AE, Hategan A, Safer D, Goldman YE, Discher DE. Cross-correlated TIRF/AFM reveals asymmetric distribution of force-generating heads along self-assembled, "synthetic" myosin filaments. *Biophys J*. 2009; 96:1952–1960. [PubMed: 19254555]

- Calvi LM, Adams GB, Weibrecht KW, Weber JM, Olson DP, Knight MC, Martin RP, Schipani E, Divieti P, Bringham FR, Milner LA, Kronenberg HM, Scadden DT. Osteoblastic cells regulate the haematopoietic stem cell niche. *Nature*. 2003; 425:841–846. [PubMed: 14574413]
- Chang JT, Palanivel VR, Kinjo I, Schambach F, Intlekofer AM, Banerjee A, Longworth SA, Vinup KE, Mrass P, Oliaro J, Killeen N, Orange JS, Russell SM, Weninger W, Reiner SL. Asymmetric T lymphocyte division in the initiation of adaptive immune responses. *Science*. 2007; 315:1687–1691. [PubMed: 17332376]
- Chasis JA, Mohandas N. Erythroblastic islands: niches for erythropoiesis. *Blood*. 2008; 112:470–478. [PubMed: 18650462]
- Chen Y, Jacamo R, Shi YX, Wang RY, Battula VL, Konoplev S, Strunk D, Hofmann NA, Reinisch A, Konopleva M, Andreeff M. Human extramedullary bone marrow in mice: a novel in vivo model of genetically controlled hematopoietic microenvironment. *Blood*. 2012; 119:4971–4980. [PubMed: 22490334]
- Chen Z, Naveiras O, Balduini A, Mammoto A, Conti MA, Adelstein RS, Ingber D, Daley GQ, Shivdasani RA. The May-Hegglin anomaly gene MYH9 is a negative regulator of platelet biogenesis modulated by the Rho-ROCK pathway. *Blood*. 2007; 110:171–179. [PubMed: 17392504]
- Cheng EC, Luo Q, Bruscia EM, Renda MJ, Troy JA, Massaro SA, Tuck D, Schulz V, Mane SM, Berliner N, Sun Y, Morris SW, Qiu C, Krause DS. Role for MKL1 in megakaryocytic maturation. *Blood*. 2009; 113:2826–2834. [PubMed: 19136660]
- Choi JS, Harley BA. The combined influence of substrate elasticity and ligand density on the viability and biophysical properties of hematopoietic stem and progenitor cells. *Biomaterials*. 2012; 33:4460–4468. [PubMed: 22444641]
- Chowdhury F, Na S, Li D, Poh YC, Tanaka TS, Wang F, Wang N. Material properties of the cell dictate stress-induced spreading and differentiation in embryonic stem cells. *Nat Mater*. 2010; 9:82–88. [PubMed: 19838182]
- Connelly JT, Gautrot JE, Trappmann B, Tan DW, Donati G, Huck WT, Watt FM. Actin and serum response factor transduce physical cues from the microenvironment to regulate epidermal stem cell fate decisions. *Nat Cell Biol*. 2010; 12:711–718. [PubMed: 20581838]
- Conti MA, Even-Ram S, Liu C, Yamada KM, Adelstein RS. Defects in cell adhesion and the visceral endoderm following ablation of nonmuscle myosin heavy chain II-A in mice. *J Biol Chem*. 2004; 279:41263–41266. [PubMed: 15292239]
- Daley GQ, Scadden DT. Prospects for stem cell-based therapy. *Cell*. 2008; 132:544–548. [PubMed: 18295571]
- Dar A, Goichberg P, Shinder V, Kalinkovich A, Kollet O, Netzer N, Margalit R, Zsak M, Nagler A, Hardan I, Resnick I, Rot A, Lapidot T. Chemokine receptor CXCR4-dependent internalization and resecretion of functional chemokine SDF-1 by bone marrow endothelial and stromal cells. *Nat Immunol*. 2005; 6:1038–1046. [PubMed: 16170318]
- Despeaux M, Chicanne G, Rouer E, De Toni-Costes F, Bertrand J, Mansat-De Mas V, Vergnolle N, Eaves C, Payrastre B, Girault JA, Racaud-Sultan C. Focal adhesion kinase splice variants maintain primitive acute myeloid leukemia cells through altered Wnt signaling. *Stem Cells*. 2012; 30:1597–1610. [PubMed: 22714993]
- DiPersio JF, Uy GL, Yasothan U, Kirkpatrick P. Plerixafor. *Nat Rev Drug Discov*. 2009; 8:105–106. [PubMed: 19180104]
- Discher DE, Janmey P, Wang YL. Tissue cells feel and respond to the stiffness of their substrate. *Science*. 2005; 310:1139–1143. [PubMed: 16293750]
- Discher DE, Mohandas N, Evans EA. Molecular maps of red cell deformation: hidden elasticity and in situ connectivity. *Science*. 1994; 266:1032–1035. [PubMed: 7973655]
- Dong C, Skalak R, Sung KL. Cytoplasmic rheology of passive neutrophils. *Biorheology*. 1991; 28:557–567. [PubMed: 1818744]
- Dulyaninova NG, Malashkevich VN, Almo SC, Bresnick AR. Regulation of myosin-IIA assembly and Mts1 binding by heavy chain phosphorylation. *Biochemistry*. 2005; 44:6867–6876. [PubMed: 15865432]

- Dupont S, Morsut L, Aragona M, Enzo E, Giulitti S, Cordenonsi M, Zanconato F, Le Digabel J, Forcato M, Bicciato S, Elvassore N, Piccolo S. Role of YAP/TAZ in mechanotransduction. *Nature*. 2011; 474:179–183. [PubMed: 21654799]
- Eckly A, Strassel C, Freund M, Cazenave JP, Lanza F, Gachet C, Leon C. Abnormal megakaryocyte morphology and proplatelet formation in mice with megakaryocyte-restricted MYH9 inactivation. *Blood*. 2009; 113:3182–3189. [PubMed: 18984861]
- Engler AJ, Sen S, Sweeney HL, Discher DE. Matrix elasticity directs stem cell lineage specification. *Cell*. 2006; 126:677–689. [PubMed: 16923388]
- Even-Faitelson L, Ravid S. PAK1 and aPKC ζ regulate myosin II-B phosphorylation: a novel signaling pathway regulating filament assembly. *Mol Biol Cell*. 2006; 17:2869–2881. [PubMed: 16611744]
- Faure-Andre G, Vargas P, Yuseff MI, Heuze M, Diaz J, Lankar D, Steri V, Manry J, Hugues S, Vascotto F, Boulanger J, Raposo G, Bono MR, Roseblatt M, Piel M, Lennon-Dumenil AM. Regulation of dendritic cell migration by CD74, the MHC class II-associated invariant chain. *Science*. 2008; 322:1705–1710. [PubMed: 19074353]
- Fernandez MI, Heuze ML, Martinez-Cingolani C, Volpe E, Donnadieu MH, Piel M, Homey B, Lennon-Dumenil AM, Soumelis V. The human cytokine TSLP triggers a cell-autonomous dendritic cell migration in confined environments. *Blood*. 2011; 118:3862–3869. [PubMed: 21772055]
- Friedenstein AJ, Gorskaja JF, Kulagina NN. Fibroblast precursors in normal and irradiated mouse hematopoietic organs. *Exp Hematol*. 1976; 4:267–274. [PubMed: 976387]
- Friedl P, Wolf K, Lammerding J. Nuclear mechanics during cell migration. *Curr Opin Cell Biol*. 2011; 23:55–64. [PubMed: 21109415]
- Fu J, Wang YK, Yang MT, Desai RA, Yu X, Liu Z, Chen CS. Mechanical regulation of cell function with geometrically modulated elastomeric substrates. *Nat Methods*. 2010; 7:733–736. [PubMed: 20676108]
- Geiger B, Spatz JP, Bershadsky AD. Environmental sensing through focal adhesions. *Nat Rev Mol Cell Biol*. 2009; 10:21–33. [PubMed: 19197329]
- Giebel B, Zhang T, Beckmann J, Spanholtz J, Wernet P, Ho AD, Punzel M. Primitive human hematopoietic cells give rise to differentially specified daughter cells upon their initial cell division. *Blood*. 2006; 107:2146–2152. [PubMed: 16249381]
- Gilbert PM, Havenstrite KL, Magnusson KE, Sacco A, Leonardi NA, Kraft P, Nguyen NK, Thrun S, Lutolf MP, Blau HM. Substrate elasticity regulates skeletal muscle stem cell self-renewal in culture. *Science*. 2010; 329:1078–1081. [PubMed: 20647425]
- Gilles L, Bluteau D, Boukour S, Chang Y, Zhang Y, Robert T, Dessen P, Debili N, Bernard OA, Vainchenker W, Raslova H. MAL/SRF complex is involved in platelet formation and megakaryocyte migration by regulating MYL9 (MLC2) and MMP9. *Blood*. 2009; 114:4221–4232. [PubMed: 19724058]
- Giordano GF, Lichtman MA. Marrow cell egress. The central interaction of barrier pore size and cell maturation. *J Clin Invest*. 1973; 52:1154–1164. [PubMed: 4512610]
- Glodek AM, Honczarenko M, Le Y, Campbell JJ, Silberstein LE. Sustained activation of cell adhesion is a differentially regulated process in B lymphopoiesis. *J Exp Med*. 2003; 197:461–473. [PubMed: 12591904]
- Gratwohl A, Baldomero H, Aljurf M, Pasquini MC, Bouzas LF, Yoshimi A, Szer J, Lipton J, Schwendener A, Gratwohl M, Frauendorfer K, Niederwieser D, Horowitz M, Koda Y, Marrow T. Worldwide Network of B. Hematopoietic stem cell transplantation: a global perspective. *JAMA*. 2010; 303:1617–1624. [PubMed: 20424252]
- Gu YC, Kortessma J, Tryggvason K, Persson J, Ekblom P, Jacobsen SE, Ekblom M. Laminin isoform-specific promotion of adhesion and migration of human bone marrow progenitor cells. *Blood*. 2003; 101:877–885. [PubMed: 12393739]
- Hare JM, Fishman JE, Gerstenblith G, DiFede Velazquez DL, Zambrano JP, Suncion VY, Tracy M, Ghersin E, Johnston PV, Brinker JA, Breton E, Davis-Sproul J, Schulman IH, Byrnes J, Mendizabal AM, Lowery MH, Rouy D, Altman P, Wong Po Foo C, Ruiz P, Amador A, Da Silva J, McNiece IK, Heldman AW. Comparison of allogeneic vs autologous bone marrow-derived

mesenchymal stem cells delivered by transendocardial injection in patients with ischemic cardiomyopathy: the POSEIDON randomized trial. *JAMA*. 2012; 308:2369–2379. [PubMed: 23117550]

- Hernandez OM, Jones M, Guzman G, Szczesna-Cordary D. Myosin essential light chain in health and disease. *Am J Physiol Heart Circ Physiol*. 2007; 292:H1643–H1654. [PubMed: 17142342]
- Hitchcock IS, Fox NE, Prevost N, Sear K, Shattil SJ, Kaushansky K. Roles of focal adhesion kinase (FAK) in megakaryopoiesis and platelet function: studies using a megakaryocyte lineage specific FAK knockout. *Blood*. 2008; 111:596–604. [PubMed: 17925492]
- Holst J, Watson S, Lord MS, Eamegdool SS, Bax DV, Nivison-Smith LB, Kondyurin A, Ma L, Oberhauser AF, Weiss AS, Rasko JE. Substrate elasticity provides mechanical signals for the expansion of hemopoietic stem and progenitor cells. *Nat Biotechnol*. 2010; 28:1123–1128. [PubMed: 20890282]
- Hooper AT, Butler JM, Nolan DJ, Kranz A, Iida K, Kobayashi M, Kopp HG, Shido K, Petit I, Yanger K, James D, Witte L, Zhu Z, Wu Y, Pytowski B, Rosenwaks Z, Mittal V, Sato TN, Rafii S. Engraftment and reconstitution of hematopoiesis is dependent on VEGFR2-mediated regeneration of sinusoidal endothelial cells. *Cell Stem Cell*. 2009; 4:263–274. [PubMed: 19265665]
- Ilani T, Vasiliver-Shamis G, Vardhana S, Bretscher A, Dustin ML. T cell antigen receptor signaling and immunological synapse stability require myosin IIA. *Nat Immunol*. 2009; 10:531–539. [PubMed: 19349987]
- Ivanovska I, Swift J, Harada T, Pajeroski JD, Discher DE. Physical plasticity of the nucleus and its manipulation. *Methods Cell Biol*. 2010; 98:207–220. [PubMed: 20816236]
- Iyer KV, Maharana S, Gupta S, Libchaber A, Tlusty T, Shivashankar GV. Modeling and experimental methods to probe the link between global transcription and spatial organization of chromosomes. *PLoS One*. 2012a; 7:e46628. [PubMed: 23049710]
- Iyer KV, Pulford S, Mogilner A, Shivashankar GV. Mechanical activation of cells induces chromatin remodeling preceding MKL nuclear transport. *Biophys J*. 2012b; 103:1416–1428. [PubMed: 23062334]
- Jacobelli J, Friedman RS, Conti MA, Lennon-Dumenil AM, Piel M, Sorensen CM, Adelstein RS, Krummel MF. Confinement-optimized three-dimensional T cell amoeboid motility is modulated via myosin IIA-regulated adhesions. *Nat Immunol*. 2010; 11:953–961. [PubMed: 20835229]
- Jansson L, Larsson J. Normal hematopoietic stem cell function in mice with enforced expression of the Hippo signaling effector YAP1. *PLoS One*. 2012; 7:e32013. [PubMed: 22363786]
- Junt T, Schulze H, Chen Z, Massberg S, Goerge T, Krueger A, Wagner DD, Graf T, Italiano JE Jr, Shivdasani RA, von Andrian UH. Dynamic visualization of thrombopoiesis within bone marrow. *Science*. 2007; 317:1767–1770. [PubMed: 17885137]
- Kaushansky K, Lok S, Holly RD, Broudy VC, Lin N, Bailey MC, Forstrom JW, Buddle MM, Oort PJ, Hagen FS, et al. Promotion of megakaryocyte progenitor expansion and differentiation by the c-Mpl ligand thrombopoietin. *Nature*. 1994; 369:568–571. [PubMed: 8202159]
- Kelley MJ, Jawien W, Ortel TL, Korczak JF. Mutation of MYH9, encoding non-muscle myosin heavy chain A, in May-Hegglin anomaly. *Nat Genet*. 2000; 26:106–108. [PubMed: 10973260]
- Kiel MJ, Yilmaz OH, Iwashita T, Terhorst C, Morrison SJ. SLAM family receptors distinguish hematopoietic stem and progenitor cells and reveal endothelial niches for stem cells. *Cell*. 2005; 121:1109–1121. [PubMed: 15989959]
- Kim KY, Kovacs M, Kawamoto S, Sellers JR, Adelstein RS. Disease-associated mutations and alternative splicing alter the enzymatic and motile activity of nonmuscle myosins II-B and II-C. *J Biol Chem*. 2005; 280:22769–22775. [PubMed: 15845534]
- Klein EA, Yin L, Kothapalli D, Castagnino P, Byfield FJ, Xu T, Levental I, Hawthorne E, Janmey PA, Assoian RK. Cell-cycle control by physiological matrix elasticity and in vivo tissue stiffening. *Curr Biol*. 2009; 19:1511–1518. [PubMed: 19765988]
- Knospe, WH. Hematopoiesis on Artificial Membranes. In: Tavassoli, M., editor. *Handbook of the Hemopoietic Microenvironment*. Clifton, NJ: The Humana Press Inc.; 1989.
- Kobayashi H, Butler JM, O'Donnell R, Kobayashi M, Ding BS, Bonner B, Chiu VK, Nolan DJ, Shido K, Benjamin L, Rafii S. Angiocrine factors from Akt-activated endothelial cells balance self-

- renewal and differentiation of haematopoietic stem cells. *Nat Cell Biol.* 2010; 12:1046–1056. [PubMed: 20972423]
- Koury ST, Koury MJ, Bondurant MC. Cytoskeletal distribution and function during the maturation and enucleation of mammalian erythroblasts. *J Cell Biol.* 1989; 109:3005–3013. [PubMed: 2574178]
- Lee J, Li M, Milwid J, Dunham J, Vinegoni C, Gorbatov R, Iwamoto Y, Wang F, Shen K, Hatfield K, Enger M, Shafiee S, McCormack E, Ebert BL, Weissleder R, Yarmush ML, Parekkadan B. Implantable microenvironments to attract hematopoietic stem/cancer cells. *Proc Natl Acad Sci U S A.* 2012; 109:19638–19643. [PubMed: 23150542]
- Lee-Thedieck C, Rauch N, Fiammengo R, Klein G, Spatz JP. Impact of substrate elasticity on human hematopoietic stem and progenitor cell adhesion and motility. *J Cell Sci.* 2012; 125:3765–3775. [PubMed: 22553208]
- Lichtman MA. Cellular deformability during maturation of the myeloblast. Possible role in marrow egress. *N Engl J Med.* 1970; 283:943–948. [PubMed: 4919083]
- Lichtman, MA.; Packman, CH.; Constine, LS. Molecular and Cellular Traffic Across the Marrow Sinuses. In: Tavassoli, M., editor. *Handbook of the Hemopoietic Microenvironment*. Clifton, NJ: The Humana Press Inc.; 1989.
- Liu J, Tan Y, Zhang H, Zhang Y, Xu P, Chen J, Poh YC, Tang K, Wang N, Huang B. Soft fibrin gels promote selection and growth of tumorigenic cells. *Nat Mater.* 2012; 11:734–741. [PubMed: 22751180]
- Lordier L, Bluteau D, Jalil A, Legrand C, Pan J, Rameau P, Jouni D, Bluteau O, Mercher T, Leon C, Gachet C, Debili N, Vainchenker W, Raslova H, Chang Y. RUNX1-induced silencing of non-muscle myosin heavy chain IIB contributes to megakaryocyte polyploidization. *Nat Commun.* 2012; 3:717. [PubMed: 22395608]
- Lu J, Sun Y, Nombela-Arrieta C, Du KP, Park SY, Chai L, Walkley C, Luo HR, Silberstein LE. Fak depletion in both hematopoietic and nonhematopoietic niche cells leads to hematopoietic stem cell expansion. *Exp Hematol.* 2012; 40:307–317. e303. [PubMed: 22155722]
- Ma X, Jana SS, Conti MA, Kawamoto S, Claycomb WC, Adelstein RS. Ablation of nonmuscle myosin II-B and II-C reveals a role for nonmuscle myosin II in cardiac myocyte karyokinesis. *Mol Biol Cell.* 2010; 21:3952–3962. [PubMed: 20861308]
- Matsumura F. Regulation of myosin II during cytokinesis in higher eukaryotes. *Trends Cell Biol.* 2005; 15:371–377. [PubMed: 15935670]
- Mazo IB, Massberg S, von Andrian UH. Hematopoietic stem and progenitor cell trafficking. *Trends Immunol.* 2011; 32:493–503. [PubMed: 21802990]
- McBeath R, Pirone DM, Nelson CM, Bhadriraju K, Chen CS. Cell shape, cytoskeletal tension, and RhoA regulate stem cell lineage commitment. *Dev Cell.* 2004; 6:483–495. [PubMed: 15068789]
- Mendez-Ferrer S, Michurina TV, Ferraro F, Mazloom AR, Macarthur BD, Lira SA, Scadden DT, Ma'ayan A, Enikolopov GN, Frenette PS. Mesenchymal and haematopoietic stem cells form a unique bone marrow niche. *Nature.* 2010; 466:829–834. [PubMed: 20703299]
- Merkel R, Simson R, Simson DA, Hohenadl M, Boulbitch A, Wallraff E, Sackmann E. A micromechanic study of cell polarity and plasma membrane cell body coupling in *Dictyostelium*. *Biophys J.* 2000; 79:707–719. [PubMed: 10920005]
- Miyake T, Kung CK, Goldwasser E. Purification of human erythropoietin. *J Biol Chem.* 1977; 252:5558–5564. [PubMed: 18467]
- Morita Y, Iseki A, Okamura S, Suzuki S, Nakauchi H, Ema H. Functional characterization of hematopoietic stem cells in the spleen. *Exp Hematol.* 2011; 39:351–359. e353. [PubMed: 21185906]
- Naveiras O, Nardi V, Wenzel PL, Hauschka PV, Fahey F, Daley GQ. Bone-marrow adipocytes as negative regulators of the haematopoietic microenvironment. *Nature.* 2009; 460:259–263. [PubMed: 19516257]
- Nilsson SK, Johnston HM, Whitty GA, Williams B, Webb RJ, Denhardt DT, Bertoncello I, Bendall LJ, Simmons PJ, Haylock DN. Osteopontin, a key component of the hematopoietic stem cell niche and regulator of primitive hematopoietic progenitor cells. *Blood.* 2005; 106:1232–1239. [PubMed: 15845900]

- North TE, Goessling W, Peeters M, Li P, Ceol C, Lord AM, Weber GJ, Harris J, Cutting CC, Huang P, Dzierzak E, Zon LI. Hematopoietic stem cell development is dependent on blood flow. *Cell*. 2009; 137:736–748. [PubMed: 19450519]
- Notta F, Doulatov S, Laurenti E, Poepl A, Jurisica I, Dick JE. Isolation of single human hematopoietic stem cells capable of long-term multilineage engraftment. *Science*. 2011; 333:218–221. [PubMed: 21737740]
- Olson EN, Nordheim A. Linking actin dynamics and gene transcription to drive cellular motile functions. *Nat Rev Mol Cell Biol*. 2010; 11:353–365. [PubMed: 20414257]
- Pajeroski JD, Dahl KN, Zhong FL, Sammak PJ, Discher DE. Physical plasticity of the nucleus in stem cell differentiation. *Proc Natl Acad Sci U S A*. 2007; 104:15619–15624. [PubMed: 17893336]
- Papayannopoulou T, Priestley GV, Nakamoto B. Anti-VLA4/VCAM-1-induced mobilization requires cooperative signaling through the kit/mkit ligand pathway. *Blood*. 1998; 91:2231–2239. [PubMed: 9516120]
- Pollard TD, Korn ED. Acanthamoeba myosin. II. Interaction with actin and with a new cofactor protein required for actin activation of Mg 2+ adenosine triphosphatase activity. *J Biol Chem*. 1973; 248:4691–4697. [PubMed: 4268864]
- Raab M, Swift J, Dingal PC, Shah P, Shin JW, Discher DE. Crawling from soft to stiff matrix polarizes the cytoskeleton and phosphoregulates myosin-II heavy chain. *J Cell Biol*. 2012; 199:669–683. [PubMed: 23128239]
- Rafii S, Shapiro F, Pettengell R, Ferris B, Nachman RL, Moore MA, Asch AS. Human bone marrow microvascular endothelial cells support long-term proliferation and differentiation of myeloid and megakaryocytic progenitors. *Blood*. 1995; 86:3353–3363. [PubMed: 7579438]
- Ragu C, Elain G, Mylonas E, Ottolenghi C, Cagnard N, Daegelen D, Passegue E, Vainchenker W, Bernard OA, Penard-Lacronique V. The transcription factor Srf regulates hematopoietic stem cell adhesion. *Blood*. 2010; 116:4464–4473. [PubMed: 20709909]
- Reichl EM, Ren Y, Morpew MK, Delannoy M, Effler JC, Girard KD, Divi S, Iglesias PA, Kuo SC, Robinson DN. Interactions between myosin and actin crosslinkers control cytokinesis contractility dynamics and mechanics. *Curr Biol*. 2008; 18:471–480. [PubMed: 18372178]
- Rey M, Valenzuela-Fernandez A, Urzainqui A, Yanez-Mo M, Perez-Martinez M, Penela P, Mayor F Jr, Sanchez-Madrid F. Myosin IIA is involved in the endocytosis of CXCR4 induced by SDF-1alpha. *J Cell Sci*. 2007; 120:1126–1133. [PubMed: 17327270]
- Sacchetti B, Funari A, Michienzi S, Di Cesare S, Piersanti S, Saggio I, Tagliafico E, Ferrari S, Robey PG, Riminucci M, Bianco P. Self-renewing osteoprogenitors in bone marrow sinusoids can organize a hematopoietic microenvironment. *Cell*. 2007; 131:324–336. [PubMed: 17956733]
- Saha K, Keung AJ, Irwin EF, Li Y, Little L, Schaffer DV, Healy KE. Substrate modulus directs neural stem cell behavior. *Biophys J*. 2008; 95:4426–4438. [PubMed: 18658232]
- Samaniego R, Sanchez-Martin L, Estechea A, Sanchez-Mateos P. Rho/ROCK and myosin II control the polarized distribution of endocytic clathrin structures at the uropod of moving T lymphocytes. *J Cell Sci*. 2007; 120:3534–3543. [PubMed: 17895369]
- Schofield R. The relationship between the spleen colony-forming cell and the haemopoietic stem cell. *Blood Cells*. 1978; 4:7–25. [PubMed: 747780]
- Scholey JM, Brust-Mascher I, Mogilner A. Cell division. *Nature*. 2003; 422:746–752. [PubMed: 12700768]
- Scotti C, Piccinini E, Takizawa H, Todorov A, Bourguine P, Papadimitropoulos A, Barbero A, Manz MG, Martin I. Engineering of a functional bone organ through endochondral ossification. *Proc Natl Acad Sci U S A*. 2013; 110:3997–4002. [PubMed: 23401508]
- Seki M. Hematopoietic colony formation in a macrophage layer provided by intraperitoneal insertion of cellulose acetate membrane. *Transplantation*. 1973; 16:544–549. [PubMed: 4356866]
- Seri M, Cusano R, Gangarossa S, Caridi G, Bordo D, Lo Nigro C, Ghiggeri GM, Ravazzolo R, Savino M, Del Vecchio M, d'Apolito M, Iolascon A, Zelante LL, Savoia A, Balduini CL, Noris P, Magrini U, Belletti S, Heath KE, Babcock M, Glucksman MJ, Aliprandis E, Bizzaro N, Desnick RJ, Martignetti JA. Mutations in MYH9 result in the May-Hegglin anomaly, and Fechtner and

- Sebastian syndromes. The May-Hegglin/Fechtner Syndrome Consortium. *Nat Genet.* 2000; 26:103–105. [PubMed: 10973259]
- Shimi T, Butin-Israeli V, Adam SA, Goldman RD. Nuclear lamins in cell regulation and disease. *Cold Spring Harb Symp Quant Biol.* 2010; 75:525–531. [PubMed: 21467145]
- Shin JW, Swift J, Spinler KR, Discher DE. Myosin-II inhibition and soft 2D matrix maximize multinucleation and cellular projections typical of platelet-producing megakaryocytes. *Proc Natl Acad Sci U S A.* 2011; 108:11458–11463. [PubMed: 21709232]
- Shin ME, He Y, Li D, Na S, Chowdhury F, Poh YC, Collin O, Su P, de Lanerolle P, Schwartz MA, Wang N, Wang F. Spatiotemporal organization, regulation, and functions of tractions during neutrophil chemotaxis. *Blood.* 2010; 116:3297–3310. [PubMed: 20616216]
- Smith CM 2nd, Burris SM, White JG. Micropipette aspiration of guinea pig megakaryocytes: absence of fragmentation and dependence on maturation stage. *Blood.* 1989; 73:1570–1575. [PubMed: 2713495]
- Song J, Kiel MJ, Wang Z, Wang J, Taichman RS, Morrison SJ, Krebsbach PH. An in vivo model to study and manipulate the hematopoietic stem cell niche. *Blood.* 2010; 115:2592–2600. [PubMed: 20110425]
- Soofi SS, Last JA, Liliensiek SJ, Nealey PF, Murphy CJ. The elastic modulus of Matrigel as determined by atomic force microscopy. *J Struct Biol.* 2009; 167:216–219. [PubMed: 19481153]
- Stiehler M, Bunger C, Baatrup A, Lind M, Kassem M, Mygind T. Effect of dynamic 3-D culture on proliferation, distribution, and osteogenic differentiation of human mesenchymal stem cells. *J Biomed Mater Res A.* 2009; 89:96–107. [PubMed: 18431785]
- Stier S, Ko Y, Forkert R, Lutz C, Neuhaus T, Grunewald E, Cheng T, Dombkowski D, Calvi LM, Rittling SR, Scadden DT. Osteopontin is a hematopoietic stem cell niche component that negatively regulates stem cell pool size. *J Exp Med.* 2005; 201:1781–1791. [PubMed: 15928197]
- Straight AF, Cheung A, Limouze J, Chen I, Westwood NJ, Sellers JR, Mitchison TJ. Dissecting temporal and spatial control of cytokinesis with a myosin II inhibitor. *Science.* 2003; 299:1743–1747. [PubMed: 12637748]
- Sugiyama T, Kohara H, Noda M, Nagasawa T. Maintenance of the hematopoietic stem cell pool by CXCL12-CXCR4 chemokine signaling in bone marrow stromal cell niches. *Immunity.* 2006; 25:977–988. [PubMed: 17174120]
- Szilvassy SJ, Humphries RK, Lansdorp PM, Eaves AC, Eaves CJ. Quantitative assay for totipotent reconstituting hematopoietic stem cells by a competitive repopulation strategy. *Proc Natl Acad Sci U S A.* 1990; 87:8736–8740. [PubMed: 2247442]
- Taichman RS, Emerson SG. Human osteoblasts support hematopoiesis through the production of granulocyte colony-stimulating factor. *J Exp Med.* 1994; 179:1677–1682. [PubMed: 7513014]
- Takeda K, Kishi H, Ma X, Yu ZX, Adelstein RS. Ablation and mutation of nonmuscle myosin heavy chain II-B results in a defect in cardiac myocyte cytokinesis. *Circ Res.* 2003; 93:330–337. [PubMed: 12893741]
- Taniguchi H, Toyoshima T, Fukao K, Nakauchi H. Presence of hematopoietic stem cells in the adult liver. *Nat Med.* 1996; 2:198–203. [PubMed: 8574965]
- Thomas ED, Lochte HL Jr, Lu WC, Ferrebee JW. Intravenous infusion of bone marrow in patients receiving radiation and chemotherapy. *N Engl J Med.* 1957; 257:491–496. [PubMed: 13464965]
- Ting SB, Deneault E, Hope K, Cellot S, Chagraoui J, Mayotte N, Dorn JF, Laverdure JP, Harvey M, Hawkins ED, Russell SM, Maddox PS, Iscove NN, Sauvageau G. Asymmetric segregation and self-renewal of hematopoietic stem and progenitor cells with endocytic Ap2a2. *Blood.* 2012; 119:2510–2522. [PubMed: 22174158]
- Titushkin I, Cho M. Modulation of cellular mechanics during osteogenic differentiation of human mesenchymal stem cells. *Biophys J.* 2007; 93:3693–3702. [PubMed: 17675345]
- Trumpp A, Essers M, Wilson A. Awakening dormant haematopoietic stem cells. *Nat Rev Immunol.* 2010; 10:201–209. [PubMed: 20182459]
- Tsai RK, Discher DE. Inhibition of "self" engulfment through deactivation of myosin-II at the phagocytic synapse between human cells. *J Cell Biol.* 2008; 180:989–1003. [PubMed: 18332220]

- Tullio AN, Accili D, Ferrans VJ, Yu ZX, Takeda K, Grinberg A, Westphal H, Preston YA, Adelstein RS. Nonmuscle myosin II-B is required for normal development of the mouse heart. *Proc Natl Acad Sci U S A*. 1997; 94:12407–12412. [PubMed: 9356462]
- Ubukawa K, Guo YM, Takahashi M, Hirokawa M, Michishita Y, Nara M, Tagawa H, Takahashi N, Komatsuda A, Nunomura W, Takakuwa Y, Sawada K. Enucleation of human erythroblasts involves nonmuscle myosin IIB. *Blood*. 2011
- Vascotto F, Lankar D, Faure-Andre G, Vargas P, Diaz J, Le Roux D, Yuseff MI, Sibarita JB, Boes M, Raposo G, Mougneau E, Glaichenhaus N, Bonnerot C, Manoury B, Lennon-Dumenil AM. The actin-based motor protein myosin II regulates MHC class II trafficking and BCR-driven antigen presentation. *J Cell Biol*. 2007; 176:1007–1019. [PubMed: 17389233]
- Visnjic D, Kalajzic I, Gronowicz G, Aguila HL, Clark SH, Lichtler AC, Rowe DW. Conditional ablation of the osteoblast lineage in Col2.3deltatg transgenic mice. *J Bone Miner Res*. 2001; 16:2222–2231. [PubMed: 11760835]
- Wang F, Kovacs M, Hu A, Limouze J, Harvey EV, Sellers JR. Kinetic mechanism of non-muscle myosin IIB: functional adaptations for tension generation and maintenance. *J Biol Chem*. 2003; 278:27439–27448. [PubMed: 12704189]
- Wang HB, Dembo M, Hanks SK, Wang Y. Focal adhesion kinase is involved in mechanosensing during fibroblast migration. *Proc Natl Acad Sci U S A*. 2001; 98:11295–11300. [PubMed: 11572981]
- Wang N, Tytell JD, Ingber DE. Mechanotransduction at a distance: mechanically coupling the extracellular matrix with the nucleus. *Nat Rev Mol Cell Biol*. 2009; 10:75–82. [PubMed: 19197334]
- Waugh RE, Mantalaris A, Bauserman RG, Hwang WC, Wu JH. Membrane instability in late-stage erythropoiesis. *Blood*. 2001; 97:1869–1875. [PubMed: 11238131]
- Weinbaum S, Cowin SC, Zeng Y. A model for the excitation of osteocytes by mechanical loading-induced bone fluid shear stresses. *J Biomech*. 1994; 27:339–360. [PubMed: 8051194]
- Weissman IL, Shizuru JA. The origins of the identification and isolation of hematopoietic stem cells, and their capability to induce donor-specific transplantation tolerance and treat autoimmune diseases. *Blood*. 2008; 112:3543–3553. [PubMed: 18948588]
- White JG, Burris SM, Tukey D, Smith C 2nd, Clawson CC. Micropipette aspiration of human platelets: influence of microtubules and actin filaments on deformability. *Blood*. 1984; 64:210–214. [PubMed: 6145458]
- Wilson CA, Tsuchida MA, Allen GM, Barnhart EL, Applegate KT, Yam PT, Ji L, Keren K, Danuser G, Theriot JA. Myosin II contributes to cell-scale actin network treadmilling through network disassembly. *Nature*. 2010; 465:373–377. [PubMed: 20485438]
- Winer JP, Janmey PA, McCormick ME, Funaki M. Bone marrow-derived human mesenchymal stem cells become quiescent on soft substrates but remain responsive to chemical or mechanical stimuli. *Tissue Eng Part A*. 2009; 15:147–154. [PubMed: 18673086]
- Winkler IG, Sims NA, Pettit AR, Barbier V, Nowlan B, Helwani F, Poulton IJ, van Rooijen N, Alexander KA, Raggatt LJ, Levesque JP. Bone marrow macrophages maintain hematopoietic stem cell (HSC) niches and their depletion mobilizes HSCs. *Blood*. 2010; 116:4815–4828. [PubMed: 20713966]
- Zeng L, Xiao Q, Margariti A, Zhang Z, Zampetaki A, Patel S, Capogrossi MC, Hu Y, Xu Q. HDAC3 is crucial in shear- and VEGF-induced stem cell differentiation toward endothelial cells. *J Cell Biol*. 2006; 174:1059–1069. [PubMed: 16982804]
- Zhang J, Niu C, Ye L, Huang H, He X, Tong WG, Ross J, Haug J, Johnson T, Feng JQ, Harris S, Wiedemann LM, Mishina Y, Li L. Identification of the haematopoietic stem cell niche and control of the niche size. *Nature*. 2003; 425:836–841. [PubMed: 14574412]
- Zhang Y, Conti MA, Malide D, Dong F, Wang A, Shmist YA, Liu C, Zerfas P, Daniels MP, Chan CC, Kozin E, Kachar B, Kelley MJ, Kopp JB, Adelstein RS. Mouse models of MYH9-related disease: mutations in nonmuscle myosin II-A. *Blood*. 2011
- Zhao B, Tumaneng K, Guan KL. The Hippo pathway in organ size control, tissue regeneration and stem cell self-renewal. *Nat Cell Biol*. 2011; 13:877–883. [PubMed: 21808241]

Zhu J, Garrett R, Jung Y, Zhang Y, Kim N, Wang J, Joe GJ, Hexner E, Choi Y, Taichman RS, Emerson SG. Osteoblasts support B-lymphocyte commitment and differentiation from hematopoietic stem cells. *Blood*. 2007; 109:3706–3712. [PubMed: 17227831]

Highlights

1. Mechanobiology is part of adult stem cell specification programs.
2. Myosin-II mediates diverse mechanobiological functions in marrow and blood.
3. Mechanical properties of the nucleus regulate cellular trafficking.

Figure 1. Myosin, lamin and mechanical factors influence differentiation and trafficking from bone marrow

Bone marrow microenvironments may provide different modes of biophysical regulation, which are principally regulated by actin-myosin at the cell membrane that generates force when cells sense matrix elasticity, followed by force transmission to the nucleus and lamins (Inset). Mesenchymal stem cells (MSCs) in marrow sense matrix elasticity and differentiate into stromal lineages, including osteoblasts and adipocytes. Hematopoietic stem cells (HSCs) are kept in quiescence (G0) near osteoblasts at the endosteum, which is stiffer than the central marrow space. HSC numbers are also suppressed by adipocytes. HSCs traffic towards the vascular niche and undergo either self-renewing expansion (symmetric division) or differentiation (asymmetric division) into progenitors (HSC/Ps) and mature cells. Cells at the vascular niche might also be exposed to shear force, which could affect differentiation. Cells also undergo selective trafficking out of the marrow due to physical properties of both cell membrane cortex and nucleus as regulated, respectively, by myosin and lamin. MSCs are additionally present as pericytes which could also regulate hematopoietic cell trafficking. MK: Megakaryocyte; EP: Erythroid Progenitor; plt: Platelet; RBC: Red Blood Cell; WBC: White Blood Cell.