Special Section on Transporters in Toxicity and Disease—Minireview

Inhibition of the Multidrug Resistance P-Glycoprotein: Time for a Change of Strategy?

Richard Callaghan, Frederick Luk, and Mary Bebawy

Division of Biomedical Science & Biochemistry, Research School of Biology, College of Medicine, Biology & Environment, The Australian National University, Canberra, New South Wales, Australia (R.C.); and School of Pharmacy, Graduate School of Health, The University of Technology, Sydney, New South Wales, Australia (F.L., M.B.)

Received November 21, 2013; accepted January 31, 2014

ABSTRACT

P-glycoprotein (P-gp) is a key player in the multidrug-resistant phenotype in cancer. The protein confers resistance by mediating the ATP-dependent efflux of an astonishing array of anticancer drugs. Its broad specificity has been the subject of numerous attempts to inhibit the protein and restore the efficacy of anticancer drugs. The general strategy has been to develop compounds that either compete with anticancer drugs for transport or act as direct inhibitors of P-gp. Despite considerable *in vitro* success, there are no compounds currently available to “block” P-gp-mediated resistance in the clinic. The failure may be attributed to toxicity, adverse drug

interaction, and numerous pharmacokinetic issues. This review provides a description of several alternative approaches to overcome the activity of P-gp in drug-resistant cells. These include 1) drugs that specifically target resistant cells, 2) novel nanotechnologies to provide high-dose, targeted delivery of anticancer drugs, 3) compounds that interfere with nongenomic transfer of resistance, and 4) approaches to reduce the expression of P-gp within tumors. Such approaches have been developed through the pursuit of greater understanding of resistance mediators such as P-gp, and they show considerable potential for further application.

Introduction

The “permeability glycoprotein” or P-glycoprotein (P-gp or ABCB1) was discovered in 1976 in rodent cells known to display reduced sensitivity to anticancer drugs (Juliano and Ling, 1976). It was soon demonstrated that selection of cultured cancer cell lines in chemotherapeutic drugs displayed a phenotype consistent with the presence of P-gp. Moreover, these drug-resistant cell lines displayed resistance to a large number of chemically, structurally, and functionally unrelated drugs; hence the moniker of “multidrug resistance” (MDR). By the 1980s, antibodies had been developed to P-gp, and it was revealed that the protein was expressed in many distinct types of cancer as well as numerous normal tissues (Kartner et al., 1985; Cordon-Cardo et al., 1989, 1990).

The overexpression of P-gp in cancer was either an inherent or acquired process: the former, a reflection of its physiologic expression, and the latter, generated by the presence of anticancer drugs. P-gp confers resistance by preventing sufficient accumulation of anticancer drugs within the cell, thereby avoiding their cytotoxic or apoptotic effects. This is achieved by its ability to mediate ATP-dependent drug translocation across the plasma membrane against considerable concentration gradients. P-gp is a member of the B-class of the eukaryotic

ATP binding cassette (ABC) superfamily of transporters. Its influence in conferring MDR was at one time considered the paramount factor in the phenotype (Steinbach and Legrand, 2007). However, the burgeoning biochemical characterization of cancer cells revealed that the protein is a member of a network of cellular factors or tissue features that produce drug resistance (Mellor and Callaghan, 2008). The influence of P-gp was apparently further diluted by the discovery of two other ABC proteins able to confer MDR, namely, MRP1 (ABCC1) and BCRP (ABCG2) (Cole et al., 1992; Doyle et al., 1998). It is worth noting that although all three mediate active drug extrusion, their substrate specificities and expression patterns in cancer are distinct but with some overlap. The present review will focus on the role of P-gp and attempts to overcome its unwanted influence in cancer.

The multiplicity of factors contributing to drug resistance and the inability to overcome the actions of P-gp and restore the sensitivity of chemotherapy have led to researchers questioning its very involvement in clinical resistance (Bradshaw and Arceci, 1998; Merino et al., 2004; Perez-Tomas, 2006). This clear overreaction should be tempered by the plethora of investigations that have described the association of P-gp with drug resistance and the positive relationship between expression and poor prognosis (Gottesman et al., 2002; Leonard et al., 2003; Modok et al., 2006; Shaffer et al., 2012). The present review will not further discuss the relative merit or influence of P-gp in drug resistance but concentrate on efforts to overcome its actions.

Originally, it was thought that the actions of P-gp were limited to conferring resistance to classic genotoxic anticancer drugs, such as

Richard Callaghan would like to acknowledge funding support from the Association for International Cancer Research [Project Grant 12-0008] and The Wellcome Trust [Project Grant 094392/Z/10/Z].

dx.doi.org/10.1124/dmd.113.056176.

ABBREVIATIONS: ABC, ATP binding cassette; CHO, Chinese hamster ovary; MDR, multidrug resistance; MP, microparticle; P450, cytochrome P450; PEG, polyethylene glycol; P-gp, P-glycoprotein; PKC, protein kinase C.

vinblastine, doxorubicin, and paclitaxel. The broad or polyspecificity of P-gp is legendary (or infamous), and the list of compounds known to interact with this transporter is well in excess of 300 (Wang et al., 2011; Chen et al., 2012). It is apparent that many of the much touted “new generation” anticancer compounds (e.g., kinase inhibitors) are also substrates for transport by P-gp (Hegedus et al., 2002; Wang and Fu, 2010). There is a clear need to generate compounds, or strategies, to overcome the actions of P-gp in 1) limiting the effectiveness of chemotherapy in cancer and 2) influencing the pharmacokinetic profile of a vast number of clinically prescribed drugs.

Overcoming Drug Resistance to Chemotherapy Caused by P-gp

The general strategy to overcome multidrug resistance has been to coadminister chemical inhibitors of P-gp with anticancer drugs. Inhibition of P-gp would thereby lead to increased accumulation of anticancer drug within the cell and produce cell cytotoxicity. Alternatively, addition of a P-gp substrate in conjunction with the anticancer drug would achieve a similar effect by competing for the transport process. The first inhibitor (or more correctly referred to as a P-gp modulator) identified was the L-type calcium channel blocker verapamil (Tsuruo et al., 1982, 1983). This drug was shown to circumvent MDR using a variety of cell cytotoxicity, transport, binding, and photolabeling assays (Cornwell et al., 1987; Safa, 1988). However, clinical trials with verapamil were beset by serious cardiac side effects, and the compound was removed as a viable option (Ozols et al., 1987; Dalton et al., 1989). Verapamil inhibits L-type calcium channels with picomolar affinity; however, the potency to block P-gp was in the micromolar range, a 10^6 higher concentration.

In the ensuing 30 years, three distinct generations of P-gp modulator have been produced, and unfortunately a clinically useable inhibitor remains elusive (McHugh and Callaghan, 2008; Crowley et al., 2010). The first generation of compound used drugs with established and unrelated pharmacological actions and relied on the polyspecificity of P-gp. The next generation employed chemically modified first generation inhibitors. Ideally, these compounds were devoid of the parent compound's activity while retaining (or improving) the potency of interaction with P-gp. The third generation produced inhibitors from de novo synthesis using a variety of combinatorial chemistry approaches and benefiting from the burgeoning database of structure-activity relationships for drug–P-gp interaction.

A large number of compounds have been examined, and several recurring themes have emerged from their preclinical and clinical characterization. The first two generations were beset with poor potency of the compounds and a number of “off-target effects” that resulted in problems with toxicity (Gottesman et al., 2002; McHugh and Callaghan, 2008).

The third generation of P-gp modulators was frequently associated with adverse drug reactions that necessitated a reduction in the dose of anticancer agents. Many of the adverse drug interactions are related to the overlapping substrate specificity between P-gp and the drug metabolizing enzyme cytochrome P450 (CYP3A4 isoform) (Wacher et al., 1995; Yu, 1999). The combined effects of these two “defender proteins” will influence the absorption, distribution, metabolism, and elimination of a large proportion of known medications. Therefore, the coadministration of two drugs (i.e., modulator and anticancer drug) that interact with both P-gp and CYP3A4 often generates unpredictable toxicity related to the emergence of unwanted pharmacokinetic parameters. Such issues with pharmacokinetic profiles have resulted in the apparent downfall of potent P-gp inhibitors, such as Tariquidar (Pusztai et al., 2005). The fall from grace of this leading P-gp inhibitor led to considerable pessimism regarding the validity of utilizing efflux

pump inhibition as a means to overcome drug resistance. However, more recent observations have demonstrated that the coadministration of Tariquidar with Vinorelbine (Abraham et al., 2009) did not produce a similar toxicity profile as demonstrated with doxorubicin/docetaxel (Pusztai et al., 2005). Moreover, a structure-activity investigation has produced derivatives of Tariquidar with near negligible ability to interact with CYP3A4, while retaining P-gp inhibition (Labrie et al., 2007). These observations provide an argument for the retention of this strategy.

The presence of “endogenous or physiological” P-gp has also become problematic to the chemical inhibition strategy. P-gp is expressed at barrier tissue to sanctuary sites (e.g., blood-brain barrier) and at secretory/absorptive tissues (e.g., gastrointestinal tract) (Cordon-Cardo et al., 1989, 1990). The protein acts as a cellular defender and is involved in establishing the overall pharmacokinetic profile for numerous drugs. Unfortunately, chemical inhibitors are not capable of discriminating between P-gp expressed at normal sites in the body and that found in cancerous tissue. Therefore, the strategy generates unwanted side effects at a number of nontumor sites in the body, for example, increased permeability at the blood-brain barrier. Increased penetration, or distribution, of genotoxic anticancer drugs into the brain is associated with a severe toxicity profile that necessitates dose reduction.

Novel Approaches to Overcoming the Actions of P-gp

The preceding sections indicated a significant role for P-gp in conferring MDR to genotoxic anticancer drugs and many novel, target specific cytostatic compounds. The controversy surrounding the quantitative extent of its role in solid tumors cannot overshadow its involvement. Moreover, the importance of chemotherapy as a primary, adjuvant, or palliative treatment in cancer remains considerable and thus justifies the need to overcome, or evade, the influence of P-gp.

Unfortunately, the strategy of developing potent and selective compounds to inhibit P-gp and overcome resistance has encountered a number of issues. The issues relate to toxicity, dose reduction of anticancer drugs, and perturbation of key barrier tissues. In recent years there is a general pessimism by the pharmaceutical sector and research funding bodies despite the consistent and demonstrable improvements in design of inhibitors. There are also numerous epiphenomena related to P-gp expression that offer the prospect of therapeutic intervention. In this review we briefly describe some of the novel approaches rooted in academic research to counter, or overcome, the phenomenon of P-gp-mediated drug resistance.

Collateral Sensitivity

The emergence of a drug-resistant phenotype may be considered the product of the influence of a number of environmental stress factors. Typically, this will involve altered patterns of gene expression to enable survival of the cancer cell. A large number of cultured cell lines have been rendered drug resistant by prolonged exposure to certain anticancer drugs, and they undoubtedly contain modified expression patterns for numerous resistance mediators or enablers. The advent of high-throughput proteomic analyses revealed that drug-resistant cells or tumors display considerably modified protein expression profiles (Righetti et al., 2005; Peng et al., 2010). For example, in a taxol-resistant A2780 ovarian cancer cell line, proteomic analysis revealed marked changes in stress response effectors, cell cycle, and apoptotic mediators and numerous alterations in pathways for bioenergetic metabolism (Cicchillitti et al., 2009). Any proteomic alterations that are specific to drug-resistant cancer cells may, therefore, provide a

target for potential therapeutic management. In acquiring a drug-resistant phenotype, cancer cells may unwittingly proffer targets for their own eradication.

Increased or even hypersensitivity of resistant cells to various drugs was first observed in bacterial cells in the early 1950s (Szybalski and Bryson, 1952). The phenomenon was referred to as collateral sensitivity and was promoted a decade later for improving the efficacy of anticancer drugs in combination chemotherapy (Paigen, 1962). In the 1980s a multitude of studies demonstrated improved efficacy of methotrexate (Herman et al., 1979), folate analogs (Diddens et al., 1983), and DNA alkylating drugs (Sladek et al., 1985) in the presence of collateral sensitizing compounds. Unfortunately, the majority of these studies were not able to provide a description of the underlying resistant phenotype. A study in 1985 (Gupta, 1985) demonstrated collateral sensitivity between a spectrum of anticancer drugs in Chinese hamster ovary (CHO) cells that were eventually shown to express high levels of P-gp. Subsequently, in 1987, Cano-Gauci and Riordan provided the first demonstration that P-gp modulators (i.e., calcium channel blockers) preferentially inhibited the growth of drug-resistant cell lines.

It appears that the phenomenon of collateral sensitivity is closely associated with all three of the multidrug ABC pumps expressed in cancer cells (Oguro et al., 1990; Jensen et al., 1992). The precise mechanism underlying collateral sensitivity remains unresolved, with many hypotheses presented. As efforts in the 1990s to purify functional P-gp began in earnest it was widely demonstrated that resistant cells were more susceptible to procedures (e.g., nitrogen cavitation, shear force) or compounds (e.g., surfactants and ionophores), leading to physical disruption of the plasma membrane (Bech-Hansen et al., 1976; Callaghan and Riordan, 1995). It was proposed that high expression of this large polytopic membrane protein conferred greater fragility to the host cell membrane. However, the vast majority of observations were gleaned from cell lines selected for drug resistance and therefore likely to display numerous functional alterations. In addition, the issue of whether the membrane biophysical changes were a cause or effect of the resistant phenotype remained unclear as did the relationship to collateral sensitivity.

Observations that metabolic inhibitors also imparted collateral sensitivity suggested that bioenergetic metabolism in resistant cells may provide another therapeutic target (Ferretti et al., 1993; Bentley et al., 1996; Goda et al., 2002). The metabolic adaptations observed in cancer cells (e.g., Warburg effect) are now considered a hallmark feature of oncogenic transformation (Gillies and Gatenby, 2007; Zhao et al., 2011). Multidrug efflux pumps such as P-gp mediate an ATP-dependent transport process and sustain a high rate of ATP hydrolysis in the presence of substrate. Cancer cells have modified their metabolic profile to ensure sufficient anabolism to generate biomass for proliferation. This is achieved by a dampening of mitochondrial oxidative phosphorylation, with a heavy reliance on glycolysis despite its lower yield of energy (Bui and Thompson, 2006; Gillies and Gatenby, 2007). Thus, resistant cancer cells, with ATP-consuming efflux pumps, may conceivably be more susceptible to conditions of low energy status.

A significant proportion of established collateral sensitivity agents is known to be a substrate for transport by P-gp or stimulate its rate of ATP hydrolysis. A study by Broxterman et al. (1988) demonstrated that the collateral sensitizing drug verapamil significantly depleted ATP levels in a drug-resistant cell line. It has been proposed that depletion of ATP necessitates increased rates of oxidative phosphorylation in drug-resistant cells and that this oxidative stress results in an increased production of reactive oxygen species (e.g., O_2^- , H_2O_2). Increased reactive oxygen species and

concomitant reductions in the antioxidant glutathione were associated with apoptosis in drug-resistant CHO cells (Karwatsky et al., 2003). Conversely, compounds that are known to inhibit ATPase activity of P-gp (e.g., Tariquidar and PSC833) can diminish the effectiveness of collateral sensitivity agents. In addition, the extent of collateral sensitivity is proportional to the expression levels of P-gp, which may also reflect the strain on ATP levels (Goldsborough et al., 2011). More systematic characterization of ATP levels and metabolic fuel utilization in drug-resistant cells is clearly warranted.

A number of compounds have been demonstrated to elicit collateral sensitivity and include Tiopronin (Goldsborough et al., 2011), Desmosumotin (Nakagawa-Goto et al., 2008), NSC73306 (Hall et al., 2011), Dp44mT (Whitnall et al., 2006), and sigma-2 receptor agonists (Niso et al., 2013). These compounds impart collateral sensitivity; however, their mechanism remains unclear to date. Although these compounds are too preliminary for clinical application, they provide proof-of-principle that targeting this metabolic “weakness” in drug-resistant cells may provide an important adjunct to conventional chemotherapy.

Cytotoxic Drug Delivery Using Particles or Polymers

Chemotherapy drugs, particularly the genotoxic class, are unfortunately associated with severe and dose-limiting side effects, particularly in proliferating tissues. Therefore, the emergence of a drug-resistant phenotype cannot simply be overcome with escalation of dose. As indicated in preceding sections, the strategy to overcome resistance by inhibiting the activity of efflux pumps, such as P-gp, is also fraught with danger. P-gp is expressed at numerous locations in healthy tissue, and its inhibition will effectively open sanctuary sites (e.g., central nervous system and testes) to cytotoxic anticancer drugs (Leslie et al., 2005; de Vries et al., 2007; Robillard et al., 2012; Ke et al., 2013).

The observation that encapsulating doxorubicin within liposomes could circumvent P-gp-mediated MDR (Thierry et al., 1989; Sadasivan et al., 1991) provided a degree of cautious optimism. Clearly, this novel drug delivery route was able to bypass the actions of P-gp, because liposomal doxorubicin would enter the cells after endocytic engulfment of the particles rather than diffusion through the bilayer. This observation was used to formulate the “vacuum cleaner” hypothesis for the mechanism of P-gp (Higgins and Gottesman, 1992; Féré, 2000). Elegant studies with the fluorescent P-gp substrate calcein-AM were used to demonstrate that drugs do not need to “enter” the cytoplasm to become translocated by P-gp (Homolya et al., 1993). A mechanism of translocation whereby P-gp extracts drugs directly from the lipid milieu (Raviv et al., 1990; Homolya et al., 1993) gained general acceptance.

The liposomal delivery system may also provide a safe mechanism for significant dose escalation, because the volume of distribution for the drug will be modified by restricting diffusion out of the circulatory system. Clearance from the circulatory system was recognized early on as a limiting factor for the use of liposome-based systems. The primary issue is the ability of the reticuloendothelial system to trap and remove liposomes from the plasma, thereby considerably reducing residence time in the circulation (Oku and Namba, 1994). The incorporation of polymers, such as PEG or sialo-gangliosides, reduces trapping by the reticuloendothelial system (Oku and Namba, 1994; Gabizon and Martin, 1997). Tumors are widely accepted as a suitable target for liposomal encapsulation strategies because of their aberrant vasculature. In particular, the intratumor vasculature is leaky and thus enables high extravasation of the liposomes.

A considerable and ongoing level of interest in this strategy remains, although some refinement remains to improve particle stability and provide targeted delivery of the payload to the tumor.

Encapsulation strategies are now considerably more elaborate than the early liposomal systems. Modern delivery systems are referred to as nanoparticles or nanocarriers that feature drugs adsorbed, internalized, conjugated, or chelated to a platform (Hall et al., 2007; McNeil, 2009; Milane et al., 2011). The platforms include emulsions (e.g., PEG), liposomes, polymers, colloidal gold, and nanocrystals; moreover, they frequently contain mixtures of these. Nanoparticles improve the therapeutic index (LD_{50}/ED_{50}) by increasing bioavailability (fraction of dose reaching circulation) and reducing toxicity (Daum et al., 2012). Nanocarriers with encapsulation render the pharmacokinetic profile of a drug to be entirely dictated by properties of the platform. In addition, the high surface-to-volume ratios enable high encapsulation or surface adsorption of the payload drug.

Nanocarriers may be readily configured to contain 1) multiple payloads (e.g., drug and DNA/RNA), 2) molecules to enable tissue-specific targeting, and 3) "optical" agents for imaging and tracking. Strategies to target nanocarriers to tumors make use of the altered surface protein profile or expression levels for receptors in cancer cells. For example, folate derivatives (Yang et al., 2011), transferrin (Bellocq et al., 2003), or epidermal growth factor receptor (EGFR)-related molecules (El-Sayed et al., 2006; Magadala and Amiji, 2008) may be surface localized on nanocarriers to facilitate interaction with their respective receptor targets, which frequently display distinct expression patterns at the tumor site. The most commonly used techniques for modern imaging of nanocarriers are positron emission tomography or single photon emission computed tomography, which make use of a growing arsenal of radiolabeled compounds (Petersen et al., 2012). However, optical approaches such as fluorescence, infrared spectroscopy, and ultrasound are also widely used in experimental or clinical settings (Kozłowska et al., 2009). Imaging techniques provide measures of plasma clearance and tumor bioavailability: the latter a largely overlooked parameter in standard clinical trial design yet integral in generating an efficacious response. After internalization of the nanocarrier, the payload must be released to reach the specific intracellular target. Typically, internalized particles will enter the endosomal pathway that leads to lysosomal degradation (Riezman et al., 1997; Wattiaux et al., 2000), which may result in irretrievable transformation or sequestration of the payload. A number of strategies have been adopted, including the use of cell-penetrating peptides (Sarko et al., 2010; Choi et al., 2011) or platforms containing drug tethers that are destabilized by the acidic intratumoral microenvironment (Andreev et al., 2010; Gao et al., 2010; Du et al., 2013).

The use of nanocarriers to circumvent MDR by P-gp has adopted a number of guises. Lipid-dextran complexes as nanocarriers were shown to increase the potency of doxorubicin and suppress the expression of P-gp in resistant cells (Susa et al., 2010). The twofold effect of these nanocarriers on resistance was achieved using a payload of high doxorubicin dose and siRNA directed against the *mdr1* gene. Another study using a dual payload strategy (paclitaxel and lonidamine) also overcame the multidrug-resistant phenotype in cultured cells (Milane et al., 2011). Paclitaxel is a tubulin-disrupting anticancer drug, whereas lonidamine is a hexokinase II inhibitor that reduces glycolytic flux that produces cellular energetic stress. The platform consisted of a combination of polycaprolactone with PEG and contained an EGFR specific peptide. The latter was included to ensure targeting to drug-resistant MDR cells, which have been demonstrated to express elevated levels of EGFR.

Super-hi-magnetization nanocarriers produced with a highly magnetic Fe_3O_4 core and a poly-[N-(1-butyric acid)]aniline shell have also been used to tackle P-gp-mediated MDR (Hua et al., 2011). The Fe_3O_4 core acted as a contrast agent to enable tracking of the nanocarriers using magnetic resonance imaging techniques. High doxorubicin loading was achieved (271 mg drug per mg nanocarrier) by virtue of conjugating the drug to the aqueous shell. The conjugated doxorubicin facilitated high-capacity delivery and markedly improved thermal stability of the drug. A recent investigation used platform comprising a copolymer of polystyrene oxide-polyethylene oxide to target drug-resistant cells expressing P-gp (Cambón et al., 2013a,b). Intriguingly the copolymer unimers were able to directly inhibit the activity of P-gp, and the iron oxide-titanium oxide core enabled magnetic resonance imaging tracking of the spherical nanocarriers. Doxorubicin was the primary payload and was bound to the shell via titanium oxide, which was labile in the acidic pH within a tumor microenvironment. These polymers were remarkably stable and delivered ~150% more doxorubicin than observed with drug alone.

This exciting area of research has limitless possibilities and provides the potential for high-dose delivery of anticancer drugs and P-gp inhibitors (chemical or nucleotide based). The delivery may be safely tracked using noninvasive procedures and specifically targeted to the tumor site.

Microparticles as Novel Therapeutic Targets for the Prevention of MDR

Microparticles (MPs), also known as microvesicles, are enclosed plasma membrane fragments of size 0.1–1 μm characterized with phosphatidylserine expression on their surface (Gyorgy et al., 2011). MPs are typically released by vesiculation from the plasma membrane of stimulated or preapoptotic eukaryotic cells and had long been deemed to be inert cell debris (Freyssinet, 2003; Freyssinet and Toti, 2010).

In the last decade however, increasing evidence suggests that MP plays an essential role in cell-to-cell communications, allowing for cell regulation and crosstalk without the need for direct cell-to-cell contact. MPs act as vectors disseminating different biologic information, depending on their donor cell membrane and cytoplasmic composition. It is clear that MPs are important intermediates in inflammation, coagulation, vascular homeostasis, chemotaxis, adhesiveness, and thrombogenicity of endothelial cells and cells of hematopoietic lineage (Morel et al., 2004; Hugel et al., 2005). MPs shed from cancer cells are associated with tumor angiogenesis, evasion of immunosurveillance, cell invasiveness, survival, apoptosis, chemoresistance, and the hypercoagulable state (Dvorak et al., 1983; Ginestra et al., 1997; Dolo et al., 1999; Angelucci et al., 2000; Kim et al., 2002; Morel et al., 2004; Bebawy et al., 2009; Jaiswal et al., 2012b). Moreover, higher concentrations of systemic MPs are linked to disease progression and development in cerebral malaria, sepsis, autoimmune disorders, atherosclerosis, HIV-1, diabetes, and cancer (Horstman and Ahn, 1999; Nomura et al., 2004, 2008; Combes et al., 2006; Bebawy et al., 2009; Burnier et al., 2009).

The emergence of MDR was long attributed to "genetic alterations" within the cancer cell after an initial chemotherapeutic insult resulting in alterations to the cellular apparatus for prosurvival mechanisms. Bebawy et al. (2009) first reported that MDR can also be acquired in a non-"genetic manner" through a process of cell-to-cell communication in the absence of drug exposure. MDR cancer cells were shown to spontaneously shed MPs from their surface. These MDR-MPs contain cargo from the donor-resistant cells, in particular, functional resistance

proteins, which when transferred to a drug-responsive recipient cancer cell confer MDR within 4 hours (Bebawy et al., 2009; Lu et al., 2013).

This phenomenon has since been shown to occur *in vivo* (Jaiswal et al., 2013), whereby these microparticles transfer resistance proteins deep within the tumor core within 24 hours of MP exposure. This process also serves to "retemplate" the transcriptional landscape of recipient cells and in doing so results in the selective dominance of deleterious traits within the cancer cell population (Jaiswal et al., 2012a). Furthermore, in addition to the transfer of resistance proteins mediating MDR, these microparticles effectively sequester anticancer drugs and reduce the available free drug concentration available to a tumor mass, hence constituting a parallel resistance pathway (Gong et al., 2012).

Modulation and/or inhibition of MP formation and release may provide a novel therapeutic strategy for the prevention of MDR (Roseblade et al., 2013). A range of inhibitors have demonstrated promising effects in diminishing MP production by targeting key players in the formation of MPs. These include compounds that modulate calpain activation, calcium channels, and other factors that indirectly stimulate MP production (Roseblade et al., 2013).

Other inhibitors such as ticlopidine, pantethine, cystamine, LMP-20, or ROCK inhibitors have also been shown to lower MP levels after treatment (Shouzu et al., 2004; Wassmer et al., 2005; Penet et al., 2008; Antonyak et al., 2012). Although some of these drugs are effective in reducing MP production in patients, they were not capable of bringing down MP levels similar to the healthy control involved in the same study (Nomura et al., 2005, 2007). Further investigations are required to identify more effective inhibitors of MP formation and production, because it would provide a promising treatment strategy in the management of tumor drug resistance.

Altered Expression of P-gp

The expression of P-gp in normal (i.e., noncancerous) tissues is regulated by the nuclear orphan receptors namely, the pregnane X, steroid and xenobiotic, and the constitutive androstane receptors (Geick et al., 2001; Synold et al., 2001; Xu et al., 2005). These receptors mediate induction of the protein in response to cellular needs, typically defined by the presence of potential substrates. In particular, these two receptor subtypes are targets for a number of clinically used medications as well as steroid-based metabolites. The receptors also induce expression of phase I (e.g., cytochrome P450) and phase II (e.g., glutathione transferase) metabolizing enzymes (Ekins and Erickson, 2002; Xu et al., 2005; Weiss and Haefeli, 2013), further reinforcing the functional link between the efflux pumps and biotransformation pathways.

Expression of P-gp in cancer tissue is a more complex, or perhaps a more convoluted, regulatory system. There have been a wide range of factors that regulate P-gp expression, including hypoxia (via hypoxia inducible factor 1 α) (Comerford et al., 2002), metabolic acidosis (Thews et al., 2010), and metabolic perturbation such as glucose deprivation (Ledoux et al., 2003) and generation of reactive oxygen species (Wartenberg et al., 2005). Many of these factors derive from the hostile intratumor microenvironment and reflect cell or tissue stress. Moreover, the ability of anticancer drugs and radiotherapy to alter P-gp expression occurs via a cellular damage/stress response rather than a classic induction mechanism. It appears that P-gp is a key "first responder" to chemical or environmental insult on cancer tissue to remove potentially lethal compounds. After induction of cellular stress, the precise pathway to invoke induction of P-gp expression is complex and likely involves a multitude of cell signaling pathways (Callaghan et al.,

2008; Sui et al., 2012). It is for this reason that an ever increasing number of investigations have targeted specific kinases in signaling pathways with a view to modulating P-gp expression and, thus, the degree of chemoresistance.

The ability of kinases or signaling pathways to regulate P-gp function and/or expression warrants a discussion of posttranslational modifications for this transporter. The seminal manuscript that gave P-gp its name relied on detecting the protein in drug-resistant CHO cell membranes by virtue of its glycosylation status (Juliano and Ling, 1976). Further studies on its glycosylation status indicated that P-gp was synthesized as a 140-kDa precursor that was glycosylated over a 2- to 4-hour period to its fully mature form that migrated as a broad band at 180 kDa in SDS-PAGE (Richert et al., 1988; Yoshimura et al., 1989). Two investigations demonstrated that in the drug-resistant human KB cell line, P-gp was markedly stable, with a half-life greater than 24 hours (Richert et al., 1988; Yoshimura et al., 1989). In multidrug-resistant CHO cells, turnover of P-gp was considerably more rapid, with a stability half-life of 17 ± 3 hours (McClellan and Hill, 1993). More recently, it was demonstrated that expression of a P-gp-EGFP fusion protein (human P-gp isoform) in KB cells displayed a total half-life of 2.2 days and a surface stability of 3.7 days (Petriz et al., 2004). It remains unclear why the human and hamster versions of P-gp display distinct stabilities; moreover, this is not the only reported difference between the human and rodent isoforms.

It was subsequently demonstrated that the stability of P-gp was increased by 4- to 6-fold by serum deprivation or growing cells at high density (Muller et al., 1995), suggesting an involvement of growth factors in stability. Similarly, the expression of P-gp is cell cycle dependent, with greatest stability in the G0/G1 phase (Zhang and Ling, 2000).

What is the "trigger" to alter the stability of membrane-bound P-gp? The most likely candidate is the phosphorylation status, and the protein has of numerous PKC consensus motifs within the linker region (Chen et al., 1986; Gros et al., 1986). Moreover, it has been demonstrated that P-gp is phosphorylated by PKC at several of these motifs within the linker region (Chambers et al., 1993; Orr et al., 1993). A number of studies examined the phosphorylation status of P-gp in cell lines selected for drug resistance and suggested that it played a key role in the resistant phenotype (Marsh and Center, 1987; Mellado and Horwitz, 1987; Meyers, 1989; Staats et al., 1990). Subsequently, a number of investigations focused on using chemical inhibitors of PKC isoforms to reduce the activity or expression of P-gp and restore drug sensitivity to the cells (Chambers et al., 1990, 1992; Bates et al., 1992). The inhibitors were highly successful and pointed to a regulatory role of phosphorylation for P-gp. However, two manuscripts in the mid-1990s using site-directed mutagenesis of the PKC motifs in the linker region deflated this strategy (Germann et al., 1996; Goodfellow et al., 1996). Disruption of the PKC-mediated phosphorylation status did not alter the ability of P-gp to confer drug resistance; this was despite clear evidence of an interaction between the two proteins (Yang et al., 1996). It is now recognized that the chemical inhibitors of PKC were substrates for transport by P-gp, and their effects were related to direct inhibition with anticancer drugs for transport rather than via altered expression (Sato et al., 1990; Germann et al., 1995; Smith and Zilfou, 1995).

Perhaps unsurprisingly, the strategy to alter P-gp phosphorylation status to circumvent drug resistance was discontinued. In the ensuing period a great deal of research has focused on defining the myriad signaling pathways that control cancer cell growth. Many of these pathways (e.g., Ras/MAPK, JNK, p38 MAPK, protein kinase A- and PKC-related proteins, and PI3K) involve cell proliferation status, mediate apoptotic signaling, or generate the stress response. These

TABLE 1
A selection of novel compounds found to reverse P-gp MDR by altering signal transduction

Compound	Probable Pathway or Target	Reference
SC-51089	Prostaglandin E2 receptor	Pekcec et al., 2009
PHIL-7 (idarubicin derivative)	JNK-phosphorylation	Peng et al., 2013
siRNA Bile extract	Wnt/ β -catenin	Shen et al., 2013
PEITC (phenethyl isothiocyanate)	Pi3k-Akt	Tang et al., 2013
Dioscin	NF- κ B	Wang et al., 2013
Parthenolide	NF- κ B and Hsp70	Xin et al., 2013
Procyanidin	NF- κ B and MAPK/ERK	Zhao et al., 2013

signaling pathways are known to modulate P-gp expression in cells, tumor spheroids, and animal models (Chen and Sikic, 2012; Sui et al., 2012; Breier et al., 2013). In addition, both chemical inhibitors and silencing strategies (e.g., siRNA) of enzymes within these pathways have been demonstrated to affect P-gp function. Table 1 provides a small and selected list (restricted to 2013) of signaling pathway inhibitors that alter P-gp function in drug-resistant cells.

Unfortunately, the vast majority of these studies have adopted a similar strategy to those involving PKC in the 1990s, namely that the reporter assay was based exclusively on measurement of overall cytotoxicity. On the basis of this previous experience, it may be argued that mechanistic information will be of paramount importance in the possibility of reviving this strategy to tackle MDR. Key parameters include whether 1) the phosphorylation (or other post-translational modification) status of P-gp, 2) the total cellular expression level of P-gp, or 3) the cellular localization of P-gp has been altered. It would also be informative to ascertain whether novel kinases or phosphatases directly interact with the transporter. Pharmacological investigations should address whether the chemical inhibitors of signaling pathways actually modulate P-gp function by competing with anticancer drugs for the transport process.

Differential Modulation of P-gp

At present, the common strategy of incorporating pharmacological inhibitors of P-gp in conventional treatment regimens is complicated with issues of specificity, toxicity, and differential modulation. It was shown previously that P-gp transport function, and subsequently MDR in cancer, is differentially modulated depending on the drug combination used and the site of interaction within the plasma membrane bilayer (Bebawy et al., 2001; Huang et al., 2007). Specifically, modulators appear to have a selective effect depending on the anticancer drug with which they are combined. For instance, it has been shown that verapamil reversed resistance of CEM/VLB100 cells to vinblastine and fluorescein-colchicine but not to colchicine (Bebawy et al., 2001; Huang et al., 2007). In contrast, chlorpromazine reversed resistance to vinblastine but not to fluorescein-colchicine and increased the degree of resistance to colchicine (Bebawy et al., 2001; Huang et al., 2007). Similarly, gypenoside extracts from *Gynnostemma pentaphyllum* effectively circumvent colchicine resistance by 15-fold in P-gp-overexpressed CEM/VLB100 cells, while having no effect on vinblastine and taxol resistance in the same cells (Huang et al., 2007). This concept of differential pharmacological modulation has since been proposed as a cellular mechanism potentially contributing to the emergence of pharmacoresistant schizophrenia (Bebawy and Chetty, 2008), supporting the significance and application of these early findings across different clinical conditions. This evidence provides us with a strategy for tailoring drug regimens to include MDR inhibitors according to the specific anticancer drug combinations used.

Perspectives and Summary

The vast majority of investigations aimed at overcoming drug resistance conferred by P-gp have employed the mantra of direct inhibition of the transporter. Unfortunately, this approach has not yet reached clinical success due to the complex array of drug toxicity, altered pharmacokinetics and adverse drug interactions. The current review has presented a number of alternative approaches to circumventing P-gp-mediated MDR. The novel strategies have been identified through the tireless efforts by researchers to gain a more complete understanding of this phenotype. These strategies will hopefully continue to develop and generate targeted, selective, and nontoxic strategies to eradicate the presence of MDR in cancer.

Authorship Contributions

Wrote or contributed to the writing of the manuscript: Callaghan, Luk, Bebawy.

References

- Abraham J, Edgerly M, Wilson R, Chen C, Rutt A, Bakke S, Robey R, Dwyer A, Goldspiel B, and Balis F, et al. (2009) A phase I study of the P-glycoprotein antagonist tariquidar in combination with vinorelbine. *Clin Cancer Res* **15**:3574–3582.
- Andreev OA, Engelman DM, and Reshetnyak YK (2010) pH-sensitive membrane peptides (pHLIPs) as a novel class of delivery agents. *Mol Membr Biol* **27**:341–352.
- Angelucci A, D'Ascenzo S, Festuccia C, Gravina GL, Bologna M, Dolo V, and Pavan A (2000) Vesicle-associated urokinase plasminogen activator promotes invasion in prostate cancer cell lines. *Clin Exp Metastasis* **18**:163–170.
- Antonyak MA, Wilson KF, and Cerione RA (2012) R(h)oads to microvesicles. *Small GTPases* **3**:219–224.
- Bates SE, Currier SJ, Alvarez M, and Fojo AT (1992) Modulation of P-glycoprotein phosphorylation and drug transport by sodium butyrate. *Biochemistry* **31**:6366–6372.
- Bebawy M and Chetty M (2008) Differential pharmacological regulation of drug efflux and pharmacoresistant schizophrenia. *Bioessays* **30**:183–188.
- Bebawy M, Combes V, Lee E, Jaiswal R, Gong J, Bonhoure A, and Grau GE (2009) Membrane microparticles mediate transfer of P-glycoprotein to drug sensitive cancer cells. *Leukemia* **23**:1643–1649.
- Bebawy M, Morris MB, and Roufogalis BD (2001) Selective modulation of P-glycoprotein-mediated drug resistance. *Br J Cancer* **85**:1998–2003.
- Bech-Hansen NT, Till JE, and Ling V (1976) Pleiotropic phenotype of colchicine-resistant CHO cells: cross-resistance and collateral sensitivity. *J Cell Physiol* **88**:23–31.
- Bellocc NC, Pun SH, Jensen GS, and Davis ME (2003) Transferrin-containing, cyclodextrin polymer-based particles for tumor-targeted gene delivery. *Bioconjug Chem* **14**:1122–1132.
- Bentley J, Bell SE, Quinn DM, Kellett GL, and Warr JR (1996) 2-deoxy-D-glucose toxicity and transport in human multidrug-resistant KB carcinoma cell lines. *Oncol Res* **8**:77–84.
- Bradshaw DM and Arceci RJ (1998) Clinical relevance of transmembrane drug efflux as a mechanism of multidrug resistance. *J Clin Oncol* **16**:3674–3690.
- Breier A, Gibalova L, Seres M, Barancik M, and Sulova Z (2013) New insight into p-glycoprotein as a drug target. *Anticancer Agents Med Chem* **13**:159–170.
- Broxterman HJ, Pinedo HM, Kuiper CM, Kaptein LC, Schuurhuis GJ, and Lankelma J (1988) Induction by verapamil of a rapid increase in ATP consumption in multidrug-resistant tumor cells. *FASEB J* **2**:2278–2282.
- Bui T and Thompson CB (2006) Cancer's sweet tooth. *Cancer Cell* **9**:419–420.
- Burnier L, Fontana P, Kwak BR, and Angelillo-Scherer A (2009) Cell-derived microparticles in haemostasis and vascular medicine. *Thromb Haemostasis* **101**:439–451.
- Callaghan R, Crowley E, Potter S, and Kerr ID (2008) P-glycoprotein: so many ways to turn it on. *J Clin Pharmacol* **48**:365–378.
- Callaghan R and Riordan JR (1995) Collateral sensitivity of multidrug resistant cells to narcotic analgesics is due to effects on the plasma membrane. *Biochim Biophys Acta* **1236**:155–162.
- Cambón A, Brea J, Loza MI, Alvarez-Lorenzo C, Concheiro A, Barbosa S, Taboada P, and Mosquera V (2013) Cytocompatibility and P-glycoprotein inhibition of block copolymers: structure-activity relationship. *Mol Pharm* **10**:3232–3241.
- Cambón A, Rey-Rico A, Barbosa S, Soltero JF, Yeates SG, Brea J, Loza MI, Alvarez-Lorenzo C, Concheiro A, and Taboada P, et al. (2013a) Poly(styrene oxide)-poly(ethylene oxide) block copolymers: From “classical” chemotherapeutic nanocarriers to active cell-response inducers. *J Control Release* **167**:68–75.

- Cambón A, Rey-Rico A, Mistry D, Brea J, Loza MI, Attwood D, Barbosa S, Alvarez-Lorenzo C, Concheiro A, and Taboada P, et al. (2013b) Doxorubicin-loaded micelles of reverse poly (butylene oxide)-poly(ethylene oxide)-poly(butylene oxide) block copolymers as efficient "active" chemotherapeutic agents. *Int J Pharm* **445**:47–57.
- Cano-Gauci DF and Riordan JR (1987) Action of calcium antagonists on multidrug resistant cells. Specific cytotoxicity independent of increased cancer drug accumulation. *Biochem Pharmacol* **36**:2115–2123.
- Chambers TC, McAvoy EM, Jacobs JW, and Eilon G (1990) Protein kinase C phosphorylates P-glycoprotein in multidrug resistant human KB carcinoma cells. *J Biol Chem* **265**:7679–7686.
- Chambers TC, Pohl J, Raynor RL, and Kuo JF (1993) Identification of specific sites in human P-glycoprotein phosphorylated by protein kinase C. *J Biol Chem* **268**:4592–4595.
- Chambers TC, Zheng B, and Kuo JF (1992) Regulation by phorbol ester and protein kinase C inhibitors, and by a protein phosphatase inhibitor (okadaic acid), of P-glycoprotein phosphorylation and relationship to drug accumulation in multidrug-resistant human KB cells. *Mol Pharmacol* **41**:1008–1015.
- Chen CJ, Chin JE, Ueda K, Clark DP, Pastan I, Gottesman MM, and Roninson IB (1986) Internal duplication and homology with bacterial transport proteins in the mdr1 (P-glycoprotein) gene from multidrug-resistant human cells. *Cell* **47**:381–389.
- Chen KG and Sikic BI (2012) Molecular pathways: regulation and therapeutic implications of multidrug resistance. *Clin Cancer Res* **18**:1863–1869.
- Chen L, Li Y, Yu H, Zhang L, and Hou T (2012) Computational models for predicting substrates or inhibitors of P-glycoprotein. *Drug Discov Today* **17**:343–351.
- Choi YS, Lee JY, Suh JS, Lee SJ, Yang VC, Chung CP, and Park YJ (2011) Cell penetrating peptides for tumor targeting. *Curr Pharm Biotechnol* **12**:1166–1182.
- Cicchilitti L, Di Michele M, Urbani A, Ferlini C, Donat MB, Scambia G, and Rotilio D (2009) Comparative proteomic analysis of paclitaxel sensitive A2780 epithelial ovarian cancer cell line and its resistant counterpart A2780TC1 by 2D-DIGE: the role of ERp57. *J Proteome Res* **8**:1902–1912.
- Cole SP, Bhardwaj G, Gerlach JH, Mackie JE, Grant CE, Almquist KC, Stewart AJ, Kurz EU, Duncan AM, and Deeley RG (1992) Overexpression of a transporter gene in a multidrug-resistant human lung cancer cell line. *Science* **258**:1650–1654.
- Combes V, Coltel N, Faillie D, Wassmer SC, and Grau GE (2006) Cerebral malaria: role of microparticles and platelets in alterations of the blood-brain barrier. *Int J Parasitol* **36**:541–546.
- Cornford KM, Wallace TJ, Karhausen J, Louis NA, Montalto MC, and Colgan SP (2002) Hypoxia-inducible factor-1-dependent regulation of the multidrug resistance (MDR1) gene. *Cancer Res* **62**:3387–3394.
- Cordon-Cardo C, O'Brien JP, Boccia J, Casals D, Bertino JR, and Melamed MR (1990) Expression of the multidrug resistance gene product (P-glycoprotein) in human normal and tumor tissues. *J Histochem Cytochem* **38**:1277–1287.
- Cordon-Cardo C, O'Brien JP, Casals D, Rittman-Grauer L, Biedler JL, Melamed MR, and Bertino JR (1989) Multidrug-resistance gene (P-glycoprotein) is expressed by endothelial cells at blood-brain barrier sites. *Proc Natl Acad Sci USA* **86**:695–698.
- Cornwell MM, Pastan I, and Gottesman MM (1987) Certain calcium channel blockers bind specifically to multidrug-resistant human KB carcinoma membrane vesicles and inhibit drug binding to P-glycoprotein. *J Biol Chem* **262**:2166–2170.
- Crowley E, McDevitt CA, and Callaghan R (2010) Generating inhibitors of P-glycoprotein: where to, now? *Methods Mol Biol* **596**:405–432.
- Dalton WS, Grogan TM, Meltzer PS, Scheper RJ, Durie BG, Taylor CW, Miller TP, and Salmon SE (1989) Drug-resistance in multiple myeloma and non-Hodgkin's lymphoma: detection of P-glycoprotein and potential circumvention by addition of verapamil to chemotherapy. *J Clin Oncol* **7**:415–424.
- Daum N, Tscheka C, Neumeyer A, and Schneider M (2012) Novel approaches for drug delivery systems in nanomedicine: effects of particle design and shape. *Wiley Interdiscip Rev Nanomed Nanobiotechnol* **4**:52–65.
- de Vries NA, Zhao J, Kroon E, Buckle T, Beijnen JH, and van Tellingen O (2007) P-glycoprotein and breast cancer resistance protein: two dominant transporters working together in limiting the brain penetration of topotecan. *Clin Cancer Res* **13**:6440–6449.
- Diddens H, Niethammer D, and Jackson RC (1983) Patterns of cross-resistance to the antifolate drugs trimetrexate, methotrexate, and CB3717 in human lymphoma and osteosarcoma cells resistant to methotrexate. *Cancer Res* **43**:5286–5292.
- Dolo V, D'Ascenzo S, Violini S, Pompucci L, Festuccia C, Ginestra A, Vittorelli ML, Canevari S, and Pavan A (1999) Matrix-degrading proteinases are shed in membrane vesicles by ovarian cancer cells in vivo and in vitro. *Clin Exp Metastasis* **17**:131–140.
- Doyle LA, Yang W, Abruzzo LV, Krogmann T, Gao Y, Rishi AK, and Ross DD (1998) A multidrug resistance transporter from human MCF-7 breast cancer cells. *Proc Natl Acad Sci USA* **95**:15665–15670.
- Du J-Z, Mao C-Q, Yuan Y-Y, Yang X-Z, and Wang J (2013) Tumor extracellular acidity-activated nanoparticles as drug delivery systems for enhanced cancer therapy. *Biotechnol Adv* DOI: 10.1016/j.biotechadv.2013.08.002.
- Dvorak HF, Van DeWater L, Bitzer AM, Dvorak AM, Anderson D, Harvey VS, Bach R, Davis GL, DeWolf W, and Carvalho AC (1983) Procoagulant activity associated with plasma membrane vesicles shed by cultured tumor cells. *Cancer Res* **43**:4434–4442.
- Ekins S and Erickson JA (2002) A pharmacophore for human pregnane X receptor ligands. *Drug Metab Dispos* **30**:96–99.
- El-Sayed IH, Huang X, and El-Sayed MA (2006) Selective laser photo-thermal therapy of epithelial carcinoma using anti-EGFR antibody conjugated gold nanoparticles. *Cancer Lett* **239**:129–135.
- Ferretti A, Chen LL, Di Vito M, Barca S, Tombesi M, Cianfriglia M, Bozzi A, Strom R, and Podo F (1993) Pentose phosphate pathway alterations in multi-drug resistant leukemic T-cells: 31P NMR and enzymatic studies. *Anticancer Res* **13**:867–872.
- Ferté J (2000) Analysis of the tangled relationships between P-glycoprotein-mediated multidrug resistance and the lipid phase of the cell membrane. *Eur J Biochem* **267**:277–294.
- Freyssinet JM (2003) Cellular microparticles: what are they bad or good for? *J Thromb Haemost* **1**:1655–1662.
- Freyssinet JM and Toti F (2010) Formation of procoagulant microparticles and properties. *Thromb Res* **125** (Suppl 1):S46–S48.
- Gabizon A and Martin F (1997) Polyethylene glycol-coated (pegylated) liposomal doxorubicin. Rationale for use in solid tumors. *Drugs* **54** (Suppl 4):15–21.
- Gao W, Chan JM, and Farokhzad OC (2010) pH-Responsive nanoparticles for drug delivery. *Mol Pharm* **7**:1913–1920.
- Geick A, Eichelbaum M, and Burk O (2001) Nuclear receptor response elements mediate induction of intestinal MDR1 by rifampin. *J Biol Chem* **276**:14581–14587.
- Germann UA, Chambers TC, Ambudkar SV, Licht T, Cardarelli CO, Pastan I, and Gottesman MM (1996) Characterization of phosphorylation-defective mutants of human P-glycoprotein expressed in mammalian cells. *J Biol Chem* **271**:1708–1716.
- Germann UA, Chambers TC, Ambudkar SV, Pastan I, and Gottesman MM (1995) Effects of phosphorylation of P-glycoprotein on multidrug resistance. *J Bioenerg Biomembr* **27**:53–61.
- Gillies RJ and Gatenby RA (2007) Adaptive landscapes and emergent phenotypes: why do cancers have high glycolysis? *J Bioenerg Biomembr* **39**:251–257.
- Ginestra A, Monea S, Seghezzi G, Dolo V, Nagase H, Mignatti P, and Vittorelli ML (1997) Urokinase plasminogen activator and gelatinases are associated with membrane vesicles shed by human HT1080 fibrosarcoma cells. *J Biol Chem* **272**:17216–17222.
- Goda K, Nagy H, Mechetter E, Cianfriglia M, and Szabó G, Jr (2002) Effects of ATP depletion and phosphate analogues on P-glycoprotein conformation in live cells. *Eur J Biochem* **269**:2672–2677.
- Goldsborough AS, Handley MD, Dulcey AE, Pluchino KM, Kannan P, Brimacombe KR, Hall MD, Griffiths G, and Gottesman MM (2011) Collateral sensitivity of multidrug-resistant cells to the orphan drug tiopronin. *J Med Chem* **54**:4987–4997.
- Gong J, Jaiswal R, Mathys JM, Combes V, Grau GE, and Bebaawy M (2012) Microparticles and their emerging role in cancer multidrug resistance. *Cancer Treat Rev* **38**:226–234.
- Goodfellow HR, Sardini A, Ruetz S, Callaghan R, Gros P, McNaughton PA, and Higgins CF (1996) Protein kinase C-mediated phosphorylation does not regulate drug transport by the human multidrug resistance P-glycoprotein. *J Biol Chem* **271**:13668–13674.
- Gottesman MM, Fojo T, and Bates SE (2002) Multidrug resistance in cancer: role of ATP-dependent transporters. *Nat Rev Cancer* **2**:48–58.
- Gros P, Croop J, and Housman D (1986) Mammalian multidrug resistance gene: complete cDNA sequence indicates strong homology to bacterial transport proteins. *Cell* **47**:371–380.
- Gupta RS (1985) Cross-resistance of vinblastine- and taxol-resistant mutants of Chinese hamster ovary cells to other anticancer drugs. *Cancer Treat Rep* **69**:515–521.
- György B, Szabó TG, Pásztói M, Pál Z, Misják P, Aradi B, László V, Pállinger E, Pap E, and Kittel A, et al. (2011) Membrane vesicles, current state-of-the-art: emerging role of extracellular vesicles. *Cell Mol Life Sci* **68**:2667–2688.
- Hall JB, Dobrovolskaia MA, Patri AK, and McNeil SE (2007) Characterization of nanoparticles for therapeutics. *Nanomedicine (Lond)* **2**:789–803.
- Hall MD, Brimacombe KR, Varonka MS, Pluchino KM, Monda JK, Li J, Walsh MJ, Boxer MB, Warren TH, and Fales HM, et al. (2011) Synthesis and structure-activity evaluation of isatin- β -thiosemicarbazones with improved selective activity toward multidrug-resistant cells expressing P-glycoprotein. *J Med Chem* **54**:5878–5889.
- Hegedus T, Orfi L, Seprodi A, Váradi A, Sarkadi B, and Kéri G (2002) Interaction of tyrosine kinase inhibitors with the human multidrug transporter proteins, MDR1 and MRP1. *Biochim Biophys Acta* **1587**:318–325.
- Herman TS, Cress AE, and Gerner EW (1979) Collateral sensitivity to methotrexate in cells resistant to adriamycin. *Cancer Res* **39**:1937–1942.
- Higgins CF and Gottesman MM (1992) Is the multidrug transporter a flippase? *Trends Biochem Sci* **17**:18–21.
- Homolya L, Holló Z, Germann UA, Pastan I, Gottesman MM, and Sarkadi B (1993) Fluorescent cellular indicators are extruded by the multidrug resistance protein. *J Biol Chem* **268**:21493–21496.
- Horstman LL and Ahn YS (1999) Platelet microparticles: a wide-angle perspective. *Crit Rev Oncol Hematol* **30**:111–142.
- Hua MY, Yang HW, Liu HL, Tsai RY, Pang ST, Chuang KL, Chang YS, Hwang TL, Chang YH, and Chuang HC, et al. (2011) Superhigh-magnetization nanocarrier as a doxorubicin delivery platform for magnetic targeting therapy. *Biomaterials* **32**:8999–9010.
- Huang TH, Bebaawy M, Tran VH, and Roufogali BD (2007) Specific reversal of multidrug resistance to colchicine in CEM/VLB(100) cells by Gynostemma pentaphyllum extract. *Phytomedicine* **14**:830–839.
- Hugel B, Martínez MC, Kunzelmann C, and Freyssinet JM (2005) Membrane microparticles: two sides of the coin. *Physiology (Bethesda)* **20**:22–27.
- Jaiswal R, Gong J, Sambasivam S, Combes V, Mathys JM, Davey R, Grau GE, and Bebaawy M (2012a) Microparticle-associated nucleic acids mediate trait dominance in cancer. *FASEB J* **26**:420–429.
- Jaiswal R, Luk F, Dalla PV, Grau GE, and Bebaawy M (2013) Breast cancer-derived microparticles display tissue selectivity in the transfer of resistance proteins to cells. *PLoS ONE* **8**:e61515.
- Jaiswal R, Luk F, Gong J, Mathys JM, Grau GE, and Bebaawy M (2012b) Microparticle conferred microRNA profiles—implications in the transfer and dominance of cancer traits. *Mol Cancer* DOI: 10.1186/1476-4598-11-37 [published ahead of print].
- Jensen PB, Roed H, Sehested M, Demant EJ, Vindelov L, Christensen IJ, and Hansen HH (1992) Doxorubicin sensitivity pattern in a panel of small-cell lung-cancer cell lines: correlation to etoposide and vincristine sensitivity and inverse correlation to carmustine sensitivity. *Cancer Chemother Pharmacol* **31**:46–52.
- Juliano RL and Ling V (1976) A surface glycoprotein modulating drug permeability in Chinese hamster ovary cell mutants. *Biochem Biophys Acta* **455**:152–162.
- Kartner N, Evernden-Porelle D, Bradley G, and Ling V (1985) Detection of P-glycoprotein in multidrug-resistant cell lines by monoclonal antibodies. *Nature* **316**:820–823.
- Karwatsky J, Lincoln MC, and Georges E (2003) A mechanism for P-glycoprotein-mediated apoptosis as revealed by verapamil hypersensitivity. *Biochemistry* **42**:12163–12173.
- Ke AB, Eyal S, Chung FS, Link JM, Mankoff DA, Muzi M, and Unadkat JD (2013) Modeling cyclosporine A inhibition of the distribution of a P-glycoprotein PET ligand, 11C-verapamil, into the maternal brain and fetal liver of the pregnant nonhuman primate: impact of tissue blood flow and site of inhibition. *J Nucl Med* **54**:437–446.
- Kim CW, Lee HM, Lee TH, Kang C, Kleinman HK, and Gho YS (2002) Extracellular membrane vesicles from tumor cells promote angiogenesis via sphingomyelin. *Cancer Res* **62**:6312–6317.
- Kozłowska D, Foran P, MacMahon P, Shelly MJ, Eustace S, and O'Kennedy R (2009) Molecular and magnetic resonance imaging: The value of immunoliposomes. *Adv Drug Deliv Rev* **61**:1402–1411.
- Labrie P, Maddafo SP, Lacroix J, Catalano C, Lee DK, Rakhit S, and Gaudreault RC (2007) In vitro activity of novel dual action MDR anthranilamide modulators with inhibitory activity on CYP-450 (Part 2). *Bioorg Med Chem* **15**:3854–3868.
- Ledoux S, Yang R, Friedlander G, and Laouari D (2003) Glucose depletion enhances P-glycoprotein expression in hepatoma cells: role of endoplasmic reticulum stress response. *Cancer Res* **63**:7284–7290.

- Leonard GD, Fojo T, and Bates SE (2003) The role of ABC transporters in clinical practice. *Oncologist* **8**:411–424.
- Leslie EM, Deeley RG, and Cole SP (2005) Multidrug resistance proteins: role of P-glycoprotein, MRP1, MRP2, and BCRP (ABCG2) in tissue defense. *Toxicol Appl Pharmacol* **204**:216–237.
- Lu JF, Luk F, Gong J, Jaiswal R, Grau GE, and Bebawy M (2013) Microparticles mediate MRP1 intercellular transfer and the re-templating of intrinsic resistance pathways. *Pharmacol Res* **76**:77–83.
- Magadala P and Amiji M (2008) Epidermal growth factor receptor-targeted gelatin-based engineered nanocarriers for DNA delivery and transfection in human pancreatic cancer cells. *AAPS J* **10**:565–576.
- Marsh W and Center MS (1987) Adriamycin resistance in HL60 cells and accompanying modification of a surface membrane protein contained in drug-sensitive cells. *Cancer Res* **47**:5080–5086.
- McClean S and Hill BT (1993) Evidence of post-translational regulation of P-glycoprotein associated with the expression of a distinctive multiple drug-resistant phenotype in Chinese hamster ovary cells. *Eur J Cancer* **29A**:2243–2248.
- McHugh K and Callaghan R (2008) Clinical trials on MDR reversal agents, in *Multidrug Resistance: Biological and Pharmaceutical Advances in Antitumour Treatment* (Colabufo NA, ed) pp 321–353. Research Signpost, Kerala.
- McNeil SE (2009) Nanoparticle therapeutics: a personal perspective. *Wiley Interdiscip Rev Nanomed Nanobiotechnol* **1**:264–271.
- Mellado W and Horwitz SB (1987) Phosphorylation of the multidrug resistance associated glycoprotein. *Biochemistry* **26**:6900–6904.
- Mellor HR and Callaghan R (2008) Resistance to chemotherapy in cancer: a complex and integrated cellular response. *Pharmacology* **81**:275–300.
- Merino V, Jiménez-Torres NV, and Merino-Sanjuán M (2004) Relevance of multidrug resistance proteins on the clinical efficacy of cancer therapy. *Curr Drug Deliv* **1**:203–212.
- Meyers MB (1989) Protein phosphorylation in multidrug resistant Chinese hamster cells. *Cancer Commun* **1**:233–241.
- Milane L, Duan Z, and Amiji M (2011) Therapeutic efficacy and safety of paclitaxel/Ionidamine loaded EGFR-targeted nanoparticles for the treatment of multi-drug resistant cancer. *PLoS ONE* **6**:e24075.
- Modok S, Mellor HR, and Callaghan R (2006) Modulation of multidrug resistance efflux pump activity to overcome chemoresistance in cancer. *Curr Opin Pharmacol* **6**:350–354.
- Morel O, Toti F, Hugel B, and Freyssinet JM (2004) Cellular microparticles: a disseminated storage pool of bioactive vascular effectors. *Curr Opin Hematol* **11**:156–164.
- Muller C, Laurent G, and Ling V (1995) P-glycoprotein stability is affected by serum deprivation and high cell density in multidrug-resistant cells. *J Cell Physiol* **163**:538–544.
- Nakagawa-Goto K, Bastow KF, Chen TH, Morris-Natschke SL, and Lee KH (2008) Antitumor agents 260. New desmosdomotin B analogues with improved in vitro anticancer activity. *J Med Chem* **51**:3297–3303.
- Niso M, Abate C, Contino M, Ferorelli S, Azzariti A, Perrone R, Colabufo NA, and Berardi F (2013) Sigma-2 receptor agonists as possible antitumor agents in resistant tumors: hints for collateral sensitivity. *ChemMedChem* **8**:2026–2035.
- Nomura S, Inami N, Kimura Y, Omoto S, Shouzu A, Nishikawa M, and Iwasaka T (2007) Effect of nifedipine on adiponectin in hypertensive patients with type 2 diabetes mellitus. *J Hum Hypertens* **21**:38–44.
- Nomura S, Ozaki Y, and Ikeda Y (2008) Function and role of microparticles in various clinical settings. *Thromb Res* **123**:8–23.
- Nomura S, Shouzu A, Omoto S, Nishikawa M, and Iwasaka T (2005) Benidipine improves oxidized LDL-dependent monocyte and endothelial dysfunction in hypertensive patients with type 2 diabetes mellitus. *J Hum Hypertens* **19**:551–557.
- Nomura S, Takahashi N, Inami N, Kajitara T, Yamada K, Nakamori H, and Tsuda N (2004) Probucol and ticlopidine: effect on platelet and monocyte activation markers in hyperlipidemic patients with and without type 2 diabetes. *Atherosclerosis* **174**:329–335.
- Oguro M, Seki Y, Okada K, and Andoh T (1990) Collateral drug sensitivity induced in CPT-11 (a novel derivative of camptothecin)-resistant cell lines. *Biomed Pharmacother* **44**:209–216.
- Oku N and Namba Y (1994) Long-circulating liposomes. *Crit Rev Ther Drug Carrier Syst* **11**:231–270.
- Orr GA, Han EK, Browne PC, Nieves E, O'Connor BM, Yang CP, and Horwitz SB (1993) Identification of the major phosphorylation domain of murine mdr1b P-glycoprotein. Analysis of the protein kinase A and protein kinase C phosphorylation sites. *J Biol Chem* **268**:25054–25062.
- Ozols RF, Cunnion RE, Klecker RW, Jr, Hamilton TC, Ostchega Y, Parrillo JE, and Young RC (1987) Verapamil and adriamycin in the treatment of drug-resistant ovarian cancer patients. *J Clin Oncol* **5**:641–647.
- Paigen K (1962) The prediction of growth-inhibitory drug combinations showing enhanced differential toxicity and collateral sensitivity. *Cancer Res* **22**:1290–1296.
- Pekcec A, Unkrüer B, Schlichtiger J, Soerensen J, Hartz AMS, Bauer B, van Vliet EA, Gorter JA, and Potschka H (2009) Targeting prostaglandin E2 EP1 receptors prevents seizure-associated P-glycoprotein up-regulation. *J Pharmacol Exp Ther* **330**:939–947.
- Penet MF, Abou-Hamdan M, Coltel N, Cornille E, Grau GE, de Reggi M, and Gharib B (2008) Protection against cerebral malaria by the low-molecular-weight thiol pantethine. *Proc Natl Acad Sci USA* **105**:1321–1326.
- Peng H, Yuan X, Shi R, Wei X, Ren S, Yan C, Ding Y, Lin Y, Fan D, and Yang M, et al. (2013) PHIL-7 inhibits cell growth and induces apoptosis in leukemia cell line K562 as well as its MDR- counterpart K562/A02 through producing reactive oxygen species. *Eur J Pharmacol* **718**:459–468.
- Peng XC, Gong FM, Wei M, Chen X, Chen Y, Cheng K, Gao F, Xu F, Bi F, and Liu JY (2010) Proteomic analysis of cell lines to identify the irinotecan resistance proteins. *J Biosci* **35**:557–564.
- Pérez-Tomás R (2006) Multidrug resistance: retrospect and prospects in anti-cancer drug treatment. *Curr Med Chem* **13**:1859–1876.
- Petersen AL, Hansen AE, Gabizon A, and Andresen TL (2012) Liposome imaging agents in personalized medicine. *Adv Drug Deliv Rev* **64**:1417–1435.
- Pétriz J, Gottesman MM, and Aran JM (2004) An MDR-EGFP gene fusion allows for direct cellular localization, function and stability assessment of P-glycoprotein. *Curr Drug Deliv* **1**:43–56.
- Pusztai L, Wagner P, Ibrahim N, Rivera E, Theriault R, Booser D, Symmans FW, Wong F, Blumenschein G, and Fleming DR, et al. (2005) Phase II study of tariquidar, a selective P-glycoprotein inhibitor, in patients with chemotherapy-resistant, advanced breast carcinoma. *Cancer* **104**:682–691.
- Raviv Y, Pollard HB, Bruggemann EP, Pastan I, and Gottesman MM (1990) Photosensitized labeling of a functional multidrug transporter in living drug-resistant tumor cells. *J Biol Chem* **265**:3975–3980.
- Richert ND, Aldwin L, Nitecki D, Gottesman MM, and Pastan I (1988) Stability and covalent modification of P-glycoprotein in multidrug-resistant KB cells. *Biochemistry* **27**:7607–7613.
- Riezman H, Woodman PG, van Meer G, and Marsh M (1997) Molecular mechanisms of endocytosis. *Cell* **91**:731–738.
- Righetti PG, Castagna A, Antonoli P, Cecconi D, Campostrini N, and Righetti SC (2005) Proteomic approaches for studying chemoresistance in cancer. *Expert Rev Proteomics* **2**:215–228.
- Robillard KR, Hoque T, and Bendayan R (2012) Expression of ATP-binding cassette membrane transporters in rodent and human sertoli cells: relevance to the permeability of antiretroviral therapy at the blood-testis barrier. *J Pharmacol Exp Ther* **340**:96–108.
- Roseblade A, Luk F, Rawling T, Ung A, Grau GE, and Bebawy M (2013) Cell-derived microparticles: new targets in the therapeutic management of disease. *J Pharm Pharm Sci* **16**:238–253.
- Sadasivan R, Morgan R, Fabian C, and Stephens R (1991) Reversal of multidrug resistance in HL-60 cells by verapamil and liposome-encapsulated doxorubicin. *Cancer Lett* **57**:165–171.
- Safa AR (1988) Photoaffinity labeling of the multidrug-resistance-related P-glycoprotein with photoactive analogs of verapamil. *Proc Natl Acad Sci USA* **85**:7187–7191.
- Sarko D, Beijer B, Garcia Boy R, Nothelfer EM, Leotta K, Eisenhut M, Altmann A, Haberkorn U, and Mier W (2010) The pharmacokinetics of cell-penetrating peptides. *Mol Pharm* **7**:2224–2231.
- Sato W, Yusa K, Naito M, and Tsuruo T (1990) Staurosporine, a potent inhibitor of C-kinase, enhances drug accumulation in multidrug-resistant cells. *Biochem Biophys Res Commun* **173**:1252–1257.
- Shaffer BC, Gillet JP, Patel C, Baer MR, Bates SE, and Gottesman MM (2012) Drug resistance: still a daunting challenge to the successful treatment of AML. *Drug Resist Updat* **15**:62–69.
- Shen D-Y, Zhang W, Zeng X, and Liu C-Q (2013) Inhibition of Wnt/β-catenin signaling downregulates P-glycoprotein and reverses multi-drug resistance of cholangiocarcinoma. *Cancer Sci* **104**:1303–1308.
- Shouzu A, Nomura S, Omoto S, Hayakawa T, Nishikawa M, and Iwasaka T (2004) Effect of ticlopidine on monocyte-derived microparticles and activated platelet markers in diabetes mellitus. *Clin Appl Thromb Hemost* **10**:167–173.
- Sladek NE, Low JE, and Landkammer GJ (1985) Collateral sensitivity to cross-linking agents exhibited by cultured L1210 cells resistant to oxazaphosphorines. *Cancer Res* **45**:625–629.
- Smith CD and Zilfou JT (1995) Circumvention of P-glycoprotein-mediated multiple drug resistance by phosphorylation modulators is independent of protein kinases. *J Biol Chem* **270**:28145–28152.
- Staats J, Marquardt D, and Center MS (1990) Characterization of a membrane-associated protein kinase of multidrug-resistant HL60 cells which phosphorylates P-glycoprotein. *J Biol Chem* **265**:4084–4090.
- Steinbach D and Legrand O (2007) ABC transporters and drug resistance in leukemia: was P-gp nothing but the first head of the Hydra? *Leukemia* **21**:1172–1176.
- Sui H, Fan ZZ, and Li Q (2012) Signal transduction pathways and transcriptional mechanisms of ABCB1/P-gp-mediated multiple drug resistance in human cancer cells. *J Int Med Res* **40**:426–435.
- Susa M, Iyer AK, Ryu K, Choy E, Hornicek FJ, Mankin H, Milane L, Amiji MM, and Duan Z (2010) Inhibition of ABCB1 (MDR1) expression by a siRNA nanoparticulate delivery system to overcome drug resistance in osteosarcoma. *PLoS ONE* **5**:e10764.
- Synold TW, Dussault I, and Forman BM (2001) The orphan nuclear receptor SXR coordinately regulates drug metabolism and efflux. *Nat Med* **7**:584–590.
- Szybalski W and Bryson V (1952) Genetic studies on microbial cross resistance to toxic agents. I. Cross resistance of *Escherichia coli* to fifteen antibiotics. *J Bacteriol* **64**:489–499.
- Tang T, Song X, Liu YF, and Wang WY (2013) PEITC reverse multi-drug resistance of human gastric cancer SGC7901/DDP cell line. *Cell Biol Int* DOI: 10.1002/cbin.10169.
- Thews O, Dillenburg W, Fellner M, Buchholz HG, Bausbacher N, Schreckenberger M, and Rösch F (2010) Activation of P-glycoprotein (Pgp)-mediated drug efflux by extracellular acidosis: in vivo imaging with 68Ga-labelled PET tracer. *Eur J Nucl Med Mol Imaging* **37**:1935–1942.
- Thierry AR, Jorgensen TJ, Forst D, Belli JA, Dritschilo A, and Rahman A (1989) Modulation of multidrug resistance in Chinese hamster cells by liposome-encapsulated doxorubicin. *Cancer Commun* **1**:311–316.
- Tsuruo T, Iida H, Nojiri M, Tsukagoshi S, and Sakurai Y (1983) Circumvention of vincristine and Adriamycin resistance in vitro and in vivo by calcium influx blockers. *Cancer Res* **43**:2905–2910.
- Tsuruo T, Iida H, Tsukagoshi S, and Sakurai Y (1982) Increased accumulation of vincristine and adriamycin in drug-resistant P388 tumor cells following incubation with calcium antagonists and calmodulin inhibitors. *Cancer Res* **42**:4730–4733.
- Wacher VJ, Wu CY, and Benet LZ (1995) Overlapping substrate specificities and tissue distribution of cytochrome P450 3A and P-glycoprotein: implications for drug delivery and activity in cancer chemotherapy. *Mol Carcinog* **13**:129–134.
- Wang L, Meng Q, Wang C, Liu Q, Peng J, Huo X, Sun H, Ma X, and Liu K (2013) Dioscin restores the activity of the anticancer agent adriamycin in multidrug-resistant human leukemia K562/adriamycin cells by down-regulating MDR1 via a mechanism involving NF-κB signaling inhibition. *J Nat Prod* **76**:909–914.
- Wang XK and Fu LW (2010) Interaction of tyrosine kinase inhibitors with the MDR- related ABC transporter proteins. *Curr Drug Metab* **11**:618–628.
- Wang Z, Chen Y, Liang H, Bender A, Glen RC, and Yan A (2011) P-glycoprotein substrate models using support vector machines based on a comprehensive data set. *J Chem Inf Model* **51**:1447–1456.
- Wartenberg M, Gronczynska S, Bekhite MM, Saric T, Niedermeier W, Hescheler J, and Sauer H (2005) Regulation of the multidrug resistance transporter P-glycoprotein in multicellular prostate tumor spheroids by hyperthermia and reactive oxygen species. *Int J Cancer* **113**:229–240.
- Wassmer SC, Cianciolo GJ, Combes V, and Grau GE (2005) Inhibition of endothelial activation: a new way to treat cerebral malaria? *PLoS Med* **2**:e245.
- Wattiaux R, Laurent N, Wattiaux-De Coninck S, and Jadot M (2000) Endosomes, lysosomes: their implication in gene transfer. *Adv Drug Deliv Rev* **41**:201–208.
- Weiss J and Haeefel WE (2013) Potential of the novel antiretroviral drug rilpivirine to modulate the expression and function of drug transporters and drug-metabolising enzymes in vitro. *Int J Antimicrob Agents* **41**:484–487.

- Whitnall M, Howard J, Ponka P, and Richardson DR (2006) A class of iron chelators with a wide spectrum of potent antitumor activity that overcomes resistance to chemotherapeutics. *Proc Natl Acad Sci USA* **103**:14901–14906.
- Xin Y, Yin F, Qi S, Shen L, Xu Y, Luo L, Lan L, and Yin Z (2013) Parthenolide reverses doxorubicin resistance in human lung carcinoma A549 cells by attenuating NF- κ B activation and HSP70 up-regulation. *Toxicol Lett* **221**:73–82.
- Xu C, Li CY, and Kong AN (2005) Induction of phase I, II and III drug metabolism/transport by xenobiotics. *Arch Pharm Res* **28**:249–268.
- Yang H, Lou C, Xu M, Wu C, Miyoshi H, and Liu Y (2011) Investigation of folate-conjugated fluorescent silica nanoparticles for targeting delivery to folate receptor-positive tumors and their internalization mechanism. *Int J Nanomedicine* **6**:2023–2032.
- Yang JM, Chin KV, and Hait WN (1996) Interaction of P-glycoprotein with protein kinase C in human multidrug resistant carcinoma cells. *Cancer Res* **56**:3490–3494.
- Yoshimura A, Kuwazuru Y, Sumizawa T, Ikeda S, Ichikawa M, Usagawa T, and Akiyama S (1989) Biosynthesis, processing and half-life of P-glycoprotein in a human multidrug-resistant KB cell. *Biochim Biophys Acta* **992**:307–314.
- Yu DK (1999) The contribution of P-glycoprotein to pharmacokinetic drug-drug interactions. *J Clin Pharmacol* **39**:1203–1211.
- Zhang W and Ling V (2000) Cell-cycle-dependent turnover of P-glycoprotein in multidrug-resistant cells. *J Cell Physiol* **184**:17–26.
- Zhao BX, Sun YB, Wang SQ, Duan L, Huo QL, Ren F, and Li GF (2013) Grape seed procyanidin reversal of p-glycoprotein associated multi-drug resistance via down-regulation of NF- κ B and MAPK/ERK mediated YB-1 activity in A2780/T cells. *PLoS ONE* **8**:e71071.
- Zhao Y, Liu H, Riker AI, Fodstad O, Ledoux SP, Wilson GL, and Tan M (2011) Emerging metabolic targets in cancer therapy. *Front Biosci (Landmark Ed)* **16**:1844–1860.

Address correspondence to: Richard Callaghan, Research School of Biology, Building 134, Linneaus Way, Australian National University, Acton, Canberra 0200, NSW, Australia. E-mail richard.callaghan@anu.edu.au