CORRESPONDENCE

The Malabsorption of Commonly Occurring Mono- and Disaccharides: Levels of Investigation and Differential Diagnoses

by Prof. Dr. med. Martin Raithel, Dr. med. Michael Weidenhiller, Dr. med. Alexander Hagel, Urban Hetterich, Prof. Dr. med. Markus F. Neurath, Prof. Dr. med. Peter C. Konturek in volume 46/2013

Correction Required

No one would doubt the fact that the number of patients seeking help for carbohydrate malabsorption problems has substantially risen over recent years. When asked, many patients express the opinion that they have an intolerance to fructose, whereas just a few questions regarding clinical symptoms will clarify that what actually affects them is a far more harmless fructose resorption disorder. The authors explicitly caution in their article (1) that “gastrointestinal malabsorption of fructose should not be confused with hereditary fructose intolerance.” But the term “gastrointestinal carbohydrate intolerance,” as used in the abstract, does not exactly help clarify matters.

The tables and figures show substantial gaps and errors when mentioning congenital defects in the monosaccharide metabolism, and these defects do not in any case relate symptomatically to malabsorptions in terms of the differential diagnosis. Although “galactosemia” refers to three congenital defects, galactokinase deficiency is the only one listed. This is a defect common in persons of Sinti/Romany origin and manifests as cataract development. The only embarrassing thing is that since 2002, all neonates in Germany are examined for another type of galactosemia: galactose 1 phosphate uridylyltransferase deficiency, the so called classic galactosemia. It would be highly desirable for this information, or a correction, to be added to this article post hoc.

DOI: 10.3238/arztebl.2014.0148a

REFERENCES

1. Raithel M, Weidenhiller M, Hagel AFK, Hetterich U, Neurath MFK, Konturek PC: The malabsorption of commonly occurring mono- and disaccharides—levels of investigation and differential diagnosis. *Dtsch Aerztebl Int* 2013; 110(46): 775–82.

Prof. Dr. med. Eberhard Mönch
Berlin
eberhard.moench@t-online.de

Conflict of interest statement

Professor Mönch has received honoraria for co-authoring a publication relating to the topic under discussion.

Restrictive Diets Are to Be Avoided

The prevalence of 30–40% reported in the article cannot be concluded from the references (1). This rate corresponds roughly to the rate of self-reported food in-

tolerances or the results of laboratory based chemical diagnostic evaluation (for example, H₂ breath test, lactase polymorphism).

Since diagnostic gold standards and non-selected patient cohorts are lacking, robust data on the prevalence rates of carbohydrate intolerance are not available.

In view of the physiological variance, unnecessary dietary restrictions on the basis of pathologically rated tolerance levels should be avoided (2). Even in 0% lactase activity, the ingestion of >12 g lactose (≈240 mL milk) may remain non-symptomatic; only a third of individuals who are symptomatic in the lactose stress test avoid milk in their daily diets.

Phylogenetically related restrictions of lactose and fructose intake are of teleological importance (for example, protection from obesity, metabolic syndrome) (3).

The non-negligible proportion of psychosomatic (functional) factors cannot be identified without using double-blinded, placebo controlled food stress testing. During such tests, the full clinical symptoms of carbohydrate malabsorption (meteorism, stomach pain, diarrhea) should be present, and the sustainability of the therapeutic effect must be monitored. (Physiological) “adult hypolactasia” can be excluded by molecular genetic testing.

In contrast to immunological (protein related) food intolerances (allergies, celiac disease), the extent of fructose and lactose restriction can be titrated at the individual level. When they are given, a sufficient intake of calcium (vitamin D, vitamins, minerals, and dietary fiber) should be ensured (4).

The aim remains to allow patients a wide range of natural foods (to prepare for themselves), rather than limiting them to restrictive diets and the associated consumption of industrially produced ready meals.

DOI: 10.3238/arztebl.2014.0148b

REFERENCES

1. Raithel M, Weidenhiller M, Hagel AFK, Hetterich U, Neurath MFK, Konturek PC: The malabsorption of commonly occurring mono- and disaccharides—levels of investigation and differential diagnosis. *Dtsch Aerztebl Int* 2013; 110(46): 775–82.
2. Gibson PR, Newnham E, Barrett JS, Shepherd SJ, Muir JG: Review article: fructose malabsorption and the bigger picture. *Alimentary pharmacology & therapeutics* 2007; 25: 349–63.
3. Disse SC, Buelow A, Boedeker RH, et al.: Reduced prevalence of obesity in children with primary fructose malabsorption: a multicentre, retrospective cohort study. *Pediatric obesity* 2013; 8: 255–8.
4. Zimmer KP: Laktose- und Fruktosemalabsorption. *Monatsschr Kinderheilkd* 2007; 155: 565–76.

Janna Riechmann

Dr. med. Jan de Laffolie

Prof. Dr. med. Klaus-Peter Zimmer
Klinik für Kinder- und Jugendmedizin,
Kindergastroenterologie

Universitätsklinikum Gießen
Janna.Riechmann@paediat.med.uni-giessen.de

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply:

We thank our correspondents for the interesting contributions in response to our article on carbohydrate malabsorption (1).

In actual fact, in routine clinical practice? the terms of hereditary fructose intolerance (metabolic defect) are often confused with fructose malabsorption. For this reason we separated these terms in our article, and gastrointestinal carbohydrate intolerance relates only to resorption disorders of the bowel.

The main focus of our article in particular was on carbohydrate mediated intolerance mechanisms. For this reason, hereditary causes need to be included in the differential diagnostic evaluation, especially as these disorders often manifest after ingestion of foodstuffs containing fructose or galactose and are then misinterpreted as allergies to milk/fruit or lactose maldigestion. We thank Professor Mönch for mentioning galactose 1 phosphate uridylyltransferase and the neonatal screening program introduced in 2002. Owing to space restrictions we listed in *Table 1* only one of the possible enzyme defects in galactosemia, as an example: the reaction catalyzed by galactokinase.

As Riechmann et al. say in their letter to the editor, standardized provocation tests for the normal population to confirm the diagnosis of carbohydrate intolerance and for reporting real prevalence rates are lacking. For this reason we included under “Definition of carbohydrate intolerance” the estimated data for the population of Europe (lactose 7–20%, fructose 15–25%, sorbite 8–20%); the variations show that substantial differences exist within Europe and that people of southern or Asian origins are far more likely to suffer from lactose maldigestion.

The “prevalence of 30–40%” cited by Riechmann and colleagues relates to all non-immunological food intolerances (carbohydrate intolerance, alcohol intolerance, fat intolerance syndrome, biogenic amines, salicylates, sulfites, etc), which cumulatively can account for 30–40% (1–3).

The gold standard for the diagnostic evaluation of carbohydrate intolerance is the oral provocation test in patients, which is best combined with an H₂ breath test. As we underlined, the mere rise in the hydrogen gas curve in asymptomatic patients is a mere sign of asymptomatic malabsorption, which by itself does not require any dietary restrictions. Overestimating the meaningfulness of the H₂ breath test alone would certainly incur a risk of panic (1, 2, 4).

We mentioned that patients with carbohydrate intolerance confirmed in such a way can tolerate a certain amount of carbohydrate up to the individual threshold level, so that individual dose adjustment is required. It

is therefore an essential therapeutic task to restore a dose reduction in carbohydrates into limited range by means of professional diet advice and training for patients.

Compared with the estimated prevalence rates of gastrointestinal malabsorption that we reported, the prevalence rates of carbohydrate intolerance are much higher in patients with functional symptoms, such as irritable bowel syndrome, and allergies (1, 4).

In this setting, the differential diagnostic evaluation should actively search for such intolerances (see also *Table 2* in [1]), because certain patient cohorts can receive crucial help thanks to this knowledge (1, 2, 4).

DOI: 10.3238/arztebl.2014.0149

REFERENCES

1. Raithel M, Weidenhiller M, Hagel AFK, Hetterich U, Neurath MFK, Konturek PC: The malabsorption of commonly occurring mono- and disaccharides—levels of investigation and differential diagnosis. *Dtsch Arztebl Int* 2013; 110(46): 775–82.
2. Stein J, Kist M, Raithel M: *Erkrankungen durch Nahrungsmittel (Food-borne diseases)* 1st edition. Stuttgart: Wissenschaftlich Verlagsgesellschaft 2011; 1–472.
3. Skypala I. Adverse food reactions – an emerging issue for adults. *J Am Diet Assoc* 2011; 111: 1877–91.
4. Halmos EP, Power VA, Shepherd SJ, Gibson PR, Muir JG: A diet low in FODMAPs reduces symptoms of Irritable Bowel Syndrome. *Gastroenterology* 2014; 146: 67–75.

On behalf of the authors

Prof. Dr. med. Peter Christopher Konturek

Klinik für Innere Medizin II, Thüringen Klinik Saalfeld
Rainweg 68, 07318 Saalfeld
pkonturek@thueringen-kliniken.de

Conflict of interest statement

The author declares that no conflict of interest exists.