CORRESPONDENCE

The Diagnosis and Treatment of Celiac Disease

by Prof. Dr. med. Dr. rer. nat. Detlef Schuppan,
Prof. Dr. med. Klaus-Peter Zimmer in volume 49/2013

Practicable Only to a Degree

The authors write that, according to the ESPEGHAN guidelines, the diagnosis can be confirmed without a biopsy in selected cases in patients with the “classic gastrointestinal manifestation of the disease” (1). The original article merely says: “Histological assessment may be omitted in symptomatic patients who have high IgA anti-TG2 levels (10 times above ULN), verified by EMA positivity, and are HLA-DQ2 and/or HLA-DQ8 heterodimer positive” (1). I cannot conclude any limitation to the “classic gastrointestinal manifestation” from the original article.

Furthermore, Schuppan and Zimmer say that “testing of first-degree relatives for the HLA-DQ2/8 genotype or for anti-TG2 antibodies is recommended.” If the ESPEGHAN criteria for the screening of at-risk patients were implemented in the way described in the article, then all first-degree relatives of patients with celiac disease would have to present to a human geneticist for diagnostic evaluation. The examination of asymptomatic relatives for HLA DQ2/8 is a predictive genetic examination; §7 (1) of the German Law on Genetic Diagnosis [Gendiagnostikgesetz] stipulates that a human geneticist is the only proper specialist to undertake this test, as far as I understand it. Whether this is really practical in view of a prevalence of celiac disease of 0.2% seems questionable to me.

DOI: 10.3238/arztebl.2014.0212a

REFERENCES

1. Schuppan D, Zimmer KP: The diagnosis and treatment of celiac disease. *Dtsch Arztebl Int* 2013; 110(49): 835–46.

Dr. med. Stefan Razeghi
Pediatrics and Youth Medicine
stefanrazeghi@yahoo.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Osteomalacia Rather Than Osteoporosis

The complications of classic celiac disease listed on pages 839 and 843 of the article include osteoporosis (1). This is correct only if it refers to the symptom of a reduction in bone minerals confirmed by densitometry, not the symptoms of osteoporosis. In adult medicine, this differentiation is important (2–4).

In my experiences in bone histology, active, untreated celiac disease leads to clear vitamin D and cal-

cium deficiencies, as a result of malabsorption. According to the extent of this deficiency, what follows is a bone mineralization disorder in the sense of volume osteoidosis. However, this is not associated with a volume deficit of the entire bone tissue. Osteoidosis alone usually heals without causing further complications, in tandem with the celiac disease.

Patients with severe, long-term calcium deficiency will experience, in addition to osteoidosis, reactive fibro-osteoclasia as a feature of secondary hyperparathyroidism. Both bone changes lead to the clinical picture of mixed intestinal osteopathy.

This can ultimately result in sustained volume deficits of the bones. In older patients with mixed intestinal osteopathy who have been treated for celiac disease and in whom the celiac disease has become inactive, a structural deficit corresponding to osteoporosis can be interpreted as “residual bone damage” if mineralization levels are normal.

Osteoporosis is a pathogenetically different disease entity and can be linked to celiac disease only in terms of a differential diagnosis.

Osteoporosis and osteomalacia should always be distinguished from one another, since they require completely different therapeutic strategies.

For this reason, the term that should be used in connection with the complications of active celiac disease is that of osteomalacia, which is pathogenetically appropriate, and not osteoporosis.

DOI: 10.3238/arztebl.2014.0212b

REFERENCES

1. Schuppan D, Zimmer KP: The diagnosis and treatment of celiac disease. *Dtsch Arztebl Int* 2013; 110(49): 835–46.
2. Abendroth K: Krankheiten des Knochens. In: Stobbe H, Baumann G (eds.): *Innere Medizin – Grundlagen und Klinik innerer Krankheiten*. 7th completely revised and extended edition. Berlin, Wiesbaden: Ullstein Mosby 1996; 1101–31.
3. Holick MF: Vitamin D: Photobiology, metabolism of action and clinical applications hypocalcemic disorders. In: *Primer on the metabolic bone diseases and disorders of mineral metabolism*, Official Publication of the American Society for Bone and Mineral Research, 4th edition, Philadelphia, Baltimore, New York: Lippincott Williams & Wilkins 1999; 92–97.
4. Braun J, Felsenberg D, Minne H: Veränderungen der Knochendichte und Knochenstruktur In: Zeidler H, Zacher J, Hiepe F (eds.): *Interdisziplinäre klinische Rheumatologie*. 1st edition, Berlin, Heidelberg, New York: Springer 2001; 1148–50.

Doz. Dr. sc. med. Klaus Abendroth
Jena
klaus.abendroth@t-online.de

Conflict of interest statement

Lecturer Dr sc med Abendroth is a representative of the osteology academy of the DVO. In the context of this activity he has received consultancy fees for continuing medical educational events. Furthermore he is a member of the German Society for Osteology (*Deutsche Gesellschaft für Osteologie*) and the German association for regional expert panels for osteoporosis (*Verein Regionale Expertenkreise Osteoporose Deutschland e.V.*).

Necessary Additional Points

Studies have confirmed an increased prevalence for developing microscopic colitis in patients with sprue, and vice versa (1–3). For collagenous colitis, the reported prevalence of sprue is 3.46% (1). In patients with confirmed sprue who continue to have symptoms in spite of adherence to their restriction diet, collagenous colitis and lymphocytic colitis should always be excluded.

DOI: 10.3238/arztebl.2014.0213a

REFERENCES

1. Green PH, Yang J, Cheng J, Lee AR, Harper JW, Bhagat G: An association between microscopic colitis and celiac disease. *Clin Gastroenterol Hepatol* 2009; 7: 1210–6.
2. Kao KT, Pedraza BA, McClune AC, et al.: Microscopic colitis: a large retrospective analysis from a health maintenance organization experience. *World J Gastroenterol* 2009; 15: 3122–7.
3. Stewart M, Andrews CN, Urbanski S, Beck PL, Storr M: The association of coeliac disease and microscopic colitis: a large population-based study. *Aliment Pharmacol Ther* 2011; 33: 1340–9.
4. Schuppan D, Zimmer KP: The diagnosis and treatment of celiac disease. *Dtsch Arztebl Int* 2013; 110(49): 835–46.

Prof. Dr. med. Andreas Tromm
Ev. Krankenhaus Hattingen gGmbH
a.tromm@krankenhaus-hattingen.de

Conflict of interest statement

Prof Tromm has been reimbursed for conference delegate fees, travel costs, and hotel expenses. He has also received honoraria from the Falk Foundation for preparing continuing medical educational events.

Two-step approach

The authors reference as the basis for their summary the different evidence based guidelines that have been developed over recent years, and a literature search, whose search strategy included, among others, all publications from the past 10 years identified by using the search terms “celiac disease” and “diagnosis” (1). Unfortunately they did not include our article that was published in 2013, even though this article was based on long years of experience and aimed to reach a definitive diagnosis using a minimum number of biopsies (2).

The best diagnostic test is that which results in the fewest false-positive and false-negative diagnoses; for this reason we’d suggest the following approach:

- The first step: simultaneous measuring of IgA and IgG antibodies specific for deamidated gliadin peptides, IgA antibodies specific for human tissue transglutaminase (in addition, total IgA). Most patients will either have a positive reaction to all three tested antigens or will test negative to all three of the specific antibody tests. In both these groups, biopsy is therefore unnecessary, since the positive predictive value (ppv) is 99% and the positive likelihood ratio (lr+) 87, whereas the negative predictive value (npv) is 98% and the negative likelihood ratio (lr-) 0.01. The results become even more meaningful (ppv 99%, lr+ 86; npv 100%, lr- 0.00) (2) if a fourth test is done for IgA endomysial-specific antibodies (2).
- The second step is small bowel biopsy. It is necessary only in patients with contradictory antibody results—that is, in patients who were positive in one or two tests only. This “two-step approach” reduces the proportion of patients requiring a biopsy to one-fifth (3, 4).

DOI: 10.3238/arztebl.2014.0213b

REFERENCES

1. Schuppan D, Zimmer KP: The diagnosis and treatment of celiac disease. *Dtsch Arztebl Int* 2013; 110(49): 835–46.
2. Bürgin-Wolff A, Buser M, Hadziselimovic F: Intestinal biopsy is not always required to diagnose celiac disease: a retrospective analysis of combined antibody tests. *BMC Gastroenterology* 2013; 23: 13–9.
3. Hadziselimovic F, Bürgin-Wolff A: Celiac disease. *N Engl J Med* 2008; 358: 747.
4. Bürgin-Wolff A, Hadziselimovic F: Two-step approach for diagnosing celiac disease. *Clin Gastroenterol Hepatol* 2008; 6: 1173.

Dr. phil A. Bürgin-Wolff
Prof. Dr. med. Faruk Hadziselimovic
Institut für Zöliakie Diagnostik
Liestal
faruk@magnet.ch

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply

S Razeghi focuses on the importance of symptoms in the ESPEGHAN recommendation for the rare case in which a duodenal biopsy is not needed (in a child or adolescent with at least tenfold raised anti-TG2-IgA and EMA confirmation, positivity for HLA-DQ2 or DQ8). If this is applied in any symptom that raises suspicions of celiac disease (for example, abdominal pain), the expected result will be diagnostic overkill. The ESPAGHAN guidelines refer, among others, to the Dahlborn study (reference 27 in our article) (1), in which a highly significant or a significant difference in the TG2-IgA titers existed between children with severe malabsorption and mild symptoms. Vivas et al (2) also showed that children with celiac disease do not only have higher TG2-IgA than adults, but are also more likely to have “classic” symptoms (malabsorption, diarrhea, failure to thrive). The law on genetic testing was correctly cited, but gastroenterologists can also acquire a qualification for “specialty related human genetic counseling” and therefore be allowed to screen asymptomatic patients at risk of celiac disease using the HLA-DQ2/8 genetic test.

The suggestion by A Bürgin-Wolff and F Hadziselimovic—to determine anti-TG-IgA as well as IgA and IgG antibodies against deamidated gliadin peptides, was not supported in the article by Giersiepen et al. (reference e5 in our article), after evidence-based evaluation of 2510 studies of the diagnostic potential of celiac serology testing. Current (prospective) studies are investigating this diagnostic approach. With all due respect for non-invasive diagnostic tests, and in the absence of a diagnostic gold standard for celiac disease, thorough histology of representative duodenal biopsy specimens according to March cannot be omitted, especially as the complication rate of diagnostic gastro-duodenoscopy is near 0%.

We thank K Abendroth for pointing out that malabsorption of vitamin D and calcium in active celiac disease results in osteomalacia, not osteoporosis. This may be the case where a sole mineralization disorder is suspected. In celiac disease and other inflammatory bowel disorders, however, bone formation is impaired in general, among other reasons due to increased breakdown of collagen type I. This is partly explained by the release and activity of proinflammatory cytokines, such as interleukin-1 α and TNF α in the context of intestinal inflammation (3). Accordingly, an increased fracture rate has been observed in

long-term untreated celiac disease in children, but especially in adults. This risk of fracture does not, as a rule, disappear after mere calcium and vitamin D substitution (4).

K Tromm mentions an important differential diagnosis for celiac disease, especially in adults: microscopic colitis, which is present as a comorbidity in 3–4% of celiac patients. In 6% of patients with “refractory celiac disease,” microscopic colitis was identified as the cause (5). We did not include this differential diagnosis in the table because it does not affect the small bowel. We concede, however, that it needs to be considered in patients whose celiac disease is in remission but who continue to suffer from diarrhea.

DOI: 10.3238/arztebl.2014.0213c

REFERENCES

- Schuppan D, Zimmer KP: The diagnosis and treatment of celiac disease. *Dtsch Arztebl Int* 2013; 110(49): 835–46.
- Vivas S, Ruiz de Morales JG, Riestra S, et al.: Duodenal biopsy may be avoided when high transglutaminase antibody titers are present. *WJG* 2009; 15: 4775–80.
- Bianchi ML, Bardella MT: Bone in celiac disease. *Osteoporos Int* 2008; 19: 1705–16.
- Tilg H, Moschen AR, Kaser A, Pines A, Dotan I: Gut, inflammation and osteoporosis: basic and clinical concepts. *Gut* 2008; 57: 684–94.
- Leffler DA, Dennis M, Hyett B, Kelly E, Schuppan D, Kelly CP: Etiologies and predictors of diagnosis in nonresponsive celiac disease. *Clin Gastroenterol Hepatol* 2007; 5: 445–50.

Prof. Dr. med. Klaus-Peter Zimmer

Department of General Pediatrics and Neonatology
Center for Pediatric and Adolescent Medicine
University Hospital Gießen and Marburg GmbH
Justus Liebig University, Gießen
klaus-peter.zimmer@paediat.med.uni-giessen.de

Prof. Dr. med. Dr. rer. nat. Detlef Schuppan

Institute of Translational Immunology and Department of Medicine I,
Johannes Gutenberg University, Mainz

Conflict of interest statement

Prof. Schuppan holds a patent for an anti-TG2 test and receives license fees for its use. He has received reimbursement of conference participation fees and of travel and accommodation expenses, and has been paid for the preparation of continuing medical education events, by the Schär, Merckle Recor-dati, and Instrumentation Laboratory companies.

Prof. Zimmer states that he has no conflict of interest.