Published in final edited form as:

Am J Addict. 2014 May ; 23(3): 226–233. doi:10.1111/j.1521-0391.2014.12098.x.

Predictors of drinking and functional outcomes for men and women following inpatient alcohol treatment

Dawn E. Sugarman, Ph.D.^{1,2}, Julia S. Kaufman, B.A.^{1,3}, Elisa M. Trucco, Ph.D.⁴, Jodi C. Brown, Ph.D., MSW⁵, and Shelly F. Greenfield, M.D., M.P.H.^{1,2}

¹Alcohol and Drug Abuse Treatment Program, McLean Hospital, 115 Mill Street, Belmont, MA

²Department of Psychiatry, Harvard Medical School, 401 Park Drive, Boston, MA

³Department of Psychology, Miami University, 100 Psychology Building, Oxford, OH

⁴Department of Psychiatry, Addiction Research Center, University of Michigan, 4250 Plymouth Road, Ann Arbor, MI

⁵Department of Social Work, College of Social and Behavioral Sciences, California State University, Northridge, 18111 Nordhoff Street, Northridge, CA

Abstract

Background and Objectives—This prospective study uses path analytic models to examine baseline characteristics associated with both functioning and drinking outcomes 12 months after inpatient alcohol treatment.

Methods—Alcohol dependent participants ($N = 101$) were recruited during inpatient alcohol treatment and assessed monthly one year after discharge.

Results—Alcohol severity was negatively associated with education and self-efficacy; marital status was positively associated with self-efficacy; and education and self-efficacy were negatively associated with drinking outcomes. Low alcohol severity, not having a depression diagnosis, and being married were associated with less social support impairment, which was in turn associated with better drinking outcomes. Having a history of sexual abuse did not influence drinking outcomes. However, having a history of sexual abuse was negatively associated with global functioning.

Conclusions and Scientific Significance—Drinking outcomes were associated with education, self-efficacy, social support, and diagnosis of depression at baseline; however, global functioning one year following treatment was primarily and negatively associated with sexual abuse history. Future treatment research should include measures of both functioning and drinking behavior outcomes.

Corresponding Author: Shelly F. Greenfield, M.D., M.P.H., McLean Hospital, Alcohol and Drug Abuse Treatment Program, 115 Mill St., Belmont, MA 02478, USA. 617-855-2241 (phone), 617-855-2699 (fax), sgreenfield@mclean.harvard.edu.

Declaration of Interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of this paper.

Keywords

Path Analysis; Gender; Alcohol Treatment; Functioning

BACKGROUND AND OBJECTIVES

Understanding predictors of alcohol treatment outcomes can help clinicians identify specific patient groups at greatest risk for poor outcomes, determine important targets for treatment success, and improve the accuracy of prognosis.¹ A comprehensive understanding of factors that contribute to prognosis may allow for gains in individualized treatment planning, thus reducing relapse rates.^{1,2}

Individual factors are among the most widely studied predictors of alcohol treatment outcomes (i.e., gender, education, marital status),³ but findings examining background characteristics are not always consistent. Research is mixed regarding gender differences,¹ with some studies suggesting minimal gender differences in alcohol treatment outcomes,⁴ and others suggesting women are more vulnerable to relapse.^{5,6} Studies diverge regarding the predictive power of marital status and education. Some findings suggest that marital status and education are protective factors associated with significant long-term improvements,⁷ whereas others conclude that these factors are poor predictors of outcomes.¹

Research has also examined individual clinical factors and their relationship to alcohol treatment outcomes. Co-occurring depression has been associated with poorer treatment outcomes, such as decreased abstinence rates and increased frequency of drinking following treatment.⁸ Another clinical factor, sexual abuse history, has received less attention despite evidence of high rates of trauma among alcohol dependent men and women.^{1,9} Recently, there has been evidence that individuals with a history of physical and sexual abuse demonstrate increased alcohol use following treatment compared to those not reporting an abuse history.¹⁰

In the past decade, alcohol-related measures have been examined more extensively and there is greater consistency across studies in their relationships to treatment outcomes. Alcohol-related self-efficacy, drinking patterns, and severity of alcohol use are considered the most consistent predictors of treatment outcomes.¹ Higher rates of alcohol severity at baseline significantly discriminate between individuals with positive and negative short- and long-term outcomes.^{3,11,12} Similarly, abstinence self-efficacy predicts better short- and long-term alcohol use outcomes.^{13,14}

Furthermore, a convergence of evidence from alcohol treatment trials demonstrates that treatment outcome is likely “multicausal” and is the effect of complex interactions among psychological, biological, and environmental factors.¹⁵ The lack of consistent findings across demographic and clinical characteristics may be partially attributed to a failure to account for other important variables in the same model.¹ Therefore, it is important to view treatment in context, examining specific treatment variables as well as demographic and clinical characteristics.^{3,15} Multivariate models may help provide information about the myriad factors affecting relapse and treatment outcomes.¹⁶

Alcohol treatment outcomes are typically assessed as changes in the quantity, frequency, and adverse consequences of use. Yet, alcohol use is also associated with adverse effects on functioning in multiple domains including health, psychological function, and interpersonal relationships.¹⁷ Thus, quantity-frequency measures of alcohol use may not adequately capture the complexity of treatment outcomes. Recently, the National Institute on Drug Abuse (NIDA) convened a panel of substance abuse treatment experts who recommended extending treatment outcomes beyond quantity-frequency measures of use to include psychosocial functioning.¹⁷ Few studies have evaluated predictors of alcohol treatment response with respect to both drinking and functional outcomes.¹⁸

In order to address this gap in the literature, a secondary analysis of a larger study^{10,19-21} was undertaken to examine prospectively the relationship of clinical and demographic characteristics on drinking and functional outcomes during the first year following inpatient treatment for alcohol dependence. We used path analysis with a mixed-gender sample to determine the relationship of several baseline characteristics to outcomes twelve months post-treatment.

Based on the literature, we chose to include demographic and clinical characteristics that have been associated with treatment outcome including gender,^{6,22} marital status,^{23,24} educational attainment,^{20,25} social support,⁷ abstinence self-efficacy,^{26,27} co-occurring major depressive disorder,^{25,28} alcohol severity,^{3,29} and history of sexual abuse.^{30,31} While many studies have examined these individual predictors, few have evaluated how they relate to one another in the context of treatment and their association with post-treatment functioning. This study adds to the literature by examining these predictors in a multivariate framework with multiple outcomes.

METHODS

Participants

A full description of the methods and demographic characteristics for this sample was previously published.^{19,21} Participants ($N = 101$) were recruited consecutively from the inpatient unit of McLean Hospital's Alcohol and Drug Abuse Treatment Program. Participants 18 years or older were eligible if they met criteria for alcohol dependence as determined by the Structured Clinical Interview for DSM-III-R (SCID).³² SCIDs were administered by a clinically-trained, master's level social worker, trained by the Principal Investigator (PI; SFG). All SCID diagnoses were reviewed by the PI. Exclusion criteria included an inability to complete follow-up visits and/or cognitive impairment that interfered with participants' adherence to the study protocol.

The majority of participants were White (94.0%) and male (59.4%) with some college education (79.2%). On average, participants were 43 years old ($SD = 11.1$). Forty-one percent of participants were married and a majority were either employed full-time (41.4%) or disabled or retired (25.3%). Women had significantly more years of education than men, and men were more likely to be employed full-time.²¹ There were no other significant gender differences in sociodemographic characteristics. Over half (57.4%) of the participants met criteria for a psychiatric disorder.²¹ The most prevalent comorbid disorders

were major depression (37.6%) and anxiety disorders (24.8%).²¹ Forty percent of participants reported a history of sexual abuse and 56% reported a history of physical abuse, with 28% reporting a history of both.¹⁰

Procedures

The study was approved by McLean Hospital's Institutional Review Board. All participants provided written informed consent. Participants were screened and recruited in the hospital after undergoing detoxification. During their hospital stay, participants received standard care, consisting of approximately 6-8 treatment groups per day and brief daily meetings with a psychiatrist, case manager, and counseling staff. Participants' average length of stay in the hospital ranged from 4-28 days ($M = 13$, $SD = 5.5$). Interview sessions were conducted when detoxification was completed. Participants completed structured interviews and self-report questionnaires at baseline and monthly for 12 months following discharge.

Measures

Alcohol use—The Timeline follow-back assessment method (TLFB)³³ assessed mean number of drinks per drinking day in the 12-month follow-up period. Breath alcohol tests and collateral informant reports were collected monthly to validate participants' self-report. The high validity of participants' self-reports of drinking during the follow-up period has been demonstrated and previously published.²¹

Educational attainment—The Drug and Alcohol Use Questionnaire assessed educational attainment at baseline.³⁴ Participants' responses were categorized to indicate their highest level of education: "Didn't graduate from high school," "Graduated from high school," "Some college," "Graduated from college," and "Postgraduate."

Self-efficacy—Abstinence self-efficacy was measured at baseline using the Situational Confidence Questionnaire (SCQ).³⁵ The SCQ is a 39-item self-report questionnaire in which participants indicate on a six-point scale how confident they feel about not drinking heavily in specified high-risk situations. Higher scores reflect a higher degree of confidence in one's ability to resist the urge to drink.

Social support impairment—The 'social and leisure' subscale of the Social Adjustment Scale Self-Report³⁶ measured social support at baseline and month 12. The full 54-item self-report scale measures role performance over the past two weeks within six areas: work, social and leisure, extended family, marital relationship, parental relationship(s), and family unit. The 'social and leisure' scale contains 11 items rated on a five-point scale with higher scores indicating greater impairment in social support. For the purpose of these analyses, we used the mean score of the 'social and leisure' subscale.

Functioning—Functioning at baseline and month 12 was measured using the Global Assessment of Functioning (GAF) Scale.³⁷ The GAF is an interval level measure ranging from 1 to 100, with higher numbers indicating better functioning. Test-retest reliability coefficients of the GAF range from .61 to .91.³⁷⁻³⁹ GAF scores were rated by the master's level social worker who administered the SCID. As there was only one rater, we were not

able to assess inter-rater reliability. However, the social worker was trained on scoring the GAF, had a background in administering SCIDs for research purposes, and all scores were reviewed with the PI. Furthermore, there is evidence that GAF scores used in research are more reliable than in clinical settings.⁴⁰ Thus, we feel confident in these ratings given the rigorous training and supervision.

Alcohol severity—Alcohol severity was measured using the Alcohol Dependence Scale (ADS) at baseline.⁴¹ The ADS is a 25-item self-report instrument designed to measure the degree of severity of alcohol dependence. A higher score on the ADS indicates greater severity of alcohol dependence.

Other demographic and clinical variables—Marital status and gender were assessed with a demographic questionnaire. History of sexual abuse was derived from the Life Experiences Questionnaire (LEQ) completed at baseline.⁴² The LEQ is a 29-item self-report measure of lifetime traumatic events. Current major depression diagnosis at baseline was assessed with the SCID.³²

Attrition

Participants who missed a monthly assessment were contacted and encouraged to return for the next assessment. Seventy-five participants completed some or all of the assessments and 26 participants dropped out of the study during the 12-month follow-up period. Forty-six participants completed all 12 assessments. Seventy-two participants completed month 12 assessments in person, and 3 participants completed month 12 assessments over the phone. Comparison analyses of completers versus dropouts were performed on all variables. The only statistically significant difference was marital status ($\chi^2 = 6.07$, $df = 1$, $p < .05$), with non-married participants more likely to drop out of the study.

Missing cases were excluded listwise. Mean number of drinks per drinking day was calculated for all 101 participants based on the number of days each participant completed data. At baseline, one participant was missing data for sexual abuse history and another for self-efficacy. In regard to GAF scores, 72 participants completed the month 12 assessment in person and were given GAF scores. Therefore, the model for drinking includes 99 participants and the model for functioning includes 70 participants.

Initial Model—We posited that demographic and clinical factors (marital status, gender, alcohol severity, education level, depression diagnosis, and sexual abuse history) would affect social support and efficacy. In turn, social support and efficacy were expected to be associated with drinking outcomes (see Figure 1). Given that alcohol use is associated with adverse effects on functioning¹⁷ and previous research shows associations between drinking reductions and functioning,¹⁸ we hypothesized a second parallel model with the same demographic, clinical factors, and exogenous variables, but with global functioning as the outcome.

Data Analyses—Path analysis was performed to test the theoretical models presented in Figures 1, 2 and 3. Due to the small sample size, the results of the path analysis should be considered exploratory. All analyses were conducted using the SAS System's CALIS

procedure.⁴³ These analyses used maximum likelihood estimation, and were performed on the variance-covariance matrix.

Marital status, gender, depression, and history of sexual abuse were included in the models as exogenous variables. Marital status, history of sexual abuse, depression diagnosis, and gender are dichotomous variables and were coded as “1” for: married, a history of sexual abuse, a baseline depression diagnosis, and female. The functioning model also includes the baseline GAF score as an exogenous variable.

RESULTS

Preliminary assumption testing was conducted to check for normality, linearity, multivariate outliers, and multicollinearity, with no serious violations noted. Means, standard deviations, and intercorrelations for study variables are presented in Table 1. To account for positive skewness in the drinks per drinking day variable, a log transformation was performed ($M = .66$, $SD = .54$). However, log transforming the variable and re-running the analyses did not meaningfully change results; therefore, the original, untransformed variable was used in all analyses.

Goodness of fit was assessed using multiple indices: the chi-square statistic, normed fit index (NFI),⁴⁴ non-normed fit index (NNI),⁴⁴ and comparative fit index (CFI).⁴⁵ The chi-square statistic provides a test of the null hypothesis that the reproduced covariance matrix has the specified model structure (i.e., that the models “fit” the data). Values on the NFI, NNFI, and CFI over 0.9 indicate an acceptable fit between model and data.⁴⁶

Drinking Model

We had posited that marital status, gender, alcohol severity, education level, depression diagnosis, and history of sexual abuse would be associated with social support and efficacy, which in turn would be associated with drinking outcomes. However, this initial model was a poor fit with the data, $\chi^2 (7, N = 99) = 25.60$, $p < .001$, $NFI = .86$, $NNI = .35$, $CFI = .87$. Previous independent results suggested a direct relationship between education level and drinking outcomes,²⁰ and LaGrange Multipliers suggested adding a path between education level and drinking outcomes to improve model fit. Positing a direct relationship between gender, alcohol severity, and education, the education level variable was shifted in the model to an endogenous position directly associated with drinking outcomes.

As seen in Figure 2, the model improves dramatically when education level is moved from an exogenous to endogenous position in the model. Additional minor adjustments to the initial model based on theory and fit indices included adding a path from baseline depression diagnosis and sexual abuse history to drinking outcome, as well as removing non-significant paths between gender and social support, and gender and self-efficacy. Using a conservative approach in modifying the model, all changes were made one at a time and the model re-estimated after each change.⁴⁶

The revised model (Figure 2) was a good fit with the data, $\chi^2 (13, N = 99) = 11.56$, $p = .56$, $NFI = .93$, $NNI = 1.02$, $CFI = 1.00$. The path model accounted for 34% of the variance in

mean number of drinks per drinking day. A history of sexual abuse, while independently related to drinking outcomes, was not statistically significant in the path model. The significant path coefficients reveal that individuals with lower education and self-efficacy, and higher social support scores (i.e., more social support impairment), had a higher mean number of drinks per drinking day in the 12-month follow-up period. Participants with depression at baseline had more social support impairment and a higher mean number of drinks per drinking day in the 12-month follow-up period. Married participants had higher self-efficacy and less social support impairment. Moreover, as seen in previous analyses using these data, women were more educated than men.²⁰ Lastly, participants with lower alcohol severity had a higher education level, higher self-efficacy, and less social support impairment.

Functioning Model

Performing a path analysis on the revised model using GAF scores as the functioning outcome at twelve months (and controlling for initial levels of functioning) revealed a good model fit and some interesting changes in the significant paths, $\chi^2 (15, N = 70) = 11.32, p = .72, NFI = .91, NNI = 1.11, CFI = 1.00$ (see Figure 3). Unlike the drinking model, marital status was not significantly related to any of the endogenous variables. Path coefficients for alcohol severity were of the same magnitude and direction in the functioning model as in the drinking outcomes model, as were the path coefficients between baseline depression diagnosis and social support. Baseline functioning was not related to education, self-efficacy, or social support, and there was no association between baseline functioning and 12-month functioning. Unlike the results for drinking outcome, education level, self-efficacy, and social support were not related to 12-month functioning. However, a history of sexual abuse was significantly and negatively associated with functioning.

DISCUSSION

The present study demonstrates that the individual characteristics associated with functioning and drinking outcomes twelve months following discharge from inpatient treatment are distinct. It is notable that sexual abuse history is the only baseline characteristic associated with functioning at twelve months while education, self-efficacy, and social support are significantly related to drinking outcomes. In this study, lower alcohol severity was associated with education level and self-efficacy, and these in turn were related to better drinking outcomes. The path analysis revealed that participants with lower educational attainment and lower self-efficacy had a higher mean number of drinks per drinking day in the year following treatment. Married participants had less impairment in social support, whereas participants with higher alcohol severity and a depression diagnosis had greater impairment in social support, which in turn predicted worse drinking outcomes. These results extend previous findings demonstrating a correlation between lower education levels and poorer drinking outcomes,^{20,47} and findings that depression^{21,48} and lower self-efficacy^{19,49} predict poorer drinking outcomes.

Importantly, sexual abuse history was not significantly associated with drinking at 12 months. These findings differ from those of Sacks and colleagues,³⁰ in which women with a

history of early trauma had worse substance use outcomes 12 months post-treatment; however, they defined early trauma as sexual, physical, or emotional abuse. In the present study, only a history of sexual abuse was examined. In previous analyses from this study,¹⁰ a history of sexual abuse (but not physical abuse) was associated with poorer drinking outcomes; however, this association was no longer significant when the analyses controlled for marital status, education, employment, and co-occurring psychiatric diagnoses. Our results highlight the importance of examining factors associated with treatment outcomes in multivariate models.

Global functioning at 12 months following discharge from inpatient alcohol treatment was not associated with the same predictors as drinking outcomes. Unlike results from the drinking model, education, self-efficacy, social support, and depression were not significantly associated with functioning outcomes. In direct contrast to the drinking model, sexual abuse was the *only* variable that had a significant effect on functioning outcomes. These results are consistent with findings from other studies that found lower functioning levels in abused than non-abused participants.^{30,31}

Our results suggest that a history of sexual abuse alone does not predict poorer drinking outcomes after treatment. However, it is possible that there are risk factors related to an abuse history that are associated with poorer drinking outcomes. For example, Rice and colleagues³¹ found that physically, sexually, or emotionally abused participants were less likely to marry than those without an abuse history. We found that marriage was significantly correlated with lower levels of social support impairment and greater self-efficacy, which in turn were related to better drinking outcomes. When these variables are accounted for, sexual abuse history no longer affects drinking outcomes. Yet, our findings also suggest that sexual abuse may have persistent negative effects on global functioning. These results highlight the importance of considering functioning as well as alcohol use in post-treatment outcomes. Furthermore, diverse patient characteristics may pose different risks for drinking and functioning in the year following treatment.

Limitations

The present study has several limitations. Our results may not be generalizable due to demographic characteristics of our sample compared to other populations seeking treatment for alcohol dependence. Namely, our sample was primarily white, male, and well-educated. The sample size is also a limitation. A larger sample size may have yielded sufficient power to detect a significant effect between sexual abuse and drinking outcomes or stronger correlations with functioning. Given that results from path analyses with relatively small sample sizes are often unreliable, the sample size also limits interpretation of the results. Moreover, we cannot rule out the possibility that differences in sample size between the two models may also account for the different pattern of findings. Therefore, until validated in a larger sample, our results must be interpreted with caution.

As is common with other alcohol treatment outcome research,^{3,16,50-52} the time points for follow-up did not exceed one year post-treatment. Examining longer-term outcomes might provide a better understanding of predictive factors for relapse and recovery. We also did not assess types of sexual abuse or participants' age when the abuse occurred, which may

account for differences in results. Also, sexual abuse history was measured with self-report instruments, which may have a recall and willingness-to-disclose bias. Furthermore, individuals were selected for this study from an inpatient treatment setting. As seen in previous research,^{53,54} comparing this sample to individuals treated in outpatient treatment settings may yield very different results. It is also important to note that the use of listwise deletion may yield biased results unless data are missing completely at random. A final consideration is our decision to move the education variable to an endogenous position in the model. This may seem counterintuitive given that a third, unmeasured variable may better explain the relationship between marital status, drinking severity, gender, and education level; however, we followed model fit suggestions and previous research²⁰ to examine the potential path correlations in this exploratory analysis.

Scientific Significance and Future Directions

The results of this study support previous research demonstrating the necessity of assessment and individualized treatment plans when trying to achieve optimal results for patients with alcohol dependence. We highlight the need for research examining complex associations of predictive and demographic factors in the substance-using population. Future analyses examining the interaction between multiple predictive factors during treatment and follow-up may provide valuable information concerning recovery and relapse, as well as treatment responses among alcohol dependent patients. These results support the recent NIDA panel recommendations¹⁷ to extend treatment research beyond quantity and frequency outcome measures to include functional outcomes. We encourage research seeking to improve alcohol treatment outcomes and life-functioning outcomes across diverse populations by examining factors that directly and indirectly influence these outcomes.

Acknowledgments

This research was supported in part by grants AA 09881 (SFG) from the National Institute on Alcohol Abuse and Alcoholism, Bethesda, MD and grants K08 DA000407 (SFG) and K24DA019855 (SFG) from the National Institute on Drug Abuse, Bethesda, MD; and the Dr. Ralph and Marian C. Falk Medical Research Trust (SFG), Chicago, Ill. The authors would like to thank Sara Wigderson, BA for her help with manuscript preparation.

REFERENCES

1. Adamson SJ, Sellman JD, Frampton CMA. Patient predictors of alcohol treatment outcome: A systematic review. *J. Subst. Abuse Treat.* 2009; 36(1):75–86. [PubMed: 18657940]
2. Kadden, RM.; Skerker, PM. Treatment decision making and goal setting. In: McCrady, BS.; Epstein, EE., editors. *Addictions: A comprehensive guidebook*. Oxford University Press; New York, NY US: 1999. p. 216-231.
3. Connors G, Maisto S, Zywiak W. Understanding relapse in the broader context of post-treatment functioning. *Addiction*. 1996; 91(Suppl):S173–S189. [PubMed: 8997791]
4. Kampman K, Volpicelli J, Alterman A, Cornish J, O'Brien C. Amantadine in the treatment of cocaine-dependent patients with severe withdrawal symptoms. *The American Journal of Psychiatry*. 2000; 157(12):2052–2054. [PubMed: 11097979]
5. Bottlender M, Soyka M. Outpatient alcoholism treatment: Predictors of outcome after 3 years. *Drug Alcohol Depend.* 2005; 80(1):83–89. [PubMed: 15878808]
6. Grella CE, Scott CK, Foss MA, Joshi V, Hser Y-I. Gender differences in drug treatment outcomes among participants in the Chicago Target Cities Study. *Eval. Program Plann.* 2003; 26(3):297–310.

7. McAweeney MJ, Zucker RA, Fitzgerald HE, Puttler LI, Wong MM. Individual and Partner Predictors of Recovery from Alcohol-Use Disorder over a Nine-Year Interval: Findings from a Community Sample of Alcoholic Married Men. *J. Stud. Alcohol.* 2005; 66(2):220–228. [PubMed: 15957673]
8. Gamble SA, Conner KR, Talbot NL, Yu Q, Tu XM, Connors GJ. Effects of pretreatment and posttreatment depressive symptoms on alcohol consumption following treatment in Project MATCH. *Journal of Studies on Alcohol and Drugs.* 2010; 71(1):71–77. [PubMed: 20105416]
9. Boles SM, Joshi V, Grella C, Wellisch J. Childhood Sexual Abuse Patterns, Psychosocial Correlates, and Treatment Outcomes Among Adults in Drug Abuse Treatment. *Journal of Child Sexual Abuse: Research, Treatment, & Program Innovations for Victims, Survivors, & Offenders.* 2005; 14(1):39–55.
10. Greenfield SF, Kolodziej ME, Sugarman DE, et al. History of abuse and drinking outcomes following inpatient alcohol treatment: A prospective study. *Drug Alcohol Depend.* 2002; 67(3): 227–234. [PubMed: 12127193]
11. Hodgins DC, el-Guebaly N, Armstrong S, Dufour M. Implications of depression on outcome from alcohol dependence: A 3-year prospective follow-up. *Alcohol. Clin. Exp. Res.* 1999; 23(1):151–157. [PubMed: 10029217]
12. McLellan AT, Lewis DC, O'Brien CP, Kleber HD. Drug dependence, a chronic medical illness. *JAMA.* 2004; 284(13):1689–1695. [PubMed: 11015800]
13. Moos RH, Moos BS. Protective resources and long-term recovery from alcohol use disorders. *Drug Alcohol Depend.* 2007; 86(1):46–54. [PubMed: 16769181]
14. Witkiewitz K. Predictors of heavy drinking during and following treatment. *Psychology of Addictive Behaviors.* 2011; 25(3):426–438. [PubMed: 21480681]
15. Moos RH, Finney JW. The expanding scope of alcoholism treatment evaluation. *Am. Psychol.* 1983; 38(10):1036–1044. [PubMed: 6314859]
16. Maisto SA, Zywiak WH, Connors GJ. Course of functioning 1 year following admission for treatment of alcohol use disorders. *Addict. Behav.* 2006; 31(1):69–79. [PubMed: 15919159]
17. Tiffany ST, Friedman L, Greenfield SF, Hasin DS, Jackson R. Beyond drug use: a systemic consideration of other outcomes in evaluations of treatments for substance use disorders. *Addiction.* 2012; 107(4):709–718. [PubMed: 21981638]
18. Moos, RH.; Finney, JW.; Cronkite, RC. *Alcoholism Treatment: Context, Process, and Outcome.* Oxford University Press; New York: 1990.
19. Greenfield SF, Hufford MR, Vagge LM, Muenz LR, Costello ME, Weiss RD. The relationship of self-efficacy expectancies to relapse among alcohol dependent men and women: A prospective study. *J. Stud. Alcohol.* 2000; 61(2):345–351. [PubMed: 10757147]
20. Greenfield SF, Sugarman DE, Muenz LR, Patterson MD, He DY, Weiss RD. The relationship between educational attainment and relapse among alcohol-dependent men and women: A prospective study. *Alcohol. Clin. Exp. Res.* 2003; 27(8):1278–1285. [PubMed: 12966322]
21. Greenfield SF, Weiss RD, Muenz LR, et al. The effect of depression on return to drinking: a prospective study. *Arch. Gen. Psychiatry.* 1998; 55(3):259–265. [PubMed: 9510220]
22. Greenfield SF, Brooks AJ, Gordon SM, et al. Substance abuse treatment entry, retention, and outcome in women: a review of the literature. *Drug Alcohol Depend.* Jan 5; 2007 86(1):1–21. [PubMed: 16759822]
23. Wilsnack SC, Wilsnack RW. Epidemiology of women's drinking. *J. Subst. Abuse.* 1991; 3(2):133–157. [PubMed: 1821278]
24. Schneider KM, Kviz FJ, Isola ML, Filstead WJ. Evaluating multiple outcomes and gender differences in alcoholism treatment. *Addict. Behav.* 1995; 20(1):1–21. [PubMed: 7785474]
25. Dodge R, Sindelar J, Sinha R. The role of depression symptoms in predicting drug abstinence in outpatient substance abuse treatment. *J. Subst. Abuse Treat.* 2005; 28(2):189–196. [PubMed: 15780549]
26. Kadden RM, Litt MD. The role of self-efficacy in the treatment of substance use disorders. *Addict. Behav.* 2011; 36(12):1120–1126. [PubMed: 21849232]

27. Gullo MJ, Dawe S, Kambouropoulos N, Staiger PK, Jackson CJ. Alcohol expectancies and drinking refusal self-efficacy mediate the association of impulsivity with alcohol misuse. *Alcohol. Clin. Exp. Res.* 2010; 34(8):1386–1399. [PubMed: 20528818]
28. Najt P, Fusar-Poli P, Brambilla P. Co-occurring mental and substance abuse disorders: A review on the potential predictors and clinical outcomes. *Psychiatry Res.* 2011; 186(2-3):159–164. [PubMed: 20728943]
29. McLellan AT, Alterman AI, Metzger DS, et al. Similarity of outcome predictors across opiate, cocaine, and alcohol treatments: Role of treatment services. *J Consult Clin Psychol.* 1994; 62(6): 1141–1158. [PubMed: 7860812]
30. Sacks JY, McKendrick K, Banks S. The impact of early trauma and abuse on residential substance abuse treatment outcomes for women. *J. Subst. Abuse Treat.* 2008; 34(1):90–100. [PubMed: 17574799]
31. Rice C, Mohr CD, Del Boca FK, et al. Self-reports of physical, sexual and emotional abuse in an alcoholism treatment sample. *J. Stud. Alcohol.* Jan; 2001 62(1):114–123. [PubMed: 11271959]
32. Spitzer, RL.; Williams, JBW.; Gibbon, M.; First, MB. Structured Clinical Interview for DSM-III-R Axis II Disorders, (SCID-II). American Psychiatric Press, Inc.; Washington, D. C.: 1990.
33. Sobell, LC.; Sobell, MB.; Litten, RZ.; Allen, JP. Measuring Alcohol Consumption: Psychosocial and Biochemical Methods. Humana Press; 1992. Timeline follow-back: A technique for assessing self-reported alcohol consumption; p. 41-72.
34. Griffin ML, Weiss RD, Mirin SM, Lange U. A comparison of male and female cocaine abusers. *Arch. Gen. Psychiatry.* 1989; 46(2):122–126. [PubMed: 2913971]
35. Annis, H.; Davis, C. Assessment of expectancies. In: Donovan, D.; Marlatt, G., editors. *Assessment of Addictive Behaviors*. Guilford Press; New York: 1988. p. 84-111.
36. Weissman MM, Bothwell S. Assessment of social adjustment by patient self-report. *Arch. Gen. Psychiatry.* 1976; 33:1111–1115. [PubMed: 962494]
37. Spitzer, RGM.; Williams, J.; Endicott, J. Global Assessment of Functioning (GAF) Scale. In: Sederer, L.; Dickey, B., editors. *Outcomes Assessment in Clinical Practice*. Williams and Wilkins; Baltimore: 1996. p. 76-78.
38. Hall RC. Global assessment of functioning. A modified scale. *Psychosomatics.* May-Jun;1995 36(3):267–275. [PubMed: 7638314]
39. Jones SH, Thornicroft G, Coffey M, Dunn G. A brief mental health outcome scale-reliability and validity of the Global Assessment of Functioning (GAF). *Br. J. Psychiatry.* May; 1995 166(5): 654–659. [PubMed: 7620753]
40. Söderberg P, Tungström S, Armelius BÅ. Reliability of Global Assessment of Functioning Ratings Made by Clinical Psychiatric Staff. *Psychiatr. Serv.* 2005; 56(4):434–438. [PubMed: 15812093]
41. Skinner HA, Allen BA. Alcohol dependence syndrome: Measurement and validation. *J. Abnorm. Psychol.* 1982; 91(3):199–209. [PubMed: 7096790]
42. Bryer JB, Nelson BA, Miller JB. Childhood sexual and physical abuse as factors in adult psychiatric illness. *Am. J. Psychiatry.* 1987; 144(11):1426–1430. [PubMed: 3674223]
43. Version 6 of the SAS System [computer program]. Cary, NC: 2000.
44. Bentler PM, Bonett DG. Significance tests and goodness of fit in the analysis of covariance structures. *Psychol. Bull.* 1980; 88(3):588–606.
45. Bentler, PM. EQS Structural Equations Program Manual. BMDP Statistical Software; Los Angeles: 1989.
46. Hatcher, L.; Stepanski, EJ. A step-by-step approach to using the SAS system for univariate and multivariate statistics. SAS Institute; Cary, NC US: 1994.
47. van Oers JAM, Bongers IMB, Van de Goor LAM, Garretsen HFL. Alcohol consumption, alcohol-related problems, problem drinking, and socioeconomic status. *Alcohol Alcohol.* 1999; 34(1):78–88. [PubMed: 10075406]
48. Holahan CJ, Moos RH, Holahan CK, Cronkite RC, Randall PK. Drinking to cope, emotional distress and alcohol use and abuse: A ten-year model. *J. Stud. Alcohol.* 2001; 62(2):190–198. [PubMed: 11327185]

49. Vielva I, Iraugi I. Cognitive and behavioural factors as predictors of abstinence following treatment for alcohol dependence. *Addiction*. 2001; 96(2):297–303. [PubMed: 11182875]
50. Moggi F, Ouimette PC, Finney JW, Moos RH. Effectiveness of treatment for substance abuse and dependence for dual diagnosis patients: A model of treatment factors associated with one-year outcomes. *J. Stud. Alcohol*. 1999; 60(6):856–866. [PubMed: 10606499]
51. Baker KD, Lubman DI, Cosgrave EM, et al. Impact of co-occurring substance use on 6 month outcomes for young people seeking mental health treatment. *Aust. N. Z. J. Psychiatry*. 2007; 41(11):896–902. [PubMed: 17924242]
52. McKellar J, Wagner T, Harris A, Oehlert M, Buckley S, Moos R. One-year outcomes of telephone case monitoring for patients with substance use disorder. *Addict. Behav*. 2012
53. Finney JW, Hahn AC, Moos RH. The effectiveness of inpatient and outpatient treatment for alcohol abuse: The need to focus on mediators and moderators of setting effects. *Addiction*. 1996; 91(12):1773–1796. [PubMed: 8997760]
54. Pettinati HM, Pierce JD Jr, Belden PP, Meyers K. The relationship of Axis II personality disorders to other known predictors of addiction treatment outcome. *The American Journal on Addictions*. 1999; 8(2):136–147. [PubMed: 10365194]

Figure 1.

Initial Path Model

N.S. = non-significant; * $p < .05$; ** $p < .01$, *** $p < .001$

Figure 2.
 Path Analytic Model with Average Number of Drinks per Drinking Day as Outcome
 Standardized path coefficients appear on single-headed straight arrows. N.S. = non-significant; * $p < .05$; ** $p < .01$, *** $p < .001$

Figure 3.
 Path Analytic Model with Functioning as Outcome
 Standardized path coefficients appear on single-headed straight arrows. N.S. = non-significant; * $p < .05$; ** $p < .01$, *** $p < .001$

Table 1
Means, Standard Deviations, and Intercorrelations for the Variables in the Models

Variable	N	M	SD	1	2	3	4	5	6	7	8	9	10
1. Mean number of drinks per drinking day	101	7.45	8.11										
2. 12 Month Functioning	71	67.31	14.71	-.44									
3. Education Level	101	3.37	1.15	-.32	.24								
4. Self-Efficacy	100	65.80	25.65	-.34	.36	.13							
5. Social Support	101	2.65	.63	.37	-.39	-.14	-.30						
6. Marital Status	101	.39	.49	-.31	.33	.27	.29	-.27					
7. Gender	101	1.41	.49	-.16	-.07	.29	-.05	.02	.07				
8. Drinking Severity	101	22.46	8.94	.35	-.42	-.21	-.44	.42	-.21	.17			
9. Baseline Depression	101	.38	.48	.26	-.28	.09	-.15	.31	-.08	.14	.25		
10. History of Sexual Abuse	100	.40	.49	.20	-.43	.04	-.16	.15	-.10	.43	.40	.24	
11. Baseline Functioning	101	50.81	10.25	-.35	.25	.22	.28	-.22	.22	.00	-.43	-.16	-.23