CASE REPORT

Septic arthritis secondary to rat bite fever: a challenging diagnostic course

Basil Budair, Karan Goswami, Vivek Dhukaram

Department of Trauma & Orthopaedics, University Hospitals Coventry and Warwickshire NHS Trust, Coventry, West Midlands, UK

Correspondence to
Basil Budair,
b.budair@nhs.net

Accepted 5 March 2014

SUMMARY

We describe a challenging diagnostic course of a case of septic arthritis secondary to rat bite fever (*Streptobacillus moniliformis*) in a 29-year-old man who presented with generalised malaise, polyarthralgia and bilateral palmoplantar rash.

BACKGROUND

Rat bite fever (RBF) is a potentially fatal infection with a mortality rate of 13% when left untreated.¹ Although the differential diagnosis for a patient presenting with non-specific symptoms is broad, social history and other factors must be kept in mind when reaching a working diagnosis. Clinicians should keep RBF on their differential diagnosis when a patient present with classic symptoms of fever, rash and joint involvement, especially if there is a history of exposure to rats.²

CASE PRESENTATION

A 29-year-old man of Polish origin presented with a 5-day history of worsening generalised malaise, fever, sore throat and polyarthralgia. He also reported a bilateral plantopalmar rash. He had no significant medical history and did not take any prescribed medications. He used over-the-counter antihistamines for seasonal allergic rhinitis. He worked as a manual labourer in a warehouse and drank 10 units of alcohol per week. He was a current smoker with a 3-pack-year history. Of note, he had direct daily physical contact with multiple rats that he kept as pets.

On examination, he had low-grade fever. He had congested tonsils, but with no visible exudates. There was generalised decreased painful range of movement in the right second metacarpophalangeal joint, right elbow, right knee and both ankles, which also had effusions. On inspection, there was a widespread, partially blanching, rash with vesicular, petechial and pustular components over both feet, hands, legs and buttocks (figure 1). His heart sounds were normal. Chest was clear. Abdominal examination was unremarkable. There was no palpable lymphadenopathy or any stigmata of endocarditis.

INVESTIGATIONS

Investigations on admission revealed raised inflammatory markers with an initial C reactive protein of 211 mg/L (normal <5 mg/L), erythrocyte sedimentation rate 36 mm/h (normal <14 mm/h) and white cell count 10.62×10^9 cells/L (normal range 3– 10×10^9 cells/L). He had normal liver and renal

function tests. Chest radiograph showed clear lung fields, and urine dipstick was unremarkable.

A broad microbiological, immunological and rheumatological screening was suggested by the treating rheumatology team for a working diagnosis of vasculitis. This was unremarkable, with normal double-stranded DNA antibodies (Abs) (<1 kIU/L), glomerular basement membrane Abs (<1 kIU/L), myeloperoxidase Abs (<0.2 kIU/L), proteinase-3 Abs (<0.2 kIU/L) and normal rheumatoid factor (13 kIU/L). No paraproteinaemia was detected on electrophoresis. Immunoglobulin levels were also within normal ranges (IgG 10.2 g/L, IgA 2.39 g/L, IgM 0.75 g/L).

Blood cultures were negative and showed no growth. Hepatitis screen was negative (hepatitis B surface antigen, hepatitis C antibody, hepatitis A IgM antibody were not detected). HIV testing was negative. Anti-streptolysin O antibody was normal (<200 IU/mL). Leptospira serology ELISA IgM was negative. Parvovirus B19 screen was unremarkable. Throat swab showed a scant growth of haemolytic *Streptococcus* group B. Virology reverse transcription-PCR showed no evidence of recent infection with respiratory syncytial virus, parainfluenza virus, adenovirus, human metapneumovirus, enterovirus, influenza A or influenza B.

Following a dermatology review, an alternative differential diagnosis of viral exanthematous pustulosis was considered, and a punch biopsy of skin was obtained. This showed a mild focal, non-specific, perivascular inflammation within the subcutaneous fat of uncertain significance, and no features specific for vasculitis. A negative direct immunofluorescence study on the skin biopsy was also obtained for IgM, IgG, IgA, C3 and fibrinogen.

Initial right ankle joint aspiration revealed a cloudy yellow fluid, with no crystals or organisms. No growth on culture or acid-fast bacilli were seen.

Following further discussion with microbiology, and mention of this patient keeping several pet rats, specific testing for RBF was recommended. Accordingly, a serum sample taken for bacterial 16S rRNA PCR was negative. However, a repeat aspirate of the right ankle joint was positive for the 16S rRNA gene, which is diagnostic for *Streptobacillus moniliformis* infection.

DIFFERENTIAL DIAGNOSIS

- RBF
- Vasculitis
- Viral exanthematous pustulosis

CrossMark

To cite: Budair B, Goswami K, Dhukaram V. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-204086

Figure 1 Medical photograph images of palmer (A) and planter (B) rash.

TREATMENT

After confirming diagnosis, the empirical broad-spectrum antibiotics were changed to intravenous benzylpenicillin as per microbiology advice. Within 2 weeks, he showed significant clinical improvement and remained non-feverish. His rash settled and inflammatory markers normalised. Although his right metacarpophalangeal, elbow, knee and ankle joints symptoms completely resolved, his left ankle remained tender, swollen and painful on movement. This patient was also unable to weight bear on the left ankle, raising suspicion of septic arthritis.

Consequently, MRI of the left ankle was performed to exclude a septic joint and osteomyelitis. This showed a moderate, thick-walled joint effusion within the ankle mortice joint, with moderate reactive inflammatory change in the surrounding soft tissues, consistent with a diagnosis of septic arthritis. In addition, the imaging featured an incidental 14×5 mm osteochondral defect of the anteromedial talar dome (figure 2).

After review of imaging, the orthopaedic team was contacted. On examination of the left ankle by the orthopaedic registrar, it remained painful and mildly swollen. It was not tender or warm on palpation. He had 10° loss of dorsiflexion compared with the contralateral ankle, and was unable to weight bear due to pain. Initial plan after discussion with the oncall orthopaedic consultant was to continue with antibiotics and list the patient for an arthroscopic left ankle washout on the next trauma theatre list. Unfortunately, for logistical reasons, the procedure was cancelled and specialist advice from a foot and ankle

surgeon was sought. At his review, 48 h from the initial orthopaedic assessment, the patient's left ankle symptoms had significantly improved. On examination, the patient was now able to independently weight bear. His pain had improved. There was slightly reduced dorsiflexion when compared with the right ankle, particularly apparent on walking. In view of the dramatic improvement in clinical findings, an ankle washout was deemed unnecessary, and the orthopaedic team advocated adhering to microbiology advice regarding further antibiotics.

OUTCOME AND FOLLOW-UP

The patient was discharged with a 3-week course of oral amoxicillin and referred for outpatient physiotherapy. On follow-up in clinic 6 weeks later, his rash had receded to a few faint spots on his feet, and he had regained full range of movement of the ankle. There was minimal pain on impact activities.

DISCUSSION

RBF is a rare bacterial infection caused by *Streptobacillus moniliformis*, a Gram-negative bacillus which colonises the nasopharynx of numerous rodents, especially rats, and can also be excreted in urine. It classically presents as a relapsing-remitting systemic illness involving fever, malaise, rash and migratory polyarthralgias and is associated with numerous complications. When left untreated, it carries a substantial risk of mortality. Owing to the non-specific nature of the presenting symptoms, the disease has a high potential for misdiagnosis, especially when exposure to rodents is not obvious or known. Previous case reports and reviews of RBF have reported various serious complications, including fulminant sepsis, meningitis, hepatitis, nephritis, amnionitis, pneumonia, myocarditis, endocarditis and pericarditis. Case reports have also described pericardial effusion, septic arthritis, suppurative polyarthritis and systemic vasculitis. In our patient, *S moniliformis* was eventually isolated from the repeat right ankle synovial fluid aspirate via 16S rRNA PCR, which is consistent with previous reports of septic arthritis.^{2 3}

Our case demonstrates the diagnostic challenges of RBF and related septic arthritis. Synovial fluid analysis, culture and blood cultures remain critical in the diagnosis of septic arthritis and determination of the microbiological aetiology.⁴ However, 18% of cases of septic arthritis are culture-negative.^{5 6} In this setting, molecular techniques such as 16S rRNA PCR amplification using universal primers are an important tool, and successfully aided the identification of *S moniliformis* septic arthritis in our patient. PCR is an effective way of isolating *S moniliformis*

Figure 2 MRI of the left ankle showing changes consistent with septic arthritis (A) and the incidental osteochondral defect of the talar dome (B).

which is normally difficult to isolate with standard tissue cultures.⁷

The treatment of choice for proven or highly suspected cases of RBF is penicillin,² and successful treatment with a 4-week regimen has been demonstrated.⁸ However, early joint washout and removal of purulent material from the joint space is advocated in the context of septic arthritis by the British Orthopaedic Association.⁹ Delayed or inadequate treatment of septic arthritis can lead to irreversible joint damage, resulting in disability.¹⁰ Since this patient presented to the orthopaedic team with a delayed, 3-week history of polyarthralgia, and evidence of improvement on antibiotics, washout was deemed unnecessary. Ravindran *et al*¹¹ show that native joint septic arthritis can be managed medically with serial closed-needle joint aspiration with similar outcomes to surgical washout.

Learning points

- ▶ A comprehensive social history of possible exposure to rodents should be elicited from patients with septic arthritis in order to facilitate diagnosis.
- ▶ Molecular diagnostics may be useful in the diagnosis of culture-negative septic arthritis.
- ▶ A close liaison with relevant specialties is necessary to ensure prompt diagnosis and management of patients with polyarthralgia.
- ▶ The need for surgical washout is based on clinical presentation, progression of symptoms and response to antibiotics.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 CDC. Rat-bite fever—New Mexico, 1996. *MMWR* 1998;47:89–91.
- 2 Elliott SP. Rat bite fever and *Streptobacillus moniliformis*. *Clin Microbiol Rev* 2007;20:13–22.
- 3 Wang TK, Wong SS. *Streptobacillus moniliformis* septic arthritis: a clinical entity distinct from rat-bite fever? *BMC Infect Dis* 2007;7:56.
- 4 Mathews CJ, Weston VC, Jones A, *et al*. Bacterial septic arthritis in adults. *Lancet* 2010;375:846–55.
- 5 Eder L, Zisman D, Rozenbaum M, *et al*. Clinical features and aetiology of septic arthritis in northern Israel. *Rheumatology (Oxford)* 2005;44:1559–63.
- 6 Fenollar F, Roux V, Stein A, *et al*. Analysis of 525 samples to determine the usefulness of PCR amplification and sequencing of the 16S rRNA gene for diagnosis of bone and joint infections. *J Clin Microbiol* 2006;44:1018–28.
- 7 Boot R, Oosterhuis A, Thuis HC. PCR for the detection of *Streptobacillus moniliformis*. *Lab Animal* 2002;36:200–8.
- 8 Holroyd KJ, Reiner AP, Dick JD. *Streptobacillus moniliformis* polyarthritits mimicking rheumatoid arthritis: an urban case of rat bite fever. *AM J Med* 1988;85:711–14.
- 9 Coakley G, Mathews C, Field M, *et al*. BSR & BHPR, BOA, RCGP and BSAC guidelines for management of the hot swollen joint in adults. *Rheumatology (Oxford)* 2006;45:1039–41.
- 10 Kaandorp CJE, Krijnen P, Moens HJB, *et al*. The outcome of bacterial arthritis: a prospective community-based study. *Arthritis Rheum* 1997;40:884–92.
- 11 Ravindran V, Logan I, Bourke BE. Medical vs surgical treatment for the native joint in septic arthritis: a 6-year, single UK academic centre experience. *Rheumatology (Oxford)* 2009;48:1320–2.

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow