Pitfalls of single lung transplantation (SLT) for chronic obstructive pulmonary disease

Anju Singhal, Robert Michael Reed

Department of Pulmonary and Critical Care Medicine,
University of Maryland,
Baltimore, Maryland, USA

Correspondence to
Dr Robert Michael Reed,
rreed@medicine.umaryland.edu

Accepted 22 March 2014

DESCRIPTION

Figure 1 demonstrates two complications of single lung transplantation (SLT) in chronic obstructive pulmonary disease (COPD). First, the image shows hyperinflation of the native right lung causing compression of the graft lung. In one study, patients with SLT showed no evidence of flow limitation at rest, but the majority of patients did show evidence of dynamic hyperinflation during exercise. Unsurprisingly, patients who underwent bilateral lung transplantation (BLT) showed no evidence of flow limitation at rest or during exercise. This finding likely accounts for the known decreased exercise tolerance seen in SLT versus BLT patients.¹

The image also shows evidence of malignancy in the native lung (red arrows). These were biopsy-proven adenocarcinomas. There is increased incidence of lung cancer found in the native lung of patients with SLT. One review showed lung cancer incidence of up to 6.9% in the native lung after the transplant. There have been only occasional reports of lung cancer in donor lungs. The presence of a native lung is the strongest risk factor for developing lung cancer in post-transplant patients.²

Figure 1 CT image demonstrating malignancy and hyperinflation of the native (right) lung in a patient who had received a single lung transplantation for chronic obstructive pulmonary disease.

These are two examples of what may account for the worse survival noted in patients with COPD under 60 years who undergo SLT versus BLT. There is also a trend towards improved survival with BLT over the age of 60 years; however, no sufficiently powered studies in these patients have been completed to show significant mortality improvements.³

Learning points

- ▶ Patients who underwent bilateral lung transplantation (BLT) have improved exercise tolerance and pulmonary function tests compared with single lung transplantation (SLT), likely secondary to dynamic hyperinflation of the native lung.
- ▶ There is increased risk of lung cancer in the native lung in patients with SLT due to chronic obstructive pulmonary disease.
- ▶ BLT has shown improved median survival compared with SLT in patients under 60 years.

Contributors All authors contributed to the research, drafting and revision of this manuscript.

Competing interests None.

Patient consent None.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Murciano D, Ferretti A, Boczkowski J, *et al*. Flow limitation and dynamic hyperinflation during exercise in COPD patients after single lung transplantation. *Chest* 2000;118:1248–54.
- 2 Mathew J, Kratzke RA. Lung cancer and lung transplantation: a review. *J Thorac Oncol* 2009;4:753–60.
- 3 Thabut G, Christie JD, Ravaud P, *et al*. Survival after bilateral versus single lung transplantation for patients with chronic obstructive pulmonary disease: a retrospective analysis of registry data. *Lancet* 2008;371:744–51.

CrossMark

To cite: Singhal A, Reed RM. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-204355

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow