Published in final edited form as:

Neuropharmacology. 2014 April ; 79: 590–595. doi:10.1016/j.neuropharm.2014.01.002.

Levamisole and cocaine synergism: a prevalent adulterant enhances cocaine's action *in vivo*

Christopher S. Tallarida^{1,2}, Erin Egan^{1,2}, Gissel D. Alejo^{1,2}, Robert Raffa^{1,3}, Ronald J. Tallarida^{1,2}, and Scott M. Rawls^{1,2}

¹Center for Substance Abuse Research, Temple University School of Medicine, Philadelphia, PA

²Department of Pharmacology, Temple University School of Medicine, Philadelphia, PA

³Department of Pharmaceutical Sciences, Temple University School of Pharmacy, Philadelphia, PA

Abstract

Levamisole is estimated by the Drug Enforcement Agency (DEA) to be present in about 80% of cocaine seized in the United States and linked to debilitating, and sometimes fatal, immunologic effects in cocaine abusers. One explanation for the addition of levamisole to cocaine is that it increases the amount of product and enhances profits. An alternative possibility, and one investigated here, is that levamisole alters cocaine's action *in vivo*. We specifically investigated effects of levamisole on cocaine's stereotypical and place-conditioning effects in an established invertebrate (planarian) assay. Acute exposure to levamisole or cocaine produced concentration-dependent increases in stereotyped movements. For combined administration of the two agents, isobolographic analysis revealed that the observed stereotypical response was enhanced relative to the predicted effect, indicating synergism for the interaction. In conditioned place preference (CPP) experiments, cocaine produced a significant preference shift; in contrast, levamisole was ineffective at all concentrations tested. For combination experiments, a submaximal concentration of cocaine produced CPP that was enhanced by inactive concentrations of levamisole, indicating synergism. The present results provide the first experimental evidence that levamisole enhances cocaine's action *in vivo*. Most important is the identification of synergism for the levamisole/cocaine interaction, which now requires further study in mammals.

Keywords

levamisole; cocaine; addiction; conditioned place preference; stereotypy; synergy; planarians; psychostimulant

© 2014 Elsevier Ltd. All rights reserved.

Corresponding Author: Scott M. Rawls, Ph.D., Associate Professor of Pharmacology, Department of Pharmacology, Center for Substance Abuse Research, Temple University School of Medicine, 3500 North Broad Street, Philadelphia, PA, 19140, USA Tel: (215) 707-4942; Fax: (215) 707-7068; scott.rawls@temple.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

An old drug called levamisole (Ergamisol) that was once used to treat parasitic worm infections in humans is exacerbating health risks for an estimated 2 million cocaine users in the United States (Auffenberg et al., 2013). The news media (*e.g.* Time Magazine), scientific publications, and government agencies are alerting the general public, health officials, and physicians about potentially life-threatening effects of cocaine laced with levamisole (Zhu et al 2009; Chang et al 2010; Ullrich et al., 2011). An example of such a warning is the public alert issued in September of 2009 by the U.S. Department of Health and Human Services Substance Abuse and Mental Health Services Administration warning that “a dangerous substance, levamisole, is showing up with increasing frequency in illicit cocaine powder and crack cocaine and can lead to a severe reduction in the number of white blood cells, a problem that is called agranulocytosis”. The Drug Enforcement Agency (DEA) estimates that about 80% of the cocaine seized in the US is laced with levamisole (Wolford et al 2012). DEA data from 2009 also noted an average concentration of approximately 10% levamisole detected in cocaine, and Buchanan et al (2010) demonstrated the presence of levamisole (as high as 10%) in a patient's crack cocaine pipe, thus confirming levamisole as a cocaine adulterant. Speculation about the addition of LVM to cocaine centers on two hypotheses. One is that LVM increases the amount of ‘product’ which increases profits. LVM is cheap, has similar physicochemical properties to cocaine, and is easily accessible as a veterinary pharmaceutical in regions in which the laced cocaine originates.

A second hypothesis is that levamisole is added to cocaine to modify the pharmacological properties of cocaine. To probe the latter possibility, we used an established planarian assay to determine if levamisole affects cocaine's action *in vivo*. Planarians are aquatic flatworms with a centralized nervous system often considered to be the simplest ‘brain’ (Raffa and Rawls, 2008; Buttarelli et al., 2008). Planarians contain neurotransmitter systems, including glutamate, dopamine, serotonin, acetylcholine, and GABA (Eriksson and Panula, 1994; Vyas et al., 2010; Nishimura et al., 2010), and to a limited extent display mammalian-equivalent behavioral responses (stereotypical activity, abstinence-related withdrawal, behavioral sensitization, cross-sensitization, and place conditioning) following exposure to addictive substances (Palladini et al., 1996; Pagán et al., 2008, 2009, 2013; Rowlands and Pagán, 2008; Rawls et al., 2010, 2011; Ramoz et al., 2012). The present experiments characterized levamisole and cocaine interactions using two behavioral endpoints. One was stereotypical activity, defined as the number of C-shape movements across a defined time interval (Venturini et al., 1996; Passarelli et al., 1999; Rawls et al., 2011). The other was conditioned place preference (CPP) (Tzschentke, 1998), an assay in which planarians exposed to a distinct environment in the presence of a positively reinforcing substance will later show preference for that same environment when given a choice (Zhang et al., 2013; Ramoz et al., 2012). Drug combination analysis employing isobolographic theory was used to quantify levamisole-cocaine interactions (Tallarida, 2011, 2012). The isobolographic method is derived from the principle of dose equivalence and is the standard pharmacological approach to analyze observed combination dose effects for comparison with expected, or additive, effects. Its aim is to assess synergistic and/or additive interactions between compounds administered simultaneously (Tallarida, 2011, 2012).

2. Experimental Procedures

2.1. Subjects and drugs

Planarians (*Dugesia dorotocephala*) were purchased from Carolina Biological Supply (Burlington, NC, USA). Upon arrival in the laboratory, planarians were maintained in the aqueous solution provided by Carolina Biological Supply, acclimated to room temperature (21 °C), and tested within 3 days of receipt. (–)-Cocaine hydrochloride was generously provided by the National Institute on Drug Abuse (Bethesda, MD, USA). Levamisole hydrochloride was purchased from Sigma-Aldrich (St. Louis, MO, USA). Stock solutions of each drug were prepared daily in a vehicle of tap water containing AmQuel® water conditioner. Treatment solutions were diluted with tap water containing AmQuel® water conditioner. Concentrations of cocaine were based on prior behavioral outcomes in planarians (Owaisat et al., 2012; Pagan et al., 2013; Rawls et al., 2010), and levamisole concentrations were determined empirically using previously reported K_i values as a guide (Anagnostou et al., 1996).

2.2. Behavioral experiments

2.2.1. Stereotypical activity—Individual planarians were placed randomly into a transparent petri dish (5.5 cm diameter) containing a solution of cocaine (0, 0.1, 1, 3, 5 mM) or levamisole (0, 0.1, 0.3, 0.75, 1 mM) for 5 min and stereotyped movements were quantified (Ramos et al., 2012). The concentration-effect data for each individual drug was used to determine the constant-potency ratio of the two drugs at a specified effect level (*i.e.* equi-effective doses of each drug). From this value the isobole, which indicates additivity for the predicted effect of the combination, was constructed and used to determine if the combination was additive, sub-additive or synergistic (super-additive) (Tallarida, 2011, 2012). Combination doses used in actual experiments were determined based on individual drug potencies.

2.2.2. Conditioned Place Preference (CPP)—CPP experiments were divided into 3 different phases: 1) pre-conditioning (pre-test); 2) conditioning; and 3) post-conditioning (post-test). Because planarians display a natural preference for a dark environment (Raffa et al., 2003), we used a biased, counterbalanced conditioning design to assess cocaine and levamisole preference (Ramos et al., 2012). In a biased design, the preference of each individual animal for a particular environment is determined prior to conditioning by placing the animal in the apparatus, and then by assessing the amount of time the animal spends in each compartment. The least-preferred compartment for each animal is then assigned to be the drug-paired compartment. For the pre-conditioning phase, dark and “ambient” light environments were created by covering half (both the top and bottom) of a petri dish containing water with black construction paper. An individual planarian was then placed at the midline of the dish and given free access to roam both the light and dark environments of the dish. The time spent in the least-preferred setting over a 5-min interval was then determined. This value is called the pre-test time. The least-preferred environment, as determined during pre-conditioning, is designated as the environment in which drug conditioning occurs and is therefore called the ‘drug-paired’ environment. For conditioning, planarians were exposed to either cocaine (0, 0.001, 0.01, 0.1, 1, 100 μ M) or levamisole (0,

0.01, 0.1, 1 μ M) for 30 min in the least-preferred (drug-paired) environment. For the situation in which the 'drug-paired' environment is ambient light, the petri dish is uncovered during the conditioning phase to allow exposure to the light. For the opposite situation in which the drug-paired environment is the dark, the entire petri dish is covered with black construction paper to enable exposure to a dark environment. Immediately following conditioning, the post-conditioning phase was performed in a manner identical to that described for pre-conditioning. Planarians were placed at the midline of a petri dish containing water and allowed free access to the light and dark sides of the dish for 5 min. Time spent in the drug-paired side (the original least-preferred environment) was determined (post-test), and a preference score was calculated as the difference between the post-test and pre-test times. A similar protocol has been used by our laboratory to demonstrate that planarians display CPP to different addictive substances including designer cathinones, nicotine, and sugar (Rawls et al., 2011; Ramoz et al., 2012; Zhang et al., 2013). Cocaine produced significant CPP whereas levamisole did not (see Results). Thus, for combined administration, inactive concentrations (0.01, 1 μ M) of levamisole were tested in combination with graded concentrations (0.001, 0.1, 100 μ M) of cocaine.

2.3. Data analysis

Comparisons of group means ($\pm$ S.E.M.) for experiments involving individual drugs were evaluated by one-way ANOVA, and in cases of significance, followed by Dunnett's test. For combined administration, isobolographic analysis, plus two-way ANOVA followed by Bonferroni's test, was used. Values of $p < 0.05$ were considered statistically significant.

3. Results

3.1. Combination of cocaine and levamisole elicits synergistic stereotypical activity

Effects of acute exposure to cocaine or levamisole on stereotyped movements are presented in Fig. 1. For cocaine (Fig. 1A), one-way ANOVA revealed a significant main effect [$F(4, 35) = 58.12, p < 0.0001$]. Post-hoc analysis indicated that 3 and 5 mM cocaine produced significantly greater stereotypical activity than water-exposed controls ($p < 0.001$). The maximal number of stereotyped movements (42.75 ± 4.11) was produced by 5 mM cocaine. For levamisole (Fig. 1B), one-way ANOVA revealed a significant main effect [$F(4, 35) = 47.78, p < 0.0001$]. Post-hoc analysis indicated that concentrations of 0.3, 0.75 and 1 mM levamisole produced significantly greater stereotypical activity compared to water-exposed controls ($p < 0.001$). The maximal number of stereotyped movements (25.88 ± 2.18) was produced by 1 mM levamisole.

The individual concentration-effect curve of each individual drug (determined from the data in Fig. 1) allows a determination of equally effective doses (*e.g.*, effect 20% is inserted into the equation of each curve and the dose that produces this effect is determined). Specifically, cocaine gives a concentration of 2.25 mM whereas levamisole gives a concentration of 0.703 mM. These ED₂₀ values for cocaine and levamisole reveal that the agents have a constant-potency ratio (cocaine:levamisole) of 3.20 with potencies of 2.25 mM for cocaine and 0.703 for levamisole. In other words, these values are the equi-effective doses of cocaine and levamisole that produce the same effect level (an effect of 20 stereotyped

movements). From those values the isobole, which indicates additivity for the combination of cocaine and levamisole, was determined and shown in Fig. 2B as the diagonal line with intercepts 2.25 ± 0.097 (horizontal) and 0.703 ± 0.125 (vertical). Four combinations of cocaine and levamisole, all of which maintained the constant-potency ratio of 3.20 (*i.e.*, cocaine concentration (mM)/levamisole concentration (mM): 2.25/0.703; 1.12/0.35; 0.56/0.17; 0.28/0.08), were tested in actual experiments and their effects on stereotyped movements are shown in Fig. 2A. As was the case with the individual agents (Fig. 1), the constant-potency combinations of cocaine and levamisole produced concentration-dependent increases in stereotyped movements (Fig. 2A). Fig 2B shows the isobole. This is a plot of all dose combinations (cocaine and levamisole) that are expected to give the specified effect when there is no interaction. The observed dose combination point (0.598 mM for cocaine, 0.186 mM for levamisole) that achieved an effect level of 20 stereotyped movements, however, is below the isobole, thereby indicating synergism, *i.e.*, the effect is achieved with this lower dose combination. (Tallarida, 2012).

3.2. Levamisole enhances cocaine-induced conditioned place preference (CPP)

The place conditioning effects of cocaine and levamisole are presented in Fig. 3. For cocaine (Fig. 3A), one-way ANOVA indicated a significant main effect on CPP [$F(5, 46) = 2.832, p < 0.05$]. Conditioning with a concentration of 0.01 μM cocaine produced a significantly greater preference shift (100 ± 15 s) than water controls (40 ± 10 s) ($p < 0.05$). For levamisole (Fig. 3B), a significant main effect was not indicated by one-way ANOVA [$F(3, 28) = 1.271, p > 0.05$].

Using the CPP data obtained for the individual drugs as a guide (Fig. 3), we combined inactive concentrations of levamisole (0.01, 1 μM) with submaximal (0.001 μM), as well as higher and more efficacious concentrations (0.1, 100 μM), of cocaine. In the case in which an inactive concentration of levamisole is administered with an active concentration of cocaine, the expected effect becomes one in which the preference shift produced by the combination is simply equal to the preference shift produced by cocaine alone. The analysis is then one in which the effect of the active drug, cocaine, is statistically compared before and after the addition of levamisole. For actual experiments (Fig. 4), two-way ANOVA revealed a significant effect of levamisole [$F(2, 60) = 3.51, p < 0.05$]. Bonferroni analysis indicated that the preference shift produced by a submaximal concentration of cocaine (0.001 μM) was enhanced in the presence of levamisole ($p < 0.05$) (Fig. 4). The shift produced by cocaine (0.001 μM) alone was 43 ± 18 s; in the presence of 0.01 and 1 μM the preference shift was 123 ± 23 s and 119 ± 13 s, respectively. Levamisole did not significantly affect the preference shift elicited by higher concentrations of cocaine (0.1, 100 μM) ($p > 0.05$).

4. Discussion

The aim of our study was to provide experimental information about effects of levamisole on cocaine's action *in vivo* and that effort revealed synergism for the cocaine and levamisole interaction. We demonstrated that two neuropharmacological effects of cocaine, stereotyped movements and CPP, were enhanced in select cases in which cocaine was administered in

combination with levamisole. As is often the case when synergism is first detected, we do not yet know the precise mechanism, but the very detection of synergism is an important first step in exploring mechanism and impact. Understanding the pharmacology of a prevalent combination such as cocaine and levamisole is critical because poly-drug use (*e.g.* cocaine + alcohol, heroin + cocaine) can increase dangers that each drug poses by itself and make it more difficult to identify efficacious medications to manage dependence, craving, and relapse. An example of the challenge posed by poly-drug addiction is the impact of alcohol dependence on the efficacy of modafinil in treating cocaine dependence (Anderson et al 2009; Vocci and Elkashef, 2005). In this case, modafinil treatment reduces cocaine use in subjects that are not dependent on alcohol but, in cocaine users who are alcohol dependent, modafinil loses its efficacy.

Effects of cocaine on the endpoints studied here (stereotypy and CPP) are established across different species, including planarians (Pagan et al., 2013; Rawls et al., 2010). Although effects of levamisole on motor and reward systems are not well defined, some of its biological actions are consistent with an agent that enhances effects of cocaine through direct or indirect augmentation of monoamine activity. For example, levamisole inhibits monoamine oxidase (MAO) and activates nicotinic receptors (*e.g.* $\alpha 3\beta 4$, $\alpha 3\beta 2$), and both MAO inhibitors and nicotinic agonists enhance cocaine effects and increase dopamine transmission (Hernando et al 2012; Levandoski et al 2003; Agarwal et al 1990). Levamisole also elevates levels of endogenous opioids associated with drug highs and attenuates the opiate withdrawal syndrome in rats (Spector et al 1998). Furthermore, levamisole, when used as adjuvant therapy for colon cancer, has been reported to cause mood-elevating effects (Goldin et al 1982). On the basis of those combined data, one explanation for the levamisole synergism with cocaine in activity and CPP assays is enhanced dopamine activity, perhaps through the simultaneous block of dopamine catabolism, or activation of nicotinic acetylcholine receptors, by levamisole and blockade of dopamine uptake by cocaine. Pharmacodynamic mechanisms are not the only potential explanations for the synergy; pharmacokinetic interactions, such as levamisole inhibiting catabolism of cocaine into inactive metabolites, or a chemical interaction between the two drugs that leads to enhancement of the pharmacological properties of cocaine, may have played a role.

When administered by themselves, cocaine and levamisole increased stereotyped movements. Although drugs possessing abuse liability often increase stereotypical activity in laboratory animals (Koob, 1992), the relationship, if any, between stereotyped movements in planarians, rats and mice is uncertain. Part of the reason is that stereotypy is broadly defined as a repetitive, ritualistic movement and is not uniformly quantified across different species. For rodents a scoring system is frequently used to quantify individual stereotypical movements, such as rearing, grooming, sniffing, and head bobbing (Tanda et al., 2007), and photocell beam interruptions are used to quantify cumulative stereotyped movements (Rasmussen et al., 2011; Hummell and Unterwald, 2003). For planarians we quantify stereotypical activity as C-shape movements, which are detected following exposure to different addictive substances including cocaine, synthetic cathinones, nicotine, and caffeine. (Zhang et al., 2013; Ramoz et al., 2012; Rawls et al., 2010; Pagan et al., 2008; Palladini et al., 2006). C-shape movements in planarians have also been described as

“seizure-like activity” based on evidence that proconvulsants induce C-shape movements that are prevented by concurrent exposure to antiepileptic agents (Ramakrishnan and Desaeer, 2011; Raffa et al., 2010). Furthermore, planarians exposed to cholinergic agonists exhibit muscle contractions resembling C-shape movements (Nishimura et al., 2010; Rawls et al., 2011; Owaisat et al., 2012). Given that seizure-like activity has been reported in mice following administration of higher doses of levamisole (Rehni and Singh, 2010), and that levamisole activates nicotinic acetylcholine receptors involved in seizures (Levandoski et al., 2003), it cannot be discounted that a proconvulsant action of levamisole was partly responsible for producing the C-shape movements in planarians.

The effectiveness of cocaine in planarian motor and CPP assays varied with concentration; lower concentrations produced the greatest preference shifts whereas 1000-fold higher concentrations were required to elicit stereotyped movements. Similar dose-related trends have been observed in corresponding rat assays and suggest that the rewarding and aversive effects of cocaine are separable by dose. For example, in rat CPP studies, cocaine displays an inverted U-shaped dose response in which the highest doses (*e.g.* 30 mg/kg) produced diminished responses due to their aversive effects (Bardo et al., 1995; Tzschentke, 2007; Zakharova et al., 2009). Doses of cocaine at the high end of the inverted U-shaped curve also produce more robust stereotypical activity (Estevez et al., 1979; Bhattacharyya et al., 1979) and, at even higher doses, eventually seizure (Zagnoni and Albano, 2002). Our results with planarians suggest a similar phenomenon for cocaine in which its rewarding effects are more robust at lower concentrations that do not elicit significant increases in stereotypical activity.

Levamisole enhanced CPP produced by submaximal concentrations of cocaine but did not alter preference shifts produced by higher, and more effective, concentrations of cocaine. Levamisole's preferential effect on submaximal concentrations of cocaine may be related to the magnitude of the preference shift produced by cocaine itself. For instance, the more effective concentrations of cocaine are likely to be associated with a ‘ceiling effect’, a phenomenon in which a drug produces a maximum effect so that increasing the drug dosage or, as in the present case, adding another type of drug does not increase overall efficacy (Lufty and Cowan, 2004; Tallarida, 2012). Indeed, when cocaine was administered to planarians by itself, the maximal preference shift was about 100 s. For combinations of cocaine and levamisole, the maximal preference shift detected was about 125 s. The relative similarity in magnitude of those preference shifts, coupled with the inverted U-shaped dose response curve of cocaine by itself, supports the interpretation that levamisole synergism with cocaine is more likely to be detected at submaximal concentrations of cocaine.

A comment is perhaps warranted on a liability of the planarian assay, which is the challenge of quantifying changes in motility following acute exposure to stimulant drugs. Planarians display slight to modest increases in motility immediately following exposure to concentrations of amphetamine and nicotine in the lower portion of their concentration curves (Raffa and Martley, 2005). However, acute exposure to modest to higher concentrations decreases motility (Pagan et al., 2009; Rawls et al., 2011).

In conclusion, we provide the first evidence that levamisole enhances cocaine's action *in vivo*. Most important is the identification of levamisole synergism with cocaine, which was detected in two different assays, including CPP, in which the environmental preference for cocaine was augmented by the presence of levamisole. The synergism between cocaine and levamisole now requires further, and more detailed, investigation in mammalian models of CPP and self-administration to determine if levamisole affects the positive reinforcing and drug seeking properties of cocaine.

Acknowledgments

Funding Sources: National Institute on Drug Abuse grants DA07237 and DA013429

References

- Agarwal A, Shukla OP, Ghatak S, Tekwani BL. Biogenic amines, metabolites and monoamine oxidase in the filarial worm *Setaria cervi*. *Int J Parasitol*. 1990; 20:873–881. [PubMed: 2276863]
- Anagnostou F, Plas C, Forest N. Ecto-alkaline phosphatase considered as levamisole-sensitive phosphohydrolase at physiological pH range during mineralization in cultured fetal calvaria cells. *J Cell Biochem*. 1996; 60:484–494. [PubMed: 8707888]
- Anderson AL, Reid MS, Li SH, Holmes T, Shemanski L, Slee A, Smith EV, Kahn R, Chiang N, Vocci F, Ciraulo D, Dackis C, Roache JD, Salloum IM, Somoza E, Urschel HC, Elkashef AM. Modafinil for the treatment of cocaine dependence. *Drug Alcohol Depend*. 2009; 104:133–139. [PubMed: 19560290]
- Auffenberg C, Rosenthal LJ, Dresner N. Levamisole: A Common Cocaine Adulterant with Life-Threatening Side Effects. *Psychosomatics*. 2013
- Bardo MT, Rowlett JK, Harris MJ. Conditioned place preference using opiate and stimulant drugs: a meta-analysis. *Neurosci Biobehav Rev*. 1995; 19:39–51. [PubMed: 7770196]
- Bhattacharyya AK, Pradhan SN. Interactions between motor activity and stereotypy in cocaine-treated rats. *Psychopharmacology (Berl)*. 1979; 63:311–312. [PubMed: 113821]
- Buchanan JA, Oyer RJ, Patel NR, Jacquet GA, Bornikova L, Thienelt C, Shriver DA, Shockley LW, Wilson ML, Hurlbut KM, Lavonas EJ. A confirmed case of agranulocytosis after use of cocaine contaminated with levamisole. *J Med Toxicol*. 2010; 6:160–164. [PubMed: 20358411]
- Buttarelli FR, Pellicano C, Pontieri FE. Neuropharmacology and behavior in planarians: translations to mammals. *Comp Biochem Physiol C Toxicol Pharmacol*. 2008; 147:399–408. [PubMed: 18294919]
- Chang A, Osterloh J, Thomas J. Levamisole: a dangerous new cocaine adulterant. *Clin Pharmacol Ther*. 2010; 88:408–411. [PubMed: 20668440]
- Eriksson KS, Panula P. gamma-Aminobutyric acid in the nervous system of a planarian. *J Comp Neurol*. 1994; 345:528–536. [PubMed: 7962699]
- Estevez VS, Ho BT, Englert LF. Metabolism correlates of cocaine-induced stereotypy in rats. *Pharmacol Biochem Behav*. 1979; 10:267–271. [PubMed: 572058]
- Goldin A, Chirigos MA, Macdonald JS, Fefer A, Mihich E. Biologic-response modifiers and adjuvant chemotherapy: consideration of selected preclinical investigations in relation to clinical potential. *Recent Results Cancer Res*. 1982; 80:351–356. [PubMed: 6173903]
- Hernando G, Bergé I, Rayes D, Bouzat C. Contribution of subunits to *Caenorhabditis elegans* levamisole-sensitive nicotinic receptor function. *Mol Pharmacol*. 2012; 82:550–560. [PubMed: 22734069]
- Hummel M, Unterwald EM. Intra-accumbens pertussis toxin sensitizes rats to the locomotor activating effects of a single cocaine challenge. *Brain Res*. 2003; 965:100–107. [PubMed: 12591125]
- Koob GF. Neural mechanisms of drug reinforcement. *Ann N Y Acad Sci*. 1992; 654:171–191. [PubMed: 1632582]

- Levandoski MM, Piket B, Chang J. The anthelmintic levamisole is an allosteric modulator of human neuronal nicotinic acetylcholine receptors. *Eur J Pharmacol.* 2003; 471:9–20. [PubMed: 12809947]
- Lutty K, Cowan A. Buprenorphine: a unique drug with complex pharmacology. *Curr Neuropharmacol.* 2004; 2:395–402. [PubMed: 18997874]
- Nishimura K, Kitamura Y, Taniguchi T, Agata K. Analysis of motor function modulated by cholinergic neurons in planarian *Dugesia japonica*. *Neuroscience.* 2010; 168:18–30. [PubMed: 20338223]
- Owaisat S, Raffa RB, Rawls SM. In vivo comparison of harmine efficacy against psychostimulants: preferential inhibition of the cocaine response through a glutamatergic mechanism. *Neurosci Lett.* 2012; 525:12–16. [PubMed: 22877698]
- Pagán OR, Deats S, Baker D, Montgomery E, Wilk G, Tenaglia M, Semon J. Planarians require an intact brain to behaviorally react to cocaine, but not to react to nicotine. *Neuroscience.* 2013; 246:265–270. [PubMed: 23684614]
- Pagán OR, Rowlands AL, Azam M, Urban KR, Bidja AH, Roy DM, Feeney RB, Afshari LK. Reversal of cocaine-induced planarian behavior by parthenolide and related sesquiterpene lactones. *Pharmacol Biochem Behav.* 2008; 89:160–170. [PubMed: 18222535]
- Pagán OR, Rowlands AL, Fattore AL, Coudron T, Urban KR, Bidja AH, Eterovi VA. A cembranoid from tobacco prevents the expression of nicotine-induced withdrawal behavior in planarian worms. *Eur J Pharmacol.* 2009; 615:118–124. [PubMed: 19490913]
- Palladini G, Ruggeri S, Stocchi F, De Pandis MF, Venturini G, Margotta V. A pharmacological study of cocaine activity in planaria. *Comp Biochem Physiol C Pharmacol Toxicol Endocrinol.* 1996; 115:41–45. [PubMed: 8983168]
- Passarelli F, Merante A, Pontieri FE, Margotta V, Venturini G, Palladini G. Opioid-dopamine interaction in planaria: a behavioral study. *Comp Biochem Physiol C Pharmacol Toxicol Endocrinol.* 1999; 124:51–55. [PubMed: 10579648]
- Raffa RB, Finno KE, Tallarida CS, Rawls SM. Topiramate-antagonism of L-glutamate-induced paroxysms in planarians. *Eur J Pharmacol.* 2010; 649:150–153. [PubMed: 20863783]
- Raffa RB, Rawls SM. A Model for drug action and abuse. Landes Bioscience; Austin, TX: 2008.
- Raffa RB, Martley AF. Amphetamine-induced increase in planarian locomotor activity and block by UV light. *Brain Res.* 2005; 1031:138–140. [PubMed: 15621023]
- Raffa RB, Dasrath CS, Brown DR. Disruption of a drug-induced choice behavior by UV light. *Behav Pharmacol.* 2003; 14:569–571. [PubMed: 14557725]
- Ramakrishnan L, Desaeer C. Carbamazepine inhibits distinct chemoconvulsant-induced seizure-like activity in *Dugesia tigrina*. *Pharmacol Biochem Behav.* 2011; 99:665–670. [PubMed: 21699913]
- Ramoz L, Lodi S, Bhatt P, Reitz AB, Tallarida C, Tallarida RJ, Raffa RB, Rawls SM. Mephedrone (“bath salt”) pharmacology: insights from invertebrates. *Neuroscience.* 2012; 208:79–84. [PubMed: 22300981]
- Rasmussen B, Unterwald EM, Rawls SM. Glutamate transporter subtype 1 (GLT-1) activator ceftriaxone attenuates amphetamine-induced hyperactivity and behavioral sensitization in rats. *Drug Alcohol Depend.* 2011; 118:484–488. [PubMed: 21524862]
- Rawls SM, Karaca F, Madhani I, Bhojani V, Martinez RL, Abou-Gharbia M, Raffa RB. β -lactamase inhibitors display anti-seizure properties in an invertebrate assay. *Neuroscience.* 2010; 169:1800–1804. [PubMed: 20600649]
- Rawls SM, Patil T, Tallarida CS, Baron S, Kim M, Song K, Ward S, Raffa RB. Nicotine behavioral pharmacology: clues from planarians. *Drug Alcohol Depend.* 2011; 118:274–279. [PubMed: 21530106]
- Rehni AK, Singh TG. Levamisole-induced reduction in seizure threshold: a possible role of nicotinic acetylcholine receptor-mediated pathway. *Naunyn Schmiedebergs Arch Pharmacol.* 2010; 382:279–285. [PubMed: 20694554]
- Rowlands AL, Pagán OR. Parthenolide prevents the expression of cocaine-induced withdrawal behavior in planarians. *Eur J Pharmacol.* 2008; 583:170–172. [PubMed: 18275955]

- Spector S, Munjal I, Schmidt DE. Effects of the immunostimulant, levamisole, on opiate withdrawal and levels of endogenous opiate alkaloids and monoamine neurotransmitters in rat brain. *Neuropsychopharmacology*. 1998; 19:417–427. [PubMed: 9778663]
- Tallarida RJ. Quantitative methods for assessing drug synergism. *Genes Cancer*. 2011; 2:1003–1008. [PubMed: 22737266]
- Tallarida RJ. Revisiting the isobole and related quantitative methods for assessing drug synergism. *J Pharmacol Exp Ther*. 2012; 342:2–8. [PubMed: 22511201]
- Tanda G, Ebbs AL, Kopajtic TA, Elias LM, Campbell BL, Newman AH, Katz JL. Effects of muscarinic M1 receptor blockade on cocaine-induced elevations of brain dopamine levels and locomotor behavior in rats. *J Pharmacol Exp Ther*. 2007; 321:334–344. [PubMed: 17255465]
- Tzschentke TM. Measuring reward with the conditioned place preference (CPP) paradigm: update of the last decade. *Addict Biol*. 2007; 12:227–462. [PubMed: 17678505]
- Ullrich K, Koval R, Koval E, Bapojé S, Hirsh JM. Five consecutive cases of a cutaneous vasculopathy in users of levamisole-adulterated cocaine. *J Clin Rheumatol*. 2011; 17:193–196. [PubMed: 21617558]
- Vocci FJ, Elkashef A. Pharmacotherapy and other treatments for cocaine abuse and dependence. *Curr Opin Psychiatry*. 2005; 18:265–270. [PubMed: 16639150]
- Vyas CA, Rawls SM, Raffa RB, Shackman JG. Glutamate and aspartate measurements in individual planaria by rapid capillary electrophoresis. *J Pharmacol Toxicol Methods*. 2011; 63:119–122. [PubMed: 20708699]
- Wolford A, McDonald TS, Eng H, Hansel S, Chen Y, Bauman J, Sharma R, Kalgutkar AS. Immune-mediated agranulocytosis caused by the cocaine adulterant levamisole: a case for reactive metabolite(s) involvement. *Drug Metab Dispos*. 2012; 40:1067–1075. [PubMed: 22393119]
- Zagnoni PG, Albano C. Psychostimulants and epilepsy. *Epilepsia* 43 Suppl. 2002; 2:28–31.
- Zakharova E, Miller J, Unterwald E, Wade D, Izenwasser S. Social and physical environment alter cocaine conditioned place preference and dopaminergic markers in adolescent male rats. *Neuroscience*. 2009; 163:890–897. [PubMed: 19580849]
- Zhang C, Tallarida CS, Raffa RB, Rawls SM. Sucrose produces withdrawal and dopamine-sensitive reinforcing effects in planarians. *Physiol Behav* 112. 2013; 113:8–13.
- Zhu NY, Legatt DF, Turner AR. Agranulocytosis after consumption of cocaine adulterated with levamisole. *Ann Intern Med*. 2009; 150:287–289. [PubMed: 19153405]

Research Highlights

- Interactions between levamisole and cocaine were investigated *in vivo*.
- Cocaine produced place conditioning and stereotypical effects in planarians.
- Levamisole produced stereotyped movements in planarians.
- Levamisole enhanced cocaine-induced conditioned place preference and stereotypy in planarians.
- Levamisole interacted synergistically with cocaine.

Fig. 1. Cocaine and levamisole produce stereotyped movements in planarians

Planarians were exposed to different concentrations of cocaine (**1A**) or levamisole (**1B**) for 5 min. The number of C-shape movements over the 5-min exposure interval were determined and presented as mean stereotypy counts $\pm$ S.E.M. $N=8$ planarians/group. *** $p < 0.001$ or * $p < 0.05$ compared to the respective water control in cocaine (**1A**) or levamisole (**1B**) experiments.

Fig. 2. Cocaine and levamisole interact synergistically to produce stereotyped movements in planarians

Cocaine and levamisole were determined to have a constant-potency ratio (cocaine:levamisole) of 3.20, with potencies of 2.25 mM for cocaine and 0.703 for levamisole, from the data in Fig. 1. **(2A)** Four combinations maintaining the constant-potency ratio of 3.20 (*i.e.*, cocaine concentration (mM)/levamisole concentration (mM): 2.25/0.703; 1.12/0.35; 0.56/0.17; 0.28/0.08) were administered to planarians for 5 min. The number of C-shape movements were determined over the 5min exposure interval and presented as mean stereotypy counts $\pm$ S.E.M. $N=8$ planarians/group. **(2B)** Isobole indicating additivity for the combination was determined from the constant-potency ratio and presented as the diagonal line with intercepts 2.25 $\pm$ 0.097 (horizontal) and 0.703 $\pm$ 0.125 (vertical). The observed dose-combination point (0.598 mM for cocaine, 0.186 mM for levamisole) determined from Fig. 2A data lies significantly below the isobole, thereby indicating synergism for the combination.

Fig. 3. Cocaine, but not levamisole, produces CPP in planarians

Data are presented as the mean preference score (s) + S.E.M. (difference between post-conditioning and pre-conditioning times) from planarians in which cocaine (**3A**) or levamisole (**3B**) was paired with ambient light during the conditioning phase. N = 8-12 planarians/group. * $p < 0.05$ compared to water control.

Fig. 4. Levamisole enhances cocaine-induced CPP

Data are presented as the mean preference score (s) + S.E.M. (difference between post-conditioning and pre-conditioning times) from planarians in which increasing concentrations of cocaine (the active agent) were co-administered with water or levamisole (LVM) (the inactive agent) in the ambient light (non-preferred environment) during the conditioning phase. N = 8 planarians/group. * $p < 0.05$ compared to water (LVM [0 μM]) + cocaine group.