Published in final edited form as:

Curr Opin Immunol. 2009 June ; 21(3): 291–297. doi:10.1016/j.coi.2009.05.008.

Diversity in CD8⁺ T cell differentiation

Ian A. Parish and Susan M. Kaech

Department of Immunobiology, Yale University School of Medicine, New Haven, CT 06520, USA

SUMMARY

CD8⁺ T cells are key effector cells of the adaptive immune system, however their activity must be tightly regulated to allow pathogen clearance while preventing immunopathology and autoimmunity. In this review, we summarise the diversity of responses that CD8⁺ T cells make to antigenic stimulation with a focus on how CD8⁺ T cell responses are regulated to achieve different immune outcomes. In particular, we discuss phenotypic diversity during tolerance induction as well as signals that drive effector and memory cell differentiation in response to infection.

INTRODUCTION

The peripheral CD8⁺ T cell repertoire is in a constant state of flux as these cells see a myriad of environmental signals that are continuously varying. In the steady-state, naïve CD8⁺ T cells must compete for homeostatic survival signals, such as MHC class I (MHC I) and interleukin-7 (IL-7), in order to survive. Moreover, antigen encounter can synergise with environmental signals to drive CD8⁺ T cells down numerous differentiation states that vary both in effector function and survival capacity. Correct co-ordination of these differentiation states is critical for both the maintenance of peripheral tolerance and the expansion and persistence of pathogen-specific clones. This review aims to summarise the diversity of CD8⁺ T cell responses to antigen, with a focus on recent work in the fields of CD8⁺ T cell tolerance, effector cell differentiation and memory (Figure 1).

MULTIPLE PATHS TO PERIPHERAL TOLERANCE

While autoreactive CD8⁺ T cells are purged from the repertoire during thymic negative selection, this is not a perfect process and rogue autoreactive CD8⁺ T cells are often released into the periphery [1]. Peripheral tolerance mechanisms have thus evolved to purge the repertoire of these dangerous self-reactive cells. Dendritic cells (DCs), a key antigen presenting cell population, are thought to drive this process by inducing tolerance within any T cells that recognise antigen presented by DCs in the steady-state.

© 2009 Elsevier Ltd. All rights reserved.

Address Correspondence: Dr. Susan Kaech, 300 Cedar Street, The Anlyan Center S641B, Yale University School of Medicine, P. O. Box 208011, New Haven, CT 06520. susan.kaech@yale.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Upon steady-state antigen recognition, peripheral tolerance induction in CD8⁺ T cells can result in two distinct differentiation states: deletion and anergy (Figure 1). In both tolerance fates, CD8⁺ T cells generally fail to develop effector functions despite proliferating in response to antigen [2,3], although cells can pass through an effector phase *en route* to tolerisation in some models [4,5]. However, despite these similarities, each tolerance fate exhibits unique features. While anergic CD8⁺ T cells persist in a hyporesponsive state caused by T cell receptor signalling defects [3], cells undergoing deletion die by apoptosis triggered by the proapoptotic BH3-only protein Bim [6]. Nevertheless, contraction does occur in models of CD8⁺ T cell anergy and this death requires Bim (I. A. Parish and W. R. Heath, unpublished observations). The biological reason for maintaining anergic cells is currently unclear, but one possibility is that anergic CD8⁺ T cells adopt an immunoregulatory state [7].

Deletion and anergy are often considered disparate cell fates, however recent data has questioned whether these cell fates are as molecularly distinct as previously thought. One of the features of anergic T cells is up-regulation of the transcription factors Egr-2 and Egr-3 by NFAT activation in the absence of AP-1 activation [8,9]. Egr-2 and Egr-3 then drive expression of immunosuppressive E3 ubiquitin ligases, such as Cbl-b, which degrade key signalling components and attenuate T cell receptor (TCR) signalling capacity [10]. Interestingly, both Egr2 and Cbl-b (as well numerous other anergy associated genes) were strongly up-regulated during deletion [11] suggesting that molecular parallels exist between deletion and anergy. Furthermore, signalling through the inhibitory signalling molecule PD-1 is required for both CD8⁺ T cell anergy [12] and deletion [13,14]. Finally, at least in CD4⁺ T cells, blocking death in a model of deletion leads to a fate of anergy [15]. Although still speculative, it is possible that common genetic pathways direct states of anergy and deletion in tolerised CD8⁺ T cells, but the outcome is largely dependent on the overall level of TCR signalling. It is known that high antigen levels typically cause anergy whereas lower antigen levels lead to deletion [16]. Thus, in situations of weak or tissue-restricted antigen presentation, deletion may prevail because the expression of genes inhibitory to TCR signalling, coupled with lower IL-7 receptor expression [11], promote cell death by “starving” cells of MHC and IL-7 survival signals. In contrast, higher antigen levels or ubiquitous antigen expression may provide a sufficiently strong and persistent TCR signal to permit survival of tolerised CD8⁺ T cells.

BREAKING TOLERANCE: THE ROAD TO AN EFFECTOR CELL

Whereas steady-state presentation of antigen by DCs causes CD8⁺ T cell tolerance by default, antigen presentation in the context of infection drives clonal expansion and CD8⁺ T cell effector and memory formation. Effector CD8⁺ T cells (often termed cytotoxic T lymphocytes, or CTL) can kill antigen-bearing target cells using granzymes and perforin, and can rapidly produce anti-viral cytokines, such as IFN- γ and TNF- α upon TCR ligation [17]. Effector cell differentiation occurs during infection partly because DC recognition of pathogen-derived products creates immunogenic DCs that bear increased amounts of co-stimulatory molecules and antigen:MHC complexes. Antigen encounter on immunogenic antigen-presenting cells for brief periods *in vitro* (as little as 1 day) can induce CD8⁺ T cell proliferation and differentiation into effectors, although longer durations of antigen

encounter *in vivo* augment clonal expansion [17,18]. Initial *in vitro* studies also suggested that TCR avidity for antigen regulates effector cell differentiation [19], however recent *in vivo* data suggest that, while avidity regulates the magnitude of the response, effector (and memory) CD8⁺ T cells can still form in response to low avidity interactions [20]. In addition, pathogen activated DCs also up-regulate Notch ligands, which can ligate Notch on CD8⁺ T cells and directly induce cytotoxicity [21].

Another key set of signals in effector cell formation during infection are innate immune system-derived inflammatory cytokines. In particular, IL-12 and the type 1 interferons (IFN α/β) promote CTL expansion and differentiation *in vitro* [22] and *in vivo* during infection [23,24], in part through induction of cytotoxic molecules and IFN- γ production [22,25]. As different infections will elicit disparate cytokine responses, the cytokines required for CTL expansion and differentiation may vary between infections. For example, CTL responses to *Listeria* infection are more IL-12 dependent [26,27], while the CTL response to LCMV infection is more IFN α/β dependent [23,28]. It should be noted that IL-12 stimulation is most effective in driving CTL differentiation when received simultaneously with a TCR stimulus [26,29], suggesting that, like CD4⁺ T cells [30], antigenic and cytokine signals must be co-delivered for optimal CTL differentiation. There is additional evidence that other inflammatory cytokines influence CTL differentiation, a topic recently reviewed in detail [22,31].

A number of transcription factors have been identified that promote CTL differentiation. The T-box transcription factor eomesodermin drives acquisition of cytotoxicity and anti-viral cytokine production [32], although the pathways responsible for eomesodermin induction are currently unclear. Another T-box transcription factor, T-bet, also appears to be a critical regulator of effector cell differentiation as T-bet deficient CD8⁺ T cells exhibit diminished cytotoxicity and altered cytokine production [33]. Moreover, IL-12 induces expression of T-bet [34,35] providing a mechanism for IL-12 driven CTL differentiation. Nevertheless, IL-12 and IFN α/β can also trigger IFN- γ production via STAT4 activation [25]. The presence of both T-bet and eomesodermin is critical for normal CTL differentiation as CD8⁺ T cells deficient in both of these factors develop an aberrant effector phenotype characterised by excessive IL-17 production [36]. In addition, the transcription factors RBP-J and CREB1 are responsible for the expression of cytotoxic molecules downstream of Notch signalling [21]. Repressors of effector function also exist; the transcriptional repressor Bcl-6 can repress expression of the cytotoxic molecule granzyme B [37], although Bcl-6 may be more important in controlling memory formation than initial effector cell differentiation (see next section). A major challenge in the field is to determine how this transcriptional network is organised to control effector cell formation, and maintain effector molecule gene expression in memory cells in the absence of infection.

While the expression of cytotoxic products in CTLs aids pathogen clearance, they may also be detrimental to the lifespan of a CTL. For example, CTLs lacking the serine protease inhibitors Spi-6 or Spi-2a, which block granzyme B and cathepsin B activity, respectively, show reduced expansion during viral infection [38,39]. Interestingly, Spi-6 is a gene upregulated by the transcription factor Id2, and Id2^{-/-} CD8⁺ T cells demonstrate a similar propensity to die and decreased clonal expansion during viral infection [40].

It should be noted that during certain chronic viral infections, CD8⁺ T cells exhibit dysfunction in their ability to mobilise effector functions, such as IFN- γ , TNF- α , IL-2 and cytotoxicity. This dysfunction (often referred to as “exhaustion”) may also be considered a unique CD8⁺ T cell differentiation state. A network of inhibitory receptors, including PD-1 [41] and LAG-3 [42], are up-regulated on exhausted T cells and appear important in driving the exhausted state. In addition, augmented IL-10 production during chronic infection also promotes an exhausted CD8⁺ T cell state [43,44] and may operate in a pathway independent of PD-1 [45]. A number of transcription factors are selectively up-regulated in exhausted CD8⁺ T cells (such as PBX3 and Blimp-1) [46] suggesting this effector cell fate is an active process directed by gene regulatory changes.

LIFE AFTER DEATH: PERSISTENCE OF CD8⁺ T CELLS INTO MEMORY

At the completion of a CD8⁺ T cell immune response, the majority of effector cells die by Bim-dependent apoptosis [47,48]. However, a proportion of cells survive this contraction process and persist as a memory T cell population that confers long-term protection against re-infection. Memory T cells, although heterogeneous in phenotype, possess a well defined set of characteristics: they exhibit enhanced proliferative capacity (relative to effector cells), augmented re-expression of effector genes upon rechallenge (relative to naïve cells) and a unique ability to self-renew and survive for long-periods of time. The molecular basis of many of these features is now emerging. Memory cells exhibit greater proliferative potential relative to effector cells because of reduced expression of the cell cycle inhibitor p27^{kip}, increased activity of the kinase CDK6 and reduced expression of the anti-proliferative transcription factor Bmi1 [49,50]. Augmented effector gene re-expression in memory cells appears due to chromatin alterations at effector gene loci [51,52]. Finally, unlike naïve or effector cells, long-lived memory CD8⁺ T cells express receptors for both IL-7 (which promotes survival) and IL-15 (which drives antigen-independent proliferation and self-renewal) [17] leading to long-term maintenance of the memory population. In addition, memory cells express higher levels of factors that promote survival, such as Bcl-2 and Spi-2a, and both enhance memory cell formation and survival [38,53].

Cells with memory potential are evident early during the immune response, although they may not necessarily display all the hallmark qualities of a mature memory CD8⁺ T cell [54]. At the peak of the T cell response to certain acute infections, such as LCMV and *Listeria*, these memory precursor cells represent a small proportion of the effector cell population, and can be distinguished by higher IL-7 receptor (IL-7R) expression and lower expression of the senescence marker KLRG1 [34,55,56]. These IL-7R^{hi} KLRG1^{lo} effector CD8⁺ T cells demonstrate the highest rate of long-term survival, homeostatic turnover (via IL-15) and secondary responses to reinfection [34,55–57]. In contrast, most of the remaining effector CD8⁺ T cells are IL-7R^{lo}, express KLRG1, and do not persist or proliferate well following reinfection [34,56,57]. A population of KLRG1^{hi} IL-7R^{hi} memory CD8⁺ T cells can additionally be found, although these cells also display a finite lifespan [34,58].

How do effector CD8⁺ T cells with memory cell potential form during infection? Our past work suggested that a gradient of T-bet may be established in effector CD8⁺ T cells, and that high T-bet levels promote terminal effector cell differentiation and senescence, while low-

levels of T-bet permit memory precursor development [34]. The appropriate balance of T-bet expression appears directly tied to effector cell survival because low levels of T-bet are sufficient for IL-15R/IL-2R β (CD122) expression, but high levels of T-bet repress IL-7R α [34,57,59]. Collectively, this suggests that cells with intermediate T-bet levels will have optimal responses to both IL-7 and IL-15, the primary cytokines that sustain memory CD8⁺ T cells [60]. Recently, it was also found that CD8⁺ T cells undergo asymmetric cell divisions during the initial cell division [61] and can be routed down either short-lived or long-lived fates. Perhaps T-bet and other cell fate determining factors are unequally partitioned during these cell divisions, leading to different cell fates. However, other studies have suggested that activated CD8⁺ T cells are not necessarily committed to one or the other fate so quickly, as longer durations of infection promote terminal differentiation of CTLs [34,56,62] and IL-12 can augment T-bet expression [34,35]. Overall, this suggests that environmental signals may amplify or refine the effector cell diversity generated by asymmetric cell division. Establishing a gradient of a lineage-determining transcription factor (such as T-bet) through infection-inducible cytokine levels does not restrict effector T cells to simply two cell fates, but instead permits the broad spectrum of effector and memory differentiation states known to exist. Furthermore, it provides a model with built-in flexibility, whereby the innate immune system can control the magnitude and potency of the effector cell response during diverse infections (which can vary in tropism, intensity and duration). Blimp-1 likely also acts in a graded fashion [57] (R. L. Rutishauser and S. M. K., unpublished data), however it is presently unclear if other transcription factors act similarly.

Other factors can promote CD8⁺ T cell memory formation. The co-stimulatory molecule receptors 4-1BB, OX40 and CD27 can augment memory cell differentiation and survival [17,63]. It is also well established that help from CD4⁺ T cells is almost always required during the primary response for optimal memory CD8⁺ T cell formation [17]. While it is unclear what factor CD4⁺ T cells supply to CD8⁺ T cells, IL-2 is one candidate as it plays a key role in promoting memory T cell recall responses to secondary infection [64] and activated CD4⁺ T cells are an abundant source of this cytokine. In addition, CD4⁺ T cells may “condition” DCs to better induce memory formation [65]. Regardless of the mechanism, CD4⁺ T cell help appears to prevent the memory precursor population from undergoing terminal differentiation; “unhelped” cells display a higher proportion of T-bet^{hi} KLRG1^{hi} cells and T-bet deficiency can partially restore their function [57].

The memory CD8⁺ T cell population that eventually forms upon pathogen clearance is heterogeneous and evolves over time. Memory cells are often subdivided based on expression of the lymph-node homing receptors CD62L and CCR7 into CD62L^{lo} CCR7^{lo} non-lymphoid tissue resident effector memory cells (T_{EM}), and CD62L^{hi} CCR7^{hi} lymphoid tissue resident central memory cells (T_{CM}). In studies in mice, most T_{EM} and T_{CM} stem from IL-7R^{hi} memory precursor cells [55,66]. However, some KLRG^{hi} senescent effector cells can persist into early memory as “short-lived T_{EM}” [34,55,56] (Figure 1) generating further heterogeneity within the memory population. While CD8⁺ T_{CM} and T_{EM} both produce IFN- γ , T_{EM} generally display immediate cytotoxicity whereas T_{CM} typically produce more IL-2 and possess a higher proliferative potential [67–69]. Importantly, the antigen-specific memory T cell population composition changes over time, with T_{EM} gradually disappearing and giving way to T_{CM}, most likely due to slow outgrowth of the

T_{CM} [67,70]. Pathogen-specific memory T cells can persist in the tissues (such as the gut and lung) long-term [71,72], although these populations may be maintained by T_{CM} that migrate into tissues and form T_{EM} [73]. Transcription factors that control the T_{CM}/T_{EM} balance have also been identified. Activation of the PI3K/mTOR pathway down-regulates CCR7 and CD62L expression via the transcription factor Klf2 [74], suggesting metabolic control of the T_{CM}/T_{EM} balance. Furthermore, the transcription factors Id2, T-bet and Blimp-1 favour T_{EM} formation (R. L. Rutishauser and S. M. K., unpublished data) [40,57] while Bcl-6 promotes T_{CM} formation [75].

CONCLUSIONS

While many cell intrinsic and extrinsic factors have been identified that shape CD8⁺ T cell fate choices, it is still unclear how the large array of lineage determining factors synergise and/or integrate to fine-tune fate choices. In particular, understanding how different cell fates are maintained (particularly in the absence of antigen) and determining the extent of fate plasticity remains a future challenge. To this end, understanding the epigenetic establishment and maintenance of the transcriptional signatures underlying different CD8⁺ T cell fates represents an exciting area of future research.

REFERENCE LIST

1. Parish IA, Heath WR. Too dangerous to ignore: self-tolerance and the control of ignorant autoreactive T cells. *Immunol Cell Biol.* 2008; 86:146–152. [PubMed: 18227854]
2. Hernandez J, Aung S, Redmond WL, Sherman LA. Phenotypic and functional analysis of CD8(+) T cells undergoing peripheral deletion in response to cross-presentation of self-antigen. *J Exp Med.* 2001; 194:707–717. [PubMed: 11560988]
3. Schwartz RH. T cell anergy. *Annu Rev Immunol.* 2003; 21:305–334. [PubMed: 12471050]
4. Hernandez J, Aung S, Marquardt K, Sherman LA. Uncoupling of proliferative potential and gain of effector function by CD8(+) T cells responding to self-antigens. *J Exp Med.* 2002; 196:323–333. [PubMed: 12163561]
5. Parish IA, Waithman J, Davey GM, Belz GT, Mintern JD, Kurts C, Sutherland RM, Carbone FR, Heath WR. Tissue destruction caused by cytotoxic T lymphocytes induces deletional tolerance. *Proc Natl Acad Sci U S A.* 2009; 106:3901–3906. [PubMed: 19234128]
6. Davey GM, Kurts C, Miller JF, Bouillet P, Strasser A, Brooks AG, Carbone FR, Heath WR. Peripheral deletion of autoreactive CD8 T cells by cross presentation of self-antigen occurs by a Bcl-2-inhibitable pathway mediated by Bim. *J Exp Med.* 2002; 196:947–955. [PubMed: 12370256]
7. Tanchot C, Guillaume S, Delon J, Bourgeois C, Franzke A, Sarukhan A, Trautmann A, Rocha B. Modifications of CD8+ T cell function during in vivo memory or tolerance induction. *Immunity.* 1998; 8:581–590. [PubMed: 9620679]
8. Macian F, Garcia-Cozar F, Im SH, Horton HF, Byrne MC, Rao A. Transcriptional mechanisms underlying lymphocyte tolerance. *Cell.* 2002; 109:719–731. [PubMed: 12086671]
9. Safford M, Collins S, Lutz MA, Allen A, Huang CT, Kowalski J, Blackford A, Horton MR, Drake C, Schwartz RH, et al. Egr-2 and Egr-3 are negative regulators of T cell activation. *Nat Immunol.* 2005; 6:472–480. [PubMed: 15834410]
10. Heissmeyer V, Macian F, Im SH, Varma R, Feske S, Venuprasad K, Gu H, Liu YC, Dustin ML, Rao A. Calcineurin imposes T cell unresponsiveness through targeted proteolysis of signaling proteins. *Nat Immunol.* 2004; 5:255–265. [PubMed: 14973438]
11. Parish IA, Rao S, Smyth GK, Juelich T, Denyer GS, Davey GM, Strasser A, Heath WR. The molecular signature of CD8+ T cells undergoing deletional tolerance. *Blood.* 2009; 113:4575–4585. [PubMed: 19204323]

12. Tsushima F, Yao S, Shin T, Flies A, Flies S, Xu H, Tamada K, Pardoll DM, Chen L. Interaction between B7-H1 and PD-1 determines initiation and reversal of T-cell anergy. *Blood*. 2007; 110:180–185. [PubMed: 17289811]
13. Goldberg MV, Maris CH, Hipkiss EL, Flies AS, Zhen L, Tudor RM, Grosso JF, Harris TJ, Getnet D, Whartenby KA, et al. Role of PD-1 and its ligand, B7-H1, in early fate decisions of CD8 T cells. *Blood*. 2007; 110:186–192. [PubMed: 17392506]
14. Keir ME, Freeman GJ, Sharpe AH. PD-1 regulates self-reactive CD8+ T cell responses to antigen in lymph nodes and tissues. *J Immunol*. 2007; 179:5064–5070. [PubMed: 17911591]
15. Barron L, Knoechel B, Lohr J, Abbas AK. Cutting edge: contributions of apoptosis and anergy to systemic T cell tolerance. *J Immunol*. 2008; 180:2762–2766. [PubMed: 18292495] * First paper demonstrating a relationship between deletion and anergy in CD4⁺ T cells
16. Redmond WL, Marincek BC, Sherman LA. Distinct requirements for deletion versus anergy during CD8 T cell peripheral tolerance in vivo. *J Immunol*. 2005; 174:2046–2053. [PubMed: 15699134] * A study definitively showing that antigen dose regulates whether a CD8⁺ T cell adopts a fate of anergy or deletion
17. Williams MA, Bevan MJ. Effector and Memory CTL Differentiation. *Annu Rev Immunol*. 2006; 25:171–192. [PubMed: 17129182]
18. Prlic M, Hernandez-Hoyos G, Bevan MJ. Duration of the initial TCR stimulus controls the magnitude but not functionality of the CD8+ T cell response. *J Exp Med*. 2006; 203:2135–2143. [PubMed: 16908626]
19. Bachmann MF, Sebzda E, Kundig TM, Shahinian A, Speiser DE, Mak TW, Ohashi PS. T cell responses are governed by avidity and co-stimulatory thresholds. *Eur J Immunol*. 1996; 26:2017–2022. [PubMed: 8814240]
20. Zehn D, Lee SY, Bevan MJ. Complete but curtailed T-cell response to very low-affinity antigen. *Nature*. 2009; 458:211–214. [PubMed: 19182777] * A study demonstrating that low affinity antigens, despite inducing less CTL expansion, can promote effector and memory CD8⁺ T cell differentiation
21. Maekawa Y, Minato Y, Ishifune C, Kurihara T, Kitamura A, Kojima H, Yagita H, Sakata-Yanagimoto M, Saito T, Taniuchi I, et al. Notch2 integrates signaling by the transcription factors RBP-J and CREB1 to promote T cell cytotoxicity. *Nat Immunol*. 2008; 9:1140–1147. [PubMed: 18724371] * A study demonstrating that Notch signaling, and the transcription factors downstream of the Notch signal, promotes the acquisition of cytotoxic function in CD8⁺ T cells
22. Mescher MF, Curtsinger JM, Agarwal P, Casey KA, Gerner M, Hammerbeck CD, Popescu F, Xiao Z. Signals required for programming effector and memory development by CD8+ T cells. *Immunol Rev*. 2006; 211:81–92. [PubMed: 16824119]
23. Cousens LP, Peterson R, Hsu S, Dorner A, Altman JD, Ahmed R, Biron CA. Two roads diverged: interferon alpha/beta- and interleukin 12-mediated pathways in promoting T cell interferon gamma responses during viral infection. *J Exp Med*. 1999; 189:1315–1328. [PubMed: 10209048]
24. Thompson LJ, Kolumam GA, Thomas S, Murali-Krishna K. Innate inflammatory signals induced by various pathogens differentially dictate the IFN- γ dependence of CD8 T cells for clonal expansion and memory formation. *J Immunol*. 2006; 177:1746–1754. [PubMed: 16849484]
25. Nguyen KB, Watford WT, Salomon R, Hofmann SR, Pien GC, Morinobu A, Gadina M, O'Shea JJ, Biron CA. Critical role for STAT4 activation by type 1 interferons in the interferon-gamma response to viral infection. *Science*. 2002; 297:2063–2066. [PubMed: 12242445]
26. Cui W, Joshi NS, Jiang A, Kaech SM. Effects of Signal 3 during CD8 T cell priming: Bystander production of IL-12 enhances effector T cell expansion but promotes terminal differentiation. *Vaccine*. 2009; 27:2177–2187. [PubMed: 19201385]
27. Pearce EL, Shen H. Generation of CD8 T cell memory is regulated by IL-12. *J Immunol*. 2007; 179:2074–2081. [PubMed: 17675465]
28. Kolumam GA, Thomas S, Thompson LJ, Sprent J, Murali-Krishna K. Type I interferons act directly on CD8 T cells to allow clonal expansion and memory formation in response to viral infection. *J Exp Med*. 2005; 202:637–650. [PubMed: 16129706] ** This paper demonstrates that different types of infection generate diverse cytokine environments that are required for pathogen-specific CTL formation

29. Curtsinger JM, Schmidt CS, Mondino A, Lins DC, Kedl RM, Jenkins MK, Mescher MF. Inflammatory cytokines provide a third signal for activation of naive CD4⁺ and CD8⁺ T cells. *J Immunol.* 1999; 162:3256–3262. [PubMed: 10092777]
30. Sporri R, Reis e Sousa C. Inflammatory mediators are insufficient for full dendritic cell activation and promote expansion of CD4(+) T cell populations lacking helper function. *Nat Immunol.* 2005; 6:163–170. [PubMed: 15654341]
31. Haring JS, Badovinac VP, Harty JT. Inflaming the CD8⁺ T cell response. *Immunity.* 2006; 25:19–29. [PubMed: 16860754]
32. Pearce EL, Mullen AC, Martins GA, Krawczyk CM, Hutchins AS, Zediak VP, Banica M, DiCioccio CB, Gross DA, Mao CA, et al. Control of effector CD8⁺ T cell function by the transcription factor Eomesodermin. *Science.* 2003; 302:1041–1043. [PubMed: 14605368] ** Study identifying eomesodermin as a key regulator of effector CD8⁺ T cell differentiation
33. Sullivan BM, Juedes A, Szabo SJ, von Herrath M, Glimcher LH. Antigen-driven effector CD8 T cell function regulated by T-bet. *Proc Natl Acad Sci U S A.* 2003; 100:15818–15823. [PubMed: 14673093] ** First evidence that T-bet plays a role in inducing effector CD8⁺ T cell functions
34. Joshi NS, Cui W, Chande A, Lee HK, Urso DR, Hagman J, Gapin L, Kaech SM. Inflammation directs memory precursor and short-lived effector CD8(+) T cell fates via the graded expression of T-bet transcription factor. *Immunity.* 2007; 27:281–295. [PubMed: 17723218] * This study showed that, as effector CD8⁺ T cells differentiate, those with higher T-bet expression terminally differentiate and lose memory cell potential
35. Takemoto N, Intlekofer AM, Northrup JT, Wherry EJ, Reiner SL. Cutting Edge: IL-12 inversely regulates T-bet and eomesodermin expression during pathogen-induced CD8⁺ T cell differentiation. *J Immunol.* 2006; 177:7515–7519. [PubMed: 17114419]
36. Intlekofer AM, Banerjee A, Takemoto N, Gordon SM, Dejong CS, Shin H, Hunter CA, Wherry EJ, Lindsten T, Reiner SL. Anomalous type 17 response to viral infection by CD8⁺ T cells lacking T-bet and eomesodermin. *Science.* 2008; 321:408–411. [PubMed: 18635804]
37. Yoshida K, Sakamoto A, Yamashita K, Arguni E, Horigome S, Arima M, Hatano M, Seki N, Ichikawa T, Tokuhisa T. Bcl6 controls granzyme B expression in effector CD8⁺ T cells. *Eur J Immunol.* 2006; 36:3146–3156. [PubMed: 17125145] * Paper demonstrating that T-bet and eomesodermin are both required for effector CD8⁺ T cells to adopt a CTL phenotype
38. Liu N, Phillips T, Zhang M, Wang Y, Opferman JT, Shah R, Ashton-Rickardt PG. Serine protease inhibitor 2A is a protective factor for memory T cell development. *Nat Immunol.* 2004; 5:919–926. [PubMed: 15311278]
39. Zhang M, Park SM, Wang Y, Shah R, Liu N, Murmann AE, Wang CR, Peter ME, Ashton-Rickardt PG. Serine protease inhibitor 6 protects cytotoxic T cells from self-inflicted injury by ensuring the integrity of cytotoxic granules. *Immunity.* 2006; 24:451–461. [PubMed: 16618603]
40. Cannarile MA, Lind NA, Rivera R, Sheridan AD, Camfield KA, Wu BB, Cheung KP, Ding Z, Goldrath AW. Transcriptional regulator Id2 mediates CD8⁺ T cell immunity. *Nat Immunol.* 2006; 7:1317–1325. [PubMed: 17086188]
41. Barber DL, Wherry EJ, Masopust D, Zhu B, Allison JP, Sharpe AH, Freeman GJ, Ahmed R. Restoring function in exhausted CD8 T cells during chronic viral infection. *Nature.* 2006; 439:682–687. [PubMed: 16382236] ** The first study identifying that the exhausted state of CD8⁺ T cells is induced by expression of an inhibitory receptor, PD-1, on virus-specific CD8⁺ T cells during chronic infection
42. Blackburn SD, Shin H, Haining WN, Zou T, Workman CJ, Polley A, Betts MR, Freeman GJ, Vignali DA, Wherry EJ. Coregulation of CD8⁺ T cell exhaustion by multiple inhibitory receptors during chronic viral infection. *Nat Immunol.* 2009; 10:29–37. [PubMed: 19043418]
43. Brooks DG, Trifilo MJ, Edelmann KH, Teyton L, McGavern DB, Oldstone MB. Interleukin-10 determines viral clearance or persistence in vivo. *Nat Med.* 2006; 12:1301–1309. [PubMed: 17041596]
44. Ejrnaes M, Filippi CM, Martinic MM, Ling EM, Togher LM, Crotty S, von Herrath MG. Resolution of a chronic viral infection after interleukin-10 receptor blockade. *J Exp Med.* 2006; 203:2461–2472. [PubMed: 17030951]

45. Brooks DG, Ha SJ, Elsaesser H, Sharpe AH, Freeman GJ, Oldstone MB. IL-10 and PD-L1 operate through distinct pathways to suppress T-cell activity during persistent viral infection. *Proc Natl Acad Sci U S A*. 2008; 105:20428–20433. [PubMed: 19075244]
46. Wherry EJ, Ha SJ, Kaech SM, Haining WN, Sarkar S, Kalia V, Subramaniam S, Blattman JN, Barber DL, Ahmed R. Molecular signature of CD8⁺ T cell exhaustion during chronic viral infection. *Immunity*. 2007; 27:670–684. [PubMed: 17950003]
47. Hildeman DA, Zhu Y, Mitchell TC, Bouillet P, Strasser A, Kappler J, Marrack P. Activated T cell death in vivo mediated by proapoptotic bcl-2 family member bim. *Immunity*. 2002; 16:759–767. [PubMed: 12121658]
48. Pellegrini M, Belz G, Bouillet P, Strasser A. Shutdown of an acute T cell immune response to viral infection is mediated by the proapoptotic Bcl-2 homology 3-only protein Bim. *Proc Natl Acad Sci U S A*. 2003; 100:14175–14180. [PubMed: 14623954]
49. Heffner M, Fearon DT. Loss of T cell receptor-induced Bmi-1 in the KLRG1(+) senescent CD8(+) T lymphocyte. *Proc Natl Acad Sci U S A*. 2007; 104:13414–13419. [PubMed: 17686974]
50. Veiga-Fernandes H, Rocha B. High expression of active CDK6 in the cytoplasm of CD8 memory cells favors rapid division. *Nat Immunol*. 2004; 5:31–37. [PubMed: 14647273]
51. Araki Y, Fann M, Wersto R, Weng NP. Histone acetylation facilitates rapid and robust memory CD8 T cell response through differential expression of effector molecules (eomesodermin and its targets: perforin and granzyme B). *J Immunol*. 2008; 180:8102–8108. [PubMed: 18523274]
52. Northrop JK, Wells AD, Shen H. Cutting edge: chromatin remodeling as a molecular basis for the enhanced functionality of memory CD8 T cells. *J Immunol*. 2008; 181:865–868. [PubMed: 18606637] * (51 & 52) First papers to illustrate the role of epigenetic modifications in the phenotype of memory CD8⁺ T cells
53. Wojciechowski S, Tripathi P, Bourdeau T, Acero L, Grimes HL, Katz JD, Finkelman FD, Hildeman DA. Bim/Bcl-2 balance is critical for maintaining naive and memory T cell homeostasis. *J Exp Med*. 2007; 204:1665–1675. [PubMed: 17591857]
54. Kaech SM, Hemby S, Kersh E, Ahmed R. Molecular and functional profiling of memory CD8 T cell differentiation. *Cell*. 2002; 111:837–851. [PubMed: 12526810]
55. Kaech SM, Tan JT, Wherry EJ, Konieczny BT, Surh CD, Ahmed R. Selective expression of the interleukin 7 receptor identifies effector CD8 T cells that give rise to long-lived memory cells. *Nat Immunol*. 2003; 4:1191–1198. [PubMed: 14625547]
56. Sarkar S, Kalia V, Haining WN, Konieczny BT, Subramaniam S, Ahmed R. Functional and genomic profiling of effector CD8 T cell subsets with distinct memory fates. *J Exp Med*. 2008; 205:625–640. [PubMed: 18316415]
57. Intlekofer AM, Takemoto N, Kao C, Banerjee A, Schambach F, Northrop JK, Shen H, Wherry EJ, Reiner SL. Requirement for T-bet in the aberrant differentiation of unhelped memory CD8⁺ T cells. *J Exp Med*. 2007; 204:2015–2021. [PubMed: 17698591]
58. Rubinstein MP, Lind NA, Purton JF, Filippou P, Best JA, McGhee PA, Surh CD, Goldrath AW. IL-7 and IL-15 differentially regulate CD8⁺ T-cell subsets during contraction of the immune response. *Blood*. 2008; 112:3704–3712. [PubMed: 18689546]
59. Intlekofer AM, Takemoto N, Wherry EJ, Longworth SA, Northrup JT, Palanivel VR, Mullen AC, Gasink CR, Kaech SM, Miller JD, et al. Effector and memory CD8⁺ T cell fate coupled by T-bet and eomesodermin. *Nat Immunol*. 2005; 6:1236–1244. [PubMed: 16273099] * First study to show that low levels of T-bet are needed for optimal memory formation as T-bet controls CD122 expression
60. Joshi NS, Kaech SM. Effector CD8 T cell development: a balancing act between memory cell potential and terminal differentiation. *J Immunol*. 2008; 180:1309–1315. [PubMed: 18209024]
61. Chang JT, Palanivel VR, Kinjo I, Schambach F, Intlekofer AM, Banerjee A, Longworth SA, Vinup KE, Mrass P, Oliaro J, et al. Asymmetric T lymphocyte division in the initiation of adaptive immune responses. *Science*. 2007; 315:1687–1691. [PubMed: 17332376] ** Paper that proposed the asymmetric cell division theory of memory cell development
62. Badovinac VP, Harty JT. Manipulating the rate of memory CD8⁺ T cell generation after acute infection. *J Immunol*. 2007; 179:53–63. [PubMed: 17579021]

63. Mousavi SF, Soroosh P, Takahashi T, Yoshikai Y, Shen H, Lefrancois L, Borst J, Sugamura K, Ishii N. OX40 costimulatory signals potentiate the memory commitment of effector CD8⁺ T cells. *J Immunol.* 2008; 181:5990–6001. [PubMed: 18941188]
64. Williams MA, Tyznik AJ, Bevan MJ. Interleukin-2 signals during priming are required for secondary expansion of CD8⁺ memory T cells. *Nature.* 2006; 441:890–893. [PubMed: 16778891]
* This study demonstrated that IL-2 signalling is required for optimal memory CD8⁺ T cell formation
65. Smith CM, Wilson NS, Waithman J, Villadangos JA, Carbone FR, Heath WR, Belz GT. Cognate CD4(+) T cell licensing of dendritic cells in CD8(+) T cell immunity. *Nat Immunol.* 2004; 5:1143–1148. [PubMed: 15475958]
66. Sarkar S, Teichgraber V, Kalia V, Polley A, Masopust D, Harrington LE, Ahmed R, Wherry EJ. Strength of stimulus and clonal competition impact the rate of memory CD8 T cell differentiation. *J Immunol.* 2007; 179:6704–6714. [PubMed: 17982060]
67. Wherry EJ, Teichgraber V, Becker TC, Masopust D, Kaech SM, Antia R, von Andrian UH, Ahmed R. Lineage relationship and protective immunity of memory CD8 T cell subsets. *Nat Immunol.* 2003; 4:225–234. [PubMed: 12563257]
68. Masopust D, Vezys V, Marzo AL, Lefrancois L. Preferential localization of effector memory cells in nonlymphoid tissue. *Science.* 2001; 291:2413–2417. [PubMed: 11264538] ** First study to demonstrate that CD8⁺ T_{EM} in non-lymphoid tissues have greater CTL function than T_{CM}
69. Wolint P, Betts MR, Koup RA, Oxenius A. Immediate cytotoxicity but not degranulation distinguishes effector and memory subsets of CD8⁺ T cells. *J Exp Med.* 2004; 199:925–936. [PubMed: 15051762]
70. Marzo AL, Klonowski KD, Le Bon A, Borrow P, Tough DF, Lefrancois L. Initial T cell frequency dictates memory CD8⁺ T cell lineage commitment. *Nat Immunol.* 2005; 6:793–799. [PubMed: 16025119]
71. Kohlmeier JE, Miller SC, Woodland DL. Cutting edge: Antigen is not required for the activation and maintenance of virus-specific memory CD8⁺ T cells in the lung airways. *J Immunol.* 2007; 178:4721–4725. [PubMed: 17404250]
72. Masopust D, Vezys V, Wherry EJ, Barber DL, Ahmed R. Cutting edge: gut microenvironment promotes differentiation of a unique memory CD8 T cell population. *J Immunol.* 2006; 176:2079–2083. [PubMed: 16455963]
73. Marzo AL, Yagita H, Lefrancois L. Cutting edge: migration to nonlymphoid tissues results in functional conversion of central to effector memory CD8 T cells. *J Immunol.* 2007; 179:36–40. [PubMed: 17579018]
74. Sinclair LV, Finlay D, Feijoo C, Cornish GH, Gray A, Ager A, Okkenhaug K, Hagenbeek TJ, Spits H, Cantrell DA. Phosphatidylinositol-3-OH kinase and nutrient-sensing mTOR pathways control T lymphocyte trafficking. *Nat Immunol.* 2008; 9:513–521. [PubMed: 18391955] * Study showing that there is metabolic regulation of the T_{EM}/T_{CM} balance via mTOR induced Klf2
75. Ichii H, Sakamoto A, Kuroda Y, Tokuhisa T. Bcl6 acts as an amplifier for the generation and proliferative capacity of central memory CD8⁺ T cells. *J Immunol.* 2004; 173:883–891. [PubMed: 15240675]

Figure 1. A model for generating diverse CD8⁺ T cell fates

(Top portion) Naïve CD8⁺ T cells (brown cell) that encounter antigen on steady-state, tolerogenic DCs (beige cell) proliferate without acquiring effector functions (dark blue cells) in a process dependent on PD-1 and the molecules Egr2, Egr3 (Egr2/3) and Cbl-b. Such tolerised T cells typically up-regulate PD-1 and down-regulate IL-7R as indicated, and either die (black cell) or become anergic (i.e., deficient in TCR signalling as denoted by the red cross). Antigen (Ag) levels control this process, with high Ag levels promoting anergy, and low Ag levels causing deletion. (Lower portion) Immunogenic DCs (red cell), which have encountered pathogen derived products, activate naïve CD8⁺ T cells to form effector cells (cytotoxic granules and production of effector cytokines are indicated). This differentiation process is regulated by molecules such as Id2, Spi-6, Notch, T-bet and eomesodermin (Eomes). These effector cells can adopt multiple cell fates, such as memory precursor cell fates (characterised by IL-7R^{hi}, KLRG1^{lo}, T-bet^{lo}, Blimp-1^{int}, Spi-2a^{hi} and Bcl-2^{hi} expression; bottom left) or short-lived effector cell fates (with IL-7R^{lo}, KLRG1^{hi}, T-bet^{hi} and Blimp-1^{hi} expression; middle). Such short-lived cells can either become senescent effectors and die by bim-dependent apoptosis, or persist into early memory as short-lived

effector memory cells that are $CCR7^{lo}$, $CD62L^{lo}$ and $KLRG1^{hi}$. In contrast, memory precursor cells are long-lived and can be $CCR7^{lo}$, $CD62L^{lo}$, $Blimp-1^{int}$ effector memory cells (promoted by Id2, Klf2 and Blimp-1), or $CCR7^{hi}$, $CD62L^{hi}$, $Blimp-1^{lo}$ central memory cells (driven by Bcl-6). Evidence exists that $KLRG1^{lo}$ effector cells can also give rise to $KLRG1^{hi}$ short-lived effector memory cells (denoted by broken arrow and question marks; N. S. Joshi, T. W. Hand and S. M. K., unpublished observations). In some chronic viral infections, PD-1, LAG-3 and IL-10 causes cells to acquire an exhausted phenotype (light blue cell) characterised by high PD-1 and LAG-3 expression and low IL-7R levels. Although the exhausted cell is depicted as sharing a common effector cell precursor with other cell fates, it is currently unclear at what point during effector cell differentiation this fate branches.