Published in final edited form as:

Angew Chem Int Ed Engl. 2013 December 16; 52(51): 13576–13579. doi:10.1002/anie.201305711.

[2+2+2] Cycloadditions of Siloxy Alkynes with 1,2-Diazines: From Reaction Discovery to Identification of an Antiglycolytic Chemotype**

Dr. Timothy J. Montavon, Dr. Yunus E. Türkmen, Noumaan A. Shamsi, Christopher Miller, Chintan S. Sumaria, Prof. Dr. Viresh H. Rawal, and Prof. Dr. Sergey A. Kozmin

Chicago Tri-Institutional Center for Chemical Methods and Library Development, Department of Chemistry, The University of Chicago, Chicago, IL 60637 (USA)

Viresh H. Rawal: vrawal@uchicago.edu; Sergey A. Kozmin: skozmin@uchicago.edu

Abstract

A newly uncovered Brønsted acid-promoted [2+2+2] cycloaddition between siloxy alkynes and 1,2-diazines produces novel polycyclic compounds with high efficiency and excellent diastereoselectivity under exceedingly mild conditions. A small-molecule library synthesized using this reaction yielded a novel chemotype, which inhibited glycolytic ATP production by blocking glucose uptake in CHO-K1 cells.

Keywords

Cycloaddition; Polycycles; Siloxyalkynes; Warburg Effect; Glycolysis Inhibitor

The synthesis of new nitrogen-containing heterocycles plays a pivotal role in chemical biology and medicinal chemistry, as reflected by their many applications in the development of pharmacological probes and drugs.^[1] Despite notable progress, there is a significant need for the identification of new nitrogen heterocycles that target previously unexplored regions of biogenic chemical space. Among the many possible synthetic strategies to such compounds, cycloadditions involving C-N multiple bonds are particularly attractive as they generate complex cyclic products via simultaneous formation of multiple bonds starting from readily available precursors.^[2] In this communication, we describe the discovery and development of a formal [2+2+2] cycloaddition of siloxy alkynes with phthalazines – a process that had not been previously described for either 1,2-diazines or electron-rich alkynes.^[3–7] This effort has not only afforded heterocyclic products with a unique pentacyclic ring system but has also enabled the identification of a novel chemotype that inhibits glycolytic ATP production by direct blockage of glucose uptake in CHO-K1 cells. Due to the prevalence of the Warburg effect in many human cancers, such compounds may

**We are grateful for financial support to the National Institutes of Health (P50 GM086145 and R01GM069990), the Chicago Biomedical Consortium with support from the Searle Funds at the Chicago Community Trust. We thank Dr. Ian Steele for X-ray crystallographic analysis of 31.

Correspondence to: Viresh H. Rawal, vrawal@uchicago.edu; Sergey A. Kozmin, skozmin@uchicago.edu.

Supporting information for this article is available on the WWW under [link place holder]

prove useful in the development of new therapeutics that target re-programmed energy metabolism of rapidly proliferating cells.^[8]

Our study began by examining the reaction of phthalazine **1** with siloxy alkyne **2** in the presence of common Brønsted acids. While no reaction between **1** and **2** was observed in the absence of such additives, even at elevated temperatures, we found that addition of simple pyridinium salts promoted the formation of a new pentacyclic product **3** (Scheme 1).

After examining a range of mono- and bispyridinium salts in various solvents, we determined the optimum protocol to entail the use of a stoichiometric amount of pyridinium trifluoromethanesulfonimide in CH₂Cl₂ at room temperature, which produced lactam **3** as a single diastereomer in 77% yield. While most of the known [2+2+2] cycloadditions typically require the presence of a transition metal catalyst,^[9] the present method promotes the condensation under remarkably mild reaction conditions, using only a simple, weak Brønsted acid. The excellent diastereoselectivity of this transformation is also highly noteworthy. The atom connectivity within the reaction product was initially determined to be that in lactam **3** based on extensive use of NMR spectroscopic methods. Ultimately, the structure was secured and the relative stereochemistry of the three newly created stereogenic centers was defined through X-ray crystallographic analysis (*vide infra*).

Interestingly, while a range of substituted mono and bispyridinium trifluoromethanesulfonimides were found to be effective as reaction promoters, the use of only HNTf₂, in the absence of pyridine, produced lactam **3** with lower efficiency (48% yield) and diminished diastereoselectivity (83:17). Furthermore, the use of either pyridinium chloride, pyridinium *p*-toluenesulfonate or pyridinium triflate substantially decreased product yields or completely abolished the reaction. These results highlight the importance of the weakly nucleophilic trifluoromethanesulfonimide counterion,^[10] an observation consistent with those made in the course of previous studies of Brønsted acid-promoted transformations of siloxy alkynes.^[3]

Having established a general reaction protocol, we began a detailed investigation of the scope of this [2+2+2] cycloaddition. With regards to siloxy alkyne substitution, we found that both alkyl and aryl substituents are well tolerated, providing the expected products in good yields with high levels of diastereoselection (Table 1, Entries 1–4). When the steric bulk of the substituent in direct proximity to the alkyne is increased, the yield of the reaction is lowered slightly but the diastereoselectivity remains relatively unaffected. For instance, siloxy alkynes **6** and **8** afforded the expected products **7** and **9** in 73% and 69% isolated yields, respectively (Table 1, Entries 2 and 3). Furthermore, the use of 1-siloxy-propyne **10** resulted in efficient formation of lactam **11** with high diastereoselectivity (d.r. > 98:2). Taken together, these results suggest that a wide range of siloxy alkyne substituents would be well tolerated in this reaction.

The scope of the reaction was next evaluated with respect to a series of substituted 1,2-diazine derivatives. For instance, benzo[g]phthalazine **12** provided the expected product **13** in good yield, with high diastereoselection (d.r. > 98:2, Table 1, Entry 5). The reaction of alkyne **2** with 6,7-dimethyl phthalazine **14** gave lactam **15** in 65% yield and 91:9

diastereoselectivity (Table 1, Entry 6). The lower efficiency of this reaction may be attributed to the decreased electrophilicity of the phthalazine moiety resulting from the presence of the two alkyl substituents. Both the halogen-containing and heteroaromatic diazines **16** and **18** served as excellent substrates in the [2+2+2] cycloaddition, providing the expected products **17** and **19** in 73% and 81% yields, respectively (Table 1, Entries 7 and 8). Finally, 1-phenyl phthalazine **20** was found to participate in this transformation in a regio- and diastereoselective manner, providing a single [2+2+2] adduct in 61% yield (Table 1, Entry 9). Simple pyridazines were unreactive under the current reaction conditions. This observation is consistent with the lower reactivity of pyridazines compared to that of phthalazines in other reactions, including [4+2] cycloadditions with electron-rich dienophiles.

We next examined a possibility of conducting a three-component reaction using two different 1,2-diazines. Reaction of siloxy alkyne **2** with phthalazine **1** and 1-phenyl phthalazine **20** delivered roughly equal quantities of all four possible products (Table 2, Entry 1), as expected given the small electronic perturbation provided by the phenyl substituent C-2. However, treatment of siloxy alkyne **2** with phthalazine **1** and 6,7-dichlorophthalazine **16** with pyridinium trifluoromethanesulfonimide afforded predominantly the hybrid product **24**, as well as smaller quantities of cycloadducts **17** and **25** (Table 2, entry 2). Finally, the use of benzo[g]phthalazine **12** in a similar three-component reaction with **1** and **2** resulted in the formation of only one of the hybrid products, **26**, as the major product along with the cycloadduct **13** (Table 2, entry 3). These results highlight the unique ability of this process to discriminate between two similar but electronically differentiated 1,2-diazine moieties.

Two mechanistic pathways can explain the formation of pentacyclic lactams originating from [2+2+2] cycloadditions of phthalazines with siloxy alkynes (Scheme 2). The reaction can proceed via initial formation of ketenium ion **28** via protonation of siloxy alkyne by the pyridinium salt (pathway A). This electrophilic intermediate can be intercepted by phthalazine **1** to give enol silane **29**, a heterodiene with nucleophilic and electrophilic carbons at opposite ends. Addition of a second phthalazine moiety to **29** would trigger cyclization of the central ring, to afford lactam **32** upon protodesilylation. Alternatively, the process can begin with protonation of phthalazine by the pyridinium catalyst (pathway B). The resulting phthalazinium ion would then react with alkyne **32** to give the ketenium ion **33**. Interception of this intermediate by another phthalazine **1** would give ammonium ion **34**, which can trigger protodesilylation of the enol silane moiety to give the observed product **31**. Both mechanistic pathways are consistent with the results of deuterium labeling experiments performed using deuterio-pyridinium triflate, which resulted in exclusive deuterium incorporation at the α carbon of the *N*-acyl hydrazone moiety.^[11] Further mechanistic studies are required in order to establish which of the two reaction pathways affords the final cycloaddition product.

To demonstrate further the synthetic utility of the novel structural types obtained by cycloadditions of siloxy alkynes and 1,2-diazines, we have developed a route to selectively functionalize the *N*-acyl hydrazone moiety of the observed [2+2+2] adducts (Scheme 3). The two diazine moieties of pentacyclic lactam **3** are chemically differentiated, one a

hydrazone and the other an *N*-acyl hydrazone. While hydrazones are notoriously unreactive towards nucleophilic additions, *N*-acyl hydrazones are quite reactive and have proven to be valuable substrates for the synthesis of diverse classes of nitrogen-containing compounds.^[12] We found that treatment of **3** with a slight excess of allyl trichlorosilane resulted in highly chemo- and diastereoselective allylation at the azomethine carbon of the *N*-acyl hydrazone moiety, providing allylated product **35** in 86% yield.^[13] Treatment of **35** with either benzoyl chloride or benzyl isocyanate afforded the expected acylation products, amide **36** or hydrazide **37**, respectively. Hydrazide **37** was highly crystalline and its structure was determined by X-ray crystallography, which provided unambiguous stereochemical assignment of the initial cycloadduct **3** and subsequent allylation product **35**.

We next employed this chemo- and diastereoselective elaboration strategy to assemble a representative collection of 56 structurally diverse compounds,^[11] which was then evaluated for their ability to inhibit aerobic glycolysis in cancer cells. To this end, we employed our recently developed screening approach, which is based on monitoring glycolytic ATP production in live cells with chemically impaired mitochondria.^[14] We performed the screen by subjecting antimycin A treated CHO-K1 cells to each member of the newly assembled chemical library^[11] and monitored production of ATP using a well-established luciferase-based protocol. The structure of the most potent inhibitor **38** is shown in Figure 1A.

This compound elicited the expected dose-dependent ATP suppression profile shown in Figure 1B. The ATP production was inhibited much more efficiently when cells were treated with a combination of **38** and antimycin A ($IC_{50} = 11.4 \mu M$). To further validate inhibition of aerobic glycolysis, we measured the effects of **38** on lactate production, which was found to decrease in a dose dependent manner (Fig. 1C). Finally, we found that **38** inhibited cellular uptake of 3H labelled 2-deoxy glucose with $IC_{50} = 2.2 \mu M$ (Fig. 1D), confirming the antiglycolytic effect of this compound.

In summary, we have described highly efficient and diastereoselective [2+2+2] cycloadditions of siloxy alkynes with phthalazines. This synthetic process is promoted under mild conditions by simple pyridinium salts and provides rapid access to new, complex heterocyclic products, richly adorned with handles for further functionalization. Cellular screen of a representative collection of such structurally diverse compounds enabled identification of a new chemical inhibitor of aerobic glycolysis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. Pozharskii, A.F.; Soldatenkov, A.; Katritzky, A.R. *Heterocycles in Life and Society*. John Wiley & Sons, Inc; Hoboken: 2011.
2. Hassner, A., editor. *Topics in Heterocyclic Chemistry*, Vol 12: Synthesis of Heterocycles via Cycloadditions I. Springer; Berlin: 2008. (b) Hassner, A., editor. *Topics in Heterocyclic Chemistry*, Vol 13: Synthesis of Heterocycles via Cycloadditions II. Springer; Berlin: 2010.
3. For other Brønsted acid-promoted reactions of siloxy alkynes, see: Zhang L, Kozmin SA. *J Am Chem Soc*. 2004; 126:10204–10205. [PubMed: 15315407] Sun J, Kozmin SA. *J Am Chem Soc*.

- 2005; 127:13512–13513. [PubMed: 16190711] Zhcang L, Sun J, Kozmin SA. *Tetrahedron*. 2006; 62:11371–11380.
4. For review of ynolates and siloxy alkynes, see: Shindo M. *Tetrahedron*. 2007; 63:10–36. For reviews of ynamides and ynamines, see: Zifcsak CA, Mulder JA, Hsung RP, Rameshkumar C, Wei LL. *Tetrahedron*. 2001; 57:7575–7606. Evano G, Coste A, Jouvin K. *Angew Chem Int Ed*. 2010; 49:2840–2859.
5. For other cycloadditions of alkoxy and siloxy alkynes, see: Danheiser RL, Gee SK. *J Org Chem*. 1984; 49:1672. Kowalski CJ, Lal GS. *J Am Chem Soc*. 1988; 110:3693–3695. Sweis RF, Schramm MP, Kozmin SA. *J Am Chem Soc*. 2004; 126:7442–7443. [PubMed: 15198582] Clark TB, Woerpel KA. *Org Lett*. 2006; 8:4109–4112. [PubMed: 16928086] Qi X, Ready JM. *Angew Chem Int Ed*. 2008; 47:7068–7070. Türkmen YE, Montavon TJ, Kozmin SA, Rawal VH. *J Am Chem Soc*. 2012; 134:9062–9065. [PubMed: 22607029]
6. For a review of cycloadditions of heterocyclic aza dienes, see: Boger DL. *Tetrahedron*. 1983; 39:2869. Boger DL. *Chem Rev*. 1986; 86:781–793.
7. For cycloadditions of 1,2-diazines with two equivalents of ynamines, see: Oishi E, Taido N, Miyashita A, Sato S, Ohta S, Ishida H, Tanji K, Higashino T. *Chem Pharm Bull*. 1991; 39:1713–1718. Oishi E, Iwamoto K-T, Okada T, Suzuki S, Tanji K, Miyashita A, Higashino T. *Chem Pharm Bull*. 1994; 42:2219–2224.
8. Vander Heiden MG, Cantley LC, Thompson CB. *Science*. 2009; 324:1029–1033. [PubMed: 19460998]
9. a) Chopade PR, Louie J. *Adv Synth Catal*. 2006; 348:2307–2327. b) Shibata T, Tsuchikama K. *Org Biomol Chem*. 2008; 6:1317–1323. [PubMed: 18385836] c) Dominguez G, Pérez-Castells J. *Chem Soc Rev*. 2011; 40:3430–3444. [PubMed: 21431173]
10. Antonietti S, Dalla V, Duñach E. *Angew Chem Int Ed*. 2010; 49:7860–7888.
11. For details, see Supporting Information.
12. For recent reviews on hydrazones and N-acyl hydrazones see Sugiura M, Kobayashi S. *Angew Chem Int Ed*. 2005; 44:5176–5186. Friestad GK. *Eur J Org Chem*. 2005:3157–3172. Brehme R, Enders D, Fernandez R, Lassaletta JM. *Eur J Org Chem*. 2007:5629–5660. Lazny R, Nodzevska A. *Chem Rev*. 2010; 110:1386–1434. [PubMed: 20000672]
13. a) Kobayashi S, Hirabayashi R. *J Am Chem Soc*. 1999; 121:6942–6943. b) Hirabayashi R, Ogawa C, Sugiura M, Kobayashi S. *J Am Chem Soc*. 2001; 123:9493–9499. [PubMed: 11572669] c) Ogawa C, Sugiura M, Kobayashi S. *J Org Chem*. 2002; 67:5359–5364. [PubMed: 12126428]
14. Ulanovskaya O, Cui J, Kron SJ, Kozmin SA. *Chem Biol*. 2011; 18:222–230. [PubMed: 21338919]

Figure 1.

a. Chemical structure of **38**. b. Inhibition of cellular ATP levels in PC3 cells by **38** in the presence or absence of Complex III inhibitor, antimycin A c. Inhibition of lactate production in A549 cells upon treatment with **38**. Cytochalasin B was used as a positive control d. Inhibition of glucose uptake in CHO-K1 cells by **38** using ^3H 2-DG scintillation experiments. All values are presented as percentage of vehicle treated samples. Each value is the mean SEM of duplicate values from a representative experiment. 2-DG = 2-deoxy glucose.

Scheme 1.

[2+2+2] Cycloaddition of Phthalazine (1) with a 1-Siloxy-1-Hexyne (2). TIPS = triisopropylsilyl; Tf = trifluoromethanesulfonyl; Me = methyl.

Scheme 3.

Subsequent Chemoselective Functionalization. Bn = benzyl; DMF = dimethylformamide.

Table 1

Scope of the [2+2+2] Cycloaddition of Phthalazines and Siloxy Alkynes

Entry	1,2-Diazine	Alkyne	Product	Yield, % ^[a]	d.r. ^[b]
1				76	98:2
2				73	98:2
3				69	98:2
4				77	>98:2

Reaction scheme showing the synthesis of a 1,2,3,4-tetrahydropyridine derivative from a 1,2-diazine, an alkyne, and a pyridine derivative, catalyzed by NTf_2^+ (100 mol%) in CH_2Cl_2 at 20 °C.

Entry	1,2-Diazine	Alkyne	Product	Yield, % ^[a]	d.r. ^[b]
5				74	>98:2
6				65	91:9
7				73	>98:2
8				81	>98:2

Entry	1,2-Diazine	Alkyne	Product	Yield, % ^[a]	d.r. ^[b]
9				61	>98:2

^[a] Yields refer to those of the major isolated diastereomer of each reaction.

^[b] Diastereomeric ratios were determined by 500 MHz ¹H NMR analysis of the crude reaction mixture prior to purification. Ph = phenyl.

Table 2

Exploring Three-Component [2+2+2] Cycloadditions of Phthalazines and Siloxy Alkynes

Entry	1,2-Diazine	Product Ratio: A:B:C:D ^a	Yield, % ^b
1		3:21:22:23 = 29:21:23:27	56
2		3:17:24:25 = 5:19:48:28	43
3		3:13:26:27 = 0:33:67:0	68

^a Molar ratios based on yields of individually isolated product isomers based on tentative structural assignments.^b Combined isolated yield (over 48 hours) for reaction performed on a 0.2 mmol scale. For all sets of products, only one diastereomer was found to be present by 500 MHz ¹H NMR analysis of the crude reaction mixture. *n*-Bu = *n*-Butyl.