Published in final edited form as:

Chem Commun (Camb). 2014 May 25; 50(40): 5391–5393. doi:10.1039/c3cc47384h.

A Dual Site Catalyst for Mild, Selective Nitrile Reduction

Zhiyao Lu and Travis J. Williams

Loker Hydrocarbon Research Institute and Department of Chemistry, University of Southern California, Los Angeles, California 90089-1661

Travis J. Williams: travisw@usc.edu

Abstract

We report a novel ruthenium bis(pyrazolyl)borate scaffold that enables cooperative reduction reactivity in which boron and ruthenium centers work in concert to effect selective nitrile reduction. The pre-catalyst compound $\{[\kappa^3\text{-(1-pz)}_2\text{HB(N=CHCH}_3\text{)}]\text{Ru(cymene)}\}^+\text{TfO}^-$ (pz = pyrazolyl) was synthesized from readily-available materials through a straightforward route, thus making it an appealing catalyst for a number of reactions.

As part of our ongoing studies of dual site ruthenium, boron-containing catalysts for the manipulation of hydride groups, we have recently reported a series of [di(pyridyl)borate]ruthenium complexes (**1**, **2**)ⁱ that exhibit remarkable reactivity in a number of applications,ⁱⁱ notably including dehydrogenation of ammonia borane.ⁱⁱⁱ Although they are successful catalysts, these di(pyridyl)dimethylborate-derived complexes are cumbersome to prepare, due largely to dependence on an expensive and reactive BrBMe₂ starting material and high water and oxygen sensitivity of intermediate complexes in their syntheses. Further, despite their catalytic utility, no direct evidence has been collected to show a mechanistic account of the cooperative role, if any, that boron and ruthenium are playing in the reactive mechanisms of **1** or **2**.^{iv} We suspect this is partially due to the robustness of the bridging $\mu\text{-OH}$ ligand between the boron and ruthenium centers, which inhibits access to a free borane in catalytic reactions.

We show here a conveniently prepared, borate-pendant ruthenium complex (**3**) that retains much of the reactivity of the original di(pyridyl)borate complexes (see ESI), show that it is an efficient and selective catalyst for nitrile reduction, and provide evidence for the cooperative role that boron and ruthenium play in the reaction, as hydride donor and activating group, respectively.

The synthesis of **3** (Scheme 1) proceeds from potassium di(pyrazolyl)borohydride^{v, vi, vii} and commercially-available (cymene)ruthenium dichloride dimer to give intermediate chloride **4**. Although **4** can be isolated, it is easily converted in situ to **3** by treatment with 1 equiv. thallium (or silver) triflate in nitrile solution. The synthesis proceeds in two smooth

steps without the need for materials that are cost-prohibitive or difficult to manipulate: all materials are amenable to handling using standard Schlenk techniques and/or a glove box. **3** can be crystallized from isopropanol and hexanes; its molecular structure was determined by single crystal X-ray diffraction (Scheme 1). The crystal structure of **3** shows the borate ligand in a tetrahedral geometry at boron, which has analogy to the popular tris(pyrazolyl)borohydride (Tp) ligand series.^{vi, vii, viii}

The synthesis of **3** revealed an important insight into the mechanism of its catalytic reactivity. In the conversion of **4** to **3**, a hydride is transferred from a ligand B-H group to **5**'s coordinated nitrile in >90% NMR yield (Scheme 1). This reaction is the first example of our envisioned cooperative reactivity of ruthenium and boron. Contrary to the design concept from which we originally prepared **2**,ⁱⁱ boron in **3** is not behaving as a Lewis acid. The structure of **3** shows that B-H addition to the nitrile proceeds in a cis fashion, and NMR evidence reveals that the selectivity for this geometry is exclusive. Thus we believe that the mechanism for this reaction involves intramolecular hydride transfer from boron to carbon, rapidly followed by (or concerted with) boron-nitrogen coordination.

The stoichiometric reduction of acetonitrile observed in the synthesis of **3** can be made catalytic by treating **3** with nitrile and sodium borohydride,^{ix} thus enabling a mild and selective approach to the synthesis of primary amines from nitriles, which remains a contemporary topic in bifunctional catalysis.^x Known methods for nitrile reduction, e.g. excess LiAlH₄, borohydride reduction of a nitrilium salt,^{xi} metal hydride reagents,^{xii} or high-pressure hydrogenation^{xiii} can be incompatible with important synthetic handles, such as aryl bromide and nitro.^{xiv} More mild conditions compatible with groups like this require a stoichiometric portions a borane reagent,^{xv} which in some cases must be independently prepared. The new catalytic mechanism reported here enables high functional group tolerance while incorporating an inexpensive reducing agent.

Table 1 shows the discovery and optimization of our conditions for nitrile reduction. In presence of 5 mol% **3**, 2.0 equiv. NaBH₄ can reduce 4-trifluoromethylbenzonitrile (**7a**) to the corresponding benzylamine (**8a**) in 50% conversion by NMR in 5 hours (entry 1). When 1.0 equiv. of NaO^tBu was added, the analogous reaction reached >90% yield (>95% conversion) in the same time. With no catalyst, the reaction was much slower, reaching completion in 7 days with a 43% overall yield (entry 5). If **3**'s Tp-ligated homologue (**6**, Scheme 1) is used in this reaction, much of the starting material decomposed under the reaction conditions. This illustrates that the μ -acetimine ligand in **3** plays an essential role in nitrile reduction.

The substrate scope of nitriles that can be reduced using our optimized conditions is broad. As shown in table 2, aromatic, aliphatic, and heterocyclic nitriles are smoothly reduced to amines under optimized conditions. Both electron poor and electron rich substrates can be reduced in high yield. For example, In presence of 5 mol% **3**, 4.0 equiv. NaBH₄ can reduce 4-trifluorobenzonitrile **7a** to the corresponding benzylamine **8a** in 82% isolated yield (entry 1). Electron rich nitrile **7b** can be reduced with similar facility in 87% yield (entry 2). Even more oxidative nitroarene **7c** can be reduced with exclusive selectivity for the nitrile over the nitro group (92%, entry 3). We believe that this startling result is attributable to selective

binding and activation of the nitrile over the nitro by the catalyst's ruthenium center. Furthermore, we see this as additional evidence that nitrile binding to ruthenium is important to the catalytic mechanism.

Despite the presence of highly reducing conditions, an aryl bromide group in nitrile **7d** is not derivitized in the course of amine synthesis. This is an important result because aryl bromides such as this are high value substrates for cross-coupling and amination reactions relevant to the synthesis of medicinally-relevant compounds.^{xvi} This example further illustrates high-yielding reduction of an alkyl nitrile. Whereas ketone groups are known to react with NaBH₄, the reduction of **7e** (entry 5) illustrates high-yielding double reduction for the synthesis of aminoalcohol **8e**. Similar double reduction is observed in the reaction of cinnamionitrile (**7g**, entry 7) to give alkyl amine **8g** in 60% yield.

Reactions of nitriles appended to aromatic heterocycles afforded complicated results. For example, pyridine **7h** is compatible with the conditions and resulted in the formation of aminomethylpyridine **8h** in 64% yield (entry 7). By remarkable contrast, more electron rich heterocycle systems are not reduced. For example, 2-cyanofuran **7i** and 2-cyanothiophene **7j** are selectively monohydrated as opposed to reduced. Thus, amides **9ij** were isolated as the main products (entries 9 and 10). We suspect that the mechanism for these reactions involves the addition of methanol (solvent) to a ruthenium-coordinated nitrile, and that the amide products are formed upon aqueous work-up. We do not currently have a proposal to account for the selectivity of hydration versus reduction.

According to the insight gained from the stoichiometric synthesis of **3** from **4**, we propose the following template mechanism for catalysis (Scheme 2). We suspect that the bridging imine **3** is reduced by borohydride to produce the amine product. We do not have direct evidence for the intermediacy of **10**; however, treatment of **3** with a stoichiometric portion of NaBH₄ in methanol-*d*₄ results in clean desymmetrization of the catalyst's *cymene* and pyrazole C—H groups, consistent with the formation of diastereotopic protons, as expected with the pyramidalization of the bridging nitrogen ligand (see ESI for graphical spectra). Still, the intermediacy of **10** remains a proposal because we have not established the kinetic role of this transient material.

We propose that the bridging amine ligand is replaced by incoming substrate, and the borohydride group of **5** is regenerated by a hydride from NaBH₄, although we do not know the details of these steps. We do observe that treatment of **3** with a stoichiometric portion of NaBH₄ in methanol-*d*₄ results in the formation of (MeO)₄B[−], unreacted BH₄[−] and a catalyst doublet, by ¹H-coupled ¹¹B NMR, which indicates a (pz)₂BH₂ intermediate, if formed, is transient. Thus, we suspect that X in Scheme 2 is methoxide. ¹H NMR studies of the working catalyst reveal that once ligated, the reductions of the nitrile groups and imine groups are very facile. Thus, the rate-determining step could be amine for nitrile substitution of the nitrile substrate to the ruthenium center. This mechanism is a subject of ongoing work in our laboratory.

In conclusion we report here a conveniently-prepared homologue of our successful di(pyridyl)borate-ligated ruthenium complexes. The new catalytic scaffold has comparable

reactivity to the old in several key reactions and introduces new reactivity to the cooperative ruthenium, boron catalytic motif by enabling selective and high yielding nitrile reduction under mild conditions. Furthermore, this platform has yielded new insight into the cooperative reactivity of ruthenium and boron by showing a plausible scenario of how these two centers can work together respectively as activating group (ruthenium) and hydride donor (boron). Ongoing work in our laboratory regards the application of this system to the reduction of other high-value pi systems and the elucidation of the mechanistic details of these reactions.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work is supported by the University of Southern California, the Hydrocarbon Research Foundation, and the National Science Foundation (CHE-1054910). We thank Ralf Haiges for X-ray crystallography. We thank the NSF (DBI-0821671, CHE-0840366, CHE-1048807) and NIH (S10 RR25432) for analytical instrumentation.

Notes and references

- i. Conley BL, Williams TJ. *J Am Chem Soc.* 2010; 132:1764–1765. [PubMed: 20088526]
- ii. Conley BL, Williams TJ. *Comments Inorg Chem.* 2011; 32:195–218.
- iii. Conley BL, Guess D, Williams TJ. *J Am Chem Soc.* 2011; 133:14212–14215. [PubMed: 21827173]
- iv. Lu Z, Malinoski B, Flores AV, Guess D, Conley BL, Williams TJ. *Catalysts.* 2012; 2:412–421.
- v. Trofimenko, S., editor. *Scorpionates: The Coordination of Polypyrazolylborate Ligands.* Imperial College Press; London: 1999.
- vi. For recent reviews of organometallic chemistry of poly(pyrazolyl)borate ligated complexes see: Caldwell LM. *Adv Organomet Chem.* 2008; 56:1–94. Lail M, Pittard KA, Gunnoe TB. *Adv Organomet Chem.* 2008; 56:95–153. Becker E, Pavli S, Kirchner K. *Adv Organomet Chem.* 2008; 56:155–197. Crossly IR. *Adv Organomet Chem.* 2008; 56:199–321.
- vii. Abernethy RJ, Hill AF, Smith MK, Willis AC. *Organometallics.* 2009; 28:6152–6159.
- viii. Bhambri S, Tocher DA. *J Chem Soc, Dalton Trans.* 1997:3367–3372.
- ix. A ruthenium-ligated acetonitrile was reduced by NaBH₄, however the system produced unidentified products. 13: Rhodes LF, Venanzi LM. *Inorg Chem.* 1987; 26:2692–2695.
- x. Vogt M, Nerush A, Iron MA, Leituss G, Diskin-Posner Y, Shimon LJW, Ben-David Y, Milstein D. *J Am Chem Soc.* 2013; 135:17004–17018. [PubMed: 24187982]
- xi. Borch RF. *Chem Commun.* 1968:442–443.
- xii. For example, stoichiometric portions of Ni^{II}, Co^{II}, Os^{IV}, Ir^{III}, Pt^{II}, are known to mediate nitrile and nitro group reduction in the presence of excess NaBH₄. Suzuki Y, Miyaji Y, Imai Z. *Tetrahedron Lett.* 1969; 52:4555–4558.
- xiii. Selected examples see: Paul H, Bhaduri S, Lahiri GK. *Organometallics.* 2003; 22:3019–3021. Chase PA, Jurca T, Stephan DW. *Chem Commun.* 2008:1701–1703. Reguillo R, Grellier M, Vautravers N, Vendier L, Sabo-Etienne S. *J Am Chem Soc.* 2010; 132:7854–7855. [PubMed: 20481582] Miao X, Fischmeister C, Bruneau C, Dubois JL, Couturier JL. *ChemSusChem.* 2012; 5:1410–1414. [PubMed: 22707476]
- xiv. Magnus P, Sane N, Fauber BP, Lynch V. *J Am Chem Soc.* 2009; 131:16045–16047. [PubMed: 19835379]
- xv. Haddenham D, Pasumansky L, DeSoto J, Eagon S, Singaram B. *J Org Chem.* 2009; 74:1964–1970. [PubMed: 19191712]

- xvi. For selected reviews of utilization of aryl bromides in medicinal synthesis see: Hassan J, Sévignon M, Gozzi C, Schulz E, Lemaire M. *Chem Rev.* 2002; 102:1359–1470. [PubMed: 11996540] Alberico D, Scott ME, Lautens M. *Chem Rev.* 2007; 107:174–238. [PubMed: 17212475] Zeni G, Larock RC. *Chem Rev.* 2006; 106:4644–4680. [PubMed: 17091931] Corbet J-P, Mignani G. *Chem Rev.* 2006; 106:2651–2710. [PubMed: 16836296] Han F-S. *Chem Soc Rev.* 2013; 42:5270–5298. [PubMed: 23460083] Aubin Y, Fischmeister C, Thomas CM, Renaud JL. *Chem Soc Rev.* 2010; 39:694–711. [PubMed: 20111788] McGlacken GP, Bateman LM. *Chem Soc Rev.* 2009; 38:2447–2464. [PubMed: 19623360] Rauws TRM, Maes BUW. *Chem Soc Rev.* 2012; 41:2463–2497. [PubMed: 22222274]

Fig. 1.

Dual site catalysts for hydride manipulation.

Scheme 1.
Syntheses of precursor **3** and complex **6**. Molecular Structure of **3**.

Scheme 2.
Mechanistic Template for Nitrile Reductions.

Table 1

Optimization of Nitrile Reduction Conditions.

Entry	Catalyst	NaO ^t Bu	5 h conversion ^a	NMR Yield ^b
1	3	None	50%	^c
2	3	1 equiv.	> 95%	> 90%, 5 hr
3	2	1 equiv.	46%	42%, 245 hr
4	6	1 equiv.	41%	trace, 245 hr
5	no catalyst	1 equiv.	38%	43%, 245 hr

^a Starting material consumption by NMR in 5 hours.^b Product formed upon consumption of nitrile and subsequent addition of water.^c Not recorded.

Table 2

Scope of **3**-Catalyzed Nitrile Reduction.

$ \begin{array}{c} \text{N}\equiv\text{C}-\text{CH}_2-\text{R} \\ \mathbf{7} \end{array} \xrightarrow[\text{MeOH, reflux}]{\begin{array}{c} \mathbf{3} \text{ (5 mol \%)} \\ \text{NaBH}_4, \text{NaO}^t\text{Bu} \end{array}} \begin{array}{c} \text{H}_2\text{N}-\text{CH}_2-\text{R} \\ \mathbf{8} \end{array} \text{ or } \begin{array}{c} \text{H}_2\text{N}-\text{C}(=\text{O})-\text{R} \\ \mathbf{9} \end{array} $				
Entry	Nitrile	Conditions	Product	Yield ^a
1	7a	5 mol% 3 4 eq NaBH ₄ , 1 eq NaO ^t Bu MeOH, reflux, 12 h	8a	82%
2	7b	5 mol% 3 8 eq NaBH ₄ , 1 eq NaO ^t Bu MeOH, reflux, 12 h	8b	87%
3	7c	5 mol% 3 4 eq NaBH ₄ , 1 eq NaO ^t Bu MeOH, reflux, 4 h	8c	92%
4	7d	5 mol% 3 8 eq NaBH ₄ , 1 eq NaO ^t Bu MeOH, reflux, 14 h	8d	85%
5	7e	5 mol% 3 4 eq NaBH ₄ , 1 eq NaO ^t Bu MeOH, reflux, 8 h	8e	80%
6	7f	5 mol% 3 4 eq NaBH ₄ , 1 eq NaO ^t Bu MeOH, reflux, 12 h	8f	84%
7	7g	5 mol% 3 8 eq NaBH ₄ , 1 eq NaO ^t Bu MeOH, reflux, 12 h	8g	60%
8	7h	5 mol% 3 4 eq NaBH ₄ , 1 eq NaO ^t Bu MeOH, reflux, 8 h	8h	64%
9	7i	5 mol% 3 4 eq NaBH ₄ , 1 eq NaO ^t Bu MeOH, reflux, 8 h	9i	56%

Entry	Nitrile	Conditions	Product	Yield ^a
10	7j	5 mol% 3 4 eq NaBH ₄ , 1 eq NaO ^t Bu MeOH, reflux, 8	9j	87%

^a Reported yields are isolated yields.