Review

Bioactive natural products from Chinese marine flora and fauna

Zhen-fang ZHOU, Yue-wei GUO*

State Key Laboratory of Drug Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China

In recent decades, the pharmaceutical application potential of marine natural products has attracted much interest from both natural product chemists and pharmacologists. Our group has long been engaged in the search for bioactive natural products from Chinese marine flora (such as mangroves and algae) and fauna (including sponges, soft corals, and mollusks), resulting in the isolation and characterization of numerous novel secondary metabolites spanning a wide range of structural classes and various biosynthetic origins. Of particular interest is the fact that many of these compounds show promising biological activities, including cytotoxic, antibacterial, and enzyme inhibitory effects. By describing representative studies, this review presents a comprehensive summary regarding the achievements and progress made by our group in the past decade. Several interesting examples are discussed in detail.

Keywords: marine natural products; biological activity; mangrove; algae; soft coral; sponges; mollusks

Acta Pharmacologica Sinica (2012) 33: 1159–1169; doi: 10.1038/aps.2012.110; published online 3 Sep 2012

Introduction

The unique ocean habitat has caused marine organisms to evolve distinctive metabolic pathways, producing remarkable secondary metabolites that differ from those of terrestrial plants. Because the compounds isolated from marine organisms are structurally and biologically intriguing, research on marine natural products has been attracting the attention of chemists as a challenging research topic. To date, more than 30 000 compounds bearing unusual structures and exhibiting various bioactivities have been isolated from marine plants and invertebrates.

In recent years, the interest in natural products from Chinese marine organisms has increased dramatically, making the annual average number of natural products isolated from marine organisms increase rapidly. As one of the most active research groups in this area, we have been devoted to the investigation of Chinese marine plants and invertebrates since 2000. In the course of our studies, a series of structurally diverse and biologically interesting compounds belonging to various structural classes, such as macrocyclic polydisulfides, spirodioxynaphthalene, and alkaloids, have been isolated and characterized. Through these productive studies, we have made important contributions to the research and develop-

ment of Chinese marine natural products. This review summarizes the progress and achievements made by our group in the study of Chinese marine flora and fauna in the past decade, and several interesting examples are discussed in detail.

Chemical studies on Chinese marine plants

Algae

Microalgae play important roles in marine biological systems. Through their photosynthetic ability, algae are the major producers of biomass and organic compounds in the ocean. More importantly, many algal metabolites present unique structures and are formed via biosynthetic routes that are quite different from those known to produce terrestrial metabolites.

Marine dinoflagellates are flagellated organisms with both photosynthetic and heterotrophic attributes. The lifestyle of these organisms is diverse, and many of them participate in symbiotic relationships. For some time, the study of dinoflagellates has focused on their toxin production and pigment composition, and only a small number of these organisms have been found to produce other secondary metabolites. As part of our ongoing research on the biologically active substances of Chinese marine organisms, three new polyhydroxypolyene compounds, lingshuiols A and B (**1** and **2**) and lingshuiol (**3**), have been isolated from the Chinese marine dinoflagellate *Amphidinium* sp collected at Lingshui Bay, Hainan Province, China^[1, 2] (Figure 1). The structures of these compounds are

* To whom correspondence should be addressed.

E-mail ywguo@mail.shcnc.ac.cn

Received 2012-06-17 Accepted 2012-07-12

Figure 1. Structures of polyhydroxypolyene compounds 1–3.

characterized by linear carbon chains, as determined by extensive analyses of 2D NMR spectroscopic data; however, the absolute configurations remain to be determined. In previous studies, polyhydroxypolyene compounds were found to exhibit antifungal, hemolytic, and antimicrobial activities^[3–5]. It was demonstrated that lingshuiol (**3**) possessed potent *in vitro* cytotoxic activities against A549 and HL-60 cells, with IC₅₀ values of 0.21 and 0.23 $\mu\text{mol/L}$, respectively.

Red algae of the genus *Laurencia* are found throughout the world, mostly in tropical and subtropical regions. Since Irie's pioneering investigations on *Laurencia* over the last four decades^[6], a number of cuparene- and laurane-derived sesquiterpenoids have been isolated from this genus. The chemical investigation of the Chinese red alga *Laurencia okamurai* Yamada led to the isolation of a new laurane sesquiterpenoid, 3 β -hydroxyaplysin (**4**), and two novel rearranged sesquiterpenes, laurokamurenes A and B (**5** and **6**)^[7] (Figure 2). Their structures, including their relative stereochemistry, were determined based on a detailed interpretation of 2D NMR spectra and comparisons with related known compounds. To the best of our knowledge, this is also the first report of the isolation of sesquiterpenes containing a 2,2,3-trimethylcyclopentenyl moiety from a natural source. The sesquiterpenoids isolated from the red algae of the genus *Laurencia* in recent years can be classified into more than 20 sesquiterpenoid skeletons^[8]. However, in most cases, the three methyl groups in the aliphatic portion are located at either positions 1,2,3

Figure 2. Structures of sesquiterpenes 4–8.

(laurene type) or 1,2,2 (cuparene type), with the exception of laurokamurenes A and B (**5** and **6**).

Two novel aromatic valerenane-type sesquiterpenes, caulerpals A and B (**7** and **8**), were isolated from the Chinese green alga *Caulerpa taxifolia*^[9] (Figure 2). Compounds **7** and **8** were evaluated for their inhibitory activity against human protein tyrosine phosphatase 1B (PTP1B), a potential drug target for the treatment of type 2 diabetes and obesity, but neither compound exhibited a potent PTP1B inhibitory activity. It is important to note that caulerpals A and B (**7** and **8**) are the only two known compounds with an aromatic valerenane-type carbon skeleton from a natural source.

Mangroves

Mangroves comprise a large number of various salt-tolerant plants growing in tropical and subtropical intertidal estuarine zones. Mangrove plants are usually categorized into two subgroups, true mangrove and semi-mangrove plants, according to their living environment. True mangrove plants are confined to the typical intertidal mangrove habitats where the seawater salinity is usually 17.0‰–36.4‰. Semi-mangrove plants, however, grow on the landward fringe of the mangrove habitats or in the terrestrial marginal zones that are subjected to irregularly high tides. In general, mangrove plants consist of 84 species globally, belonging to 24 genera and 16 families; 14 are semi-mangrove species^[10,11].

Most of our studies involve plants belonging to the Rhizophoraceae, Sonneratiaceae, Euphorbiaceae, and Meliaceae families. A series of unusual compounds bearing unprecedented structures were identified in these taxa, some of which showed marked biological activities and are currently in pre-clinical studies.

For instance, an unusual novel macrocyclic polydisulfide gymnorhizol (**9**) was isolated from the mangrove plant *Bru-guiera gymnorhiza* (Rhizophoraceae family). Compound **9** possesses an uncommon carbon skeleton that is characteristic of a 15-membered macrocycle composed of 3 repeated 1,3-dimercaptopropan-2-ol units^[12] (Figure 3). The structure of compound **9** was established by extensive spectroscopic studies and confirmed using X-ray crystallography^[13]. The first total synthesis of gymnorhizol (**9**) was completed by our group with a high overall yield^[14] (Scheme 1).

Prompted by the unusual structure of gymnorhizol (**9**) and its potent inhibitory activity against PTP1B (an IC₅₀ value of 14.9 μmol/L), we investigated the same mangrove species collected from different areas, which led to the isolation of two new macrocyclic polydisulfides, **10** and **11**^[15]. The structures of **10** and **11** were unequivocally determined through single-crystal X-ray diffraction analysis.

A plausible biosynthetic pathway for these cyclic polydisulfides was also proposed in which the 1,3-dimercaptopropan-2-ol unit should be the common building block for the biosynthesis of all of these macrocyclic polydisulfides; the self-cyclization of the 1,3-dimercaptopropan-2-ol unit of different

Figure 3. Structures of macrocyclic polydisulfides **9–11** and their X-rays.

sizes gave rise to macrocyclic polydisulfides ranging from 5-membered to 30-membered polydisulfides or even to larger

Scheme 1. Total synthesis of gymnorhizol (**9**).

polydisulfides. Marine disulfide- and multisulfide-containing metabolites are a special and important class of natural products. Due to the increasing interest in these compounds, the progress in this area was previously summarized by our group^[16], encompassing the literature published between 1971 and 2010.

To obtain sufficient amounts of the polydisulfides for pharmacological research, the mangrove plant *B. gymnorrhiza* collected from Zhanjiang, Guangdong Province, was investigated. Unexpectedly, instead of the ubiquitous macrocyclic polydisulfides, seven new spirodioxynaphthalene compounds, named palmarumycins BG1–BG7 (**12–18**) (Figure 4), were found^[17]. It is noteworthy that these intriguing compounds were generally considered to be fungal metabolites, even though there were four reports on the isolation of palmarumycins (or preussomerins) from plants^[18–21]. The fact that compounds **12–18** were isolated in quite an appreciable quantity indicated that these compounds were produced by *B. gymnorrhiza*; however, we could not rule out the possibility that they were the metabolites of an endophytic or epiphytic fungus. The absolute configurations of metabolites **12–18** were determined using time-dependent density functional theory electronic circular dichroism (TDDFTECD) calculations of the solution conformers. It was found that compound **16** displayed an inhibitory activity against the HL-60 and MCF-7 cell lines. Additionally, we published a report summarizing the literature on the isolation, structure elucidation, biological activities, biosynthesis, and chemical synthesis of spirodioxynaphthalenes over the last 20 years^[22].

Although the chemical investigation of the plants belonging to the Sonneratiaceae family was not as fruitful as that of the Rhizophoraceae family, a novel α -alkylbutenolide dimer, paracaseolide A (**19**), was isolated from the mangrove plant *Sonneratia paracaseolaris* (Sonneratiaceae family); this compound is characterized by an unusual tetraquinane oxa-cage bislactone skeleton bearing two linear alkyl chains^[23] (Figure 5). Moreover, paracaseolide A (**19**) exhibited a potent inhibitory activity against the dual-specificity phosphatase Cdc25B, a key enzyme for cell cycle progression, with an IC₅₀ value of 6.44 μ mol/L. A retrosynthetic pathway for paracaseolide A (**19**) was proposed in which **19** should be biosynthesized through a Diels–Alder [4+2] cycloaddition, starting from two molecules of **20** (Scheme 2). Paracaseolide A (**19**) represents the first example of an α -alkylbutenolide dimer from a natural source. Upon publication, the article was immediately covered in the

Figure 4. Structures of spirodioxynaphthalene compounds **12–18**.

Figure 5. Structures of compounds **19** and **21–25**.

journal of *Natural Product Reports* as “hot off the press”^[24].

In addition, three known compounds, integracins A and B (**21** and **22**) and 15-dehydroxy-integracin B (**24**), were isolated for the first time from the Chinese mangrove plant *Sonneratia hainanensis*^[25] (Figure 5). Although compounds **21–24** had been reported previously, their absolute configurations were not resolved until their absolute chemistry was determined using Mosher’s method, the specific rotation analysis of the alcohols (**24** and **25**) obtained from integracin A (**21**) in two steps, and chemical correlation. Integracin A (**21**) exhibited a cytotoxic effect against the tumor cell lines HepG2 and NCI-H460, both of which were 100% inhibited at 25 mg/mL.

The Euphorbiaceae family is composed of various species that are distributed widely throughout China. Although poisonous, most of the plants in this family are utilized in folk

Scheme 2. Plausible retrosynthetic pathway for paracaseolide A (**19**).

medicine. Thus, the substances of the mangrove plant *Excoecaria agallocha* L (Euphorbiaceae family for the first time) were investigated, which led to the isolation of four diterpenes with an unprecedented skeleton, named excoagallochaols A–D (**26–29**)^[26] (Figure 6). In addition to these terpenes, a novel octacyclic alkaloid with an unusual scaffold, suffruticosine (**30**), was isolated from *Securinega suffruticosa* (Pall) Rehd in the same family^[27] (Figure 6). The absolute configuration of suffruticosine (**30**) was determined on the basis of CD spectral analysis.

Figure 6. Structures of diterpenoids and the alkaloid **26–36**.

Plants of the Meliaceae family are a rich source of limonoids, which are a structurally diverse group of natural products with highly complex polycyclic skeletons. These unusual structural features have attracted the attention of chemists as

challenging targets for total synthesis, bioactivity evaluation, and biosynthetic studies. Chemical investigation on the stem bark of the plant *Toona ciliata* var *pubescens* resulted in the isolation of nine new compounds, including toonapubesin A (**31**), toonapubesic acid A (**32**), and their derivatives^[28] (Figure 6). The proposed structure of **32** was confirmed by the X-ray diffraction analysis of its methyl ester, whereas the absolute configuration was determined by a novel solid-state TDDFT ECD approach on its methyl ester.

More importantly, macrolide (*R*)-de-O-methylasiodiplodin (**33**), isolated from the semi-mangrove *Cerbera manghas*, was found to be a potent nonsteroidal antagonist of the mineralocorticoid receptor (MR), a drug target for the treatment of hypertension and other cardiovascular diseases, showing an IC₅₀ value of 8.93 μmol/L. Due to its low natural yield, further study on its structural derivatization/modification and the structure-activity relationship (SAR) was conducted. Consequently, a series of analogs were synthesized and evaluated for their MR antagonistic activities^[29], and compounds **34–36** were found to exhibit a better selectivity and more potent activity against MR than compound **33**, with IC₅₀ values ranging from 0.58 to 1.11 μmol/L.

Chemical studies on Chinese invertebrates Sponges

Sponges are animals of the phylum Porifera, which consists of approximately 5000–10000 species that can be classified mainly according to the composition of their skeletons as calcarea, glass sponges, and demosponges. Their chemical variability results from the individual species, particular metabolism and complex environment. In the past half century, a large number of compounds, including terpenoids, alkaloids, and polyacetylenes, were isolated and characterized from various sponges collected from different regions of the world.

For instance, five new sesquiterpenes, lingshuiolides A and B (**37** and **38**), lingshuiperoxide (**39**), isodysetherin (**40**), and spirolingshuidolide (**41**) were isolated from the Hainan sponge *Dysidea septosa*^[30] (Figure 7). The absolute configuration of lingshuiolide B (**38**) was established by using a modified Mosher's method. The chemical study on another *Dysidea* sp led to the discovery of a new sesquiterpene quinone (**42**), along with a known related analog dysidine (**43**)^[31] (Figure 7). A bioassay showed that dysidine (**43**) had a potent PTP1B inhibitory activity, with an IC₅₀ value of 6.70 μmol/L, whereas the new compound **42** exhibited a moderate PTP1B inhibitory activity and cytotoxicity, with IC₅₀ values of 39.50 and 19.45

Three unprecedented diastereoisomeric dimers, *cis*-dimers A (**44**) and B (**45**) and *trans*-dimer C (**46**), representing bis-bisabolene skeletons, and their potential precursor dehydrotheonelline (**47**) have been isolated from the South China Sea sponges *Axinyssa variabilis* and *Lipastrotethya ana*^[33] (Figure 7). The three diastereoisomers (**44–46**) were possibly biosynthesized through an intermolecular [4+2] Diels-Alder cycloaddition involving two molecules of dehydrotheonelline (**47**). The *trans*- and *cis*-dimers were possibly formed through *exo*- and *endo*-Diels-Alder coupling, respectively (Scheme 3). Furthermore, a mixture of *cis*-dimers **44** and **45** and the pure *trans*-dimer **46** were active at 50 mg/cm² in a feeding-deterrence test against the gold fish *Carassius auratus*, suggesting a possible defensive role for these compounds.

Soft corals

Soft corals from the South China Sea have been extensively studied by Chinese chemists and have yielded a plethora of steroids and terpenoids. It has been suggested that such secondary metabolites produced by soft corals are most likely involved in the defensive mechanisms of the animals; indeed, the animals appear to be relatively free from predation.

As expected, the chemical study of the soft coral *Lobophytum cristatum* Tixier-Durivault led to the isolation of two new prenylgermacrane-type diterpenoids, lobophytumins A and B (**48** and **49**), two new prenyleudesmane-type diterpenoids, lobophytumins C and D (**50** and **51**), and two new spatane-type

diterpenoids, lobophytumins E and F (**52** and **53**)^[34] (Figure 8). Their structures, including the relative configurations, were determined via a detailed analysis of the spectroscopic data and comparison with related known compounds. In addition,

tion, the absolute configuration of lobophytumin C (**50**) was tentatively assigned by comparing its specific rotation with that of the closely related model compound (-)- β -selinene; in contrast, the stereochemistry of C-11 in compounds **48–49** and **52–53** remains undefined. This is the first report of spatane-type diterpenoids from a soft coral source^[34], and it supports Faulkner's proposal of prenylgermacrene as the precursor of many diterpenes, including quite different but biogenetically related ones. In a bioactivity evaluation, lobophytumins C and D (**50** and **51**) showed weak *in vitro* cytotoxicities against the tumor cell lines A549 and HCT-116.

Casbane diterpenes are characterized by the presence of a dimethyl-cyclopropyl moiety fused to the 14-membered ring, structures that are extremely rare in nature and marine organisms and previously found only in *Sinularia microclavata*^[35]. From the Hainan soft coral *Sinularia depressa*, we isolated nine casbane diterpenes (**54–62**), either *cis* or *trans* ring junctions, including two pairs of epimers: depressin (**54**) and 1-*epi*-depressin (**55**) and 10-hydroxydepressin (**56**) and 1-*epi*-10-hydroxydepressin (**57**)^[36] (Figure 9). Compounds **56** and **57** played a key role in the establishment of the absolute stereochemistry, as the *S* configuration at C-10 of both compounds could be easily assigned using Mosher's method. Three further casbanes (**59–61**) displayed identical structures, with the exception of the stereochemistries at the ring junction, which were *cis* in **59** and *trans* in both **60** and **61**. Compound **58** was similar to **57**, except for the oxidation of the hydroxyl; nevertheless, compound **62** had many similarities with 10-hydroxydepressin (**56**), with the exception of an additional oxygen atom. 10-Hydroxydepressin (**56**) showed a cytotoxic activity against the tumor cell lines HepG2 and SW-1990, with IC₅₀ values of 61 and 37 μ mol/L, respectively, and an antimicrobial activity against *Staphylococcus aureus* and *Escherichia coli* at the concentration of 17 μ mol/L.

The genus *Sarcophyton* is a productive source for unusual tetraterpenoids, which are constructed by coupling two cembranoid units through the Diels-Alder reaction. Five novel biscembranoids, ximaolides A–E (**64–68**), together with their proposed biogenetic precursor, methyl tortuosoate (**63**), were isolated from the Hainan soft coral *Sarcophyton tortuosum*^[37] (Figure 10). The relative stereochemistries of ximaolides A and E (**64** and **68**) were acquired by X-ray diffraction analysis, whereas the stereochemistries of the other compounds were suggested by both biosynthetic considerations and NOESY experiments. Furthermore, ximaolide D (**67**) is the first example of a biscembranoid possessing a tetrahydrofuran ring

between the A and B rings.

The study of *Sarcophyton lactum* also led to the discovery of two unprecedented biscembranoids, bislatumlides A and B (**69** and **70**), for which the dienophilic double bond belonged to an α,β -conjugated- γ -lactone, along with a new cembranolide diterpene, isosarcophytonolide D (**71**)^[38] (Figure 10). Compounds **69** and **70** differ in the configuration of the double bond at C-11: *E* for **69** and *Z* for **70**. However, it was proven that the isomerization of the double bond in **69** readily occurred in the presence of trace amount of acid. Compounds **69** and **70** could be biosynthesized by the Diels-Alder addition of two cembranoid units. Compound **71** and the known compound sarcophytonolide D (**72**) may comprise half of the biosynthetic precursors, as shown in Scheme 4.

Scheme 4. Plausible Diels-Alder reaction leading to compounds **69** and **70**.

Steroids are a large group of compounds encountered in marine organisms and are reported to display various bioactivities. For instance, a novel steroid with an uncommon 21-oic acid methyl ester moiety, designated methyl spongoate (**73**), was isolated from the Sanya soft coral *Spongodes* sp; this steroid exhibited a potent cytotoxicity against BEL-7402 tumor cells *in vitro*^[39]. Later, its stereoselective synthesis was achieved in our laboratory^[40]. Owing to its unusual structure and potent cytotoxicity, a number of analogs with different C-20 side chains were synthesized and evaluated for their cytotoxic activities, and some of the analogs were found to be more potent than compound **73** against the A549, HCT-116, HepG2, SW-1990, MCF-7, and NCI-H460 tumor cell lines. The preliminary SAR study suggested that the unsaturated carbonyl moiety, a Michael acceptor in ring A, and the C-20 side chain played important roles in the cytotoxic effect of these derivatives^[41] (Figure 10).

Mollusks

Mollusks are a large phylum of invertebrate animals that are

Figure 9. Structures of casbane diterpenes **54–62**.

Figure 10. Structures of bisembranoids **63–73**.

highly diverse in size and anatomical structure and also in their behavior and habitat. Mollusks show such a variety in body structures that it is difficult to find definite characteristics that apply to all of the classification groups. The two most universal features are a mantle with an important cavity used for breathing and excretion and the structure of the nervous system. However, most of the animals in the Opisthobranchia family lose their mantles and require other defensive system for protection. Thus, these animals usually produce structurally diverse substances to form a chemical defensive system, representing a different source of bioactive compounds from marine organisms. As expected, a series of metabolites with complex structures and intriguing biological activities have been isolated from mollusks.

The study on the South China Sea nudibranch *Tritoniopsis elegans* led to the isolation of four diterpenes, tritoniopsins A–D (**74–77**), which display an unprecedented pyran ring in the cladiellane framework and represent a novel cladiellane-based diterpene family^[42] (Figure 11). These compounds were also found in the nudibranch's prey, *Cladiella krempfi*. The structure of tritoniopsin A (**74**) was elucidated by X-ray diffraction

analysis, whereas the *S* configuration of the hydroxyl at C-6 in tritoniopsin C (**76**) was assigned by applying Mosher's method. Both tritoniopsins A and B (**74** and **75**) were discovered from *T. elegans* and *C. krempfi*, but, surprisingly, the major compound in this mollusk was tritoniopsin A (**74**). Such an uneven metabolite distribution could be explained by the fact that the mollusk either modified dietary tritoniopsin B (**75**) or selectively accumulated compound **74**. The presence of these unique metabolites in both the nudibranch and the soft coral *C. krempfi* also confirmed the trophic relationship between the predator and prey.

During the investigation of the nudibranch *Glossodoris rufomarginata* and its unidentified prey sponge from the South China Sea, it was observed that scalaradial (**78**) and its deacetyl derivative (**79**) were the main components of the extract of the dietary sponge, whereas a series of related scalaranes (**80–84**) (Figure 11), most likely derived from dietary scalaradial, were isolated from the mollusk, thus proving the ability of *Glossodoris rufomarginata* to modify dietary molecules^[43]. The finding of these molecules (**80**, **81**, and **84**) with a keto functionalization at C-12 is particularly interesting.

Figure 11. Structures of the compounds 74–92 from mollusks.

This structural feature is quite common in the homoscalarane series but unusual in the scalaranes, and in particular it has been encountered only in nudibranch metabolites. In fact, in addition to 12-deacetoxy-12-oxo-deoxoscalarin (**80**), previously isolated from Indian *Glossodoris atromarginata*^[44], only one 12-keto-scalarane, 12-deacetoxy-12-oxo-18-*epi*-scalaradial, from Pacific nudibranch *Chromodoris youngbleuthi*^[45], has been previously reported.

In our research on the sea hare *Aplysia dactylomela*, two novel triterpenes, aplysiols A and B (**85** and **86**), characterized as squalene-derived polyethers, were isolated^[46] (Figure 11). The absolute stereochemistry of aplysiol A (**85**) was determined using Mosher's method and biogenetic considerations, whereas the configuration of aplysiol B (**86**) was established by an integrated NMR-QM (quantum mechanical) approach on the basis of ¹³C NMR chemical shifts and ^{2,3}J_{C-H} coupling-constant DFT (density functional theory) calculations.

Marine pulmonata of the Onchidiidae family are shell-less mollusks living on sheltered intertidal shores. Previous chemical studies on distinct species in this family mostly focused on polypropionates. In our study, onchidione (**87**), as the main metabolite, was characterized from the first collection of the Chinese marine pulmonate *Onchidium* sp (Figure 11), which

was present both in the mucus and in the mantle^[47]. The structure of the polypropionate was suggested by an analysis of spectral data. In particular, the relative stereochemistry of the central ring was supported by a series of nuclear overhauser enhancement (NOE) experiments, and the complete structure was confirmed by X-ray diffraction analysis. A recent study on the second collection of the mollusk led to the isolation of onchidiol (**88**) and 4-*epi*-onchidiol (**89**), related alcohols of onchidione (**87**)^[48] (Figure 11). The absolute configurations of onchidiol (**88**) and 4-*epi*-onchidiol (**89**) were established by TDDFTECD. Moreover, the absolute chemistry of onchidione (**87**) was corroborated by X-ray diffraction analysis, with the final refinement using Cu-Kα data.

In the final section of this review, we describe the nitrogen-containing compounds of marine mollusks. The first example consists of phidianidines A and B (**90** and **91**) obtained from *Phidiana militaris*, belonging to the Glaucidae family^[49] (Figure 11). The two novel alkaloids were characterized with an unprecedented carbon skeleton featuring the presence of a 1,2,4-oxadiazole ring linking the indole system through a methylene bridge and displaying an aminoalkylguanidine group at C-3'. The 1,2,4-oxadiazole ring is extremely rare in nature. The phidianidines were tested in a panel of tumor

cell lines (C6 rat glioma cells, HeLa human epithelial cervical cancer cells and CaCo-2 human epithelial colorectal adenocarcinoma cells) and non-tumor cell lines (H9c2 rat embryonic cardiac myoblasts and 3T3-L1 murine embryonic fibroblasts) and showed a strong and selective activity against C6 and 3T3-L1 cells by IC_{50} values within the nanomolar range, as shown in Table 1.

Table 1. Cytotoxicity profile of compounds **90** and **91** against tumor and non-tumor cell lines, IC_{50} ($\mu\text{mol/L}$)^a.

Cell line	Phidianidine A (90)	Phidianidine B (91)
C6	0.642±0.2	0.98±0.3
HeLa	1.52±0.3	0.417±0.4
CaCo-2	35.5±4	100.2±8.5
3T3-L1	0.14±0.2	0.786±0.3
H9c2	2.26±0.6	5.42±0.8

^a IC_{50} values are expressed as mean±SEM ($n=24$) of the three independent experiments. Bold values show IC_{50} of less than 1 $\mu\text{mol/L}$.

The second example is an unusual molecule, actisonitrile (**92**), found in the mantle of *Actinocyclus papillatus*^[50] (Figure 11), belonging to the Actinocyclus family that is another family of opisthobranch, which was never studied thus far. Despite its structure is quite simple, the determination of the absolute stereochemistry of **92** was a challenging task, yet its absolute configuration was recently solved by a synthetic approach. Thus, both the *S* and *R* enantiomers were synthesized, and the comparison of optical rotation and CD profile of the natural product with those of the pair of the synthetic enantiomers resulted in the assignment of the *R* configuration at the C-2 position of compound **92**.

Conclusions

This review summarizes representative substances from Chinese marine plants and invertebrates found by our group, indicating that marine organisms are an inexhaustible source of new molecules that often display unique structures and sometimes have very interesting pharmacological properties, such as antifungal, antibacterial, enzyme-inhibitory, and other activities. Some of the compounds isolated from marine organisms even show promising activities within the nanomolar range and can be developed as drug candidates or at least lead compounds in the near future. Although our research on marine natural products is limited, it has established the foundation for further studies.

Because our understanding of the biosynthetic origin and the real ecological role of these bioactive compounds remains incomplete, further research to clarify these important issues is necessary.

Acknowledgements

The authors wish to acknowledge all of their colleagues who have contributed to the studies presented herein and

to thank Prof T Kurtán (University of Debrecen, Hungary) for his contribution to the establishment of absolute stereochemistry by CD methodology. The reported research was financially supported by the National Marine '863' Project (No 2011AA09070102), the National Natural Science Foundation of China (No 21072204, 21021063, 40976048, 31070310, and 81072572), the SKLDR/SIMM Projects (No SIMM 1203ZZ-03 and SIMM 1105KF-04), and Syngenta-SIMM-PhD Studentship Project, and partially funded by the National S&T Major Project (2011ZX09307-002-03) and the EU 7th Framework Programme-IRSES Project (2010-2014).

References

- Huang XC, Zhao D, Guo YW, Wu HM, Trivellone E, Cimino G. Lingshuiols A and B, two new polyhydroxy compounds from the Chinese marine dinoflagellate *Amphidinium* sp. *Tetrahedron Lett* 2004; 45: 5501–4.
- Huang XC, Zhao D, Guo YW, Wu HM, Lin LP, Wang ZH, et al. Lingshuiol, a novel polyhydroxyl compound with strongly cytotoxic activity from the marine dinoflagellate *Amphidinium* sp. *Bioorg Med Chem Lett* 2004; 14: 3117–20.
- Doi Y, Ishibashi M, Nakamichi H, Kosaka T, Ishikawa T, Kobayashi J. Luteophanol A, a new polyhydroxyl compound from symbiotic marine dinoflagellate *Amphidinium* sp. *J Org Chem* 1997; 62: 3820–3.
- Paul GK, Matsumori N, Murata M, Tachibana K. Isolation and chemical structure of amphidinol 2, a potent hemolytic compound from marine dinoflagellate *Amphidinium klebsii*. *Tetrahedron Lett* 1995; 36: 6279–82.
- Satake M, Murata M, Yasumoto T, Fujita T, Naoki H. Amphidinol, a polyhydroxy-polyene antifungal agent with an unprecedented structure, from a marine dinoflagellate, *Amphidinium klebsii*. *J Am Chem Soc* 1991; 113: 9859–61.
- Irie T, Suzuki M, Masamune T. Laurencin, a constituent from laurencia species. *Tetrahedron Lett* 1965; 6: 1091–9.
- Mao SC, Guo YW. A laurane sesquiterpene and rearranged derivatives from the Chinese red alga *Laurencia okamurai* Yamada. *J Nat Prod* 2006; 69: 1209–11.
- Mao SC, Guo YW. Sesquiterpenes from algae of the Genus *Laurencia*: chemistry and biological activities. *Chin Tradit Herb Drugs* 2004; 35: 8–18.
- Mao SC, Guo YW, Shen X. Two novel aromatic valerenane-type sesquiterpenes from the Chinese green alga *Caulerpa taxifolia*. *Bioorg Med Chem Lett* 2006; 16: 2947–50.
- Wang BS, Liang SC, Zhang WY, Zan QJ. Mangrove Flora of the world. *Acta Bot Sin* 2003; 45: 644–53.
- Li MY, Xiao Q, Pan JY, Wu J. Natural products from semi-mangrove flora: source, chemistry and bioactivities. *Nat Prod Rep* 2009; 26: 281–98.
- Sun YQ, Guo YW. Gymnorrhizol, an unusual macrocyclic polydisulfide from the Chinese mangrove *Bruguiera gymnorrhiza*. *Tetrahedron Lett* 2004; 45: 5533–5.
- Sun YQ, Guo YW. Crystal structure of 1,2,6,7,11,12-hexathia-cyclopentadecane-4,9,14-triol, $C_{29}H_{48}O_3S_6$. *Z Kristallogr NCS* 2004; 219: 121–3.
- Gong JX, Shen X, Yao LG, Jiang H, Krohn K, Guo YW. Total synthesis of gymnorrhizol, an unprecedented 15-membered macrocyclic polydisulfide from the Chinese mangrove *Bruguiera gymnorrhiza*. *Org Lett* 2007; 9: 1715–6.
- Huang XY, Wang Q, Liu HL, Zhang Y, Xin GR, Shen X, et al. Diastereoisomeric macrocyclic polydisulfides from the mangrove *Bruguiera*

- gymnorhiza*. *Phytochemistry* 2009; 70: 2096–100.
- 16 Jiang CS, Muller WE, Schroder HC, Guo YW. Disulfide- and multisulfide-containing metabolites from marine organisms. *Chem Rev* 2012; 112: 2179–207.
- 17 Cai YS, Kurtan T, Miao ZH, Mandi A, Komaromi I, Liu HL, et al. Palmarumycins BG1-BG7 and preussomerin BG1: establishment of their absolute configurations using theoretical calculations of electronic circular dichroism spectra. *J Org Chem* 2011; 76: 1821–30.
- 18 Wang L, Zhang CF, Wang ZT, Zhang M, Xu LS. Five new compounds from *Dendrobium crystallinum*. *J Asian Nat Prod Res* 2009; 11: 903–11.
- 19 Prajoubklang A, Sirithunyalug B, Charoenchai P, Suvannakad R, Sriubolmas N, Piyamongkol S, et al. Bioactive deoxypreussomerins and dimeric naphthoquinones from *Diospyros ehretioides* fruits: deoxypreussomerins may not be plant metabolites but may be from fungal epiphytes or endophytes. *Chem Biodivers* 2005; 2: 1358–67.
- 20 Ravindranath N, Ravinder Reddy M, Mahender G, Ramu R, Ravi Kumar K, Das B. Deoxypreussomerins from *Jatropha curcas*: are they also plant metabolites? *Phytochemistry* 2004; 65: 2387–90.
- 21 Kouam TNM, Lavaud C, Massiot G, Nuzillard JM, Connolly JD, Ryeroff DS. Bipendensin, an unusual phenolic acetal from *Azela bipendensis*. *Nat Prod Lett* 1993; 3: 299–303.
- 22 Cai YS, Guo YW, Krohn K. Structure, bioactivities, biosynthetic relationships and chemical synthesis of the spirodioxynaphthalenes. *Nat Prod Rep* 2010; 27: 1840–70.
- 23 Chen XL, Liu HL, Li J, Xin GR, Guo YW. Paracaseolide A, first alpha-alkylbutenolide dimer with an unusual tetraquinane oxa-cage bis-lactone skeleton from Chinese mangrove *Sonneratia paracaseolaris*. *Org Lett* 2011; 13: 5032–5.
- 24 Hill RA, Sutherland A. Hot off the press. *Nat Prod Rep* 2011; 28: 1879–82.
- 25 Liu HL, Huang XY, Li J, Xin GR, Guo YW. Absolute configurations of integracins A, B, and 15'-dehydroxy-integracin B. *Chirality* 2012; 24: 459–62.
- 26 Wang JD, Zhang W, Li ZY, Xiang WS, Guo YW, Krohn K. Elucidation of excogallochaols A–D, four unusual diterpenoids from the Chinese mangrove *Excoecaria agallocha*. *Phytochemistry* 2007; 68: 2426–31.
- 27 Qin S, Liang JY, Gu YC, Guo YW. Suffruticosine, a novel octacyclic alkaloid with an unprecedented skeleton from *Securinega suffruticosa* (Pall) Rehd. *Tetrahedron Lett* 2008; 49: 7066–9.
- 28 Wang JR, Liu HL, Kurtan T, Mandi A, Antus S, Li J, et al. Protolimonoids and norlimonoids from the stem bark of *Toona ciliata* var *pubescens*. *Org Biomol Chem* 2011; 9: 7685–96.
- 29 Jiang CS, Zhou R, Gong JX, Chen LL, Kurtan T, Shen X, et al. Synthesis, modification, and evaluation of (R)-de-O-methylasiodiplodin and analogs as nonsteroidal antagonists of mineralocorticoid receptor. *Bioorg Med Chem Lett* 2011; 21: 1171–5.
- 30 Huang XC, Li J, Li ZY, Shi L, Guo YW. Sesquiterpenes from the Hainan Sponge *Dysidea septosa*. *J Nat Prod* 2008; 71: 1399–403.
- 31 Li Y, Zhang Y, Shen X, Guo YW. A novel sesquiterpene quinone from Hainan sponge *Dysidea villosa*. *Bioorg Med Chem Lett* 2009; 19: 390–2.
- 32 Zhang Y, Li Y, Guo YW, Jiang HL, Shen X. A sesquiterpene quinone, dysidine, from the sponge *Dysidea villosa*, activates the insulin pathway through inhibition of PTPases. *Acta Pharmacol Sin* 2009; 30: 333–45.
- 33 Mao SC, Manzo E, Guo YW, Gavagnin M, Mollo E, Ciavatta ML, et al. New diastereomeric bis-sesquiterpenes from Hainan marine sponges *Axinyssa variabilis* and *Lipastrotethya ana*. *Tetrahedron* 2007; 63: 11108–13.
- 34 Li L, Sheng L, Wang CY, Zhou YB, Huang H, Li XB, et al. Diterpenes from the Hainan soft coral *Lobophytum cristatum* Tixier-Durivault. *J Nat Prod* 2011; 74: 2089–94.
- 35 Zhang CX, Yan SJ, Zhang GW, Lu WG, Su JY, Zeng LM, et al. Cytotoxic diterpenoids from the soft coral *Sinularia microclavata*. *J Nat Prod* 2005; 68: 1087–9.
- 36 Li Y, Carbone M, Vitale RM, Amodeo P, Castelluccio F, Sicilia G, et al. Rare casbane diterpenoids from the Hainan soft coral *Sinularia depressa*. *J Nat Prod* 2010; 73: 133–8.
- 37 Jia R, Guo YW, Chen P, Yang YM, Mollo E, Gavagnin M, et al. Biscembranoids and their probable biogenetic precursor from the Hainan soft coral *Sarcophyton tortuosum*. *J Nat Prod* 2007; 70: 1158–66.
- 38 Yan XH, Gavagnin M, Cimino G, Guo YW. Two new biscembranes with unprecedented carbon skeleton and their probable biogenetic precursor from the Hainan soft coral *Sarcophyton latum*. *Tetrahedron Lett* 2007; 48: 5313–6.
- 39 Yan XH, Lin LP, Ding J, Guo YW. Methyl spongoate, a cytotoxic steroid from the Sanya soft coral *Spongodes* sp. *Bioorg Med Chem Lett* 2007; 17: 2661–3.
- 40 Gong JX, Miao ZH, Yao LG, Ding J, Kurtan T, Guo YW. Stereoselective synthesis of methyl spongoate, a new steroid with potent antitumor activities. *Synlett* 2010; 2010: 480–2.
- 41 Jiang CS, Huang CG, Feng B, Li J, Gong JX, Kurtan T, et al. Synthesis and antitumor evaluation of methyl spongoate analogs. *Steroids* 2010; 75: 1153–63.
- 42 Ciavatta ML, Manzo E, Mollo E, Mattia CA, Tedesco C, Irace C, et al. Tritoniopsins A–D, cladiellane-based diterpenes from the South China Sea nudibranch *Tritoniopsis elegans* and its prey *Cladiella krempfi*. *J Nat Prod* 2011; 74: 1902–7.
- 43 Gavagnin M, Mollo E, Docimo T, Guo YW, Cimino G. Scalarane metabolites of the nudibranch *Glossodoris rufomarginata* and its dietary sponge from the South China Sea. *J Nat Prod* 2004; 67: 2104–7.
- 44 Fontana A, Cavaliere P, Ungur N, D'Souza L, Parameswaram PS, Cimino G. New scalaranes from the nudibranch *Glossodoris atromarginata* and its sponge prey. *J Nat Prod* 1999; 62: 1367–70.
- 45 Terem B, Scheuer PJ. Scalaradial derivatives from the nudibranch chromodoris youngbleuthi and the sponge spongia oceanica. *Tetrahedron* 1986; 42: 4409–12.
- 46 Manzo E, Gavagnin M, Bifulco G, Cimino P, Di MS, Ciavatta ML, et al. Aplysiols A and B, squalene-derived polyethers from the mantle of the sea hare *Aplysia dactylomela*. *Tetrahedron* 2007; 63: 9970–8.
- 47 Carbone M, Gavagnin M, Mattia CA, Lotti C, Castelluccio F, Pagano B, et al. Structure of onchidione, a bis-γ-pyrone polypropionate from a marine pulmonate mollusk. *Tetrahedron* 2009; 65: 4404–9.
- 48 Wang JR, Carbone M, Gavagnin M, Mandi A, Antus S, Yao LG, et al. Assignment of absolute configuration of bis-γ-pyrone polypropionates from marine pulmonate molluscs. *Eur J Org Chem* 2012; 2012: 1107–11.
- 49 Carbone M, Li Y, Irace C, Mollo E, Castelluccio F, Di Pascale A, et al. Structure and cytotoxicity of phidianidines A and B: first finding of 1,2,4-oxadiazole system in a marine natural product. *Org Lett* 2011; 13: 2516–9.
- 50 Manzo E, Carbone M, Mollo E, Irace C, Di Pascale A, Li Y, et al. Structure and synthesis of a unique isonitrile lipid isolated from the marine mollusk *Actinocyclus papillatus*. *Org Lett* 2011; 13: 1897–9.