Published in final edited form as:

J Allergy Clin Immunol. 2014 May ; 133(5): 1246–1254. doi:10.1016/j.jaci.2013.10.039.

Asthma, Allergy, and Responses to Methyl Donor Supplements and Nutrients

Sunita Sharma, MD, MPH^{1,2} and **Augusto Litonjua, MD, MPH^{1,2}**

¹Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, and Harvard Medical School, Boston, MA, USA

²Division of Pulmonary and Critical Care Medicine, Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA

Abstract

After a brief period of stabilization, recent data has shown that the prevalence of asthma and allergic diseases continue to increase. Atopic diseases are major public health problems resulting in significant disability and resource utilization globally. While environmental factors influence the development of atopic disease, dietary changes may partially explain the high burden of atopic disease. One mechanism by which diet is suspected to impact asthma and allergy susceptibility is through epigenetic mechanisms including DNA methylation. Dietary methyl donors are important in the one-carbon metabolism pathway that is essential for DNA methylation. Findings from both observational studies and interventional trials of dietary methyl donor supplementation on the development and treatment of asthma and allergy have produced mixed results. While issues related to the differences in study design partially explain the heterogeneous results, two other issues have been largely overlooked in these studies. Firstly, these nutrients affect one of many pathways and occur in many of the same foods. Secondly, it is now becoming clear that the human intestinal microbiome is involved in the metabolism and production of the B vitamins and other methyl donor nutrients. Future studies will need to account for both the interrelationships between these nutrients and the effects of the microbiome.

Keywords

Allergy; Asthma; Betaine; Choline; Vitamin B2; Vitamin B6; Vitamin B12; Folate; Methyl Donor; Microbiome

© 2013 American Academy of Allergy, Asthma and Immunology. Published by Mosby, Inc. All rights reserved.

Correspondence: Sunita Sharma, MD, MPH Channing Division of Network Medicine Brigham and Women's Hospital 181 Longwood Avenue Boston, MA 02115 sunita.sharma@channing.harvard.edu Tel: 617-525-1234 Fax: 617-525-0958.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

Asthma and allergic diseases including allergic rhinitis and atopic dermatitis continue to be major public health problems resulting in significant disability and resource utilization globally^{1, 2}. Although the trend in asthma prevalence in the United States had initially stabilized, since 2003 the prevalence of asthma and allergic disease in the United States and abroad is rising³⁻⁷. There is substantial evidence for the genetic predisposition to complex diseases like asthma and allergy with multiple genetic loci demonstrating reproducible associations with these diseases, but genetics alone is unable to explain the overall burden of atopic disease. While the etiology of the rising prevalence is not definitively known, several factors have been proposed to account for this increase including environmental etiologies^{8, 9}. Environmental exposures such as tobacco smoke have been shown to influence the incidence of both asthma and allergy in children¹⁰⁻¹² and in adults^{13, 14}. Other environmental exposures including traffic related air pollution have also shown modest associations with atopic diseases in children¹⁵⁻¹⁷. However, even the contribution of these environmental exposures to asthma and allergy susceptibility is unlikely to fully explain the increasing prevalence of asthma and allergic diseases worldwide.

Epidemiologic data suggests that changes in dietary patterns may also partially explain the rising prevalence of asthma and allergy in industrialized countries¹⁸⁻²⁰. However, the role of certain dietary nutrients in the development of atopic diseases has been conflicting in part due to the fact that the role of diet in the development of atopy is complex, difficult to measure, and is limited by a dearth of interventional trials investigating the impact of diet on atopic disease susceptibility.

Methyl Donors

One of the proposed mechanisms by which diet is suspected to impact asthma and allergy susceptibility is through epigenetic mechanisms including DNA methylation, which results in heritable changes in gene expression without alteration of the original DNA sequence²¹. Methylation of CpG islands within regulatory regions of DNA is an important mechanism for the control of gene expression and has been shown to be transmissible across generations²². DNA methylation results from the transfer of methyl groups to the cytosine in the CpG dinucleotide resulting in a reversible modification that can alter both chromosome stability and suppress gene transcription. Environmental exposures, including air pollution and dietary factors, have been shown to influence the epigenome²³. Furthermore, recent evidence from the Normative Aging Study suggests that prior allergen sensitization is associated with increased methylation of the retrotransposon-derived element Alu, which is a proxy for global DNA methylation, ($\beta = 0.32$ for sensitized vs. nonsensitized patients; $p = 0.003$) even after adjustment for smoking status and air pollutants²⁴.

Dietary methyl donors including folate and choline are necessary for the one-carbon metabolism pathway that produces S-adenosylmethionine (SAM), which is the universal methyl donor that is essential for the DNA methylation process to occur (Figure 1). Vitamin B12 and betaine, which are additional agents derived from the diet, are also necessary for methionine synthesis, which is another step that is important in DNA methylation.

Therefore, differential intake of these nutrients may lead to differences in DNA methylation and ultimately alter gene expression. In an agouti mouse model, mothers on a diet supplemented with methyl donors resulted in hypermethylation of the metastable epiallele in the progeny resulting in a brown coat color, lower incidence of obesity, and increased longevity²⁵. In a mouse model of allergic airway disease, *in utero* supplementation of a diet that is high in methyl donors (including folic acid, choline, and vitamin B12) resulted in increased atopy (IgE and eosinophils) and airway responsiveness in the exposed progeny²⁶. In addition, *in utero* exposure to a high methyl donor diet was associated with increased methylation of the Runt-related transcription factor (*RUNX3*) gene in the progeny, which resulted in decreased expression of the gene. *Runx3* is known to negatively regulate allergic airway disease, suggesting that the methylation changes induced by a diet high in methyl donors may result in increased atopic diseases. Given the emerging evidence from animal models demonstrating a role for dietary methyl donors in the development of atopic diseases, dietary methyl donors have been investigated in human subjects as a possible etiology of the rising prevalence of atopic disease.

In humans dietary methyl groups such as folate, vitamin B12, and choline are a source of methyl donors for DNA methylation (Table 1). Methyl donors, especially folate, have been extensively studied in asthma and to a lesser degree in allergy. However, the data for the association of dietary methyl donors with the development of asthma and allergy in humans remains unclear. In this review we examine the existing evidence for the role of methyl donors in the development and treatment of asthma and allergy. We have limited this review to articles that have examined specific nutrients rather than foods or diets that contain the nutrient. Food intake and dietary patterns and asthma and allergies have recently been reviewed and the readers are directed to those reviews^{18, 20, 27, 28}

Folate

Folate is a water-soluble B vitamin that is naturally present in a variety of foods and is available in nutritional supplements. It is naturally occurring in the diet as folate, and folic acid is the a fully oxidized monoglutamate form that is used as a dietary supplement²⁹. Folic acid consists of a p-aminobenzoic molecule linked to a pteridine ring and one molecule of glutamic acid. Dietary folates are polyglutamates, which include additional glutamate residues³⁰. Folate is found naturally in foods including dark leafy vegetables, fruits, nuts, beans, poultry, meat, eggs, seafood, and dairy products. In addition, it is found (in folic acid form) in fortified food products including breads, cereals, and grain products. Dietary supplements including multivitamins, however, are the major source of folic acid in humans. Unlike naturally-occurring folate, which is hydrolyzed in the intestines and reduced to tetrahydrofolate upon absorption, folic acid requires reduction and methylation in the liver via dihydrofolate reductase. It should be noted that controversy exists regarding how best to measure folate status in humans³¹⁻³³, but this is beyond the scope of the current review.

Folate is necessary for many biologic processes including purine and pyrimidine synthesis and amino acid production. Folate is also needed for the methylation of deoxyuridylate to thymidylate in the formation of DNA, which is required for normal cell division. Folate in the form of 5-methyl-tetrahydrofolate (5-methyl-THF) provides the methyl groups necessary

for the formation of S-adenosyl-methionine (SAM), which is the universal methyl donor required for DNA methylation. In addition to its role in DNA methylation, folate may have antioxidant effects, which is relevant to asthma and allergies^{34, 35}. The use of folic acid supplements before and during the first trimester of pregnancy reduces the risk of neural tube defects and other congenital anomalies and is, therefore, recommended as part of public health guidelines worldwide³⁶⁻³⁹. Because asthma and allergies develop in early childhood, prenatal exposures are thought to play a role in the development of these disorders, and folate intake has been studied as a potential factor. The association between folate and asthma was recently extensively reviewed by Blatter and colleagues⁴⁰. With regard to the question of prenatal folate intake, folic acid supplementation, or maternal folate status and the development of asthma and allergies in children, a few studies showed increased risks of early wheeze and lower respiratory illnesses,⁴¹ asthma,^{42, 43} and eczema (but not asthma)⁴⁴. However, the bulk of birth cohort studies have shown no effect of folate intake or folic acid supplement use at various points in pregnancy⁴⁵⁻⁵¹ and asthma or allergies. We agree with Blatter and colleagues that given the weight of the evidence against a moderate to strong effect of maternal folate status on asthma and allergies, there is no reason to change the current recommendations regarding prenatal folic acid supplementation.

The effect of maternal folate intake on DNA methylation and the development of atopic disease in the offspring is a topic of recent interest⁵². In a study of 24 pregnant women, maternal folic acid intake during pregnancy and cord blood folate levels were correlated with cord blood LINE-1 methylation, a measure of global DNA methylation⁵³. Furthermore, among 107 pregnant Korean women, serum folate levels were significantly associated with percent methylation of the placenta⁵⁴. Additionally, peri-conceptional use of folic acid supplements was associated with higher methylation of insulin-like growth factor II (*IGF2*) in young children⁵⁵. However, in a prospective cohort study of 534 pregnant women participating in Project Viva, there was no significant association found between maternal dietary intake of folate as assessed by questionnaire in both the first and second trimesters of pregnancy with cord blood LINE-1 methylation⁵⁶. The discrepancy in these results may in part be due to relatively small sample sizes in the studies and in the assessment of DNA methylation. Studies using newer and more comprehensive methods^{57, 58} of assessing DNA methylation (using genome-wide techniques) are underway, and using these methods together with folate intake assessment can better answer the question of whether DNA methylation of specific regions in the genome may modify the risk of asthma and allergies.

Post-natal studies of the association of folate and asthma have also resulted in mixed findings. Two studies have shown an increased risk with folate. In a small study of 138 at-risk children, those with higher folate levels in early childhood (<6yrs old) had increased risks of sensitization in later childhood to foods and aeroallergens compared with children who had lower levels.⁵⁹ However, there were no associations of increased folate level with asthma or wheezing. In a large cross-sectional study of 1601 Australian adults, dietary intake of folate was significantly associated with asthma, but not with markers of airway hyper- responsiveness or atopy.⁶⁰ Three large cross-sectional studies have found a protective effect. In a study of 6,784 Danish adults, higher serum folate levels were found to be associated with lower risks for an asthma diagnosis, but no associations were found with

lung function or atopy⁶¹. On the other hand, in the US NHANES cohort (both children and adults), higher serum folate levels were associated with decreased risks for wheeze, total serum IgE, and atopy, but no association was noted for physician-diagnosed asthma⁶². In a study of 1,030 English adults, dietary folate intake was associated with lower risks for physician-diagnosed asthma⁶³. Finally, four studies have found no associations between either serum folate levels^{34, 64, 65} or folate intake⁶⁶ with asthma or allergies.

As can be seen above, results of post natal studies of folate and asthma and allergy have been inconsistent. Furthermore, randomized controlled trials investigating folic acid supplementation on asthma and allergy susceptibility are lacking. We agree with the conclusion of Blatter and colleagues⁴⁰ that there is insufficient data to exclude an effect of folate on either asthma and allergy development or modification of disease course. Reasons for the heterogeneous results of these studies may include differences in study design and analytic approach, the difficulty of folate status measurements, timing of folate supplementation, confounding due to other exposures, sample size, age, differences in the phenotypic assessment of allergy and atopy, or differences in genetic predisposition. In addition to these study design considerations, Figure 1 shows that other methyl donor nutrients are involved in the pathway leading to DNA methylation. Thus, perturbations in folate intake may not be sufficient to derail the pathway unless there are also changes to the levels of the other methyl donor nutrients. While some of the reports reviewed above^{47, 50, 51, 60, 61, 63, 66} investigated other nutrients or foods, these were studied individually and no consideration was given to the relationships between each of these nutrients. While studies of diet patterns^{67, 68} address this issue to a certain extent, pathway-based or systems-based analyses⁶⁹ might help dissect the complex interrelationships between individual nutrients.

Studies investigating the association of genetic variants in genes in the folate metabolism pathway with asthma and allergy have been limited to date. The vast majority of studies investigating the role of genetic associations have been performed on genetic variants in the methylene-tetrahydrofolate reductase (*MTHFR*) gene due to the association of its common variant C677T with reduced enzymatic function of the gene⁷⁰. The results of studies investigating genetic associations of the C677T *MTHFR* genotype with asthma and atopy have been variable. In a cross-sectional study of 1,671 subjects from Denmark, the *MTHFR* genotype was associated with specific IgE⁷¹. However, in a population-based birth cohort that included genotypic information on 7,356 mothers and 5,364 children from the Avon Longitudinal Study of Parents and Children (ALSPAC) there was no genetic association of the *MTHFR* C677T genotype with atopy, allergy, or asthma in the mothers or children⁴⁶. In addition, a smaller population-based study of 1,189 adults from Denmark also showed no association of the *MTHFR* polymorphism with allergy (IgE, skin-tests) or asthma⁷². Interestingly, in a population-based prospective cohort study of 2,001 children that investigated the interaction between *MTHFR* genetic variants and methyl donor levels on asthma and eczema, a significant interaction of the *MTHFR* C677T mutation and higher folate levels on an increased risk of eczema was identified⁷³. Although there may be a difference in genetic predisposition that underlies the association of methyl donors with asthma and allergy disease susceptibility, these results suggest that there may also be

complex gene-by-environment or gene-gene interactions that have yet to be fully elucidated. Furthermore, genetic studies investigating other genes in this pathway may help elucidate the genetic predisposition to these allergy and asthma.

An additional reason for the heterogeneous effects of folate may be differences in intestinal microbiota in the study populations. It has been shown that bacteria commonly found in human large intestines can produce folate,⁷⁴ and whereas folate is mainly absorbed in the small intestines, absorption also occurs in the large intestines⁷⁵. Several strains of bacteria from the genus *Bifidobacteria* have been shown to produce folate both in vitro⁷⁶ and in vivo.⁷⁷ Bacteria from the genus *Lactobacillus* generally need folate for growth, but one strain, *Lactobacillus plantarum*, has been shown to be a folate producer⁷⁸. It is easy to see how folate production by intestinal microbiota can modify the effects of folate ingested in the diet, and has already been considered in studies of diet and colon cancer risk^{78, 79}. While there is keen interest in the gut microbiome and asthma and allergy risk,^{80, 81} these studies are in their infancy and future studies in cohorts that are collecting both diet and microbiome data will be needed to investigate this link.

Choline

Choline is a tri-methylated water soluble B vitamin that is important in many biologic functions⁸². Although it is synthesized in small quantities by the liver, choline is present in many foods and is a component of nutritional supplements and multivitamins. Choline is naturally found in many food sources including soybeans, egg, peanuts, cauliflower, lentils, and flax seeds⁸³. Choline is a precursor of phosphatidylcholine and sphingomyelin, which are phospholipids that are components of biologic membranes and are involved in maintaining cell structure. Choline has also been shown to aid in the movement of lipids in and out of cells^{84, 85}. In addition, it is a precursor to platelet-activating factor and acetylcholine. Choline can be oxidized into betaine, which provides the methyl groups needed to resynthesize methionine from homocysteine thereby providing methionine for methylation reactions⁸⁶. This pathway is an alternative to one that uses the co-factor 5-methyltetrahydrofolate (Figure 1).

Animal models have demonstrated the anti-inflammatory properties of choline supplementation in inflammatory arthritis⁸⁷. In addition, an in vitro model showed that choline deficiency resulted in the loss of membrane phosphatidyl choline and resulted in the induction of apoptosis⁸⁸. Mice fed with a choline deficient diet showed altered antioxidant properties and an increase in oxidative stress⁸⁹. Furthermore, dietary choline supplementation resulted in decreased reactive oxygen species and a suppressed the inflammatory response to ovalbumin challenge in a murine model of allergic airway disease⁹⁰.

Human data for the association of choline with the development of asthma and allergy is limited. A small study of 23 asthmatic subjects with documented aspirin sensitivity treated with choline magnesium trisalicylate (CMT), demonstrated the safety of choline supplementation, but did not result in an improvement in pulmonary function, nasal congestion, or rhinorrhea⁹¹. However, studies of the treatment of asthmatic subjects with tricholine citrate demonstrated symptomatic improvement in their asthma with decreased

symptom scores, increased symptom-free days, and decreased rescue medication use^{92, 93}. Furthermore, individuals treated with high-dose choline were also noted to have improvements in lung function⁹² and had decreased airway responsiveness⁹³. In a treatment trial of 76 patients with asthma treated with choline or standard pharmacotherapy, choline supplementation resulted in improved quality of life, decreased asthma medication use, improvements in airway responsiveness, and significantly reduced markers of inflammation including interleukin-4 (IL-4), IL-5, and tumor necrosis factor alpha (TNF- α)⁹⁴. Interestingly, in a case-control study investigating the metabolomic signature of the serum of asthmatics compared to non-asthmatic controls, choline was significantly lower in asthmatic subjects⁹⁵. In addition, this metabolomic profile was associated with decreased lung function including the forced expiratory volume in one second (FEV₁) percent predicted⁹⁵, demonstrating a possible role for choline in the pathogenesis of asthma.

More studies are needed to elucidate the effects of choline in the development of asthma and allergies. As with folate, there appears to be a role of the intestinal microbiome in mediating the effects of dietary choline. Through metabolomics studies, the role of gut microbiome metabolism of dietary choline in atherosclerosis has been elucidated. Gut microbiome metabolism of choline and phosphatidylcholine lead to the formation of trimethylamine N-oxide (TMAO),^{96, 97} which was recently shown to increase the risk of major cardiovascular events such as death, myocardial infarction, or stroke.⁹⁸ The role of TMAO in asthma and allergies is unknown, however, the metabolomic study⁹⁵ cited above linking lower choline levels in serum with asthma suggests that this line of research should be investigated further.

Vitamin B12

Vitamin B12 (B12) is a water-soluble vitamin that plays a key role in one-carbon metabolism. Vitamin B12 is naturally found in several foods including fish, meat, poultry, eggs, and dairy products. Vitamin B12 is also found in fortified foods including cereals and grains as well as in dietary supplements²⁹. B12 exists in several forms and contains the mineral cobalt, giving them the name “cobalamins”. Methylcobalamin and 5-deoxyadenosylcobalamin are the forms of B12 that are important in human metabolism⁹⁹. Vitamin B12 is required for red blood cell formation, DNA synthesis, and proper neurologic function¹⁰⁰. Folate metabolism is influenced by vitamin B12 levels¹⁰¹. B12 acts as a co-factor for folate, and provides methyl groups for the synthesis of methionine and its derivative SAM, which is a universal methyl donor for multiple substrates: DNA, RNA, proteins, lipids, and hormones. B12 deficiency results in decreased methylation of deoxyuridylic acid (dUMP) to deoxythymidylic (dTTP), leading to an excessive incorporation of uracil into DNA as well as DNA hypomethylation¹⁰². Animal models have shown that increased B12 is associated with an increased risk of allergic airway disease that is mediated through epigenetic mechanisms including changes in DNA methylation²⁶.

The results of studies investigating B12 intake and the development of atopy in humans have been variable. In a single birth cohort, maternal serum B12 concentrations during pregnancy were significantly associated with an overall increased prevalence of atopic dermatitis in children⁴⁴. However, a study of 763 asthmatic children in Japan, showed that maternal vitamin B12 intake during pregnancy was not associated with the development of

wheeze or eczema⁵⁰. In addition, serum levels and dietary intake of B12 were not associated with asthma and atopy in a large, adult cohort study⁶¹. These studies have been limited by small sample sizes and a lack of interventional studies. Therefore, the role of vitamin B12 in the development of asthma and allergy remains unclear.

Vitamins B2 and B6

Vitamin B2, or riboflavin, is required for several cellular processes. There are a variety of dietary sources of vitamin B2 including eggs, meat, milk and cheese. In addition, it is found in many fortified foods including cereals and rice as well as in nutritional supplements including multivitamins. Vitamin B6 is a water-soluble vitamin that is naturally occurring in many foods. Vitamin B6 is the generic name for six compounds with vitamin B6 activity: pyridoxine, pyridoxal, pyridoxamine, and their respective 5'-phosphate esters. The vitamin B6 compounds are naturally occurring in fish, beef liver, fruits, vegetables, and grains²⁹. In the United States, most adults obtain most of their vitamin B6 from fortified foods including cereals, beef, and poultry. Furthermore, vitamin B6 is found in multivitamins and supplements containing other B complex vitamins. Vitamin B2 and B6 perform a wide variety of biologic functions including assisting in the metabolic pathway of one-carbon compounds (Figure 1).

Although the data for the role of vitamins B2 and B6 in asthma pathogenesis is limited, several small studies have investigated the role of vitamins B2 and B6 on the risk of atopy. In a study investigating maternal vitamin B intake during pregnancy and the subsequent risk of wheeze and eczema in infants, Miyake and colleagues demonstrated no evidence of association of maternal vitamin B2 or B6 intakes with the subsequent risk of wheeze or eczema up to two years of age in Japanese children⁵⁰. In a small cross-sectional study evaluating the prevalence of allergy in school children in Vietnam, vitamin B2 intake was found to be associated with a reduced risk of allergy ($p=0.038$)¹⁰³. Using the European Respiratory Health Survey (ECRHS) data to determine the role of dietary intake on the prevalence of allergic sensitization, Heinrich and colleagues showed no association of vitamin B2 intake and the prevalence of allergy¹⁰⁴. Finally, in a cross-sectional population based study of 1671 subjects with and without atopy in Denmark, dietary intake of vitamins B2 and B6 were not significantly associated with atopy⁷¹. However, there was noted to be at least a borderline interaction between *MTHFR* genotype (C677T) and the dietary intake of vitamins B2 and B6 with a decreased risk of atopy⁷¹.

While there is data to suggest that vitamin B6 level is associated with homocysteine metabolism¹⁰⁵ and that increased dietary intake of vitamin B6 results in decreased plasma homocysteine levels¹⁰⁶, its role in asthma and allergy is not yet clear. A few studies have investigated these vitamins in established asthma. In a small study of 31 patients with asthma participating in a double-blind trial, vitamin B6 supplementation in steroid-dependent asthma was not associated with improvement in lung function, asthma symptom scores, or 24 hour urinary 5-HIAA levels¹⁰⁷. However, in a different study of 15 adult patients with asthma, erythrocyte pyridoxal phosphate concentrations were significantly lower in asthmatics compared to non-asthmatic controls¹⁰⁸. Furthermore, supplementation with vitamin B6 in these subjects was associated with decreased asthma symptoms and

decreased exacerbation rates¹⁰⁸. In a double-blind study of 76 asthmatic children supplementation with vitamin B6 was associated with improvement in asthma symptoms resulting in a reduction in the dosage of asthma-treatment medications¹⁰⁹.

Although definitive clinical trials are lacking, a review of the current literature only supports a limited effect of vitamins B2 and B6 on atopic disease susceptibility and disease treatment due to the discrepancy of the results obtained to date. Some of the variability in the results of these studies may be due to study design, sample size, analytic approach, the difficulty of B vitamin status assessment, and age.

Betaine

Betaine is a water soluble molecule that is involved in a variety of biologic processes including DNA methylation. Betaine is found in a variety of food sources including beets, broccoli, spinach, seafood, and grains. In addition, it is found in small quantities in nutritional supplements and multivitamins¹¹⁰. It is essential in one-carbon metabolism playing an essential role in the transition from homocysteine to methionine. Studies investigating the role of betaine in the development of asthma and allergy are lacking. Furthermore, studies investigating the role of betaine in DNA methylation in humans is limited. In a large cohort study of 534 pregnant women in Project Viva, there was no significant association found between maternal dietary intake of betaine in both the first and second trimesters of pregnancy with cord blood LINE-1 methylation⁵⁶. However, periconceptional intake of betaine was inversely associated with cord-blood methylation, but the effect of betaine on methylation was attenuated after adjustment for cadmium intake⁵⁶. Given its importance in one-carbon metabolism, studies investigating the role of dietary betaine intake in the development of asthma and allergy are warranted.

Summary and Future Directions

Dietary changes are a proposed mechanism underlying the increasing prevalence of asthma and allergic disease worldwide. Several observational studies and a few small interventional trials have attempted to assess the role of dietary methyl donors in the development and treatment of asthma and allergy. Due to the interest in the early origins of asthma and allergic diseases, and because of recommendations for folic acid supplementation in pregnancy for the prevention of neural tube defects, folate intake has been investigated as a potentially modifiable exposure in pregnancy. However, the results of the studies investigating prenatal folate intake or folate status published to date have been mixed, limiting any definitive conclusions on the role of prenatal folate with regards to its association with asthma and allergy. Studies investigating postnatal folate intake or status in childhood or adulthood have also been mixed, and studies of other dietary methyl donors have been limited.

While more studies are needed to allow for more definitive conclusions regarding the effect of these nutrients on asthma and allergies, appropriate attention to study design is essential for future investigations. In addition, two other issues need to be considered in either analyzing existing datasets or designing new studies – broader approaches such as pathway or systems analyses and incorporation of the effects of the intestinal microbiome. Few of the

studies reviewed here have investigated more than one nutrient. Given that these nutrients are intricately related in pathways (Figure 1), future studies need to include analyses of multiple nutrients. Furthermore, given the complexities of the one-carbon metabolism pathway, analyses including a systems-based approach or pathway analyses may further our understanding of the role of dietary methyl donors in the development of atopic disease⁶⁹.

Finally, the role of the microbiome in health is gaining wider recognition. The intestinal microbiome has been shown to be actively involved in the production and metabolism of these nutrients. As reviewed, several strains of the genus *Bifidobacteria* have the ability to produce folate. The microbiome also has a role in either the production or metabolism of other B vitamins (reviewed in LeBlanc et al.⁷⁴). B vitamin production by intestinal bacteria are mediating or modifying the effects of ingested nutrients, and may partly explain the heterogeneity in the results of the studies investigating these nutrients to date. The design of future studies will need to incorporate analyses of the intestinal microbiome to fully understand the effects of these nutrients on human health in general, and in the development of asthma and allergies in particular. For example, in the ongoing Vitamin D Antenatal Asthma Reduction Trial (VDAART, U01 HL091528 and [ClinicalTrials.gov # NCT00920621](https://clinicaltrials.gov/ct2/show/study/NCT00920621)), we are collecting both pre- and postnatal dietary information (including vitamins other than vitamin D) and stool samples (R01HL108818) to enable such studies. Integration of these types of datasets (i.e. dietary data and microbiome data) will lead to novel insights into the biologic basis of health and disease^{111, 112}.

Acknowledgments

Funding Sources: R01 HL111108 and Dr. Sharma is funded by K08 HL096833. Dr. Litonjua is funded by U01 HL091528 and R01 HL111108.

Abbreviations

(CMT)	Choline Magnesium Trisalicylate
(SAH)	S-Adenosyl-homocysteine
(SAM)	S-Adenosyl-methionine
(THF)	Tetrahydrofolate
(B2)	Vitamin B2
(B6)	Vitamin B6
(B12)	Vitamin B12

References

1. Ozdoganoglu T, Songu M. The burden of allergic rhinitis and asthma. Therapeutic advances in respiratory disease. 2012; 6:11–23. [PubMed: 22179899]
2. To T, Stanojevic S, Moores G, Gershon AS, Bateman ED, Cruz AA, et al. Global asthma prevalence in adults: findings from the cross-sectional world health survey. BMC public health. 2012; 12:204. [PubMed: 22429515]

3. Vital signs: asthma prevalence, disease characteristics, and self-management education: United States, 2001--2009. *MMWR. Morbidity and mortality weekly report*. 2011; 60:547--52. [PubMed: 21544044]
4. Pearce N, Ait-Khaled N, Beasley R, Mallol J, Keil U, Mitchell E, et al. Worldwide trends in the prevalence of asthma symptoms: phase III of the International Study of Asthma and Allergies in Childhood (ISAAC). *Thorax*. 2007; 62:758--66. [PubMed: 17504817]
5. Masoli M, Fabian D, Holt S, Beasley R. The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy*. 2004; 59:469--78. [PubMed: 15080825]
6. Williams H, Stewart A, von Mutius E, Cookson W, Anderson HR. Is eczema really on the increase worldwide? *The Journal of allergy and clinical immunology*. 2008; 121:947--54. e15. [PubMed: 18155278]
7. Bjorksten B, Clayton T, Ellwood P, Stewart A, Strachan D. Worldwide time trends for symptoms of rhinitis and conjunctivitis: Phase III of the International Study of Asthma and Allergies in Childhood. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology*. 2008; 19:110--24. [PubMed: 17651373]
8. Eder W, Ege MJ, von Mutius E. The asthma epidemic. *The New England journal of medicine*. 2006; 355:2226--35. [PubMed: 17124020]
9. Lewis S, Butland B, Strachan D, Bynner J, Richards D, Butler N, et al. Study of the aetiology of wheezing illness at age 16 in two national British birth cohorts. *Thorax*. 1996; 51:670--6. [PubMed: 8882071]
10. Kerkhof M, Boezen HM, Granell R, Wijga AH, Brunekreef B, Smit HA, et al. Transient early wheeze and lung function in early childhood associated with chronic obstructive pulmonary disease genes. *The Journal of allergy and clinical immunology*. 2013
11. Mitchell EA, Beasley R, Keil U, Montefort S, Odhiambo J. The association between tobacco and the risk of asthma, rhinoconjunctivitis and eczema in children and adolescents: analyses from Phase Three of the ISAAC programme. *Thorax*. 2012; 67:941--9. [PubMed: 22693180]
12. Lin SY, Reh DD, Clipp S, Irani L, Navas-Acien A. Allergic rhinitis and secondhand tobacco smoke: a population-based study. *American journal of rhinology & allergy*. 2011; 25:e66--71. [PubMed: 21679502]
13. Strachan DP, Butland BK, Anderson HR. Incidence and prognosis of asthma and wheezing illness from early childhood to age 33 in a national British cohort. *BMJ*. 1996; 312:1195--9. [PubMed: 8634562]
14. Withers NJ, Low L, Holgate ST, Clough JB. The natural history of respiratory symptoms in a cohort of adolescents. *American journal of respiratory and critical care medicine*. 1998; 158:352--7. [PubMed: 9700106]
15. Gruzdeva O, Bergstrom A, Hulchiy O, Kull I, Lind T, Melen E, et al. Exposure to air pollution from traffic and childhood asthma until 12 years of age. *Epidemiology*. 2013; 24:54--61. [PubMed: 23222555]
16. Gehring U, Wijga AH, Brauer M, Fischer P, de Jongste JC, Kerkhof M, et al. Traffic-related air pollution and the development of asthma and allergies during the first 8 years of life. *American journal of respiratory and critical care medicine*. 2010; 181:596--603. [PubMed: 19965811]
17. Nicolai T, Carr D, Weiland SK, Duhme H, von Ehrenstein O, Wagner C, et al. Urban traffic and pollutant exposure related to respiratory outcomes and atopy in a large sample of children. *The European respiratory journal*. 2003; 21:956--63. [PubMed: 12797488]
18. Allan K, Devereux G. Diet and asthma: nutrition implications from prevention to treatment. *Journal of the American Dietetic Association*. 2011; 111:258--68. [PubMed: 21272700]
19. McKeever TM, Britton J. Diet and asthma. *American journal of respiratory and critical care medicine*. 2004; 170:725--9. [PubMed: 15256393]
20. Nurmatov U, Devereux G, Sheikh A. Nutrients and foods for the primary prevention of asthma and allergy: systematic review and meta-analysis. *The Journal of allergy and clinical immunology*. 2011; 127:724--33. e1--30. [PubMed: 21185068]
21. Ptashne M. On the use of the word 'epigenetic'. *Current biology : CB*. 2007; 17:R233--6. [PubMed: 17407749]

22. Anway MD, Cupp AS, Uzumcu M, Skinner MK. Epigenetic transgenerational actions of endocrine disruptors and male fertility. *Science*. 2005; 308:1466–9. [PubMed: 15933200]
23. Edwards TM, Myers JP. Environmental exposures and gene regulation in disease etiology. *Environmental health perspectives*. 2007; 115:1264–70. [PubMed: 17805414]
24. Sordillo JE, Lange NE, Tarantini L, Bollati V, Zanobetti A, Sparrow D, et al. Allergen sensitization is associated with increased DNA methylation in older men. *International archives of allergy and immunology*. 2013; 161:37–43. [PubMed: 23257623]
25. Wolff GL, Kodell RL, Moore SR, Cooney CA. Maternal epigenetics and methyl supplements affect agouti gene expression in Avy/a mice. *FASEB J*. 1998; 12:949–57. [PubMed: 9707167]
26. Hollingsworth JW, Maruoka S, Boon K, Garantzotis S, Li Z, Tomfohr J, et al. In utero supplementation with methyl donors enhances allergic airway disease in mice. *The Journal of clinical investigation*. 2008; 118:3462–9. [PubMed: 18802477]
27. Litonjua AA. Dietary factors and the development of asthma. *Immunology and allergy clinics of North America*. 2008; 28:603–29. ix. [PubMed: 18572110]
28. Varraso R. Nutrition and asthma. *Current allergy and asthma reports*. 2012; 12:201–10. [PubMed: 22407651]
29. Dietary Fact Sheet: Vitamin B6. 2011.
30. Bailey, L.; Gregory, J. Folate. International Life Sciences Institute; Washington DC: 2006.
31. Wright AJ, Dainty JR, Finglas PM. Folic acid metabolism in human subjects revisited: potential implications for proposed mandatory folic acid fortification in the UK. *The British journal of nutrition*. 2007; 98:667–75. [PubMed: 17617936]
32. Powers HJ. Folic acid under scrutiny. *The British journal of nutrition*. 2007; 98:665–6. [PubMed: 17697404]
33. Yetley EA, Pfeiffer CM, Phinney KW, Bailey RL, Blackmore S, Bock JL, et al. Biomarkers of vitamin B-12 status in NHANES: a roundtable summary. *The American journal of clinical nutrition*. 2011; 94:313S–21S. [PubMed: 21593512]
34. Lin JH, Matsui W, Aloe C, Peng RD, Diette GB, Breyse PN, et al. Relationships between folate and inflammatory features of asthma. *The Journal of allergy and clinical immunology*. 2013; 131:918–20. [PubMed: 23237419]
35. Moens AL, Champion HC, Claeys MJ, Tavazzi B, Kaminski PM, Wolin MS, et al. High-dose folic acid pretreatment blunts cardiac dysfunction during ischemia coupled to maintenance of high-energy phosphates and reduces postreperfusion injury. *Circulation*. 2008; 117:1810–9. [PubMed: 18362233]
36. Richard-Tremblay AA, Sheehy O, Berard A. Annual trends in use of periconceptional folic Acid and birth prevalence of major congenital malformations. *Current drug safety*. 2013; 8:153–61. [PubMed: 23845112]
37. Hure A, Young A, Smith R, Collins C. Diet and pregnancy status in Australian women. *Public health nutrition*. 2009; 12:853–61. [PubMed: 18647425]
38. Recommendations for the use of folic acid to reduce the number of cases of spina bifida and other neural tube defects. MMWR. Recommendations and reports : Morbidity and mortality weekly report. Recommendations and reports / Centers for Disease Control. 1992; 41:1–7.
39. From the Centers for Disease Control and Prevention. Recommendations for use of folic acid to reduce number of spina bifida cases and other neural tube defects. *JAMA : the journal of the American Medical Association*. 1993; 269:1233, 6–8. [PubMed: 8437292]
40. Blatter J, Han YY, Forno E, Brehm J, Bodnar L, Celedon JC. Folate and asthma. *American journal of respiratory and critical care medicine*. 2013; 188:12–7. [PubMed: 23650899]
41. Haberg SE, London SJ, Stigum H, Nafstad P, Nystad W. Folic acid supplements in pregnancy and early childhood respiratory health. *Archives of disease in childhood*. 2009; 94:180–4. [PubMed: 19052032]
42. Haberg SE, London SJ, Nafstad P, Nilsen RM, Ueland PM, Vollset SE, et al. Maternal folate levels in pregnancy and asthma in children at age 3 years. *The Journal of allergy and clinical immunology*. 2011; 127:262–4. 4, e1. [PubMed: 21094522]

43. Whitrow MJ, Moore VM, Rumbold AR, Davies MJ. Effect of supplemental folic acid in pregnancy on childhood asthma: a prospective birth cohort study. *American journal of epidemiology*. 2009; 170:1486–93. [PubMed: 19880541]
44. Kiefte-de Jong JC, Timmermans S, Jaddoe VW, Hofman A, Tiemeier H, Steegers EA, et al. High circulating folate and vitamin B-12 concentrations in women during pregnancy are associated with increased prevalence of atopic dermatitis in their offspring. *The Journal of nutrition*. 2012; 142:731–8. [PubMed: 22399526]
45. Bekkers MB, Elstgeest LE, Scholtens S, Haveman-Nies A, de Jongste JC, Kerkhof M, et al. Maternal use of folic acid supplements during pregnancy, and childhood respiratory health and atopy. *The European respiratory journal*. 2012; 39:1468–74. [PubMed: 22034647]
46. Granell R, Heron J, Lewis S, Davey Smith G, Sterne JA, Henderson J. The association between mother and child MTHFR C677T polymorphisms, dietary folate intake and childhood atopy in a population-based, longitudinal birth cohort. *Clinical and experimental allergy : journal of the British Society for Allergy and Clinical Immunology*. 2008; 38:320–8. [PubMed: 18070159]
47. Litonjua AA, Rifas-Shiman SL, Ly NP, Tantisira KG, Rich-Edwards JW, Camargo CA Jr. et al. Maternal antioxidant intake in pregnancy and wheezing illnesses in children at 2 y of age. *The American journal of clinical nutrition*. 2006; 84:903–11. [PubMed: 17023719]
48. Magdelijns FJ, Mommers M, Penders J, Smits L, Thijs C. Folic acid use in pregnancy and the development of atopy, asthma, and lung function in childhood. *Pediatrics*. 2011; 128:e135–44. [PubMed: 21690114]
49. Martinussen MP, Risnes KR, Jacobsen GW, Bracken MB. Folic acid supplementation in early pregnancy and asthma in children aged 6 years. *American journal of obstetrics and gynecology*. 2012; 206:72, e1–7. [PubMed: 21982024]
50. Miyake Y, Sasaki S, Tanaka K, Hirota Y. Maternal B vitamin intake during pregnancy and wheeze and eczema in Japanese infants aged 16-24 months: the Osaka Maternal and Child Health Study. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology*. 2011; 22:69–74. [PubMed: 20561231]
51. Nwaru BI, Erkkola M, Ahonen S, Kaila M, Kronberg-Kippila C, Ilonen J, et al. Intake of antioxidants during pregnancy and the risk of allergies and asthma in the offspring. *European journal of clinical nutrition*. 2011; 65:937–43. [PubMed: 21559033]
52. Karmaus W, Ziyab AH, Everson T, Holloway JW. Epigenetic mechanisms and models in the origins of asthma. *Current opinion in allergy and clinical immunology*. 2013; 13:63–9. [PubMed: 23242116]
53. Fryer AA, Nafee TM, Ismail KM, Carroll WD, Emes RD, Farrell WE. LINE-1 DNA methylation is inversely correlated with cord plasma homocysteine in man: a preliminary study. *Epigenetics : official journal of the DNA Methylation Society*. 2009; 4:394–8. [PubMed: 19755846]
54. Park BH, Kim YJ, Park JS, Lee HY, Ha EH, Min JW, et al. [Folate and homocysteine levels during pregnancy affect DNA methylation in human placenta]. *Journal of preventive medicine and public health = Yebang Uihakhoe chi*. 2005; 38:437–42. [PubMed: 16358830]
55. Steegers-Theunissen RP, Obermann-Borst SA, Kremer D, Lindemans J, Siebel C, Steegers EA, et al. Periconceptional maternal folic acid use of 400 microg per day is related to increased methylation of the IGF2 gene in the very young child. *PloS one*. 2009; 4:e7845. [PubMed: 19924280]
56. Boeke CE, Baccarelli A, Kleinman KP, Burris HH, Litonjua AA, Rifas-Shiman SL, et al. Gestational intake of methyl donors and global LINE-1 DNA methylation in maternal and cord blood: prospective results from a folate-replete population. *Epigenetics : official journal of the DNA Methylation Society*. 2012; 7:253–60. [PubMed: 22430801]
57. Devries A, Vercelli D. Epigenetics of human asthma and allergy: promises to keep. *Asian Pacific journal of allergy and immunology / launched by the Allergy and Immunology Society of Thailand*. 2013; 31:183–9. [PubMed: 24053700]
58. Kim HP, Lee YS, Park JH, Kim YJ. Transcriptional and epigenetic networks in the development and maturation of dendritic cells. *Epigenomics*. 2013; 5:195–204. [PubMed: 23566096]

59. Okupa AY, Lemanske RF Jr, Jackson DJ, Evans MD, Wood RA, Matsui EC. Early-life folate levels are associated with incident allergic sensitization. *The Journal of allergy and clinical immunology*. 2013; 131:226–8. e1–2. [PubMed: 23040888]
60. Woods RK, Walters EH, Raven JM, Wolfe R, Ireland PD, Thien FC, et al. Food and nutrient intakes and asthma risk in young adults. *The American journal of clinical nutrition*. 2003; 78:414–21. [PubMed: 12936923]
61. Thuesen BH, Husemoen LL, Ovesen L, Jorgensen T, Fenger M, Gilderson G, et al. Atopy, asthma, and lung function in relation to folate and vitamin B(12) in adults. *Allergy*. 2010; 65:1446–54. [PubMed: 20456312]
62. Matsui EC, Matsui W. Higher serum folate levels are associated with a lower risk of atopy and wheeze. *The Journal of allergy and clinical immunology*. 2009; 123:1253–9. e2. [PubMed: 19409604]
63. Patel BD, Welch AA, Bingham SA, Luben RN, Day NE, Khaw KT, et al. Dietary antioxidants and asthma in adults. *Thorax*. 2006; 61:388–93. [PubMed: 16467075]
64. Farres MN, Shahin RY, Melek NA, El-Kabarity RH, Arafa NA. Study of folate status among Egyptian asthmatics. *Internal medicine*. 2011; 50:205–11. [PubMed: 21297321]
65. Shaheen MA, Attia EA, Louka ML, Bareedy N. STUDY OF THE ROLE OF SERUM FOLIC ACID IN ATOPIC DERMATITIS: A CORRELATION WITH SERUM IgE AND DISEASE SEVERITY. *Indian journal of dermatology*. 2011; 56:673–7. [PubMed: 22345769]
66. Bueso AK, Berntsen S, Mowinckel P, Andersen LF, Lodrup Carlsen KC, Carlsen KH. Dietary intake in adolescents with asthma--potential for improvement. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology*. 2011; 22:19–24. [PubMed: 20961335]
67. Garcia-Marcos L, Castro-Rodriguez JA, Weinmayr G, Panagiotakos DB, Priftis KN, Nagel G. Influence of Mediterranean diet on asthma in children: a systematic review and meta-analysis. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology*. 2013; 24:330–8. [PubMed: 23578354]
68. Lange NE, Rifas-Shiman SL, Camargo CA Jr, Gold DR, Gillman MW, Litonjua AA. Maternal dietary pattern during pregnancy is not associated with recurrent wheeze in children. *The Journal of allergy and clinical immunology*. 2010; 126:250–5. 5, e1–4. [PubMed: 20584543]
69. Moore JB, Weeks ME. Proteomics and systems biology: current and future applications in the nutritional sciences. *Advances in nutrition*. 2011; 2:355–64. [PubMed: 22332076]
70. Frosst P, Blom HJ, Milos R, Goyette P, Sheppard CA, Matthews RG, et al. A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. *Nature genetics*. 1995; 10:111–3. [PubMed: 7647779]
71. Husemoen LL, Toft U, Fenger M, Jorgensen T, Johansen N, Linneberg A. The association between atopy and factors influencing folate metabolism: is low folate status causally related to the development of atopy? *International journal of epidemiology*. 2006; 35:954–61. [PubMed: 16766537]
72. Thuesen BH, Husemoen LL, Fenger M, Linneberg A. Lack of association between the MTHFR (C677T) polymorphism and atopic disease. *The clinical respiratory journal*. 2009; 3:102–8. [PubMed: 20298385]
73. van der Valk RJ, Kiefte-de Jong JC, Sonnenschein-van der Voort AM, Duijts L, Hafkamp-de Groen E, Moll HA, et al. Neonatal folate, homocysteine, vitamin B12 levels and methylenetetrahydrofolate reductase variants in childhood asthma and eczema. *Allergy*. 2013; 68:788–95. [PubMed: 23692062]
74. LeBlanc JG, Milani C, de Giori GS, Sesma F, van Sinderen D, Ventura M. Bacteria as vitamin suppliers to their host: a gut microbiota perspective. *Current opinion in biotechnology*. 2013; 24:160–8. [PubMed: 22940212]
75. Aufreiter S, Gregory JF 3rd, Pfeiffer CM, Fazili Z, Kim YI, Marcon N, et al. Folate is absorbed across the colon of adults: evidence from cecal infusion of (13)C-labeled [6S]-5-formyltetrahydrofolic acid. *The American journal of clinical nutrition*. 2009; 90:116–23. [PubMed: 19439459]

76. D'Aimmo MR, Mattarelli P, Biavati B, Carlsson NG, Andlid T. The potential of bifidobacteria as a source of natural folate. *Journal of applied microbiology*. 2012; 112:975–84. [PubMed: 22335359]
77. Strozzi GP, Mogna L. Quantification of folic acid in human feces after administration of *Bifidobacterium* probiotic strains. *Journal of clinical gastroenterology*. 2008; 42(Suppl 3 Pt 2):S179–84. [PubMed: 18685499]
78. Rossi M, Amaretti A, Raimondi S. Folate production by probiotic bacteria. *Nutrients*. 2011; 3:118–34. [PubMed: 22254078]
79. O'Keefe SJ, Ou J, Aufreiter S, O'Connor D, Sharma S, Sepulveda J, et al. Products of the colonic microbiota mediate the effects of diet on colon cancer risk. *The Journal of nutrition*. 2009; 139:2044–8. [PubMed: 19741203]
80. Ly NP, Litonjua A, Gold DR, Celedon JC. Gut microbiota, probiotics, and vitamin D: interrelated exposures influencing allergy, asthma, and obesity? *The Journal of allergy and clinical immunology*. 2011; 127:1087–94. quiz 95-6. [PubMed: 21419479]
81. McLoughlin RM, Mills KH. Influence of gastrointestinal commensal bacteria on the immune responses that mediate allergy and asthma. *The Journal of allergy and clinical immunology*. 2011; 127:1097–107. quiz 108-9. [PubMed: 21420159]
82. Zeisel SH, da Costa KA. Choline: an essential nutrient for public health. *Nutrition reviews*. 2009; 67:615–23. [PubMed: 19906248]
83. Konstantinova SV, Tell GS, Vollset SE, Ulvik A, Drevon CA, Ueland PM. Dietary patterns, food groups, and nutrients as predictors of plasma choline and betaine in middle-aged and elderly men and women. *The American journal of clinical nutrition*. 2008; 88:1663–9. [PubMed: 19064529]
84. Li Z, Vance DE. Phosphatidylcholine and choline homeostasis. *J Lipid Res*. 2008; 49:1187–94. [PubMed: 18204095]
85. Blusztajn JK. Choline, a vital amine. *Science*. 1998; 281:794–5. [PubMed: 9714685]
86. McNeil SD, Nuccio ML, Ziemak MJ, Hanson AD. Enhanced synthesis of choline and glycine betaine in transgenic tobacco plants that overexpress phosphoethanolamine N-methyltransferase. *Proc Natl Acad Sci U S A*. 2001; 98:10001–5. [PubMed: 11481443]
87. Ganley O, Graessle O, Robinson H. Anti-inflammatory activity of components obtained from egg-yolk, peanut oil, and soya-bean lecithin. *J Lab Clin Med*. 1958; 51:709–14. [PubMed: 13539486]
88. Yen CL, Mar MH, Zeisel SH. Choline deficiency-induced apoptosis in PC12 cells is associated with diminished membrane phosphatidylcholine and sphingomyelin, accumulation of ceramide and diacylglycerol, and activation of a caspase. *FASEB J*. 1999; 13:135–42. [PubMed: 9872938]
89. Yoshida Y, Itoh N, Hayakawa M, Habuchi Y, Inoue R, Chen ZH, et al. Lipid peroxidation in mice fed a choline-deficient diet as evaluated by total hydroxyoctadecadienoic acid. *Nutrition*. 2006; 22:303–11. [PubMed: 16500556]
90. Mehta AK, Arora N, Gaur SN, Singh BP. Choline supplementation reduces oxidative stress in mouse model of allergic airway disease. *Eur J Clin Invest*. 2009; 39:934–41. [PubMed: 19563446]
91. Szczeklik A, Nizankowska E, Dworski R. Choline magnesium trisalicylate in patients with aspirin-induced asthma. *The European respiratory journal*. 1990; 3:535–9. [PubMed: 2198165]
92. Gupta SK, Gaur SN. A placebo controlled trial of two dosages of LPC antagonist--choline in the management of bronchial asthma. *The Indian journal of chest diseases & allied sciences*. 1997; 39:149–56. [PubMed: 9357148]
93. Gaur SN, Agarwal G, Gupta SK. Use of LPC antagonist, choline, in the management of bronchial asthma. *The Indian journal of chest diseases & allied sciences*. 1997; 39:107–13. [PubMed: 9339609]
94. Mehta AK, Singh BP, Arora N, Gaur SN. Choline attenuates immune inflammation and suppresses oxidative stress in patients with asthma. *Immunobiology*. 2010; 215:527–34. [PubMed: 19897276]
95. Jung J, Kim SH, Lee HS, Choi GS, Jung YS, Ryu DH, et al. Serum metabolomics reveals pathways and biomarkers associated with asthma pathogenesis. *Clinical and experimental allergy : journal of the British Society for Allergy and Clinical Immunology*. 2013; 43:425–33. [PubMed: 23517038]
96. al-Waiz M, Mikov M, Mitchell SC, Smith RL. The exogenous origin of trimethylamine in the mouse. *Metabolism: clinical and experimental*. 1992; 41:135–6. [PubMed: 1736035]

97. Wang Z, Klipfell E, Bennett BJ, Koeth R, Levison BS, Dugar B, et al. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature*. 2011; 472:57–63. [PubMed: 21475195]
98. Tang WH, Wang Z, Levison BS, Koeth RA, Britt EB, Fu X, et al. Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *The New England journal of medicine*. 2013; 368:1575–84. [PubMed: 23614584]
99. Klee GG. Cobalamin and folate evaluation: measurement of methylmalonic acid and homocysteine vs vitamin B(12) and folate. *Clinical chemistry*. 2000; 46:1277–83. [PubMed: 10926922]
100. Combs, G. Vitamin B12 in *The Vitamins*. Academic Press, Inc; New York: 1992.
101. Anderson OS, Sant KE, Dolinoy DC. Nutrition and epigenetics: an interplay of dietary methyl donors, one-carbon metabolism and DNA methylation. *The Journal of nutritional biochemistry*. 2012; 23:853–9. [PubMed: 22749138]
102. Okun JG, Horster F, Farkas LM, Feyh P, Hinz A, Sauer S, et al. Neurodegeneration in methylmalonic aciduria involves inhibition of complex II and the tricarboxylic acid cycle, and synergistically acting excitotoxicity. *The Journal of biological chemistry*. 2002; 277:14674–80. [PubMed: 11847233]
103. Quyen DT, Irei AV, Sato Y, Ota F, Fujimaki Y, Sakai T, et al. Nutritional factors, parasite infection and allergy in rural and suburban Vietnamese school children. *The journal of medical investigation : JMI*. 2004; 51:171–7. [PubMed: 15460903]
104. Heinrich J, Holscher B, Bolte G, Winkler G. Allergic sensitization and diet: ecological analysis in selected European cities. *The European respiratory journal*. 2001; 17:395–402. [PubMed: 11405517]
105. Ubbink JB, van der Merwe A, Delport R, Allen RH, Stabler SP, Riezler R, et al. The effect of a subnormal vitamin B-6 status on homocysteine metabolism. *The Journal of clinical investigation*. 1996; 98:177–84. [PubMed: 8690790]
106. Tucker KL, Olson B, Bakun P, Dallal GE, Selhub J, Rosenberg IH. Breakfast cereal fortified with folic acid, vitamin B-6, and vitamin B-12 increases vitamin concentrations and reduces homocysteine concentrations: a randomized trial. *The American journal of clinical nutrition*. 2004; 79:805–11. [PubMed: 15113718]
107. Sur S, Camara M, Buchmeier A, Morgan S, Nelson HS. Double-blind trial of pyridoxine (vitamin B6) in the treatment of steroid-dependent asthma. *Annals of allergy*. 1993; 70:147–52. [PubMed: 8430923]
108. Reynolds RD, Natta CL. Depressed plasma pyridoxal phosphate concentrations in adult asthmatics. *The American journal of clinical nutrition*. 1985; 41:684–8. [PubMed: 3984921]
109. Collipp PJ, Goldzier S 3rd, Weiss N, Soleymani Y, Snyder R. Pyridoxine treatment of childhood bronchial asthma. *Annals of allergy*. 1975; 35:93–7. [PubMed: 1096686]
110. Craig SA. Betaine in human nutrition. *The American journal of clinical nutrition*. 2004; 80:539–49. [PubMed: 15321791]
111. Claus SP, Swann JR. Nutrimetabonomics: applications for nutritional sciences, with specific reference to gut microbial interactions. *Annual review of food science and technology*. 2013; 4:381–99.
112. Francavilla R, Calasso M, Calace L, Siragusa S, Ndagijimana M, Vernocchi P, et al. Effect of lactose on gut microbiota and metabolome of infants with cow's milk allergy. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology*. 2012; 23:420–7. [PubMed: 22435727]

Key Messages

- Results of studies investigating the effect of dietary methyl donors on asthma and allergy susceptibility have been variable.
- The current evidence does not support the routine supplementation of dietary methyl donors for the primary prevention or treatment of asthma and allergy.
- Future studies need to incorporate pathway or systems-based analyses and the effects of the microbiome on these nutrients.

Capsule Summary

Dietary changes in methyl donors may explain the increase in atopic disease. Studies investigating methyl donor supplementation with atopy have had variable results. Insufficient evidence exists to recommend dietary methyl donor supplementation to treat atopy.

Figure 1.

Figure 1 represents a simplified diagram of the one-carbon metabolic pathway. The figure demonstrates where dietary methyl donors, identified in red, are involved in the one-carbon metabolic pathway and how they influence DNA methylation. DNA methylation depends upon the availability of methyl groups from *S*-adenosylmethionine, which is derived from methionine. This figure highlights the connection between folate, choline, methionine, betaine, and vitamins B2, B6, and B12 on DNA methylation. The pathways intersect at the formation of methionine from homocysteine.

Table 1

Methyl donor sources, function, and mechanisms of action

Methyl Donor	Dietary Sources[*]	Function	Mechanism of Action in Methylation Reactions
Folate	Green leafy vegetables, fruits, nuts, beans, grains, fortified cereals, eggs, dairy	Purine and pyrimidine production, amino acid production, and cell division	Production of S-adenosylmethionine (SAM)
Choline	Soybeans, egg, peanuts, cauliflower, lentils, and flax seeds	Precursor to phosphatidylcholine and sphingomyelin, Aids in the movement of lipids into and out of cells	Production of SAM
Vitamin B12	Fish, meat, poultry, eggs, and dairy products	Red blood cell formation, DNA synthesis, and neurologic function	Methionine Synthesis
Vitamin B2 (riboflavin)	Eggs, meat, milk and cheese	Energy production, antioxidant, red blood cell production	Methionine production
Vitamin B6 (pyridoxine)	Fish, beef liver, fruits, vegetables, and grains	Homocysteine metabolism, metabolism of fats and protein, nervous system development	Folate metabolism
Betaine	Beets, broccoli, spinach, seafood, and grains	Homocysteine metabolism, immune modulation	Methionine production

^{*} Dietary sources other than nutritional supplements or fortified foods.

Table 2

Results of studies investigating the role of non-folate methyl donors in the primary prevention or treatment of asthma and allergy

Methyl donor	Potential mechanism of action	Observational studies*	Randomized-control trials
Choline	Animal models demonstrate anti-inflammatory properties and antioxidant properties of choline.	No observational studies.	No primary prevention studies.
			RCT of asthmatic subjects showed no improvement in asthma and allergy symptoms or lung function ⁹¹ . (n=23)
			RCT of asthmatic subjects showed choline supplementation resulted in decreased asthma symptom scores, increased symptom-free days, and decreased medication use ⁹³ .
			RCT of asthmatic patients showed choline supplementation resulted in improved quality of life, decreased airway responsiveness, and decreased inflammatory markers ⁹² .
Vitamin B12	A co-factor for folate. Provides methyl groups for the synthesis of methionine and S-adenosyl-methionine.	Maternal serum vitamin B12 was associated with increased atopic dermatitis in a birth cohort ⁴⁴ . (n=8,742)	No randomized controlled trials.
		Japanese birth cohort showed maternal vitamin B12 intake during pregnancy was not associated with wheeze or eczema ⁵⁰ . (n=763)	
		Serum vitamin B12 level and dietary intake was not associated with asthma or atopy in an adult cohort study ⁶¹ . (n=6,784)	
Vitamin B2	Important in methionine synthesis and its role as a methyl donor.	Japanese birth cohort showed maternal vitamin B2 intake during pregnancy was not associated with wheeze or eczema ⁵⁰ . (n=763)	No RCTs.
		Riboflavin intake was associated with a reduced risk of allergy ¹⁰³ . (n=195)	
		Adult cohort study showed no association with vitamin B2 with the prevalence of allergy ¹⁰⁴ . (n=3,872)	
		Cross-sectional study of adult subjects with and without atopy showed no association of vitamin B2 with atopy ⁷¹ . (n=1,671)	

Methyl donor	Potential mechanism of action	Observational studies*	Randomized-control trials
Vitamin B6	Important in methionine synthesis and its role as a methyl donor.	Japanese birth cohort showed maternal vitamin B6 intake during pregnancy was not associated with wheeze or eczema ⁵⁰ . (n=763)	RCT of asthmatic patients showed that pyridoxine supplementation was not associated with asthma symptoms or lung function ¹⁰⁷ . (n=31)
		Cross-sectional study of adult subjects with and without atopy showed no association of vitamin B2 with atopy ⁷¹ (n=1,671)	RCT of adult asthmatics vitamin B6 levels were associated with decreased asthma symptoms and exacerbation rates ¹⁰⁸ . (n=15)
			RCT of childhood asthmatics treated with pyridoxine showed significant improvement in use of bronchodilators and cortisone use ¹⁰⁹ (n=76)
Betaine	Important in methionine synthesis and its role as a methyl donor. Involved in global methylation	No observational studies.	No RCTs.

* Observational studies include birth cohort studies and cross-sectional studies.