Published in final edited form as:

J Allergy Clin Immunol. 2014 May ; 133(5): 1390–1399.e6. doi:10.1016/j.jaci.2014.01.021.

Exposure to food allergens through inflamed skin promotes intestinal food allergy via the TSLP-basophil axis

Mario Noti, PhD^{a,b,e}, Brian S. Kim, MD^{a,b,d}, Mark C. Siracusa, PhD^{a,b}, Gregory D. Rak, VMD, PhD^{a,b}, Masato Kubo, PhD^{f,g}, Amin E. Moghaddam, MD^h, Quentin A. Sattentau, PhD^h, Michael R. Comeau, BScⁱ, Jonathan M. Spergel, MD, PhD^j, and David Artis, PhD^{a,b,c}

^aDepartment of Microbiology, University of Pennsylvania, Philadelphia, PA 19104, USA ^bInstitute for Immunology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA ^cDepartment of Pathobiology, School of Veterinary Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA ^dDepartment of Dermatology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA ^eDivision of Experimental Pathology, Institute of Pathology, University of Bern, Bern, CH-3010, Switzerland ^fLaboratory for cytokine regulation, Research Center for Integrative Medical Science, RIKEN Yokohama Institute, Kanagawa 230-0045, Japan ^gDivision of Molecular Pathology, Research Institute for Biomedical Science, Tokyo University of Science, Chiba, 278-0022, Japan ^hThe Sir William Dunn School of Pathology, The University of Oxford, Oxford OX13RE, UK ⁱInflammation Research, Amgen Inc., Seattle, WA 98119, USA ^jDepartment of Pediatrics, Division of Allergy and Immunology, Children's Hospital of Philadelphia, Philadelphia, PA 19104, USA

Abstract

Background—Exposure to food allergens through a disrupted skin barrier has been recognized as a potential factor in the increasing prevalence of food allergy.

Objective—To test the immunological mechanisms by which epicutaneous sensitization to food allergens predisposes to intestinal food allergy.

Methods—Mice were epicutaneously sensitized with ovalbumin (OVA) or peanut on an atopic dermatitis-like skin lesion followed by intragastric antigen challenge. Antigen-specific serum IgE levels and Th2 cytokine responses were measured by ELISA. Expression of type-2 cytokines and mast cell proteases in the intestine were measured by real-time PCR. Accumulation of basophils in

© 2014 American Academy of Allergy, Asthma and Immunology. Published by Mosby, Inc. All rights reserved.

Corresponding author: David Artis, Department of Microbiology, Institute for Immunology, Perelman School of Medicine, University of Pennsylvania, 421 Curie Boulevard, BRB II/III, room 356, Philadelphia, PA 19104-6160, USA, dartis@mail.med.upenn.edu.

Disclosure of potential conflict of interest: M.R.C is an employee and shareholder of Amgen.

AUTHOR CONTRIBUTIONS

M.N., B.S.K., M.C.S., G.D.R., and D.A. designed and performed experiments. M.K. provided Baso-DTR mice and M.R.C. provided TSLPR-deficient mice, murine rTSLP and cDNA TSLP plasmid. M.N., B.S.K., M.C.S., M.R.C., J.M.S and D.A. analyzed the data. M.N. and D.A. wrote the manuscript and all authors critically reviewed the manuscript.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

the skin and mast cells in the intestine was examined by flow cytometry. *In vivo* basophil depletion was achieved by diphtheria toxin treatment of Baso-DTR mice. For cell transfer studies, the basophil population was expanded *in vivo* by hydrodynamic tail vein injection of thymic stromal lymphopoietin cDNA plasmid.

Results—Sensitization to food allergens through an atopic dermatitis-like skin lesion is associated with an expansion of TSLP-elicited basophils in the skin that promote antigen-specific Th2 cytokine responses, elevated antigen-specific serum IgE levels and the accumulation of mast cells in the intestine promoting the development of intestinal food allergy. Critically, disruption of TSLP responses or depletion of basophils reduced the susceptibility to intestinal food allergy while transfer of TSLP-elicited basophils into intact skin promoted disease.

Conclusion—Epicutaneous sensitization on a disrupted skin barrier is associated with the accumulation of TSLP-elicited basophils that are necessary and sufficient to promote antigen-induced intestinal food allergy.

Keywords

Food allergy; atopic dermatitis; epicutaneous sensitization; TSLP; basophils; mast cells; IgE

INTRODUCTION

The prevalence of food allergies has increased in the past several decades with an estimated 5% of children and 3–4% of adults in industrialized countries living with the daily concern that exposure to certain foods may trigger a life-threatening allergic reaction.^{1,2} Food allergies are defined by an adverse immune response that occurs repeatedly following exposure to a given food and can manifest in symptoms ranging from itching, hives and diarrhea to acute anaphylaxis with a life-threatening drop in blood pressure and airway constriction.^{3,4,5} As a result, individuals must avoid certain foods to guard against accidental exposures.⁶ Currently, there is no treatment to prevent or cure food allergies, and available medications only treat symptoms after the allergic reaction occurs.^{4,7–9} As the public health and economic impact of food allergies continue to grow, there is an urgent need to develop new intervention strategies to prevent and treat this debilitating condition.

Cross-linking of allergen-specific IgE on mast cells has been shown to play a critical role in the pathogenesis of food allergy and therapeutic strategies to target the IgE-mast cell axis in patients have shown their potential for the prevention of anaphylactic responses.^{10–12} Although studies in animal models and patients have demonstrated a clear role for the IgE-mast cell axis in the pathogenesis of food allergies, little is known about the early immunological events that initiate these effector responses. Epidemiological studies have demonstrated that cutaneous inflammation associated with atopic dermatitis (AD) is a significant risk factor for the development of food allergies.^{8,13} The concept of skin barrier defects contributing to allergic sensitization is further supported by animal studies demonstrating that epicutaneous sensitization can predispose to antigen-induced allergic inflammation at other barrier sites.^{14–17} These data indicate that the skin may be a highly relevant site of food allergen sensitization.^{18–22} However, the immunological mechanisms

through which antigen sensitization in the skin can predispose to allergic inflammation in the intestine are incompletely understood.

Employing a new model of food allergy, we demonstrate that epicutaneous sensitization to food antigens on an AD-like skin lesion is associated with the infiltration of thymic stromal lymphopoietin (TSLP)-elicited basophils that promote the development of IgE-mediated intestinal food allergy in response to oral antigen exposure. In this model, TSLP-elicited basophils were necessary and sufficient for the development of antigen-induced intestinal food allergy indicating that the TSLP-basophil axis may offer a new therapeutic target in the treatment and prevention of food allergy.

RESULTS

Antigen sensitization in the context of AD-like skin inflammation promotes allergic intestinal inflammation

To study the potential mechanisms by which intestinal food allergy develops following epicutaneous sensitization, we established a mouse model of food allergy in which mice were sensitized with food antigens on a developing AD-like skin lesion induced by topical treatment with the vitamin D analog, MC903.^{24,25} WT BALB/c mice were epicutaneously sensitized to a model antigen, ovalbumin (OVA), on a developing AD-like skin lesion associated with elevated skin TSLP production (Fig 1. *A,B*). Following epicutaneous sensitization, mice were challenged intra-gastrically (i.g) with OVA resulting in the manifestation of allergic reactions and inflammation characterized by an elevated clinical allergy score (Fig 1. *C*), antigen-specific IgE responses (Fig. 1*D*), type-2 inflammation in the intestine (Fig 1. *E,F*), the accumulation of intestinal mast cells (Fig 1. *G*) and expression of mast cell-specific proteases (Fig 1. *H*) in the small intestinal lamina propria. Control vehicle treated mice or mice that were sensitized to OVA and challenged i.g. with a control antigen (bovine serum albumin, BSA) exhibited minimal or no signs of intestinal food allergy (Fig 1. *C–H*, Fig E1). A similar pattern of disease was observed in mice that were epicutaneously sensitized to peanut, another clinically relevant food allergen (Fig E2). Clinical symptoms including episodes of scratching, piloerection and reduced activity resolved within 60 min after each oral antigen exposure.

To test whether epicutaneous sensitization predisposing to intestinal food allergy may also occur through a mechanically disrupted skin barrier, mice were sensitized by means of repeated application of OVA to tape-stripped skin (Fig E3A). Similar to the MC903 based model, oral antigen exposure resulted in an increased allergy score (Fig E3B), elevated antigen-specific serum IgE (Fig E3C) and the accumulation of mast cells in the small intestinal lamina propria (Fig. E3D). Collectively, employing two different models of epicutaneous sensitization, we demonstrate that antigen sensitization through a compromised skin barrier can promote the development of antigen-induced intestinal food allergy.

Intestinal food allergy in this model was associated with systemic antigen-specific IgE responses (Fig 1. *D*) and the accumulation of IgE-bearing mast cells in the intestine (Fig 1. *G*), suggesting that the IgE-mast cell axis may play a critical role. To test the role of IgE in

this model, mice deficient in IgE (*Igh-7^{-/-}*) were epicutaneously sensitized on an AD-like skin lesion and subsequently challenged i.g. with OVA. While oral antigen challenge of *Igh-7^{+/+}* BALB/c mice resulted in an elevated clinical allergy score and antigen-specific serum IgE levels, *Igh-7^{-/-}* mice did not manifest allergic symptoms (Fig E4). Consistent with other models of food allergy and data from studies in patients, these findings demonstrate IgE as an important effector molecule in the pathogenesis of food allergy.^{26–28}

Antigen-induced food allergy is dependent on TSLP-TSLPR interactions

Skin barrier dysfunction is associated with elevated expression of TSLP in the skin and the development of type-2 cytokine-associated allergic inflammation.^{14,25,29–31} Further, gain-of-function polymorphisms in *TSLP* and elevated TSLP responses have been associated with increased susceptibility to multiple allergic diseases in humans and mice.^{25,32–37} Recent studies demonstrated that mice with AD-like skin lesions are more susceptible to the development of airway inflammation following aero-antigen challenge in a TSLP-dependent manner.³⁸ Taken together, these studies provoked the hypothesis that TSLP may be a key mechanism by which cutaneous sensitization to food antigens promotes the development of intestinal food allergy. To test the role of TSLP in this model of epicutaneous sensitization to intestinal food allergy, WT BALB/c (*Tslpr^{+/+}*) mice or mice deficient in the receptor for TSLP (*Tslpr^{-/-}*) were epicutaneously sensitized and subsequently challenged i.g. with OVA (Fig 2. A). Following oral antigen challenge, *Tslpr^{+/+}* mice exhibited an elevated clinical allergy score (Fig 2. B), enhanced antigen-specific serum IgE levels (Fig 2. C), increased intestinal inflammation associated with type-2 cytokine responses (Fig 2. D,E), accumulation of mast cells in the intestinal lamina propria (Fig 2. F) and elevated expression of mast cell-specific proteases (Fig 2. G). In striking contrast, *Tslpr^{-/-}* mice exhibited limited antigen-induced food allergy by all parameters examined (Fig 2. B–G). Collectively, these data indicate that TSLP-TSLPR interactions are necessary for the development of intestinal food allergy in this model.

TSLP is sufficient to promote antigen-induced intestinal food allergy

To test whether TSLP alone is sufficient to promote antigen-induced food allergy, WT BALB/c mice were treated intradermally (i.d.) with recombinant TSLP (rTSLP) in the presence of OVA or either PBS or OVA alone (Fig 3. A).¹⁴ Oral antigen challenge of mice sensitized to OVA in the skin in the presence of rTSLP exhibited an increased allergy score (Fig 3. B), elevated antigen-specific serum IgE levels (Fig 3. C), increased intestinal inflammation associated with type-2 cytokine responses (Fig 3. D,E) and the accumulation of mast cells in the intestine (Fig 3. F) that was associated with elevated expression levels of mast cell-specific proteases (Fig 3. G) compared to controls (Fig 3. B–G). Of note, i.d. injection of OVA alone resulted in episodes of scratching illustrated in an elevated allergy score, but no differences in all other parameters assessed. Collectively, these data demonstrate that TSLP-TSLPR interactions are necessary (Fig 2) and sufficient (Fig 3) to promote allergic responses in the intestine upon oral antigen exposure.

TSLP-elicited basophils promote antigen-specific Th2 cytokine responses in the skin

TSLP is known to act on multiple cellular targets including T cells, dendritic cells, mast cells and basophils^{33,39}. We recently demonstrated that AD-like skin lesions are associated with the infiltration of TSLP-dependent basophils that promote Th2 cytokine-associated inflammation in the skin.⁴⁰ We therefore hypothesized that TSLP-elicited basophils may promote cutaneous antigen-specific Th2 cytokine responses that contribute to the activation of IgE-mast cell responses and the subsequent development of intestinal food allergy.

Consistent with previous studies,^{21,40} AD-like skin lesions in this model were associated with the accumulation of TSLP-dependent basophils in the skin, but not the intestine (Fig E5, and data not shown). To test whether TSLP-elicited basophils influenced the development of antigen-specific Th2 cytokine responses and associated skin inflammation, we employed a genetic approach in which basophils are selectively depleted by diphtheria toxin due to the expression of the DT receptor (DTR) on their surface.⁴¹ Using this strategy, Baso-DTR^{pos} mice and Baso-DTR^{neg} littermate controls were epicutaneously sensitized with MC903 + OVA while simultaneously being treated with DT (Fig 4. A). DT treatment resulted in efficient depletion (~90 %) of basophils in Baso-DTR^{pos} mice (Fig 4. B) that was associated with a reduction in skin inflammation (Fig 4. C,D). To test whether basophil depletion affected Th2 cytokine responses, skin-draining lymph node cells were re-stimulated *ex vivo* with antigen. While Baso-DTR^{neg} mice treated with MC903 + OVA in the presence of DT exhibited elevated antigen-specific Th2 cytokine responses, this response was significantly reduced in mice depleted of basophils (Fig 4. E). Taken together, these data demonstrate that TSLP-elicited basophils in the context of an AD-like skin lesion promote skin inflammation and antigen-specific Th2-cytokine responses.

TSLP-elicited basophils regulate antigen-induced food allergy

Given that TSLP-dependent basophils promoted local antigen-specific Th2 cytokine responses (Fig 4. E), we sought to test the role of basophils in mediating intestinal food allergy. Using the same approach described above to deplete basophils, Baso-DTR^{pos} or Baso-DTR^{neg} mice were epicutaneously sensitized and orally challenged while simultaneously being treated with DT (Fig 5. A). Strikingly, while skin sensitized control Baso-DTR^{neg} mice exhibited intestinal food allergy in response to oral antigen exposure (Fig 5. B–G), depletion of basophils in Baso-DTR^{pos} mice limited antigen-induced food allergy by all parameters examined (Fig 5. B–G). Collectively, these findings demonstrate that the TSLP-basophil axis regulates antigen-specific Th2 cytokine responses, systemic antigen-specific IgE levels and the accumulation of intestinal mast cells promoting the development of intestinal food allergy.

To test whether basophils are sufficient to promote antigen-induced intestinal food allergy, gain-of-function studies employing adoptive transfer of basophils were performed. To expand the basophil population *in vivo*, BALB/c mice received hydrodynamic tail vein injections of cDNA plasmid encoding TSLP.⁴⁰ Three weeks after TSLP plasmid injections, basophils were sort-purified from the periphery and injected i.d. into the ears of naive BALB/c mice in the presence of OVA on various days prior to oral antigen exposure (Fig 6. A). While control mice treated with PBS or OVA alone did not manifest signs of intestinal

food allergy, mice adoptively transferred with TSLP-elicited basophils in the presence of OVA exhibited signs of antigen-induced intestinal food allergy by all parameters assessed (Fig 6. B–G). Collectively, these data demonstrate that TSLP-elicited basophils are both necessary and sufficient to promote IgE-mediated intestinal food allergy in this model.

DISCUSSION

AD in infancy is a risk factor for developing food allergies later in life.^{8,13} However, the immunological mechanisms through which antigen sensitization in the skin can predispose to allergic inflammation in the intestine are incompletely understood. Employing a new model of intestinal food allergy initiated by epicutaneous antigen sensitization, we demonstrate that TSLP-elicited basophils orchestrate the development of IgE-mediated intestinal food allergy. We show that in the context of an AD-like skin lesion, basophils infiltrate the skin in a TSLP-dependent manner and promote antigen-specific Th2 cytokine responses. Further, we demonstrate that TSLP-elicited basophils are both necessary and sufficient for the development of IgE-mediated intestinal food allergy. These findings support the hypothesis that antigen sensitization through a disrupted skin barrier is a risk factor for the development of food allergy and provide a cellular mechanism by which TSLP-elicited basophils promote allergic inflammation in the intestine.

Accurately modeling the progression of allergic disease in patients using murine model systems presents numerous challenges, particularly regarding the time scale of the events that lead from sensitization to an allergen at one mucosal site to allergic disease at multiple mucosal barrier surfaces. Despite these challenges, previously described murine model systems have provided significant insight into the mechanisms that mediate the pathogenesis of intestinal food allergy. Consistent with other food allergy models,^{16,42} antigen-induced allergic reactions in this model were mediated by the IgE-mast cell axis, as mice deficient in IgE did not manifest allergic symptoms in response to oral antigen exposure. While other models of epicutaneous sensitization reported anaphylactic reactions in response to oral allergen exposure¹⁶, no drop in temperature after oral antigen exposure was observed in the described model (data not shown). Further, epicutaneous sensitization in absence of antigen did not result in increased mucosal mast cell frequencies as compared to other studies.¹⁶ Differences in the schedules of sensitization of the various models may account for the observed discrepancies in models of epicutaneous sensitization. Importantly, epicutaneous antigen sensitization in the model described here resulted in the simultaneous development of intestinal food allergy and, as previously reported, eosinophilic esophagitis (EoE)-like disease, two commonly co-existing food-induced allergic disorders.²⁵ Eosinophil accumulation in this model was not only restricted to the esophagus, but was evident in the skin, the stomach, intestine and the periphery of mice with intestinal food allergy/EoE-like disease.²⁵ Taken together, the described model may provide a new research tool that can be used to test new hypotheses and potential treatment protocols in the context of food allergy/EoE.

While the effector mechanisms in the pathogenesis of food allergies are well known, the sites of sensitization to food allergens and the upstream immunological mechanisms that initiate and regulate subsequent effector responses in the intestine are poorly understood. We

previously demonstrated that TSLP promotes the population expansion of basophils and that gain-of function polymorphisms in the gene encoding for TSLP are associated with atopy and peripheral basophilia.^{25,40} In the present model system, AD-like skin lesions were associated with a significant infiltration of TSLP-elicited basophils that were necessary and sufficient to promote the development of antigen-induced intestinal food allergy. While topical treatment with MC903 results in continuous TSLP expression and a high clinical allergy score upon oral antigen exposure of previously sensitized mice, adoptive transfer of basophils into an environment with low TSLP levels may affect survival and activation status of transferred basophils resulting in a lower, but still significant clinical allergy score compared to controls. Importantly, similar to the model system used, increased expression of TSLP and infiltration of basophils into lesional skin has been reported previously in AD patients. (Kim et al., personal communication and⁴³) Together, these data indicate that TSLP-elicited basophil responses may be a key upstream mechanism that precedes the induction of effector antigen-specific IgE responses and the accumulation of mast cells in the intestine promoting allergic intestinal inflammation.

The mechanisms by which TSLP-elicited basophils promote efficient antigen sensitization and CD4⁺ Th2 cell polarization in this model remain to be elucidated. Previous studies have shown that in some settings major histocompatibility complex class II (MHC-II)-dependent interactions between basophils and CD4⁺ T cells can augment Th2-cytokine responses.⁴⁴⁻⁴⁶ However, it is likely that dendritic cells (DCs) and other antigen presenting cells contribute in antigen sampling, processing and presentation. Basophil-derived mediators may upregulate OX40L on DCs creating a Th2-permissive environment.^{39,44,46} Further studies are necessary to investigate the complex interplay between TSLP-elicited basophils, DCs and T cells in mediating local antigen sensitization and Th2 polarization.

Recent studies suggest that several cardinal features of food allergy including Th2 sensitization to allergens, IgE responses and the expansion of intestinal mast cells result from enhanced IL-4 signaling in humans and mice.^{42,47} The critical role of IL-4 in mediating intestinal-food allergy is further substantiated by the observation that IL-4 deficient mice are protected from antigen-induced food allergy in this model (data not shown). Importantly, TSLP-elicited basophils have been attributed a critical role in the induction and propagation of type-2 immune responses by providing an early innate source of IL-4.^{40,48} Our findings that TSLP-elicited basophils are recruited early in the context of AD-like inflammation and are critical for the development of subsequent intestinal food allergy provoke the hypothesis that basophil-derived IL-4 may be a key mechanism by which allergic inflammation at one epithelial barrier surface can promote the development of allergic inflammation at another barrier site. Given the short life-time of a mature basophil, it is likely that basophils accumulate in inflamed skin in response to elevated TSLP levels, and prior to degranulation, release IL-4 into the periphery that may account for optimal Th2 polarization, IgE class switching, the expansion of intestinal mast cells and subsequent intestinal allergic inflammation. In this context, adoptive transfer of IL-4 eGFP⁺ basophils into skin of WT recipients was not associated with an accumulation of IL-4 eGFP⁺ basophils in the periphery 24 h after transfer, indicating that transferred cells may not leave the skin (data not shown). However, further studies are necessary to interrogate fate of TSLP-elicited basophils recruited to the skin and the role of basophil-derived IL-4 that may

regulate the susceptibility to intestinal food allergy. Importantly, recent clinical trials using anti-IL4 antibodies (dupilumab) in patients with AD demonstrated promising results in the improvement of eczema severity⁴⁹. Given that AD patients are at a higher risk to suffer from food allergies, targeting IL-4 responses in patients with AD may be a previously unrecognized therapeutic pathway to limit the development of intestinal food allergy.

In conclusion, we establish an experimental model of food allergy that recapitulates many characteristics of human disease in that the skin may be a relevant source of allergen sensitization. AD is a known risk factor for food allergy and human AD skin lesions are associated with elevated TSLP expression and basophil infiltration. These responses, coupled with systemic allergen-specific IgE responses and the accumulation of IgE-bearing mast cells in the intestine likely act to fuel intestinal allergic inflammation. Employing this model we provide a previously unrecognized cellular mechanism by which the TSLP-basophil axis promotes early phase local antigen-specific Th2 cytokine responses, systemic antigen-specific serum IgE levels and the accumulation of IgE-bearing intestinal mast cells promoting the development of IgE-mediated intestinal food allergy. Targeting the TSLP-basophil axis in the context of AD may offer a novel therapeutic approach for early intervention and therapeutic strategies to prevent the development of intestinal allergic inflammation.

METHODS

Allergy scoring

Starting 20 min after oral antigen challenge, mice were individually observed for a period of 30 min and a clinical allergy score was quantified as follows: 0: no symptoms; 1: less than 4 episodes of scratching and rubbing around nose and head; 2: 4–10 episodes of scratching and rubbing around nose and head; 3: more than 10 episodes of scratching and rubbing around nose and head; 4: hunching and piloerection; 5: immobility (unresponsive to non-harmful, tactile disturbance); 6: death). Investigators were blinded to the treatment group conditions when scoring for allergy symptoms.

Flow cytometry

Small intestinal lamina propria cells or cells from the skin were isolated as described previously.^{27,28} Single cell suspensions were incubated with Aqua Live/Dead Fixable Dye (Invitrogen) for dead cell exclusion and stained with anti-mouse fluorochrome-conjugated monoclonal antibodies against B220, CD3e, CD4, CD5, CD8, CD11c, NK1.1, CD19, IgE, CD45, CD49b or CD117 (c-kit) (BD Bioscience, BioLegend, eBioscience). Mast cells were identified as live, lin⁻ (CD3,CD5,CD19,CD11c,NK1.1), CD45⁺, c-kit⁺, IgE⁺ cells and basophils as live, lin⁻, CD45⁺, c-kit⁻, CD49b⁺, IgE⁺ cells.

Adoptive basophil transfer

WT BALB/c mice were injected with a cDNA plasmid encoding TSLP as previously described.²⁹ Three weeks after TSLP cDNA injection, basophils were sort purified, and 5–10×10⁴ cells were transferred i.d (ears) in the presence of 100 µg OVA on various days. As control, mice were injected with PBS or 100 µg OVA alone.

ELISA, histology and real-time PCR

To measure skin TSLP levels, ears were incubated overnight in culture media, and cell-free supernatants were stored. Measurement of TSLP was performed using a commercially available DuoSet ELISA kit (eBioscience). Antigen-specific serum IgE responses were measured as described before.³⁰ For histological analysis, tissues were fixed in 4% paraformaldehyde and embedded in paraffin, and sections were stained with hematoxylin and eosin (H&E). For real-time PCR analysis, RNA was isolated using an RNeasy mini kit (QIAGEN) according to the manufacturer's instructions. cDNA was generated using a SuperscriptII reverse transcription kit (Invitrogen). Real-time quantitative PCR was performed on cDNA using SYBR green master mix (Applied Biosystems) and commercially available primer sets from Qiagen (Quantitect primer assays). Samples were normalized to β -actin and displayed as fold induction over controls.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Research in the Artis lab is supported by the National Institutes of Health (AI061570, AI087990, AI074878, AI106679, AI095466, AI095608, AI102942 and AI097333 to D.A.; the Crohn's and Colitis Foundation of America (D.A.); the Burroughs Wellcome Fund Investigator in Pathogenesis of Infectious Disease Award (D.A.); the Swiss National Science Foundation Prospective and Advanced Research Fellowships (PBBEP3_130438 and PA00P3_136468 to M.N.); T32-AR007465 and KL2-RR024132 to B.S.K., and F32-AI085828 to M.C.S. The ACC Flow Cytometry and Cell Sorting Shared Resource is partially supported by NCI Comprehensive Cancer Center Support Grant (2-P30-CA016520). This work was supported by the NIH/NIDDK P30 Center for Molecular Studies in Digestive and Liver Diseases (P30-DK050306), its pilot grant program and scientific core facilities (Molecular Pathology and Imaging, Molecular Biology, Cell Culture and Mouse).

We thank members of the Artis laboratory for discussions and critical reading of the manuscript. We thank the Matthew J. Ryan Veterinary Hospital Pathology Lab and the Abramson Cancer Center Flow Cytometry and Cell Sorting Resource Laboratory for technical advice and support.

Abbreviations used

AD	atopic dermatitis
CPE	crude peanut extract
DT	diphtheria toxin
IgE	immunoglobulin E
IL	interleukin
i.d	intradermal
i.g	intragastrically
i.p.	intraperitoneally
lin⁻	lineage negative
Mcpt	mouse mast cell protease
OVA	ovalbumin

Th2	T-helper type 2
TSLP	thymic stromal lymphopoietin
WT	wild-type

References

1. Sicherer SH, Sampson HA. Food allergy. *The Journal of allergy and clinical immunology*. 2010; 125(2 Suppl 2):S116–25. [PubMed: 20042231]
2. Burks AW, Tang M, Sicherer S, Muraro A, Eigenmann PA, Ebisawa M, et al. ICON: food allergy. *The Journal of allergy and clinical immunology*. 2012; 129(4):906–20. [PubMed: 22365653]
3. Fiocchi A, Schunemann HJ, Brozek J, Restani P, Beyer K, Troncone R, et al. Diagnosis and Rationale for Action Against Cow's Milk Allergy (DRACMA): a summary report. *The Journal of allergy and clinical immunology*. 2010; 126(6):1119–28. e12. [PubMed: 21134569]
4. Burks AW, Jones SM, Boyce JA, Sicherer SH, Wood RA, Assa'ad A, et al. NIAID-sponsored 2010 guidelines for managing food allergy: applications in the pediatric population. *Pediatrics*. 2011; 128(5):955–65. [PubMed: 21987705]
5. Berin MC, Sampson HA. Food allergy: an enigmatic epidemic. *Trends in immunology*. 2013
6. Lieberman JA, Sicherer SH. Quality of life in food allergy. *Current opinion in allergy and clinical immunology*. 2011; 11(3):236–42. [PubMed: 21464708]
7. Berin MC, Sicherer S. Food allergy: mechanisms and therapeutics. *Current opinion in immunology*. 2011; 23(6):794–800. [PubMed: 21943957]
8. Boyce JA, Assa'ad A, Burks AW, Jones SM, Sampson HA, Wood RA, et al. Guidelines for the diagnosis and management of food allergy in the United States: summary of the NIAID-sponsored expert panel report. *Nutr Res*. 2011; 31(1):61–75. [PubMed: 21310308]
9. Wang J, Sampson HA. Treatments for food allergy: how close are we? *Immunologic research*. 2012; 54(1–3):83–94. [PubMed: 22434517]
10. Brandt EB, Strait RT, Hershko D, Wang Q, Muntel EE, Scribner TA, et al. Mast cells are required for experimental oral allergen-induced diarrhea. *The Journal of clinical investigation*. 2003; 112(11):1666–77. [PubMed: 14660743]
11. Sampson HA, Leung DY, Burks AW, Lack G, Bahna SL, Jones SM, et al. A phase II, randomized, doubleblind, parallelgroup, placebocontrolled oral food challenge trial of Xolair (omalizumab) in peanut allergy. *The Journal of allergy and clinical immunology*. 2011; 127(5):1309–10. e1. [PubMed: 21397314]
12. Khoriaty E, Umetsu DT. Oral immunotherapy for food allergy: towards a new horizon. *Allergy, asthma & immunology research*. 2013; 5(1):3–15.
13. Rance F. Food allergy in children suffering from atopic eczema. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology*. 2008; 19(3):279–84. quiz 85. [PubMed: 18397414]
14. Han H, Xu W, Headley MB, Jessup HK, Lee KS, Omori M, et al. Thymic stromal lymphopoietin (TSLP)-mediated dermal inflammation aggravates experimental asthma. *Mucosal immunology*. 2012; 5(3):342–51. [PubMed: 22354320]
15. Jiang H, Hener P, Li J, Li M. Skin thymic stromal lymphopoietin promotes airway sensitization to inhalant house dust mites leading to allergic asthma in mice. *Allergy*. 2012; 67(8):1078–82. [PubMed: 22687045]
16. Bartnikas LM, Gurish MF, Burton OT, Leisten S, Janssen E, Oettgen HC, et al. Epicutaneous sensitization results in IgE-dependent intestinal mast cell expansion and food-induced anaphylaxis. *The Journal of allergy and clinical immunology*. 2013; 131(2):451–60. e1–6. [PubMed: 23374269]
17. Vaali K, Puumalainen TJ, Lehto M, Wolff H, Rita H, Alenius H, et al. Murine model of food allergy after epicutaneous sensitization: role of mucosal mast cell protease-1. *Scandinavian journal of gastroenterology*. 2006; 41(12):1405–13. [PubMed: 17101571]

18. Bieber T. Atopic dermatitis. *The New England journal of medicine*. 2008; 358(14):1483–94. [PubMed: 18385500]
19. De Benedetto A, Kubo A, Beck LA. Skin barrier disruption: a requirement for allergen sensitization? *The Journal of investigative dermatology*. 2012; 132(3 Pt 2):949–63. [PubMed: 22217737]
20. Kubo A, Nagao K, Amagai M. Epidermal barrier dysfunction and cutaneous sensitization in atopic diseases. *The Journal of clinical investigation*. 2012; 122(2):440–7. [PubMed: 22293182]
21. Leyva-Castillo JM, Hener P, Jiang H, Li M. TSLP produced by keratinocytes promotes allergen sensitization through skin and thereby triggers atopic march in mice. *The Journal of investigative dermatology*. 2013; 133(1):154–63. [PubMed: 22832486]
22. Spergel JM, Mizoguchi E, Brewer JP, Martin TR, Bhan AK, Geha RS. Epicutaneous sensitization with protein antigen induces localized allergic dermatitis and hyperresponsiveness to methacholine after single exposure to aerosolized antigen in mice. *The Journal of clinical investigation*. 1998; 101(8):1614–22. [PubMed: 9541491]
23. Zhou B, Comeau MR, De Smedt T, Liggitt HD, Dahl ME, Lewis DB, et al. Thymic stromal lymphopoietin as a key initiator of allergic airway inflammation in mice. *Nature immunology*. 2005; 6(10):1047–53. [PubMed: 16142237]
24. Li M, Hener P, Zhang Z, Kato S, Metzger D, Chambon P. Topical vitamin D3 and low- calcemic analogs induce thymic stromal lymphopoietin in mouse keratinocytes and trigger an atopic dermatitis. *Proceedings of the National Academy of Sciences of the United States of America*. 2006; 103(31):11736–41. [PubMed: 16880407]
25. Noti M, Wojno ED, Kim BS, Siracusa MC, Giacomini PR, Nair MG, et al. Thymic stromal lymphopoietin-elicited basophil responses promote eosinophilic esophagitis. *Nature medicine*. 2013; 19(8):1005–13.
26. Helm RM, Burks AW. Animal models of food allergy. *Current opinion in allergy and clinical immunology*. 2002; 2(6):541–6. [PubMed: 14752339]
27. Finkelman FD. Anaphylaxis: lessons from mouse models. *The Journal of allergy and clinical immunology*. 2007; 120(3):506–15. quiz 16–7. [PubMed: 17765751]
28. Lemon-Mule H, Nowak-Wegrzyn A, Berin C, Knight AK. Pathophysiology of food-induced anaphylaxis. *Current allergy and asthma reports*. 2008; 8(3):201–8. [PubMed: 18589838]
29. Comeau MR, Ziegler SF. The influence of TSLP on the allergic response. *Mucosal immunology*. 2010; 3(2):138–47. [PubMed: 20016474]
30. Ziegler SF. The role of thymic stromal lymphopoietin (TSLP) in allergic disorders. *Current opinion in immunology*. 2010; 22(6):795–9. [PubMed: 21109412]
31. Ziegler SF. Thymic stromal lymphopoietin and allergic disease. *The Journal of allergy and clinical immunology*. 2012; 130(4):845–52. [PubMed: 22939755]
32. Rothenberg ME, Spergel JM, Sherrill JD, Annaiah K, Martin LJ, Cianferoni A, et al. Common variants at 5q22 associate with pediatric eosinophilic esophagitis. *Nature genetics*. 2010; 42(4):289–91. [PubMed: 20208534]
33. Ziegler SF, Artis D. Sensing the outside world: TSLP regulates barrier immunity. *Nature immunology*. 2010; 11(4):289–93. [PubMed: 20300138]
34. Sherrill JD, Gao PS, Stucke EM, Blanchard C, Collins MH, Putnam PE, et al. Variants of thymic stromal lymphopoietin and its receptor associate with eosinophilic esophagitis. *The Journal of allergy and clinical immunology*. 2010; 126(1):160–5. e3. [PubMed: 20620568]
35. Harada M, Hirota T, Jodo AI, Doi S, Kameda M, Fujita K, et al. Functional analysis of the thymic stromal lymphopoietin variants in human bronchial epithelial cells. *American journal of respiratory cell and molecular biology*. 2009; 40(3):368–74. [PubMed: 18787178]
36. Blazquez AB, Mayer L, Berin MC. Thymic stromal lymphopoietin is required for gastrointestinal allergy but not oral tolerance. *Gastroenterology*. 2010; 139(4):1301–9. [PubMed: 20600023]
37. Al-Shami A, Spolski R, Kelly J, Keane-Myers A, Leonard WJ. A role for TSLP in the development of inflammation in an asthma model. *The Journal of experimental medicine*. 2005; 202(6):829–39. [PubMed: 16172260]
38. Zhang Z, Hener P, Frossard N, Kato S, Metzger D, Li M, et al. Thymic stromal lymphopoietin overproduced by keratinocytes in mouse skin aggravates experimental asthma. *Proceedings of the*

- National Academy of Sciences of the United States of America. 2009; 106(5):1536–41. [PubMed: 19188585]
39. Siracusa MC, Wojno ED, Artis D. Functional heterogeneity in the basophil cell lineage. *Advances in immunology*. 2012; 115:141–59. [PubMed: 22608258]
 40. Siracusa MC, Saenz SA, Hill DA, Kim BS, Headley MB, Doering TA, et al. TSLP promotes interleukin-3-independent basophil haematopoiesis and type 2 inflammation. *Nature*. 2011; 477(7363):229–33. [PubMed: 21841801]
 41. Sawaguchi M, Tanaka S, Nakatani Y, Harada Y, Mukai K, Matsunaga Y, et al. Role of mast cells and basophils in IgE responses and in allergic airway hyperresponsiveness. *J Immunol*. 2012; 188(4):1809–18. [PubMed: 22250079]
 42. Mathias CB, Hobson SA, Garcia-Lloret M, Lawson G, Poddighe D, Freyschmidt EJ, et al. IgE-mediated systemic anaphylaxis and impaired tolerance to food antigens in mice with enhanced IL-4 receptor signaling. *The Journal of allergy and clinical immunology*. 2011; 127(3):795–805. e1–6. [PubMed: 21167580]
 43. Ito Y, Satoh T, Takayama K, Miyagishi C, Walls AF, Yokozeki H. Basophil recruitment and activation in inflammatory skin diseases. *Allergy*. 2011; 66(8):1107–13. [PubMed: 21371044]
 44. Yoshimoto T, Yasuda K, Tanaka H, Nakahira M, Imai Y, Fujimori Y, et al. Basophils contribute to T(H)2-IgE responses in vivo via IL-4 production and presentation of peptide- MHC class II complexes to CD4+ T cells. *Nature immunology*. 2009; 10(7):706–12. [PubMed: 19465908]
 45. Sokol CL, Chu NQ, Yu S, Nish SA, Laufer TM, Medzhitov R. Basophils function as antigen-presenting cells for an allergen-induced T helper type 2 response. *Nature immunology*. 2009; 10(7):713–20. [PubMed: 19465907]
 46. Perrigoue JG, Saenz SA, Siracusa MC, Allenspach EJ, Taylor BC, Giacomini PR, et al. MHC class II-dependent basophil-CD4+ T cell interactions promote T(H)2 cytokine-dependent immunity. *Nature immunology*. 2009; 10(7):697–705. [PubMed: 19465906]
 47. Burton OT, Darling AR, Zhou JS, Noval-Rivas M, Jones TG, Gurish MF, et al. Direct effects of IL-4 on mast cells drive their intestinal expansion and increase susceptibility to anaphylaxis in a murine model of food allergy. *Mucosal immunology*. 2012
 48. Giacomini PR, Siracusa MC, Walsh KP, Grencis RK, Kubo M, Comeau MR, et al. Thymic stromal lymphopoietin-dependent basophils promote Th2 cytokine responses following intestinal helminth infection. *J Immunol*. 2012; 189(9):4371–8. [PubMed: 23024277]
 49. Simpson, E. Treatment of moderate-to-severe atopic dermatitis with dupilumab. *American Academy of Dermatology*; 2013. Meeting abstract

Clinical implications

Targeting the TSLP-basophil axis may offer a novel therapeutic approach for early intervention and therapeutic strategies to prevent the development of food allergy.

FIG. 1.

Epicutaneous sensitization promotes antigen-induced food allergy. **A**, Experimental protocol. **B**, TSLP protein in the skin. **C**, Allergy score. **D**, OVA-specific serum IgE levels. **E**, H&E staining from the jejunum. Scale bar: 100 μ m. Arrows: tissue-infiltrating granulocytes. **F**, Type-2 cytokine expression in the jejunum. **G**, Frequencies of mast cells in the lamina propria. **H**, Jejunal *Mcpt1* and *Mcpt2* expression. $N = 3-4$ per group. Data are representative of three independent experiments. ND: no disease.

FIG. 2.

TSLP-TSLPR responses mediate antigen-induced food allergy. **A**, Experimental protocol. **B**, Allergy score. **C**, OVA-specific serum IgE levels. **D**, H&E staining from the jejunum. Scale bar: 100 μ m. Arrows: tissue-infiltrating granulocytes. **E**, Type-2 cytokine expression in the jejunum. **F**, Frequencies of mast cells in the lamina propria. **G**, Jejunal *Mcpt1* and *Mcpt2* expression. $N = 3-4$ per group. Data are representative of three independent experiments. ns: not significant.

FIG. 3.

TSLP is sufficient to promote antigen-induced intestinal food allergy. **A**, Experimental protocol. **B**, Allergy score. **C**, OVA-specific serum IgE levels. **D**, H&E staining from the jejunum. Scale bar: 100 μ m. Arrows: tissue-infiltrating granulocytes. **E**, Type-2 cytokine expression in the jejunum. **F**, Frequencies of mast cells in the lamina propria. **G**, Jejunal *Mcpt1* and *Mcpt2* expression. *N* = 3 mice per group. Data are representative of three independent experiments.

FIG. 4.

AD-like skin lesions are associated with the accumulation of TSLP-elicited basophils in the skin. **A**, Experimental protocol. **B**, Frequencies of basophils in the skin. **C**, H&E staining from the skin. Scale bar: 100 μ m. **D**, Ear thickness measurement. **E**, Th2 cytokine responses in antigen re-stimulated skin-draining lymph nodes. $N = 3-4$ per group. Data are representative of three independent experiments. nd: not detected.

FIG. 5.

TSLP-elicited basophil responses regulate antigen-induced food allergy. **A**, Experimental protocol. **B**, Allergy score. **C**, OVA-specific serum IgE levels. **D**, H&E staining from the jejunum. Scale bar: 100 μ m. Arrows: tissue-infiltrating granulocytes. **E**, Type-2 cytokine expression in the jejunum. **F**, Frequencies of mast cells in the lamina propria. **G**, Jejunal *Mcpt1* and *Mcpt2* expression. $N = 3-5$ per group. Data are representative of three independent experiments.

FIG. 6.

TSLP-elicited basophils are sufficient to promote intestinal food allergy. **A**, Experimental protocol. **B**, Allergy score. **C**, OVA-specific serum IgE levels. **D**, H&E staining from the jejunum. Scale bar: 100 μ m. Arrows: tissue-infiltrating granulocytes. **E**, Type-2 cytokine expression in the jejunum. **F**, Frequencies of mast cells in the lamina propria. **G**, Jejunal *Mcpt1* and *Mcpt2* expression. $N = 3$ per group. Data are representative of three independent experiments.