Published in final edited form as:

Bull Entomol Res. 2014 June ; 104(3): 323–333. doi:10.1017/S0007485314000091.

Increased production of mitochondrial reactive oxygen species and reduced adult life span in an insecticide-resistant strain of *Anopheles gambiae*

Dennis Otali^{a,*}, Robert J. Novak^{b,c}, Wen Wan^d, Su Bu^e, Douglas R. Moellering^e, and Maria De Luca^e

^aDepartment of Biology, University of Alabama at Birmingham, 1720 2nd Ave. South, Birmingham, AL 35294-1170, USA

^bWilliam C Gorgas Center for Geographic Medicine, Division of Infectious Diseases, Department of Medicine, University of Alabama at Birmingham, 845 19th St. South, Birmingham, AL 35294-2170, USA

^cDepartment of Global Health, University of South Florida, 13201 Bruce B. Downs Blvd., Tampa, FL 33612, USA

^dDepartment of Biostatistics, Virginia Commonwealth University Medical Center, P.O. Box 980032, Richmond, VA 23298-0032, USA

^eDepartment of Nutrition Sciences, University of Alabama at Birmingham, 1720 2nd Ave. South, Birmingham, AL 35294-3360, USA

Abstract

Control of the malaria vector *An. gambiae* is still largely obtained through chemical intervention using pyrethroids, such as permethrin. However, strains of *An. gambiae* that are resistant to the toxic effects of pyrethroids have become widespread in several endemic areas over the last decade. The objective of this study was to assess differences in five life-history traits (larval developmental time and the body weight, fecundity, hatch rate, and longevity of adult females) and energy metabolism between a strain of *An. gambiae* that is resistant to permethrin (RSP), due to knockdown resistance and enhanced metabolic detoxification, and a permethrin susceptible strain reared under laboratory conditions. We also quantified the expression levels of five antioxidant enzyme genes: *GSTe3*, *CAT*, *GPXH1*, *SOD1*, and *SOD2*. We found that the RSP strain had a longer developmental time than the susceptible strain. Additionally, RSP adult females had higher wet body weight and increased water and glycogen levels. Compared to permethrin susceptible females, RSP females displayed reduced metabolic rate and mitochondrial coupling efficiency and higher mitochondrial ROS production. Furthermore, despite higher levels of *GSTe3* and *CAT* transcripts, RSP females had a shorter adult life span than susceptible females. Collectively, these results suggest that permethrin resistance alleles might affect energy metabolism, oxidative stress, and adult survival of *An. gambiae*. However, because the strains used in this study differ in their genetic backgrounds, the results need to be interpreted with

*Corresponding Author: Dennis Otali, Department of Biology, University of Alabama at Birmingham, Campbell Hall 464, 1720 2nd Ave S, Birmingham AL 35294-1170, Phone: (+1) 205-975-6205, Fax: (+1) 205-975-7128, otalid@uab.edu.

caution and replicated in other strains in order to have significant implications for malaria transmission and vector control.

Keywords

Anopheles gambiae; permethrin; ROS; mitochondria; insecticide resistance

Background

Permethrin is an insecticide that belongs to the pyrethroid family of neurotoxic agents designed to kill insects by altering the permeability of the voltage-gated sodium channel (VGSC). Pyrethroids are highly preferred to other insecticides in vector control management practices due to their low toxicity to humans, rapidity of action, and easy and fast degradability in the environment (Ray, 2001). As such, permethrin-based insecticide treated nets and indoor residual spraying have been extensively used to control *An. gambiae* populations in malaria endemic areas and have been effective in reducing the transmission of the parasites from infective mosquito females to humans (Enayati & Hemingway, 2006; WHO, 2007). However, over the last decade, strains of *An. gambiae* that are resistant to the toxic effects of permethrin and other pyrethroids have become widespread in several endemic areas in Africa (Casimiro et al., 2006; Adasi & Hemingway, 2008). The most established source of permethrin resistance is represented by point-mutations within the gene encoding the VGSC that have been associated with knockdown resistance to different insecticides (Ranson et al., 2000). But, enhanced metabolic detoxification mechanisms through up-regulation of cytochrome P450 monooxygenase (P450), glutathione S-transferase (GST), and non-specific esterase genes have also been reported in *Anopheles* mosquitoes (Vulule et al., 1999; Hemingway & Ranson, 2000; Djouaka et al., 2008; Lumjuan et al., 2011). While significant progress has been made in understanding the molecular mechanisms underlying the resistance to permethrin in malaria vectors (Li et al., 2007; Soderlund, 2008), there have been few reports of how alleles that confer resistance to insecticides affect other fitness characteristics of the malarial vectors in insecticide free environments.

Life history theory applied to this problem suggests that we should expect alleles influencing resistance to insecticide to come at the cost of other fitness traits, especially in an insecticide free environment. The life-history theory is based on the idea that physiological traits such as reproduction, storage, somatic maintenance, growth, and development are energetically costly traits. Because resources are limited, differential allocation of energy among these competing demands can produce trade-offs among traits and natural selection is thought to have shaped the way organisms partition their limiting resources to these fitness components, balancing the costs and benefits (Wiley, 1974). Fitness costs associated with resistance mechanisms have been reported in insecticide-free environments reviewed in (Brooke & Koekemoer, 2010; Kliot & Ghanim, 2012). For example, work in *An. gambiae* and *An. stephensi* showed that strains resistant to dieldrin had reduced fecundity compared to susceptible individuals despite similar longevity (Rowland, 1991). Reduced fecundity and shorter reproductive period and lifespan were also observed in carbofuran resistant aphids

(Roberto & Omoto, 2006). However, loss of fitness has not been observed in other studies (Okoye et al., 2007), suggesting that a cost of resistance may not always occur (Coustau et al., 2000; Rigby et al., 2002). This is particularly true if different molecular mechanisms of resistance exist and/or ecological factors are involved (Coustau et al., 2000). Because evolutionary fitness costs are the cornerstones of economic optimal models of malaria vector insecticide resistance (Brown et al., 2013); more research in this area is necessary. Such research could elucidate whether new resistance management strategies for insecticide use are needed to maintain or restore its efficacy (Read et al., 2009). The employment of mechanisms of detoxification in insects can be energetically costly (Coustau et al., 2000). Thus, it is conceivable that increased metabolic detoxification in pyrethroid resistance would result in a resource allocation trade-off between the detoxification mechanisms and other energetically demanding physiological functions, such as growth or reproduction (Rivero et al., 2010). In the present study, we compared life-history traits and energy metabolism between a wild-derived permethrin resistant strain of *An. gambiae*, and a permethrin susceptible strain reared under laboratory conditions. The permethrin resistant strain used in our study is characterised for a *knockdown resistance* (*kdr*) mutation and enhanced levels of P450 and esterase enzyme activities (Vulule et al., 1999) (<http://www.mr4.org/>). We focused on this resistant strain of *An. gambiae* because growing evidence suggests that pyrethroid resistance in the wild is likely due to a combination of target-site insensitivity and metabolic-based mechanisms (Brooke & Koekemoer, 2010).

Materials and Methods

Strains and colony maintenance

A strain of *An. gambiae* with reduced susceptibility to permethrin (RSP) and a permethrin susceptible (ASEMBO1) strain were used. The strains were obtained from the Malaria Research and Reference Reagent Resource Center (MR4) (<http://www.mr4.org/>).

RSP has the East African form of the *kdr* sodium channel mutation allele, L1014S, and increased P450 and beta-esterase activities (Vulule et al., 1999). RSP was originally isolated from a permethrin impregnated bed net study carried out in four adjacent villages in North West of Kisumu, Kenya (Vulule et al., 1999). The strain was established from a colony representative of *An. gambiae* selected for permethrin tolerance (Vulule et al., 1999) and subsequently deposited in the MR4 collection, where it has been selected for permethrin resistance for over ten years by treating a cohort of fourth-instar larvae with 1ppm permethrin for 24 hours every three generations (<http://www.mr4.org/>).

The permethrin susceptible strain ASEMBO1 was originally collected 50 km west of Kisumu in a control area for the United States Agency for International Development cohort bed net project approximately 4.5 km from the study area (<http://www.mr4.org/>).

The stocks were obtained twice from the MRC4 (2009 and 2012) and the bioenergetics experiments described below were performed within five months of arrival.

Experimental conditions

Mosquitoes were maintained in insectary rooms at $25^{\circ}\text{C} \pm 2^{\circ}\text{C}$, $80\% \pm 10\%$ relative humidity, and 16 hr light/8 hr dark cycle. Larvae were provided abundant food consisting of ground Tetramin (fish food). Adult mosquitoes were fed on 10% honey solution soaked in cotton sticks. Unless otherwise stated, experimental groups of each strain were three to five day old adults reared in 30 cm^3 cages. For egg production, young adults of the respective strains were allowed to freely mate and feed on blood from ears of a restrained rabbit. Moist filter paper was placed in cages for oviposition and the eggs collected were incubated overnight in a Petri dish and were hatched in deionised (DI) water. The larvae were reared in plastic pans ($29.5 \times 23.5 \times 15\text{ cm}$) containing approximately 1000 mL of water in environmental chambers (Thermo Scientific, Dubuque, Iowa, USA). Each pan contained groups of 50 larvae which fed *ad libitum* on Tetramin. Emerging adults were reared in the environmental chamber and fed *ad libitum* on 10% honey solution.

Developmental time

Groups of 16 first-instar (L1) larvae of each strain were randomly selected and transferred into 12 emergence containers (11 cm in diameter) each holding 100 mL DI water and approximately 4.8 mg of Tetramin added daily and maintained in environmental chambers. Larvae were counted daily until they reached the pupa stage when they were transferred into individual test tubes containing 3 ml of DI water.

Body weight and glycogen levels

To determine wet weight, female mosquitoes were anaesthetised with CO_2 , transferred into vials in groups of five, and weighed to 0.1 mg accuracy with an analytical balance (OHAUS corp. Pine Brook, NJ). Mosquitoes were then dried in a bath incubator (Fischer Scientific) at 60°C for 1 hour and the dry weight was measured as a proxy for their size (Siegel et al., 1994). Water content was calculated by subtracting the dry weight from the wet weight.

Glycogen content was assessed in three-five day old females fed on 10% honey using the protocol described in (Jumbo-Lucioni et al., 2010). Briefly, for each strain, 10 independent replicates, each containing a group of 10 mosquitoes were assayed. Groups of mosquitoes were homogenised on ice using 40 μL of homogenization buffer (0.01 μM KH_2PO_4 and 1 mM EDTA pH 7.4). The homogenates were centrifuged in a microcentrifuge at 2000 rpm for 2 minutes at 4°C . Aliquots of 1.67 μL of homogenate were added to 250 μL of a reagent containing 0.1 U/ml of amyloglucosidase. After 30-minute incubation period at 37°C , OD_{540} was measured. The concentration of glycogen was determined from a glycogen standard run with each replicate. An independent set of 10 independent replicates was assayed for triacylglycerol content spectrophotometrically using a commercially available kit (Sigma-Triglyceride kit) following manufactures protocol. Each sample was assayed twice and the mean was used in the analyses.

Fecundity

After blood feeding, 40 mosquitoes of each strain were randomly aspirated and transferred into individual cages with moist filter paper for egg laying. The cages were provided with

10% honey solution and egg production monitored daily. The eggs produced were counted and incubated overnight in Petri dishes to determine hatch rate after which the filter paper was replaced. Eggs were hatched in rearing pans in 1000 mL of DI water. Care was taken to ensure all eggs stuck on the sides of the rearing pan were in contact with DI water. Hatched eggs were counted on the second day.

Adult life span

One hundred and twenty pupae of each strain were randomly collected in pupa cups and transferred to three cages (40 per cage) in an environmental chamber. Pupa cups were removed from the cages after 24 hours so that pupae that failed to emerge were excluded. The emerged adults were fed on 10% honey solution *ad libitum*. The cages were monitored daily and dead individuals were counted and removed until all individuals had died.

Metabolic rate measurement

Metabolic rate was determined as described in (De Luca et al., 2010). Briefly, metabolic rate as CO₂ production was measured using a flow-through respirometry system (Qubit System Research, Kingston, Ontario, Canada). Groups of five females of each strain were anaesthetised with CO₂ and gently transferred to the respirometry chamber. CO₂ was then measured for 10 min/chamber with a 30 second flush period between measurements at a flow rate of 30ml/min. The amount of CO₂ produced by each group of mosquitoes was calculated using C950 Data Acquisition software (Qubit System Research, Kingston, Ontario, Canada).

Mitochondrial respiration rate assay

Mosquitoes were anaesthetised with CO₂ and thoraces dissected from 30 females per replicate. All mitochondrial isolation steps in the six replicates were performed on ice. Mosquitoes were chilled briefly on ice and thoraces were separated from the heads and abdomens. Dissected thoraces were placed into 200µl of ice-cold isolation buffer [250 mM sucrose, 5 mM Tris-HCl, 2 mM EDTA, 1% (w/v) bovine serum albumin (BSA), pH 7.4 at 4°C; (Miwa et al., 2003) supplemented with protease inhibitors (leupeptin 1mg/ml, aprotinin 1mg/ml and pepstatin 1mg/ml) in a 1.5 ml Eppendorf tube. The samples were pounded gently 126 times over a 2 minute period, using a custom-built, motorised micromortar. Mashed mosquitoes were filtered through a 5 micron nylon mesh, and the volume was raised to 400µl by washing the nylon membrane with additional isolation buffer. A cycle of low-speed centrifugation (1 min centrifugation at 1000g) was followed by centrifugation of the filtered solution for 10 min at 3000 g at 4°C, and the pellet re-suspended in 100µl of isolation buffer. Protein concentrations in the mitochondrial fractions were determined using a Lowry assay.

Using freshly isolated mitochondria, mitochondrial respiration assays were performed using a polarographic oxygen sensor (Oroboros oxygraph, OROBOROS® INSTRUMENTS, Innsbruck, Austria) with 0.2 mg/ml of freshly isolated mitochondria incubated in respiration medium (120mM KCl, 5mM KH₂PO₄, 3mM Hepes, 1mM EDTA, 1mM MgCl₂, and 0.2% BSA, pH 7.2; (Ferguson et al., 2005). Oxygen consumption rates were measured at 25°C (Sacktor & Sanborn, 1956). As implemented by (Miwa et al., 2003), we measured state 3

and state 4 respiration rates using the NAD^+ -linked substrates pyruvate 5mM/proline 5mM to deliver electrons into mitochondrial complex I predominately. NAD^+ -linked substrates were added to the chamber and allowed to equilibrate for 1 min, followed by the addition of ADP at a concentration of 400 μM to elicit ADP-dependent state 3. This was followed by the determination of the state 4 respiration rate, once all the added ADP had been exhausted and a steady state is reached (Affourtit et al., 2012). Mitochondrial coupling efficiency (P: O ratio), e.g. the relationship between ATP synthesis and oxygen consumption, was calculated as the amount of ADP consumed per oxygen being reduced during state 3 (Jumbo-Lucioni et al., 2012). All assays were performed within three hours of mitochondrial isolation. Data were analyzed using the software *DatLab* Version 4.1.0.8.

Detection of ROS

Mitochondria for ROS analysis were stored in ice cold conditions and used within 5 hrs after isolation. ROS levels were measured using 10-acetyl -3, 7 dihydroxyphenoazine (Amplex Red, AR; Molecular Probes, Eugene, OR), to detect hydrogen peroxide (H_2O_2) in the presence of horseradish peroxidase, producing the red-fluorescent oxidation product (excitation/emission = 571/585 nm), resorufin. The ROS production of isolated mitochondria was measured as a basal rate (state 2 respiration) initiated by the reaction of saturated levels of substrates (pyruvate and proline), which evaluates ROS produced in a leak-like state without ADP and no oxidative phosphorylation. ROS production was also measured in the presence of rotenone and antimycin A, known inhibitors of complex I and complex III of the electron transport chain (ETC), respectively. Complexes I and III have been reported as the primary sites of mitochondrial ROS production (Hinkle et al., 1967; Boveris et al., 1972; Cadenas et al., 1977; Raha et al., 2000) and these inhibitors would assess that ROS production capacity of the ETC at these complexes. All these reactions were performed in triplicate for both strains at the same time in the same plate. In addition, a vehicular control was included to provide values for non-mitochondrial ROS. The data in arbitrary units are mean activity of the reaction for 30 minutes measured in a Synergy 2 plate reader at 600 nm.

Quantitative (q) PCR

Groups of 10 permethrin susceptible and RSP females each in six replicates were homogenised using TRI REAGENT (Promega) as described in the manufacturer's instruction. First strand cDNA was synthesised from 500 ng/ μL of total RNA using SuperScript III, RNaseOUT, reverse transcriptase buffer, MgCl_2 and an Oligo(dT)₂₀ (Invitrogen) as described in the manufacturer's protocol. Thirty five cycles of amplification were performed in PCR machine. The amplification cycle was as follows: 95°C for 5 min, 95°C for 30 sec, 53°C for 30 sec and 72°C for 45 sec and 72°C 7 min. We performed quantitative qPCR using a SYBR Green Master mix and 50 ng total of cDNA per reaction and run in a Stratagene Mx3000P® qPCR machine. The primers used for qPCR on the same total RNA are listed in Table 2.

Data analysis

A χ^2 test was used to compare mean longevity between strains. A log-rank test was used to compare median longevity between strains and a Kaplan-Meier curve was provided to illustrate the difference between strains. The data analysis for metabolic rate and glycogen levels was performed using analysis of covariance, with body weight used as a covariate. A two-sample t -test was used to compare differences between means for the other phenotypes. Statistical significance was set at $\alpha < 0.05$ for each test.

Results

Developmental time

We found that the median developmental time from L1 to pupa was significantly longer (4%) in the RSP strain compared to the permethrin susceptible strain, (log-rank test $\chi^2_1 = 8.9$, $p = 0.003$) (Figure 1). Significantly longer (6%) was also the median developmental time from L1 to adult emergence (log-rank test $\chi^2_1 = 13.6$, $p = 0.0002$) (Figure 1).

Body weight, energy storage, and metabolic rate

Evidence exists of a strong trade-off between larval developmental time and adult weight in insects (Santos et al., 1994). Therefore, we measured both wet body weight and dry weight of 3–5 day old females. RSP females had significantly higher wet weight than permethrin susceptible females ($p = 0.044$) (Figure 2A). On the other hand, there were no significant differences in dry body weight between the strains ($p = 0.250$) (Figure 2B). As expected considering the higher wet weight, the RSP females had on average 5% more water content ($p = 0.026$) than permethrin susceptible female (Figure 2C).

The capacity of glycogen to bind water is 3–5 times its own weight (Schmidt-Nielsen, 1997). To explore whether the higher water content of RSP females was accompanied by an increase in glycogen level, we measured the glycogen content of whole-body sugar-fed females. Compared to permethrin susceptible females, RSP females had on average 9.5% higher levels of body weight-adjusted glycogen in ($p = 0.014$) (Figure 2 D). No statistical difference in triacylglycerol levels ($p = 0.27$) was, however, observed between the two strains (Figure 2 E). The mean triacylglycerol content for the RSP and permethrin-susceptible females was $11.36 \mu\text{l/mg} \pm 1.71$ and $14.51 \mu\text{l/mg} \pm 2.17$, respectively.

Previous studies of desiccation resistance and water balance in natural populations of *Drosophila* (Gibbs & Matzkin, 2001) have shown a positive correlation between metabolic rates and water-loss rates (Gibbs & Matzkin, 2001). The relationship between metabolic rate and water loss has been observed in a variety of other insects as reviewed by (Chown & Gaston, 1999) and is explained by the fact that reducing metabolic rates can help the insect to conserve water by reducing the need for gas exchange. Thus, we measured CO_2 production in the two strains. There was a statistical difference in metabolic rate between the strains ($p = 0.034$). As predicted, the mean VCO_2 in RSP was lower than in permethrin susceptible females, $20.48 \mu\text{l/h} \pm 1.27$ and $25.07 \mu\text{l/h} \pm 1.48$, respectively (Figure 2 F).

Fecundity

The mean number of eggs produced in one gonotrophic cycle between the two strains was not significantly different (Table 1). No differences were also observed in the number of eggs hatched and the time from blood feeding to laying eggs between the two strains (Table 1).

Longevity and mitochondrial bioenergetics

We observed a significantly reduced adult life span of RSP females compared to permethrin susceptible females (log-rank test $\chi^2_1 = 10.49$, $p = 0.0012$) (Figure 3).

Previous work in *Drosophila* reported a positive correlation between inter-individual variability in survival and mitochondrial bioenergetics (Melvin & Ballard, 2006). Thus, we next measured mitochondrial coupling efficiency (P: O ratio) and H₂O₂ production rate using NADH-linked substrates. Notably, mitochondria isolated from thoraces of RSP females showed a significantly lower coupling efficiency than those isolated from thoraces of permethrin susceptible females ($p = 0.050$) (Figure 4A). Furthermore, the RSP females' mitochondria produced significantly more ROS with only substrates present and no ADP (no oxidative phosphorylation or ATP production occurring), reflective of the leakiness of the system to proton re-entry (state 2; $p = 0.010$), and also produced significantly more ROS from complex I (rotenone; $p = 0.017$) than those from susceptible females (Figure 4B). There was no statistically significant difference in ROS production from complex III between the strains (antimycin A; $p = 0.096$) (Figure 4B).

Antioxidant gene expression

To control for the oxidative stress induced by the enhanced production of ROS in mitochondria, cells produce different enzymes that scavenge ROS, such as superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPX), and GST (Hayes & McLellan, 1999). Previous work in *An. gambiae* showed that the expression of *GSTe3* is significantly enhanced following H₂O₂ exposure (Kumar et al., 2003; Ding et al., 2005). Additionally, it has been reported that CAT is the primary antioxidant enzyme involved in oocyte protection by ROS damage in *An. gambiae* (DeJong et al., 2007). To assess whether the increased mitochondrial ROS production in the RSP strain is associated with the induction of components of the antioxidant system, we measured the expression of *GSTe3*, *CAT*, *GPXH1*, *SOD1*, *SOD2* genes using mRNA extracted from whole-body of females of each strain. We found a statistically significant difference in *GSTe3* ($p = 0.011$) and catalase ($p = 0.0068$) expression levels between the two strains, with RSP females showing on average 80% *GSTe3* (Figure 5A) and 92% *CAT* (Figure 5B) respectively higher transcript levels than permethrin susceptible females. No significant difference was observed in the expression levels of either *GPXH1* ($p = 0.650$) (Figure 5C), *SOD1* ($p = 0.430$) (Figure 5D) or *SOD2* ($p = 0.480$) (Figure 5E).

Discussion

Here, we compared several life history and energy metabolism traits of a permethrin susceptible strain of *An. gambiae* and a strain harboring *kdr* and metabolic-based

mechanisms conferring permethrin resistance (RSP strain). We observed that the RSP strain had lower metabolic rate, slower developmental time, and shorter adult life span than the permethrin susceptible strain. Additionally, consistent with previous evidence in different organisms of a relationship between enhanced levels of ROS and age-associated decline (reviewed in (Marchi et al., 2012)), mitochondria isolated from the thoraces of the RSP females were found to have lower coupling efficiency and higher ROS production rates. Collectively, these results suggest that permethrin resistance alleles could affect energy metabolism, oxidative stress, and adult survival of *An. gambiae* and therefore impose strong fitness costs to the malaria vector. However, one major limitation of this study is that the RSP and permethrin susceptible strains have different genetic background and thus we cannot exclude the possibility that different loci may be responsible for the phenotypic differences observed between the strains. Also, although our strains of *An. gambiae* were originally isolated from the wild in the same geographic area in Kenya, they were subsequently reared under laboratory conditions for over ten years and may suffer from inbreeding depression. Therefore, our results need to be interpreted with caution and replicated in independent studies in order to have significant implications for malaria transmission and vector control.

In our study, we found that both developmental times from L1 to pupation and from L1 to adult emergence were longer in RSP when compared to the permethrin susceptible strain. However, despite the longer developmental time, we did not observe a difference in dry body weight between females of the two *An. gambiae* strains. This result is interesting considering the strong relationship between molecular and physiological mechanisms that regulate the duration of developmental time and rate of growth and final body size (Shingleton, 2011). Compared to females of the permethrin susceptible strain, adult females of the RSP strain also showed higher water and glycogen levels, which is in agreement with the high energetic investment that is required to support the body's detoxifying apparatus (Coustau et al., 2000).

P450 monooxygenase-mediated detoxification is a major mechanism of resistance to insecticides due to the wide variety of substrates that P450s can metabolise (Félix & Silveira, 2012). The P450 monooxygenase system consists of two main components: the cytochrome P450, which acts as the substrate binding protein (and terminal oxidase), and the NADPH-cytochrome P450 reductase (P450 reductase), which transfers electrons from NADPH to cytochrome P450 (Kawano et al., 1987; Scott & Wheelock, 1991; Guzov et al., 1996). The overproduction of detoxifying enzymes in insects can be energetically costly (Coustau et al., 2000), and investment in resistance can be expected to produce reduced investment in competing organismal functions.

Notably, our study did not show evidence of a reproductive fitness cost associated with the presence of both target-site insensitivity and metabolic-based mechanisms of permethrin resistance in *An. gambiae*. However, the lifespan of RSP females was significantly reduced compared to the permethrin susceptible females. A potential explanation for these findings is that the resistant strain might invest more energy in enhancing the antioxidative and detoxification mechanisms and maintaining reproductive functions at the cost of somatic maintenance and repair, allowing faster rates of aging and decreased longevity (Rose &

Charlesworth, 1981a; Rose & Charlesworth, 1981b). Consistent with this idea, we found lower mitochondrial coupling efficiency and higher ROS production rates in the mitochondria isolated from the thoraces of the RSP females. ROS, including superoxide and its dismutation product H_2O_2 , are essential as signaling molecules in defense against infection and in reproduction (Sanz & Stefanatos, 2008). But, if produced in excess, they can oxidise and damage various cellular components, including mitochondrial proteins, membranes, lipids, and nuclear and mitochondrial genomes, and thus have been implicated in the aging process of a variety of species (Sanz & Stefanatos, 2008), including *Anopheles* (Monaghan et al., 2009). In insects, increased levels of P450 activity has been associated with high levels of ROS production as byproducts of the detoxification processes (Murataliev et al., 2008). As such, it is tempting to speculate that the RSP strain may exhibit higher levels of ROS produced by P450 than the permethrin-susceptible strain. These ROS may in turn oxidise and damage mitochondrial DNA, protein, and lipids, possible leading to mitochondrial dysfunction and excessive production of ROS by redox-coupled reactions within the ETC, as suggested by our mitochondrial bioenergetic results.

Antioxidant enzymes are produced by the cell in response to the oxidative stress induced by the enhanced production of ROS in mitochondria. We observed an 80% and 90% increase in expression of *GSTe3* and *CAT*, respectively, in young RSP females compared to females of the susceptible strain. As mentioned above, previous work reported that the antioxidant enzyme catalase is responsible for oocyte protection by ROS damage in *An. gambiae* (DeJong et al., 2007). Based on this observation, our gene expression data not only suggest that RSP mosquitoes might activate a strong antioxidant response to counterbalance an increased oxidative environment, but also support our hypothesis of an investment of energy resources to maintain fecundity.

Acknowledgments

We thank Michelle Moses Chambers and Gin Chuang for helping with metabolic rate and mitochondrial ROS production measurements, respectively. We also thank Jeff Leips for providing insightful comments on the manuscript. We are grateful to Edward W. Hook III and Sally Fried of the Division of Infectious Diseases, William C. Gorgas Center for Geographic Medicine, for granting permission to use the mosquito facility. The RSP strain was obtained through the MR4 as part of the BEI Resources Repository, NIAID, NIH: *Anopheles gambiae* RSP, MRA-334, deposited by J Vulule, MQ Benedict. The ASEMBO1 strain was obtained through the MR4 as part of the BEI Resources Repository, NIAID, NIH: *Anopheles gambiae* ASEMBO1 [AB], MRA-186, deposited by MQ Benedict. This study was supported by the Department of Biology at the University of Alabama at Birmingham, the Bioanalytical Redox Biology Core (DRTC P60 DK079626), and NIH grant R01 DK084219 to MD.

Abbreviations

ADP	adenosine diphosphate
AR	Amplex red
ATP	adenosine triphosphate
BSA	bovine serum albumen
CAT	catalase
DI	deionised

EDTA	ethylene diamine tetraacetic acid
ETC	electron transport chain
GPX	glutathione peroxidase
GST	glutathione S-transferase
<i>kdr</i>	knockdown resistance
MR4	Malaria Research and Reference Reagent Resource Center
NAD	nicotinamide adenine dinucleotide
ROS	reactive oxygen species
RSP	reduced susceptibility to permethrin
SOD	super oxide dismutase
VGSC	voltage-gated sodium channel

References

- Adasi K, Hemingway J. Susceptibility to three pyrethroids and detection of knockdown resistance mutation in Ghanaian *Anopheles gambiae* sensu stricto. *Journal of Vector Ecology*. 2008; 33:255–262. [PubMed: 19263844]
- Affourtit C, Quinlan CL, Brand MD. Measurement of proton leak and electron leak in isolated mitochondria. *Methods in Molecular Biology*. 2012; 810:165–182. [PubMed: 22057567]
- Ayyadevara S, Engle MR, Singh SP, Dandapat A, Lichti CF, Benes H, Shmookler Reis RJ, Liebau E, Zimniak P. Lifespan and stress resistance of *Caenorhabditis elegans* are increased by expression of glutathione transferases capable of metabolizing the lipid peroxidation product 4-hydroxynonenal. *Aging Cell*. 2005; 4:257–271. [PubMed: 16164425]
- Boveris A, Oshino N, Chance B. The cellular production of hydrogen peroxide. *Biochemical Journal*. 1972; 128:617–630. [PubMed: 4404507]
- Brooke BD, Koekemoer LL. Major effect genes or loose confederations? The development of insecticide resistance in the malaria vector *Anopheles gambiae*. *Parasities & Vectors*. 2010; 3:74.
- Brown ZS, Dickinson KL, Kramer RA. Insecticide resistance and malaria vector control: the importance of fitness cost mechanisms in determining economically optimal control trajectories. *Journal of Economic Entomology*. 2013; 106:366–374. [PubMed: 23448053]
- Cadenas E, Boveris A, Ragan CI, Stoppani AO. Production of superoxide radicals and hydrogen peroxide by NADH-ubiquinone reductase and ubiquinol-cytochrome c reductase from beef-heart mitochondria. *Archives of Biochemistry and Biophysics*. 1977; 180:248–257. [PubMed: 195520]
- Casimiro S, Coleman M, Hemingway J, Sharp B. Insecticide resistance in *Anopheles arabiensis* and *Anopheles gambiae* from Mozambique. *Journal of Medical Entomology*. 2006; 43:276–282. [PubMed: 16619611]
- Chown SL, Gaston KJ. Exploring links between physiology and ecology at macro-scales: the role of respiratory metabolism in insects. *Biological Reviews of the Cambridge Philosophical Society*. 1999; 74:87–120.
- Coustau C, Chevillon C, ffrench-Constant R. Resistance to xenobiotics and parasites: can we count the cost? *Trends in Ecology & Evolution*. 2000; 15:378–383. [PubMed: 10931680]
- De Luca M, Klimentidis YC, Casazza K, Chambers MM, Cho R, Harbison ST, Jumbo-Lucioni P, Zhang S, Leips J, Fernandez JR. A conserved role for syndecan family members in the regulation of whole-body energy metabolism. *PLoS One*. 2010; 5:e11286. [PubMed: 20585652]
- DeJong RJ, Miller LM, Molina-Cruz A, Gupta L, Kumar S, Barillas-Mury C. Reactive oxygen species detoxification by catalase is a major determinant of fecundity in the mosquito *Anopheles gambiae*.

- Proceedings of the National Academy of Sciences of the United States of America. 2007; 104:2121–2126. [PubMed: 17284604]
- Ding Y, Hawkes N, Meredith J, Eggleston P, Hemingway J, Ranson H. Characterization of the promoters of Epsilon glutathione transferases in the mosquito *Anopheles gambiae* and their response to oxidative stress. *Biochemical Journal*. 2005; 387:879–888. [PubMed: 15631620]
- Djouaka RF, Bakare AA, Coulibaly ON, Akogbeto MC, Ranson H, Hemingway J, Strode C. Expression of the cytochrome P450s, CYP6P3 and CYP6M2 are significantly elevated in multiple pyrethroid resistant populations of *Anopheles gambiae* s.s. from Southern Benin and Nigeria. *BMC Genomics*. 2008; 9:538. [PubMed: 19014539]
- Enayati AA, Hemingway J. Pyrethroid insecticide resistance and treated bednets efficacy in malaria control. *Pesticide Biochemistry and Physiology*. 2006; 84:116–126.
- Félix, R.; Silveira, H. The Role of *Anopheles gambiae* P450 Cytochrome in Insecticide Resistance and Infection. In: Perveen, F., editor. *Insecticides - Pest Engineering*. InTech; 2012. p. 503-518.
- Ferguson M, Mockett RJ, Shen Y, Orr WC, Sohal RS. Age-associated decline in mitochondrial respiration and electron transport in *Drosophila melanogaster*. *Biochemical Journal*. 2005; 390:501–511. [PubMed: 15853766]
- Gibbs AG, Matzkin LM. Evolution of water balance in the genus *Drosophila*. *The Journal of Experimental Biology*. 2001; 204:2331–2338. [PubMed: 11507115]
- Guzov VM, Houston HL, Murataliev MB, Walker FA, Feyereisen R. Molecular cloning, overexpression in *Escherichia coli*, structural and functional characterization of house fly cytochrome b5. *Journal of Biological Chemistry*. 1996; 271:26637–26645. [PubMed: 8900138]
- Hayes JD, McLellan LI. Glutathione and glutathione-dependent enzymes represent a co-ordinately regulated defence against oxidative stress. *Free Radical Research*. 1999; 31:273–300. [PubMed: 10517533]
- Hemingway J, Ranson H. Insecticide resistance in insect vectors of human disease. *Annual Review of Entomology*. 2000; 45:371–391.
- Hinkle PC, Butow RA, Racker E, Chance B. Partial resolution of the enzymes catalyzing oxidative phosphorylation. XV. Reverse electron transfer in the flavin-cytochrome beta region of the respiratory chain of beef heart submitochondrial particles. *The Journal of Biological Chemistry*. 1967; 242:5169–5173. [PubMed: 4294331]
- Jumbo-Lucioni P, Ayroles JF, Chambers MM, Jordan KW, Leips J, Mackay TF, De Luca M. Systems genetics analysis of body weight and energy metabolism traits in *Drosophila melanogaster*. *BMC Genomics*. 2010; 11:297. [PubMed: 20459830]
- Jumbo-Lucioni P, Bu S, Harbison ST, Slaughter JC, Mackay TF, Moellering DR, De Luca M. Nuclear genomic control of naturally occurring variation in mitochondrial function in *Drosophila melanogaster*. *BMC Genomics*. 2012; 13:659. [PubMed: 23171078]
- Kawano S, Kamataki T, Yasumori T, Yamazoe Y, Kato R. Purification of human liver cytochrome P-450 catalyzing testosterone 6 beta-hydroxylation. *Journal of Biochemistry*. 1987; 102:493–501. [PubMed: 3429444]
- Kliot A, Ghanim M. Fitness costs associated with insecticide resistance. *Pest Management Science*. 2012; 68:1431–1437. [PubMed: 22945853]
- Kumar S, Christophides GK, Cantera R, Charles B, Han YS, Meister S, Dimopoulos G, Kafatos FC, Barillas-Mury C. The role of reactive oxygen species on *Plasmodium* melanotic encapsulation in *Anopheles gambiae*. *Proceedings of the National Academy of Sciences of the United States of America*. 2003; 100:14139–14144. [PubMed: 14623973]
- Li X, Schuler MA, Berenbaum MR. Molecular mechanisms of metabolic resistance to synthetic and natural xenobiotics. *Annual Review of Entomology*. 2007; 52:231–253.
- Lumjuan N, Rajatileka S, Changsom D, Wicheer J, Leelapat P, Prapanthadara LA, Somboon P, Lycett G, Ranson H. The role of the *Aedes aegypti* Epsilon glutathione transferases in conferring resistance to DDT and pyrethroid insecticides. *Insect Biochemistry and Molecular Biology*. 2011; 41:203–209. [PubMed: 21195177]
- Marchi S, Giorgi C, Suski JM, Agnoletto C, Bononi A, Bonora M, De Marchi E, Missiroli S, Patergnani S, Poletti F, Rimessi A, Duszynski J, Wieckowski MR, Pinton P. Mitochondria-ros

- crosstalk in the control of cell death and aging. *Journal of Signal Transduction*. 2012; 2012:329635. [PubMed: 22175013]
- McElwee JJ, Schuster E, Blanc E, Thomas JH, Gems D. Shared transcriptional signature in *Caenorhabditis elegans* Dauer larvae and long-lived daf-2 mutants implicates detoxification system in longevity assurance. *The Journal of Biological Chemistry*. 2004; 279:44533–44543. [PubMed: 15308663]
- Melvin RG, Ballard JW. Intraspecific variation in survival and mitochondrial oxidative phosphorylation in wild-caught *Drosophila simulans*. *Aging Cell*. 2006; 5:225–233. [PubMed: 16842495]
- Miwa S, St-Pierre J, Partridge L, Brand MD. Superoxide and hydrogen peroxide production by *Drosophila* mitochondria. *Free Radical Biology & Medicine*. 2003; 35:938–948. [PubMed: 14556858]
- Monaghan P, Metcalfe NB, Torres R. Oxidative stress as a mediator of life history trade-offs: mechanisms, measurements and interpretation. *Ecology Letters*. 2009; 12:75–92. [PubMed: 19016828]
- Murataliev MB, Guzov VM, Walker FA, Feyereisen R. P450 reductase and cytochrome b5 interactions with cytochrome P450: effects on house fly CYP6A1 catalysis. *Insect Biochemistry and Molecular Biology*. 2008; 38:1008–1015. [PubMed: 18930820]
- Okoye PN, Brooke BD, Hunt RH, Coetzee M. Relative developmental and reproductive fitness associated with pyrethroid resistance in the major southern African malaria vector, *Anopheles funestus*. *Bulletin of Entomological Research*. 2007; 97:599–605. [PubMed: 17997873]
- Raha S, McEachern GE, Myint AT, Robinson BH. Superoxides from mitochondrial complex III: the role of manganese superoxide dismutase. *Free Radical Biology & Medicine*. 2000; 29:170–180. [PubMed: 10980405]
- Ranson H, Jensen B, Vulule JM, Wang X, Hemingway J, Collins FH. Identification of a point mutation in the voltage-gated sodium channel gene of Kenyan *Anopheles gambiae* associated with resistance to DDT and pyrethroids. *Insect Biochemistry and Molecular Biology*. 2000; 9:491–497.
- Ray, DE. Pyrethroid insecticides: mechanisms of toxicity, systemic poisoning syndromes, paresthesia, and therapy. In: Krieger, R.; Doull, J.; Ecobichon, D., editors. *Agents*. San Diego: Academic Press; 2001. p. 1289–1303.
- Read AF, Lynch PA, Thomas MB. How to make evolution-proof insecticides for malaria control. *PLoS Biology*. 2009; 7:e1000058. [PubMed: 19355786]
- Rigby MC, Hechinger RF, Stevens L. Why should parasite resistance be costly? *Trends in Parasitology*. 2002; 18:116–120. [PubMed: 11854088]
- Rivero A, Vezilier J, Weill M, Read AF, Gandon S. Insecticide control of vector-borne diseases: when is insecticide resistance a problem? *PLoS Pathogens*. 2010; 6:e1001000. [PubMed: 20700451]
- Roberto HK, Omoto C. Fitness cost associated with carbosulfan resistance in *Aphis gossypii* Glover (Hemiptera: Aphididae). *Neotropical Entomology*. 2006; 35:246–250. [PubMed: 17348137]
- Rose MR, Charlesworth B. Genetics of life history in *Drosophila melanogaster*. I. Sib analysis of adult females. *Genetics*. 1981a; 97:173–186. [PubMed: 6790340]
- Rose MR, Charlesworth B. Genetics of life history in *Drosophila melanogaster*. II. Exploratory selection experiments. *Genetics*. 1981b; 97:187–196. [PubMed: 6790341]
- Rowland M. Behaviour and fitness of gamma HCH/dieldrin resistant and susceptible female *Anopheles gambiae* and *An. stephensi* mosquitoes in the absence of insecticide. *Medical and Veterinary Entomology*. 1991; 5:193–206. [PubMed: 1722728]
- Sacktor B, Sanborn R. The effect of temperature on oxidative phosphorylation with insect flight muscle mitochondria. *Journal of Biophysical and Biochemical Cytology*. 1956; 2:105–107. [PubMed: 13295315]
- Santos M, Fowler K, Partridge L. Gene-environment interaction for body size and larval density in *Drosophila melanogaster*: an investigation of effects on development time, thorax length and adult sex ratio. *Heredity (Edinb)*. 1994; 72 (Pt 5):515–521. [PubMed: 8014062]
- Sanz A, Stefanatos RK. The mitochondrial free radical theory of aging: a critical view. *Current Aging Science*. 2008; 1:10–21. [PubMed: 20021368]

- Schimdt-Nielsen, K. Animal Physiology: Adaptation and Environment. Schimdt-Nielsen, K., editor. Cambridge University Press; 1997. p. 169-216.
- Scott, GJ.; Wheelock, GD. Characterization of a Cytochrome P450 responsible for pyrethroid resistance in the housefly. 2nd National Meeting of the American Chemical Society; New York. 25 August – 30 August 1991; NY: American Chemical Society; 1991. p. 16-20.
- Shingleton, A. Evolution and the regulation of growth and body size. In: Flat, T.; Heyland, A., editors. Mechanisms of Life History Evolution. The Genetics and Physiology of Life History Traits and Trade-Offs. Oxford University Press; 2011. p. 43-55.
- Siegel JP, Novak RJ, Ruesink WG. Relationship between Wing Length and Dry-Weight of Mosquitos. Journal of the American Mosquito Control Association. 1994; 10:186–196.
- Soderlund DM. Pyrethroids, knockdown resistance and sodium channels. Pest Management Science. 2008; 64:610–616. [PubMed: 18383430]
- Vulule JM, Beach RF, Atieli FK, McAllister JC, Brogdon WG, Roberts JM, Mwangi RW, Hawley WA. Elevated oxidase and esterase levels associated with permethrin tolerance in *Anopheles gambiae* from Kenyan villages using permethrin-impregnated nets. Medical and Veterinary Entomology. 1999; 13:239–244. [PubMed: 10514048]
- WHO. Insecticide-Treated Mosquito Nets: A WHO Position Statement. World Health Organization; 2007.
- Wiley RH. Evolution of social organization and life-history patterns among grouse. Quarterly Review of Biology. 1974; 49:201–227.

Figure 1.

Developmental time from first-instar larva to pupa (Panel A; $n = 465$) and to adult emergence (Panel B, $n = 337$) in permethrin susceptible and RSP strains.

Figure 2.

Adult wet weight (Panel A; $n = 30$), dry weight (Panel B; $n = 30$), water content (Panel C; $n = 30$), glycogen levels (Panel D; $n = 20$), triacylglycerol storage (Panel E; $n = 20$) and metabolic rate (Panel F; $n = 10$) in permethrin susceptible and RSP females.

Glycogen and triacylglycerol levels are normalised for body weight. Values are expressed as mean $\pm$ SE. * $p < 0.05$

Figure 3.

Adult female survivorship curves for the permethrin susceptible and RSP strains of *An. gambiae*. Survival assays were carried out using population cages with initial population sizes of 46–50 individuals.

Figure 4.

Coupling efficiency (or P:O ratio) (Panel A) and ROS production (Panel B) of mitochondria isolated from thoraces of permethrin susceptible and RSP females using NAD^+ -linked substrates (pyruvate and proline). Values are given as means $\pm$ SE of six independent replicates. * $p < 0.05$.

Figure 5.

Relative mRNA expression of antioxidant enzymes in whole body extracts of permethrin susceptible and RSP females. (Panel A) *GSTe3*: Glutathione *S* Transferase *e3*; (Panel B) *CAT*: Catalase; (Panel C) *GPXH1*: Glutathione Peroxidase; (Panel D) *SOD1*: Superoxide Dismutase 1; and (Panel E) *SOD2*: Superoxide Dismutase 2. Transcript levels of the five genes were normalised to *RSP7 Ribosomal Protein S7*. Values are given as means $\pm$ SE of six independent replicates. * $p < 0.05$.

Table 1

Reproductive characteristics for the permethrin susceptible and RSP strains of *An. gambiae*.

	Susceptible	RSP	<i>p</i> value
Eggs laid	66.07±54.46 (24–182)	56.35±63.85 (8–211)	0.46
Eggs hatched	47.92±42.54 (2–129)	50.08±35.95 (2–139)	0.84
Duration (days) to lay eggs	3.66	3.55	0.66

Mean ± SD (range)

Table 2

Nucleotide sequence of primers used for qPCRs

Gene	Primer	Sequence
<i>GPXH1</i>	GPXH1-FF	CCGGAAGGCGACGGAGAGGA
	GPXH1-RR	TCCACCGGCTGACCGTCCTT
<i>SOD1</i>	SOD1-FF	TCACTAACAATAATGTTGGCCGTA
	SOD1-RR	AACTTGATCGCATTGCCGAG
<i>RPS7</i>	RPS7-FF	TCATCATCTACGTGCCGGTG
	RPS7-RR	TACACGGCGGTGACGTTC
<i>GSTe3</i>	GSTe3-FF	GAGCTGACGGCAAAGATGATCG
	GSTe3-RR	CCTGCTTCACTAGATCCTTCGC
<i>SOD2</i>	SOD2-FF	CATCAACGTTGTGGGGCTGA
	SOD2-RR	ATGTTACCGAGATCGCCGAC
<i>CAT</i>	CAT-FF	AATTCCCGCTCATTCCGGTT
	CAT-RR	TCAGATAGTTGGCACCCACG