Research Highlight

Methylation of FoxO3 regulates neuronal cell death

Wei-guo ZHU^{1, 2, *}

Acta Pharmacologica Sinica (2012) 33: 577; doi: 10.1038/aps.2012.48

Oxidative stress plays an essential role in the development of several neurodegenerative disorders^[1]. The evolutionally conserved FoxO (FOX family O class) transcription factors, including FoxO1, FoxO3, FoxO4, and FoxO6 in mammals, induce the expression of downstream pro-apoptotic genes, thus act as critical regulators during the oxidative stress-induced neuronal cell death^[2, 3].

FoxO family proteins are regulated by a variety of post-translational modifications, such as phosphorylation, acetylation, ubiquitination and arginine methylation^[4]. For instance, serine/threonine kinase PKB/Akt is well known FoxO inhibitor. Phosphorylation of FoxO by Akt at three key residues increases its binding activity to 14-3-3 protein and results in its translocation from nucleus to cytoplasm^[5]. On the other hand,

phosphorylation of FoxO by Hippo/MST1 kinase dampens its binding to 14-3-3 protein and leads to nuclear accumulation of FoxO1/3^[3]. Similar to phosphorylation, acetylation/deacetylation of FoxO proteins dynamically regulates their biological functions through affecting their DNA-binding activity, stability and interaction with other proteins^[6].

Recently, Yuan's group of Institute of Biophysics at Chinese Academy of Sciences identified a novel post-translational modification of FoxO3 transcription factor^[7]. They found that a lysine methyltransferase, Set9, methylated FoxO3 at lysine 270 that led to inhibition of DNA binding activity and downregulation of transactivation without affecting Akt-mediated phosphorylation. They showed that lysine methylation of FoxO3 reduced oxidative stress-induced neuronal apoptosis in cerebellar granule neurons through transcriptional downregulation of its pro-apoptotic target, *Bim*. In addition, they observed that the lysine methylation of FoxO3 did not affect its protein stability, Akt-mediated phosphorylation and subcellular localization. Interestingly, they showed that oxidative stress reduced the methylation level of FoxO3, indicating that the

methylation of FoxO3 might be dynamically regulated even though they failed to identify the demethylase of FoxO3.

In short, Yuan's group identifies a novel modification of FoxO3 and shows that lysine methylation negatively regulates FoxO3-mediated transcription and neuronal cell death, which implicates a variety of neurodegenerative diseases.

¹Key laboratory of Carcinogenesis and Translational Research (Ministry of Education), Department of Biochemistry and Molecular Biology, Peking University Health Science Center, Beijing 100191, China; ²Peking University -Tsinghua University Center for Life Sciences, Beijing 100084, China

Correspondence: Prof Wei-guo ZHU
(zhuweiguo@bjmu.edu.cn)

- 1 Chong ZZ, Li F, Maiese K. Oxidative stress in the brain: novel cellular targets that govern survival during neurodegenerative disease. *Prog Neurobiol* 2005; 75: 207–46.
- 2 Calnan DR, Brunet A. The FoxO code. *Oncogene* 2008; 27: 2276–88.
- 3 Lehtinen MK, Yuan Z, Boag PR, Yang Y, Villen J, Becker EB, et al. A conserved MST-FOXO signaling pathway mediates oxidative-stress responses and extends life span. *Cell* 2006; 125: 987–1001.
- 4 van der Horst A, Burgering BM. Stressing the role of FoxO proteins in lifespan and disease. *Nat Rev Mol Cell Biol* 2007; 8: 440–50.
- 5 Brunet A, Bonni A, Zigmond MJ, Lin MZ, Juo P, Hu LS, et al. Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell* 1999; 96: 857–68.
- 6 Zhao Y, Wang Y, Zhu WG. Applications of post-translational modifications of FoxO family proteins in biological functions. *J Mol Cell Biol* 2011; 3: 276–82.
- 7 Xie Q, Hao Y, Tao L, Peng S, Rao C, Chen H, et al. Lysine methylation of FOXO3 regulates oxidative stress-induced neuronal cell death. *EMBO Rep* 2012; 13: 371–7.