Published in final edited form as:

Environ Microbiol. 2014 August ; 16(8): 2623–2634. doi:10.1111/1462-2920.12322.

Non-native acylated homoserine lactones reveal that LuxIR quorum sensing promotes symbiont stability

Sarah V. Studer^{#†,*}, Julia A. Schwartzman^{#†}, Jessica S. Ho^{†,§}, Grant D. Geske^{‡,¥}, Helen E. Blackwell[‡], and Edward G. Ruby^{†,§}

[†]Department of Medical Microbiology and Immunology, University of Wisconsin-Madison, 1550 Linden Drive, Madison WI 53706

[‡]Department of Chemistry, University of Wisconsin-Madison, 1101 University Avenue, Madison WI 53706

[#] These authors contributed equally to this work.

SUMMARY

Quorum sensing, a group behavior coordinated by a diffusible pheromone signal and a cognate receptor, is typical of bacteria that form symbioses with plants and animals. LuxIR-type acyl homoserine-lactone (AHL) quorum sensing is common in Gram-negative proteobacteria, and many members of this group have additional quorum-sensing networks. The bioluminescent symbiont *Vibrio fischeri* encodes two AHL signal synthases: AinS and LuxI. AinS-dependent quorum sensing converges with LuxI-dependent quorum sensing at the LuxR regulatory element. Both AinS- and LuxI-mediated signaling are required for efficient and persistent colonization of the squid host, *Euprymna scolopes*. The basis of the mutualism is symbiont bioluminescence, which is regulated by both LuxI- and AinS-dependent quorum sensing, and is essential for maintaining a colonization of the host. Here, we used chemical and genetic approaches to probe the dynamics of LuxI- and AinS-mediated regulation of bioluminescence during symbiosis. We demonstrate that both native AHLs and non-native AHL analogs can be used to non-invasively and specifically modulate induction of symbiotic bioluminescence *via* LuxI-dependent quorum sensing. Our data suggest that the first day of colonization, during which symbiont bioluminescence is induced by LuxIR, is a critical period that determines the stability of the *V. fischeri* population once symbiosis is established.

[#]Corresponding author, egruby@wisc.edu.

^{*}Current address: Fuel Cell Technologies Office, Office of Energy Efficiency and Renewable Energy (EERE), Department of Energy, 1000 Independence Ave. SW, Washington, DC 20009, USA

[§]Current address: Laboratory of Immune Cell Epigenetics and Signaling, The Rockefeller University, 1230 York Avenue, New York, New York 10065, USA

[¥]Current address: Cushing Academy, 39 School St., Ashburnham, MA, 01430, USA

Author contributions: SVS and JAS performed the experiments, and JAS performed the data analysis. JSH and GDG performed preliminary experiments. JSH, HEB, and EGR helped with experimental design. JAS, SVS, and EGR prepared the manuscript, with help from, and editing by, all co-authors.

INTRODUCTION

A common thread among microbes that initiate symbioses with plant and animal hosts is the quorum sensing-dependent regulation of colonization factors (Parsek and Greenberg, 2000; González and Venturi, 2012). Quorum sensing relies on perception of an endogenously synthesized secreted pheromone signal molecule, called an autoinducer, by a cognate receptor in a concentration-dependent manner. LuxIR quorum-sensing systems are widespread among Gram-negative bacteria, which use a LuxR-type quorum-sensing receptor to perceive an *N*-acyl L-homoserine lactone (AHL) chemical signal(s) (Churchill and Chen, 2010). The importance of AHL-based quorum sensing in pathogens such as *Pseudomonas aeruginosa*, or mutualists such as *Sinorhizobium meliloti* (Teplitski et al., 2011; Galloway et al., 2012), has led to significant interest in developing methods to manipulate this regulatory circuit *via* interception of the native AHL signal molecule (Rasmussen and Givskov, 2006; Amara et al., 2011; Galloway et al., 2011; Praneenarat et al., 2012).

Despite this interest, only a few studies (Hentzer et al., 2003; Wu et al., 2004; Palmer et al., 2011a) have chemically modulated bacterial AHL quorum-sensing in a host model to ask whether signaling affects colonization robustness in the host environment, and all of these studies have focused on pathogenic associations (Bjarnsholt and Givskov, 2007). Pathogens represent only a small fraction of the microbes that both encode LuxIR-type systems and colonize animal or plant hosts; thus, we chose to apply a chemical approach, in combination with existing strains of *V. fischeri* carrying mutations in AinS-LitR and LuxIR branches of quorum sensing, to study the role of the LuxIR signal circuit in the maintenance of stable, beneficial host-microbe associations.

The symbiosis between the marine bacterium *Vibrio fischeri* and the squid *Euprymna scolopes* is a model system to study the initiation and maintenance of a natural, two-partner mutualism (Mandel, 2010). A monospecific, and extracellular population of *V. fischeri* is maintained in a specialized host structure called the light organ, where, as the name would suggest, symbionts produce light in exchange for the habitat provided by the host. Bioluminescence, and other behaviors that promote the stable association of a microbe and its host, are regulated by quorum sensing in *V. fischeri* (Stabb and Visick, 2013).

The principal quorum-sensing circuit in *V. fischeri* is composed of the AHL signal molecule *N*-(3-oxo-hexanoyl)-L-homoserine lactone (3-oxo C6 HL), the signal synthase LuxI, and the AHL-binding transcriptional regulatory element LuxR. The LuxIR circuit is auto-regulated; LuxR bound to an AHL signal activates transcription of *luxI*, and promotes accumulation of the LuxI-synthesized 3-oxo-C6 HL (Fig. 1a). In addition to LuxIR, *V. fischeri* encodes a second AHL-based quorum-sensing system, which is mediated by the *N*-octanoyl L-homoserine lactone (C8 HL) signal synthesized by AinS (Fig. 1a). The C8 HL signals through the histidine kinase AinR in a concentration-dependent manner (Gilson et al., 1995; Lupp et al., 2003; Kimbrough and Stabb, 2013). A third, non-AHL, quorum signal is expressed in *V. fischeri*: a furanosyl borate diester (AI-2) that is synthesized by LuxS, and likely signals through LuxP and LuxQ, similar to the homologous set of genes in *Vibrio harveyi* (Lupp and Ruby, 2004; Neiditch et al., 2006).

All quorum-sensing pathways in *V. fischeri* intersect at LuxR (Fig. 1a). We have previously shown that in culture, both C8 HL and AI-2 accumulation contribute to activation of *luxR* transcriptional activator LitR (Fig. 1a) (Lupp et al., 2003; Lupp and Ruby, 2004). C8 HL may also weakly bind to the non-cognate receptor LuxR, and contribute to an additional overlap between signaling systems (Fig. 1a) (Dunlap, 1999; Lupp et al., 2003). In addition to the downstream targets of LuxR regulation (Lupp and Ruby, 2005; Antunes et al., 2007), C8 HL controls an extensive set of genes, independent of LuxR (Lupp and Ruby, 2005; Antunes et al., 2007).

These convergent signal cascades culminate with the transcriptional regulation of the *luxICDABEG* operon, which encodes the light-producing luciferase enzyme complex, as well as LuxI itself. Previous studies suggest that regulation by AHL quorum sensing, mediated by AinS and LuxI, is necessary for colonization and bioluminescence of *V. fischeri* in the squid host, while the contribution of LuxS signaling is not essential for either process (Lupp and Ruby, 2004). The bioluminescence of *V. fischeri* is required to maintain a stable, and long-term partnership between host and symbiont, and possibly to signal the host (Heath-Heckman et al., 2013; Koch et al., 2013). A recently recognized role for quorum signals is as effectors of cross-kingdom communication (Rumbaugh and Kaufmann, 2012); notably, the *E. scolopes* transcriptome responds to the presence of LuxI signal 3-oxo-C6 HL (Chun et al., 2008).

Despite the centrality of quorum sensing in the ‘conversation’ between squid and vibrio, much work remains to decipher the contribution of this regulatory network and its signals to the establishment and maintenance of a stable and robust symbiosis. To bring a new perspective to studies of quorum regulation and signaling in the squid-vibrio association, we took a combined chemical and genetic approach to disentangle the signaling cascade upstream and downstream of the regulatory element LitR, and to assess the relative contributions of each branch of the cascade to colonization of *E. scolopes*. We have identified a series of synthetic, non-native AHLs that are capable of strongly modulating luminescence (presumably *via* interaction with LuxR) in laboratory cultures of *V. fischeri* ES114, a squid light-organ isolate (Geske et al., 2007a; Geske et al., 2007b; Geske et al., 2008). Amongst these compounds, the phenylacetanoyl homoserine lactone (PHL) derivatives were found to be the most potent as both agonists and antagonists of luminescence. Two PHLs of particular interest were 3-nitro PHL, an agonist that, at a comparable molarity, induced higher levels of luminescence than the native LuxR ligand 3-oxo-C6 HL, and 4-iodo PHL, an antagonist that strongly inhibited luminescence induction by 3-oxo-C6 HL (structures shown in Fig. 1b).

In the current study, we show that in *V. fischeri* (i) the effects of these two PHL derivatives are specific to targets of the signal synthesized by LuxI, and do not perturb known LuxI-independent quorum signaling, and (ii) these compounds can be used, at concentrations comparable to those observed in culture studies (Geske et al., 2007a), to manipulate bacterial activities within the squid host. Our chemical studies highlight an intriguing and novel role for LuxIR quorum sensing during the initial stages of the symbiotic ‘conversation’. Specifically, we demonstrate that the first day of colonization is a critical period in which regulation by LuxIR quorum sensing predicates the long-term stability of

the symbiotic partners. These studies underscore the value of chemical methods to probe mechanisms of quorum sensing and, to our knowledge, do so for the first time in the context of a beneficial eukaryotic-prokaryotic symbiosis.

RESULTS

Non-native analogs of 3-oxo-C6 HL do not affect AinS phenotypes

Certain native AHL autoinducers or non-native antagonists are able to signal through two AHL receptors (Schaefer et al., 1996; Swem et al., 2009; Palmer et al., 2011a; Palmer et al., 2011b). To determine whether the PHL analogs interact only with the LuxI-dependent quorum-sensing pathway, or if the molecules also modulate quorum signaling upstream of LuxI, we examined the effects of added native autoinducer and non-native PHLs on two phenotypes that are regulated by LitR and AinS, but that are independent of the LuxI branch of quorum sensing in *V. fischeri* (Fig. 1a). Using concentrations of native and non-native AHLs sufficient to modulate LuxI signaling in previous studies (Geske et al., 2007a), we examined motility on soft agar (repressed by LitR and AinS), and the metabolic shift known as the acetate switch (activated by LitR and AinS) (Figs. 1a, 2).

We first performed motility assays to determine the effects of the native AHLs and the PHL analogs on AinS and LitR-dependent signaling. As previously reported (Lupp and Ruby, 2005), cells of an *ainS* mutant spotted onto soft-agar medium swam faster, and produced significantly larger outer rings of migration, than those of their wild-type parent, consistent with AinS- and LitR- dependent repression of motility (Fig. 2a). Previous work has demonstrated that none the molecules used in the present study alter growth of *V. fischeri* (Geske et al., 2007a). As anticipated, *ainS*⁻ cells spotted onto agar containing C8 HL migrated the same distance as their similarly treated parent (Fig 1a), although neither 3-oxo-C6 nor the PHL analogs influenced migration. The addition of 3-nitro PHL slightly increased the motility of the *litR* mutant but not the *ainS* mutant relative to wild type, while 4-iodo-PHL had no effect on the migration of either mutant. The effect of 3-nitro PHL on motility was slight compared to that of C8 HL, however it is possible that this analog may also contribute to LitR-dependent signaling. Notably, the effect of 3-nitro PHL was only apparent in the absence of C8 HL, suggesting that the signaling interaction would not be biologically relevant in an environment where C8 HL accumulates, such as the squid light organ. This hypothesis is consistent with the finding that 3-oxo-C6 HL may also have the capacity to signal via the LitR-dependent pathway, although with much less sensitivity than 3-oxo-C6 HL-LuxR signaling (Kimbrough and Stabb, 2013).

The transcription of the acetyl-CoA synthetase gene, *acs*, is a measure of the alteration of the AinS-dependent induction of the acetate switch (Fig. 1a). Wild-type and *ainS*-mutant strains, each carrying an *acs'-lacZ*⁺ promoter-reporter plasmid, were grown in liquid medium supplemented with the native autoinducer or non-native analog, and assayed for *acs* expression (Fig. 2b). Consistent with previous results (Studer et al., 2008), without any additions, *ainS*⁻ *V. fischeri* had a 41% lower level of *acs* expression relative to wild-type cells. Neither the native autoinducer nor the PHL analogs significantly altered the expression of *acs* in the wild-type cultures. In contrast, the addition of C8 HL to the medium raised *acs* expression in the *ainS* mutant to 110% of wild-type levels, as expected (Studer et al., 2008).

Further, 3-oxo-C6 HL, 3-nitro PHL and 4-iodo PHL did not significantly alter the expression by *ainS*⁻ *V. fischeri*, indicating that 3-nitro PHL is unable to induce the AinSR system in the absence of C8 HL, and that 4-iodo PHL does not suppress this system in wild-type cells. Together with the motility data, these results indicate that, although 3-nitro PHL and 4-iodo PHL interact with the 3-oxo-C6 HL-regulated LuxIR system, they do not alter C8 HL-regulated phenotypes through LitR, and therefore do not target this branch of quorum sensing in *V. fischeri* in an environment with active C8 HL signaling.

Native and non-native LuxR-binding molecules can signal during symbiosis

To determine whether experimentally added autoinducer signals affect *V. fischeri* during colonization of its natural host, we first examined whether the presence of 4-iodo PHL, an antagonist of 3-oxo-C6 HL in culture (Geske et al., 2007a), was also effective in *V. fischeri* within the host. Specifically, we asked whether 4-iodo PHL could antagonize LuxIR quorum sensing in the host, and reduce the level of wild-type bioluminescence to that of mutants deficient in LuxIR signaling. Wild-type colonized animals maintained in seawater containing 50 or 100 μ M 4-iodo PHL had 20- to 40-fold lower levels of luminescence at 48 h of colonization than in seawater without added antagonist (Fig. 3a). As little as 5 μ M 4-iodo PHL was sufficient to reduce the bioluminescence of *V. fischeri* cells to the level of the *luxI* mutant, which attains approximately 5% of the bioluminescence of wild-type cells in symbiosis, while a concentration of 0.5 μ M had no significant effect on symbiotic bioluminescence (Fig. 3a). Notably, the amount of 4-iodo PHL observed to reduce bioluminescence is within the range of the concentration for 50% inhibition (IC₅₀) for this molecule obtained in culture studies (IC₅₀ = 0.9 mM) (Geske et al., 2007a), and suggests that the delivery of the analogs to the symbiont population is not greatly hindered by host tissues.

We next measured the contribution of native and non-native LuxR agonists to the induction of bioluminescence. As expected (Visick et al., 2000), in the absence of added signal, the bioluminescence of animals colonized by the *luxI* mutant was lower than that of wild-type colonized animals (Fig. 3b, c). In contrast, addition of as little as 0.5 mM of either 3-oxo-C6 HL (Fig. 3b) or 3-nitro PHL (Fig. 3c) to animals colonized by *luxI*⁻ *V. fischeri* restored induction of bioluminescence. Higher concentrations of the experimentally added signals did not further increase the bioluminescence of either wild-type *V. fischeri*, or the *luxI* mutant, and we noted no significant difference in the maximum level of bioluminescence achieved by the superactivator 3-nitro PHL when compared to the native LuxR agonist, 3-oxo-C6 (concentration for 50% activation, EC₅₀ values = 0.3 and 3.0 mM, respectively, in *V. fischeri* culture; (Geske et al., 2007a)). It is still possible that the concentration of exogenously added signal experienced by the symbionts inside the squid's light organ is lower than that present in the surrounding seawater due to diffusion limitation; however, addition of either 3-oxo-C6 HL (Fig. 3b) or 3-nitro PHL (Fig. 3c) resulted in induction of luminescence in wild-type colonized animals 1 h earlier than it occurred without the added compounds, suggesting that both compounds swiftly and effectively diffuse into the light organ, reaching concentrations that result in induction of LuxIR activity.

LuxIR signaling is required for *V. fischeri* to maintain a colonization of the light organ; thus, we also examined the effect of exogenously added autoinducers on colonization levels (*i.e.*, CFUs/squid) of LuxIR pathway mutants. Addition of 5 μ M of the LuxR antagonist 4-iodo PHL to wild type *V. fischeri* diminished the host's maintenance of a light-organ colonization to a level typical of *luxI*⁻ *V. fischeri* (Fig. 3d). However, 4-iodo PHL had no effect on the colonization of the *luxI* mutant, supporting the conclusion that the molecule suppresses colonization by wild-type cells solely through its inhibition of the LuxIR pathway. Colonization levels of the *luxI* mutant in the squid light organ were fully restored to those of wild-type *V. fischeri* when either 3-oxo-C6 HL or 3-nitro PHL were added (Fig. 3d). Additions of 5 μ M 3-oxo-C6, 3-nitro PHL, or 4-iodo PHL (Fig. 3d) had no impact on the colonization defect of the *luxR* mutant, indicating that the distinct effects of both these autoinducer analogs are dependent on a functioning LuxIR signal pathway. Taken together, these results show that exogenously added autoinducers, both the native signal and the non-native agonist, induce expected LuxIR-system responses, even when the bacteria are within host tissues.

LuxR ligands do not perturb AinS-dependent symbiosis factors

To assess the effect of exogenous autoinducers on AinS-dependent quorum sensing during squid colonization by *V. fischeri*, animals colonized by wild-type, *ainS*⁻ or *litR*⁻ strains were exposed to either the native autoinducers or the non-native analogs. In the absence of added compounds, animals colonized by the *ainS* mutant had both a lower level of luminescence (Fig. 4a) and a lower number of CFU/squid (Fig. 4b) when compared to wild-type colonized animals. Luminescence, and colonization by *litR*⁻ *V. fischeri* were indistinguishable from wild-type *V. fischeri* in the absence of added compounds (Fig. 4a & b). The addition of C8 HL, but not 3-oxo-C6 HL, restored the luminescence and CFU/squid of animals colonized by the *ainS* mutant to wild-type levels, demonstrating that the mutant's defect in symbiosis is not caused solely by a repression of signal synthesis by LuxI. The addition of the antagonist, 4-iodo PHL, decreased the luminescence and colonization levels of all strains, further indicating that the LuxIR system is active, and subject to inhibition, during colonization of squid by *ainS*-deficient symbionts.

Because there are regulatory targets of LuxIR quorum sensing other than the bioluminescence gene cluster *luxCDABEG* (Antunes et al., 2007), we reasoned that if a target of LuxIR quorum sensing apart from bioluminescence contributed to host colonization, the symbiotic population would be sensitive to the addition of LuxR antagonist 4-iodo PHL, even in the absence of the *luxCDABEG* genes. Accordingly, we observed that the level of the symbiotic population of a *luxCDABEG* mutant (Bose et al., 2008), which lacks the genes that encode the luciferase complex, was significantly lower than that of wild-type *V. fischeri* at 48 hours post-colonization (Fig. S1). While the population level of *luxCDABEG* cells was not altered by 5 μ M 4-iodo PHL, the addition of 4-iodo PHL to a wild-type colonization did not completely decrease the level of symbiotic bioluminescence to that of *luxCDABEG* (data not shown), and this small amount of residual bioluminescence correlated with a slightly larger population (Fig. S1). Together, these data suggest that bioluminescence is the key LuxIR-regulated factor that contributes to initial

light-organ colonization, and that even a low level of light production during symbiosis can contribute to the maintenance of a normal symbiont population level.

An initial, yet transitory, appearance of bioluminescence is sufficient to result in robust and persistent colonization

LuxIR induction of bioluminescence is required for *V. fischeri* to maintain colonization of the light organ, but the population levels of *luxCDABEG*, *luxI* and *luxR* mutants only begin to decrease after the first 24 h of symbiosis (Visick et al., 2000; Bose et al., 2008). Surprisingly, we found that chemically inhibiting LuxIR activation of bioluminescence by the addition of 5 μ M 4-iodo-PHL after 24 h of light-organ colonization did not significantly alter colonization of *V. fischeri* at 48 h (Fig. 5), even though addition of the antagonist was sufficient to lower symbiotic bioluminescence to levels comparable to the colonization-deficient *luxI*⁻ strain. More unexpected was the observation that, after 24 hours or colonization, even doubling the length of the symbiont's exposure to 4-iodo PHL had no significant impact on colonization of the light organ (Fig. 5). Collectively, these results suggest that induction of bioluminescence by LuxIR within the first 24 h of symbiosis is sufficient to maintain robust symbiont colonization more than a day later.

To better define the window in which LuxIR-regulated bioluminescence is critical for persistent colonization of the host, we perturbed the LuxIR circuit by addition of PHL analogs in the symbiosis after 18 h of infection. We reasoned that by this point in the symbiosis: (i) the symbiont has reached its peak population in the light organ (Ruby, 1996), and (ii) the host has initiated irreversible, symbiont-dependent, morphogenic programs (Nyholm and McFall-Ngai, 2004). We first modulated the loss or gain of bioluminescence at 18 h either by adding 4-iodo PHL to squid colonized by wild-type *V. fischeri*, or by adding 3-nitro PHL to squid colonized by *luxI*⁻ *V. fischeri* from 18-48 h post-colonization. The rapid loss of animal bioluminescence upon addition of 4-iodo-PHL at 18 h (Fig. 6a), which occurred within a few hours of exposure to the antagonist (Fig. S2), was sufficient to decrease the population of wild-type *V. fischeri* to levels comparable to *luxI* by 48 h of colonization (Fig. 6b). Additionally, the rapid gain of bioluminescence achieved by exposing squid colonized by the *luxI* mutant to 3-nitro PHL after 18 h of infection (Fig. 6a) did not lead to a significant increase in the colonization level of *luxI*⁻ *V. fischeri* at 48 h (Fig. 6b). Taken together, these data suggest that continual activation of bioluminescence by LuxR, throughout the first day of colonization, rather than the induction of bioluminescence at a specific stage during the initiation of symbiosis (Nyholm and McFall-Ngai, 2004), is required to maintain a robust population over subsequent days.

If the level of bioluminescence, rather than simply its presence, is the critical determinant of colonization stability, then we reasoned that a gradual decrease in LuxR activation of bioluminescence after 18 h should not impair colonization as severely as the complete loss of bioluminescence at that juncture. To test this hypothesis, we exposed squid colonized by wild-type *V. fischeri* to 4-iodo PHL (to diminish luminescence), or *luxI*⁻ *V. fischeri* to 3-nitro PHL (to enhance luminescence) from 0-18 h of colonization. The small change in the mean luminescence levels of the animals is consistent with the retention of some of the molecule by the animals (Fig. 6a). The slight increase of wild-type bioluminescence

observed after removal of 4-iodo-PHL at 18 h was not sufficient to support symbiotic colonization at 48 h, consistent with the defect in colonization observed in squid colonized by non-bioluminescent symbionts from 0-18 h (Fig. 6b). In contrast, the gradual decrease in bioluminescence of the *luxI* mutant following 18 h exposure to 3-nitro PHL only slightly impaired symbiont colonization at 48 h, compared to the colonization level of wild-type *V. fischeri* in which bioluminescence had been severely impaired by addition of 4-iodo PHL from 18-48 h (Fig. 6b). These results suggest that the continued maintenance of a stable symbiont population requires that the symbionts achieve a threshold level of bioluminescence during the first day of colonization.

DISCUSSION

Synthetic small-molecule probes can help illuminate biological phenomena by providing both temporal and spatial levels of control to experimental manipulations and, thereby, complement traditional biochemical and genetic approaches (O' Connor et al., 2011). These tools are particularly powerful for the study of both intercellular signaling and developmental processes, where the timing and sequence of extracellular trigger molecules play crucial roles. As highlighted above, the design and application of such chemical probes to interrogate bacterial quorum sensing has recently become a target of intense study (Rasmussen and Givskov, 2006; Amara et al., 2011; Galloway et al., 2011; Praneenarat et al., 2012). Much of the practical interest in quorum-signal manipulation comes from its potential to create a new class of anti-pathogenic agents designed, for example, to inhibit bacterial virulence (Njoroge and Sperandio, 2009; Lowery et al., 2010). However, the use of such therapies to modulate bacterial pathogens within animal tissues requires a much clearer understanding of the response of the co-occurring normal microbiota to any unintended collateral effects of the treatment on their quorum-sensing processes.

The remarkable specificity that the light-organ environment shows for *V. fischeri* makes the squid-vibrio symbiosis ideal for the study of the chemical dialog that promotes a long-term and stable association. In the current study, we report for the first time the use of non-native, synthetic AHLs (3-nitro and 4-iodo PHLs) to probe this symbiosis. As in the squid-vibrio system (Lupp and Ruby, 2005; Antunes et al., 2007), the legume-rhizobium mutualism also promotes the establishment of a stable colonization by activating genes required for the earliest stages of nodulation in the host (González and Marketon, 2003). In contrast, quorum sensing by the pathogens *Vibrio cholerae* (Liu et al., 2008) and *Pectobacterium carotovora* (Palmer et al., 2011a) represses production of the bacterium's virulence factors, presumably to evade host defenses and/or delay their deployment until the population is dense enough to sustain or transmit an infection (von Bodman et al., 2003). Thus, in both mutualism and pathogenesis the timing of signaling during host colonization is key, and addition of chemical signal analogs can help probe this temporal pattern. In addition, it is tempting to speculate that differences in the 'wiring' of quorum-sensing systems among pathogens and mutualists of the same genus, such as *V. cholerae* and *V. fischeri*, contribute in some way to the dichotomy of host responses.

Specific and sensitive chemical modulation of LuxIR quorum sensing

To interpret experiments designed to chemically manipulate quorum sensing, it is necessary for any small-molecule analogs to be specific. *V. fischeri*, like many Gram-negative microbes that encode LuxIR quorum-signaling circuits, encodes an additional AHL quorum signal, synthesized by AinS (Gilson et al., 1995; Stabb et al., 2007). Although the putative receptor, AinR, is not structurally related to LuxR, both may bind acyl-homoserine lactone signals, including 3-oxo-C6 HL, and C8 HL (Miyashiro and Ruby, 2012; Kimbrough and Stabb, 2013). Furthermore, LuxR has previously been shown to react to the AinS signal, C8 HL, and AinR to the LuxR signal 3-oxo-C6 HL, although neither receptor is as sensitive to the non-cognate signal (Fig. 1a) (Lupp et al., 2003; Kimbrough and Stabb, 2013). Despite this indication of cross-talk between the natural AHLs of *V. fischeri*, our phenotypic, transcriptional, and symbiotic measurements reported herein indicate that, in the context of functional C8 signaling, the two non-native PHLs do not influence any ainS-dependent phenotypes. In the context of the symbiosis, C8 HL signaling does not function solely as a mechanism by which to induce bioluminescence at low cell density; in fact, we noted that activation of LuxIR by the exogenous addition of a LuxR agonist was not sufficient to restore normal colonization of the host by an *ainS* mutant. This finding supports the hypothesis that LuxIR-independent targets of AinSR regulation, such as motility and acetyl-CoA synthase (Lupp and Ruby, 2005; Studer et al., 2008), are required for persistent colonization of the light organ. Perhaps the key to quorum signaling in the context of symbiosis lies in the relative abundance of each AHL signal, and the regulation surrounding these levels.

We also considered the characteristics of LuxIR-mediated quorum sensing by the PHL analogs during light-organ colonization. Both PHL molecules were able to modulate LuxIR signaling at concentrations as low as those previously reported in laboratory-culture experiments (Geske et al., 2007a). In addition to LuxR activation being highly sensitive to the PHLs, the transient addition of the analogs had a long-lived effects; that is, the symbionts of animals that had been removed from seawater containing PHL continued to respond for up to one day as though the analog were present. This observation is consistent with the light-organ crypts being a diffusion-limited environment, and suggests that the ability to experimentally reverse the action of these small chemical probes will be limited in the context of this, and perhaps other, animal models.

Interestingly, while the analogs retained their efficacy in the symbiotic environment, in the squid we did not observe that the 3-nitro PHL agonist had a higher specific activity of bioluminescence induction than the natural agonist, as had been found in culture (Geske et al., 2007b). Because the host's control of bioluminescence in the light organ is likely to rely on several physiological tissue conditions, such as oxygen availability (Miyashiro and Ruby, 2012), it is possible that these environmental factors, rather than LuxIR signaling, determine the maximum level of bioluminescence produced by *V. fischeri* cells in the light organ.

Bioluminescence leaves a persistent 'first impression'

The chemical manipulation of LuxIR quorum sensing described here offers new insight into the process by which a robust and stable mutualism is established between *V. fischeri* and *E.*

scolopes. We show that LuxIR quorum signaling enhances the transition of symbiotic *V. fischeri* from an initial colonization of the host to a robust, and long-term association, and that the presence of animal bioluminescence during the first 24 h of colonization is predictive of symbiont population dynamics over subsequent days of the symbiosis. Our results define a preliminary framework on which to map existing and future studies of microbial colonization factors and, thereby, suggest principles of host selection and maintenance of a bioluminescent symbiont during initiation of the squid-vibrio mutualism.

The contribution of AHL molecules to the chemical dialogue between host and symbiont cannot be ignored (Chun et al., 2008) (Fig. 1a); however, our data suggest that direct chemical signaling does not itself contribute to the host's control of the number of symbionts in the light organ. Rather, we propose that the host's perception of bioluminescence, or of a product of the bioluminescence reaction itself, is the signal or cue that governs maintenance of symbiont population levels. The centrality of light as a cue for maintenance of a stable symbiosis is consistent with the structural and physiological attributes of the light organ: symbiont bioluminescence is perceived by the tissues of this specialized structure (Tong et al., 2009), and these tissues also have the capacity to entrain physiological rhythms based on this stimulus (Heath-Heckman et al., 2013).

Our data suggest that, below a minimum threshold of bioluminescence, the benefit of bioluminescence to the symbiont is proportional to the amount of light produced. For example, we show that, even in the absence of LuxIR signaling, there is a small amount of light produced by the basal level of luciferase, and this activity may enhance the fitness of *luxI* and *luxR* mutants, relative to a completely 'dark' strain missing the genes encoding luciferase itself. Further, our temporal manipulation of light production during the first 24 h after symbiotic infection indicates the critical contribution of light to the persistent colonization of the light organ. However, our data also demonstrate that even a slowly diminishing bioluminescence output, if it does not fall below a minimum threshold, is sufficient for maintenance of the symbiont population by the host. Finally, we show for the first time that bioluminescence produced by *V. fischeri* within the first day of colonization appears to be a critical determinant of symbiotic population dynamics over the longer term. It will be interesting to elucidate the mechanism(s) by which this initial bioluminescent 'conversation' between host and symbiont is predictive of the long-term stability of the partnership.

CONCLUSIONS

Regulation of microbial colonization factors by quorum signaling is rarely studied within the host itself. Nevertheless, it is perhaps from this vantage that the contribution of quorum sensing to bacterial population dynamics may be best understood. In this study, we demonstrate that native and non-native HL analogs can be used to modulate LuxIR quorum sensing without perturbing either (i) the AinS-LitR quorum-signaling circuitry of *V. fischeri*, or (ii) non-LuxIR dependent factors required for symbiosis with *E. scolopes*. In addition to these observations, our chemical approach for perturbing quorum-sensing circuitry suggests a novel role for LuxIR regulation of bioluminescence not revealed by previous genetic approaches. That is, we show that the LuxIR quorum-sensing circuit, from the first hours

through the first day of colonization, is required to establish the stability of the mutualism, and that the success of this initial signaling has long-term consequences for the maintenance of the symbiont population by the host. Using longitudinal studies that are becoming feasible in the squid-vibrio model (Koch et al., 2013), future work will determine how long the impact of the first LuxIR-mediated bioluminescent ‘conversation’ is reflected in the dynamics of the symbiont population.

EXPERIMENTAL PROCEDURES

Bacterial strains and growth conditions

Strains and plasmids used in this work are listed in Table S1. *V. fischeri* strains are derivatives of ES114 (Boettcher and Ruby, 1990), and were grown at 28° C in either a seawater-tryptone medium (SWT) (Boettcher and Ruby, 1990) or Luria-Bertani salt medium (LBS) (Graf et al., 1994). *Escherichia coli* strains were grown in Luria-Bertani (LB) medium (Sambrook, 1989). Media were solidified with 1.5% agar as needed. When appropriate, antibiotics were added to LB and LBS media at the following concentrations: chloramphenicol, 5 µg/ml for *V. fischeri*, and 25 µg/ml for *E. coli*; erythromycin, 5 µg/ml for *V. fischeri*; kanamycin 100 µg/ml for *V. fischeri*. Medium reagents were purchased from ThermoFisher Scientific.

β-galactosidase assay

The β-galactosidase activity assays were performed as described previously (Studer et al., 2008). Briefly, strains were cultured in SWT, the optical density (OD) at 600 nm was measured, and β-galactosidase activity was determined after 4 h, using a microtiter-dish method modified from Slauch and Silhavy (Slauch, 1991). Cell-pellets were frozen at -20° C before resuspension, and the SDS/chloroform step was omitted. The A₄₂₀ values of the microtiter wells were read every 30 s for 1 h using a GENiosPro 96-well plate reader (TECAN, Research Triangle Park, NC). Three replicates of two dilutions were tested for each cell pellet. The relative units of β-galactosidase activity were calculated using the formula: rate [V_{max}] / ([OD] × volume [mL]).

Motility assay

The swimming behavior of *V. fischeri* was determined by growing strains to approximately 1.0 OD in SWT with 0.25% dimethylsulfoxide (DMSO) containing the appropriate signal-molecule additions. The cells were collected by centrifugation of 1 ml of the culture, the cell pellet washed with SWT, and resuspended to 0.6 OD units in SWT containing the additions. Three microliters of a suspension of each strain was spotted onto the surface of a 0.25% agar plate containing defined seawater minimal medium (MM) base (Lupp and Ruby, 2005) supplemented with 0.3% casamino acids and 0.25% DMSO containing the appropriate signal molecules. The ring diameter was determined at 16 h as a measure of the relative rate of bacterial migration.

Chemical synthesis

Procedures for synthesizing 3-oxo-C6 HL, C8 HL, 3-nitro PHL and 4-iodo PHL have been previously described (Geske et al., 2007b). All compounds were suspended as stock

solutions in DMSO, and stored sealed in the freezer when not in use. In biological experiments, DMSO was added to a final concentration of 0.25% in all conditions, including those with no added compound.

Squid colonization assays

For animal colonization experiments, newly hatched squid were placed in filter-sterilized seawater (FSW) (Instant Ocean, Aquarium Systems) containing the appropriate *V. fischeri* strain at approximately 3000 CFU/ml; a control group of animals was placed in uninoculated FSW, supplemented with 0.25% DMSO containing either the natural AHL or non-native analog, or no added compound, as indicated. All animals were transferred to fresh, uninoculated FSW at 3 h, and again at approximately 16 h, after inoculation. There was no evidence that additions of up to 100 μ M of the AHLs or analogs had any adverse effects on the animals. At 48 or 72 h post-inoculation, individual animal bioluminescence was determined using a TD 20/20 (Turner Designs) luminometer, and the animals were sacrificed to determine the CFU/squid. To monitor animal bioluminescence over time, measurements were acquired with Light a Packard Tri-Carb 2100TR scintillation counter (Packard Instruments, Meriden, CT) modified to operate as an automated photometer. In all experiments, uninoculated animals had no detectable bioluminescence or CFUs.

Statistics

Statistical analysis was performed using the GraphPad Prism software package (Version 5.0, 2010). The experimental data sets were assessed for normal distribution using the D'Agostino and Pearson omnibus test (D'agostino et al., 1990). Data that fit a normal distribution were tested for significant variation among experimental groups by one- or two-factor analysis of variance (ANOVA), with post-hoc Bonferroni T-testing. Data that failed to fit a normal distribution were assessed by the Kruskal-Wallis test with Post-Hoc Dunn's multiple comparisons to identify groups of statistically equivalent means.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors thank E. Stabb, C. Brennan, J. Campbell, J.P. Gerdt, and E. Heath-Heckman for contributive discussions, and M. E. Mattmann and P. M. Shipway for compound synthesis. This work was supported by NSF (IOS 9817232) and NIH (RR12294/OD11024) to M. McFall-Ngai and EGR, and NIH (AI063326) to MM-N, as well as grants from the Greater Milwaukee Foundation, Burroughs Welcome Foundation, and Johnson & Johnson to HEB. JAS was supported by an NSF Graduate Research Fellowship, and the Chemical Biology Training Program (UW-Madison, funded by NIH grant NIGMS T32 GM008505), and SVS was supported by fellowships from the Wisconsin Alumni Research Foundation, the Molecular Biology Training Grant (UW-Madison, funded by NIH grant T32 GM07215), and the UW-Madison Graduate School. GDG was supported by a fellowship from the American Chemical Society Division of Medicinal Chemistry. JSH was supported by the Agency for Science, Technology and Research (A*STAR), Singapore.

REFERENCES

Amara N, Krom BP, Kaufmann GF, Meijler MM. Macromolecular inhibition of quorum sensing: enzymes, antibodies, and beyond. *Chem Rev.* 2011; 111:195–208. [PubMed: 21087050]

- Antunes LC, Schaefer AL, Ferreira RB, Qin N, Stevens AM, Ruby EG, Greenberg EP. Transcriptome analysis of the *Vibrio fischeri* LuxR-LuxI regulon. *J Bacteriol.* 2007; 189:8387–8391. [PubMed: 17827287]
- Bjarnsholt T, Givskov M. Quorum-sensing blockade as a strategy for enhancing host defences against bacterial pathogens. *Phil Trans R Soc B.* 2007; 362:1213–1222. [PubMed: 17360273]
- Boettcher KJ, Ruby EG. Depressed light emission by symbiotic *Vibrio fischeri* of the sepiolid squid *Euprymna scolopes*. *J Bacteriol.* 1990; 172:3701–3706. [PubMed: 2163384]
- Bose JL, Rosenberg CS, Stabb EV. Effects of luxCDABEG induction in *Vibrio fischeri*: enhancement of symbiotic colonization and conditional attenuation of growth in culture. *Arch Microbiol.* 2008; 190:169–183. [PubMed: 18521572]
- Chun CK, Troll JV, Koroleva I, Brown B, Manzella L, Snir E, et al. Effects of colonization, luminescence, and autoinducer on host transcription during development of the squid-vibrio association. *Proc Natl Acad Sci U S A.* 2008; 105:11323–11328. [PubMed: 18682555]
- Churchill ME, Chen L. Structural basis of acyl-homoserine lactone- dependent signaling. *Chem Rev.* 2010; 111:68–85. [PubMed: 21125993]
- D'agostino RB, Belanger A, D'Agostino RB Jr. A suggestion for using powerful and informative tests of normality. *Am Stat.* 1990; 44:316–321.
- Dunlap PV. Quorum regulation of luminescence in *Vibrio fischeri*. *J Mol Microbiol Biotechnol.* 1999; 1:5–12. [PubMed: 10941779]
- Galloway WR, Hodgkinson JT, Bowden S, Welch M, Spring DR. Applications of small molecule activators and inhibitors of quorum sensing in Gram- negative bacteria. *Trends Microbiol.* 2012; 20:449–458. [PubMed: 22771187]
- Galloway WRJD, Hodgkinson JT, Bowden SD, Welch M, Spring DR. Quorum sensing in Gram-negative bacteria: small-molecule modulation of AHL and AI-2 quorum sensing pathways. *Chem Rev.* 2011; 111:28–67. [PubMed: 21182299]
- Geske GD, O'Neill JC, Blackwell HE. *N*-phenylacetanoyl-L- homoserine lactones can strongly antagonize or superagonize quorum sensing in *Vibrio fischeri*. *ACS Chem Biol.* 2007a; 2:315–319. [PubMed: 17480049]
- Geske GD, O'Neill JC, Miller DM, Mattmann ME, Blackwell HE. Modulation of bacterial quorum sensing with synthetic ligands: systematic evaluation of *N*-acylated homoserine lactones in multiple species and new insights into their mechanisms of action. *J Am Chem Soc.* 2007b; 129:13613–13625. [PubMed: 17927181]
- Geske GD, O'Neill JC, Miller DM, Wezeman RJ, Mattmann ME, Lin Q, Blackwell HE. Comparative analyses of *N*-acylated homoserine lactones reveal unique structural features that dictate their ability to activate or inhibit quorum sensing. *Chembiochem.* 2008; 9:389–400. [PubMed: 18224645]
- Gilson L, Kuo A, Dunlap PV. AinS and a new family of autoinducer synthesis proteins. *J Bacteriol.* 1995; 177:6946–6951. [PubMed: 7592489]
- González JE, Marketon MM. Quorum sensing in nitrogen-fixing rhizobia. *Microbiol Mol Biol Rev.* 2003; 67:574–592. [PubMed: 14665677]
- González JF, Venturi V. A novel widespread interkingdom signaling circuit. *Trends Plant Sci.* 2012; 18:167–174. [PubMed: 23089307]
- Graf J, Dunlap PV, Ruby EG. Effect of transposon-induced motility mutations on colonization of the host light organ by *Vibrio fischeri*. *J Bacteriol.* 1994; 176:6986–6991. [PubMed: 7961462]
- Heath-Heckman EA, Peyer SM, Whistler CA, Apicella MA, Goldman WE, McFall-Ngai MJ. Bacterial bioluminescence regulates expression of a host cryptochrome gene in the squid-vibrio symbiosis. *mBio.* 2013; 4 doi:pii: e00167-13.
- Hentzer M, Wu H, Andersen JB, Riedel K, Rasmussen TB, Bagge N, et al. Attenuation of *Pseudomonas aeruginosa* virulence by quorum sensing inhibitors. *EMBO J.* 2003; 22:3803–3815. [PubMed: 12881415]
- Kimbrough JH, Stabb EV. Substrate specificity and function of the pheromone receptor AinR in *Vibrio fischeri* ES114. *J Bacteriol.* 2013 (Epub ahead of print).
- Koch EJ, Miyashiro T, McFall-Ngai MJ, Ruby EG. Features governing symbiont persistence in the squid-vibrio association. *Mol Ecol.* 2013 (doi:10.1111/mec.12474).

- Liu Z, Miyashiro T, Tsou A, Hsiao A, Goulian M, Zhu J. Mucosal penetration primes *Vibrio cholerae* for host colonization by repressing quorum sensing. *Sci Signal*. 2008; 105:9769.
- Lowery CA, Salzedo NT, Sawada D, Kaufmann GF, Janda KD. Medicinal chemistry as a conduit for the modulation of quorum sensing. *J Med Chem*. 2010; 53:7467–7489. [PubMed: 20669927]
- Lupp C, Ruby EG. *Vibrio fischeri* LuxS and AinS: comparative study of two signal synthases. *J Bacteriol*. 2004; 186:3873–3881. [PubMed: 15175301]
- Lupp C, Ruby EG. *Vibrio fischeri* uses two quorum-sensing systems for the regulation of early and late colonization factors. *J Bacteriol*. 2005; 187:3620–3629. [PubMed: 15901683]
- Lupp C, Urbanowski M, Greenberg EP, Ruby EG. The *Vibrio fischeri* quorum-sensing systems *ain* and *lux* sequentially induce luminescence gene expression and are important for persistence in the squid host. *Mol Microbiol*. 2003; 50:319–331. [PubMed: 14507383]
- Mandel MJ. Models and approaches to dissect host-symbiont specificity. *Trends Microbiol*. 2010; 18:504–511. [PubMed: 20729086]
- Miyashiro T, Ruby EG. Shedding light on bioluminescence regulation in *Vibrio fischeri*. *Mol Microbiol*. 2012; 84:795–806. [PubMed: 22500943]
- Neiditch MB, Federle MJ, Pompeani AJ, Kelly RC, Swem DL, Jeffrey PD, et al. Ligand-induced asymmetry in histidine sensor kinase complex regulates quorum sensing. *Cell*. 2006; 126:1095–1108. [PubMed: 16990134]
- Njoroge J, Sperandio V. Jamming bacterial communication: new approaches for the treatment of infectious diseases. *EMBO Mol Med*. 2009; 1:201–210. [PubMed: 20049722]
- Nyholm SV, McFall-Ngai MJ. The winnowing: establishing the squid- *vibrio* symbiosis. *Nat Rev Microbiol*. 2004; 2:632–642. [PubMed: 15263898]
- O' Connor CJ, Laraia L, Spring DR. Chemical genetics. *Chem Soc Rev*. 2011; 40:4332–4345. [PubMed: 21562678]
- Palmer AG, Streng E, Blackwell HE. Attenuation of virulence in pathogenic bacteria using synthetic quorum-sensing modulators under native conditions on plant hosts. *ACS Chem Biol*. 2011a; 6:1348–1356. [PubMed: 21932837]
- Palmer AG, Streng E, Jewell KA, Blackwell HE. Quorum sensing in bacterial species that use degenerate autoinducers can be tuned by using structurally identical non-native ligands. *Chembiochem*. 2011b; 12:138–147. [PubMed: 21154995]
- Parsek MR, Greenberg EP. Acyl-homoserine lactone quorum sensing in gram-negative bacteria: a signaling mechanism involved in associations with higher organisms. *Proc Natl Acad Sci U S A*. 2000; 97:8789–8793. [PubMed: 10922036]
- Praneenarat T, Palmer AG, Blackwell HE. Chemical methods to interrogate bacterial quorum sensing pathways. *Org Biomol Chem*. 2012; 10:8189–8199. [PubMed: 22948815]
- Rasmussen TB, Givskov M. Quorum sensing inhibitors: a bargain of effects. *Microbiology*. 2006; 152:895–904. [PubMed: 16549654]
- Ruby EG. Lessons from a cooperative, bacterial-animal association: the *Vibrio fischeri*-*Euprymna scolopes* light organ symbiosis. *Annu Rev Microbiol*. 1996; 50:591–624. [PubMed: 8905092]
- Rumbaugh KP, Kaufmann GF. Exploitation of host signaling pathways by microbial quorum sensing signals. *Curr Opin Microbiol*. 2012; 15:162–168.
- Sambrook, JF.; E.F.; Maniatis, T. *Molecular Cloning*. Cold Spring Harbor. Cold Spring Harbor Press; New York: 1989.
- Schaefer AL, Hanzelka BL, Eberhard A, Greenberg EP. Quorum sensing in *Vibrio fischeri*: probing autoinducer-LuxR interactions with autoinducer analogs. *J Bacteriol*. 1996; 178:2897–2901. [PubMed: 8631679]
- Slauch JM, Silhavy TJ. cis-acting ompF mutations that result in OmpR- dependent constitutive expression. *J Bacteriol*. 1991; 173:4039–4048. [PubMed: 1648075]
- Stabb, EV.; Visick, KL. *Vibrio fischeri*: A bioluminescent light-organ symbiont of the bobtail squid *Euprymna scolopes*. In: Rosenberg, E.; DeLong, EF.; Lory, S.; Stackebrandt, E.; Thompson, F., editors. *The Prokaryotes*. Springer-Verlag; Berlin: 2013. p. 497-532.

- Stabb, EV.; Schaefer, A.; Bose, JL.; Ruby, EG. Quorum signaling and symbiosis in the marine luminous bacterium *Vibrio fischeri*.. In: Winans, S.; Bassler, B., editors. Cell-cell signaling in bacteria. ASM Press; Washington, DC: 2007. p. 233-250.
- Studer SV, Mandel MJ, Ruby EG. AinS quorum sensing regulates the *Vibrio fischeri* acetate switch. J Bacteriol. 2008; 190:5915–5923. [PubMed: 18487321]
- Swem LR, Swem DL, O'Loughlin CT, Gatmaitan R, Zhao B, Ulrich SM, Bassler BL. A quorum-sensing antagonist targets both membrane-bound and cytoplasmic receptors and controls bacterial pathogenicity. Mol Cell. 2009; 35:143–153. [PubMed: 19647512]
- Teplitski M, Mathesius U, Rumbaugh KP. Perception and degradation of N-acyl homoserine lactone quorum sensing signals by mammalian and plant cells. Chem Rev. 2011; 111:100–116. [PubMed: 20536120]
- Tong D, Rozas NS, Oakley TH, Mitchell J, Colley NJ, McFall-Ngai MJ. Evidence for light perception in a bioluminescent organ. Proc Natl Acad Sci U S A. 2009; 106:9836–9841. [PubMed: 19509343]
- Visick KL, Foster J, Doino J, McFall-Ngai M, Ruby EG. *Vibrio fischeri* lux genes play an important role in colonization and development of the host light organ. J Bacteriol. 2000; 182:4578–4586. [PubMed: 10913092]
- von Bodman SB, Bauer WD, Coplin DL. Quorum sensing in plant- pathogenic bacteria. Annu Rev Phytopathol. 2003; 41:455–482. [PubMed: 12730390]
- Wu H, Song Z, Hentzer M, Andersen JB, Molin S, Givskov M, Høiby N. Synthetic furanones inhibit quorum-sensing and enhance bacterial clearance in *Pseudomonas aeruginosa* lung infection in mice. J Antimicrob Chemother. 2004; 53:1054–1061. [PubMed: 15117922]

Figure 1.

The core AHL-dependent pathways of quorum signaling in *V. fischeri*. **(a)** AinS synthesizes C8 HL, which is released into the environment and, when at a sufficiently high concentration, diffuses back into the cell. Subsequent interaction of C8 HL with LuxR may potentiate a weak, but potentially priming, induction of the *lux* operon (*luxICDABEG*). When a threshold concentration of C8 HL is reached, C8 HL is thought to bind its cognate receptor AinR, initiating a multi-step signaling cascade leading to an increase in LitR translation (Kimbrough and Stabb, 2013). LitR further activates transcription of *luxR*,

increases the amount of LuxR. When it binds 3-oxo C6, LuxR strongly activates transcription of the *luxICDABEG* operon. Activation of *luxI* transcription increases the synthesis of 3-oxo C6, and amplifies induction of the *lux* operon, leading to an exponential increase (autoinduction) in the synthesis of the luciferase complex and light production. 3-nitro PHL and 4-iodo PHL are structural analogs of the HL family of quorum-sensing signals, and specifically enhance or depress LuxR function, respectively. The presence of native AHL molecules, C8 HL and 3-oxo C6 HL have been shown to also alter host gene expression. **(b)** Structures of the natural autoinducers (1 & 2) and non-native autoinducer analogs (3 & 4) used in this study: (1) octanoyl homoserine lactone; (2) 3-oxo-hexanoyl homoserine lactone; (3) 3-nitrophenyl homoserine lactone; and, (4) 4-iodophenyl homoserine lactone.

Figure 2.

AinS-dependent transcriptional and phenotypic regulation is unaffected by non-native PHL derivatives. (a) The migration of the indicated mutant strains of *ainS*⁻ and *litR*⁻ *V. fischeri* was measured, relative to migration of a wild-type control, after 16 h on motility agar supplemented with either a solvent control (DMSO), or 120 nM of the indicated HL or PHL analog. Error bars represent S.E.M. of nine replicate plates. 'a', 'b' and 'c' indicate distinct treatment groups with statistically equivalent means; Two-way ANOVA (interaction $F(4,80) = 10.1$, $P < 0.001$; HL addition $F(4,80) = 20.8$, $P < 0.0001$; strain $F(1,80) = 4.0$, $P = 0.049$),

with Post-hoc Bonferroni T-tests. **(b)** The level of *acs* expression in wild-type and *ainS*⁻ strains of *V. fischeri* containing the *acs*'-*lacZ*⁺ reporter plasmid were determined after 4 h of growth in SWT supplemented with 40 nM of the indicated AHL or PHL analog. Error bars represent S.E.M. of 4 biological replicates. Significance was determined by Two-way ANOVA (interaction $F(4,20) = 7.2$, $P = 0.0009$; strain $F(1,20) = 28.7$, $P < 0.0001$; addition $F(4,20) = 5.3$, $P = 0.0045$), with Post-hoc Bonferroni comparison testing: 'a' and 'b' indicate distinct treatment groups with statistically equivalent means.

Figure 3.

Contribution of native and non-native HL analogs to symbiotic colonization and bioluminescence at 48 h post-inoculation. (a) Bioluminescence of wild-type (open boxes) was compared to *luxI*⁻ *V. fischeri* (diagonal boxes) in the presence of different concentrations of 4-iodo PHL. Significant differences in mean bioluminescence were determined by the Kruskal-Wallis test ($H = 36.2$, 10 d.f., $P < 0.0001$), with Post-Hoc Dunn's multiple comparisons ('a' and 'b' indicate distinct groups with equivalent means; $P < 0.05$). Inner fences determined by Tukey's method, dashed line indicates background level of

luminescence. Induction of bioluminescence of wild type (closed symbols), or *luxI* (open symbols) was monitored in the presence of **(b)** no added HL (circles), 0.5 μ M C6 HL (triangles), or 5 μ M C6 HL (squares), or **(c)** no added HL (circles), 0.5 μ M 3-nitro PHL (triangles), or 5 μ M 3-nitro PHL (squares). Bars represent S.E.M. of $n=5$; dashed line indicates level of detectable luminescence. **(d)** Animal colonization at 48 h was compared between inoculations with wild-type (open boxes), *luxI*⁻ (diagonal boxes), or *luxR*⁻ (dotted boxes) *V. fischeri* in the presence of 5 μ M added HL, as indicated. Significant differences in mean colony-forming units (CFU) per animal were determined by the Kruskal-Wallis test ($H = 172.4$, 15 d.f., $P < 0.0001$), with Post-Hoc Dunn's multiple comparisons ('a' and 'b' indicate distinct groups with equivalent means; $P < 0.05$). Inner fences determined by Tukey's method; dots represent outliers.

Figure 4.

AinS⁻ and *LitR*⁻ dependent colonization dynamics are not perturbed by analogs of 3-oxo C6 homoserine lactone. Squid colonized by wild-type (white), *ainS*⁻ (diagonal) or *litR*⁻ (dotted) *V. fischeri* were treated with either a DMSO solvent control (No addition), or 5 μ M additions of C8 HL, 3-oxo-C6 HL, 3-nitro PHL or 4-iodo PHL as indicated. (a) Differences in bioluminescence among treatments and strains at 48 h were assessed by the Kruskal-Wallis test ($H = 78.8$, 15 d.f., $P < 0.0001$), with Post-Hoc Dunn's multiple comparisons ('a' and 'b' indicate distinct groups with equivalent means, $P < 0.05$); dashed line indicates background

level of luminescence. **(b)** Differences in colonization levels at 48 h were assessed by the Kruskal-Wallis test ($H = 149.7$, 15 d.f., $P < 0.0001$) with Post-Hoc Dunn's multiple comparisons ('a' and 'b' indicate distinct groups with equivalent means, $P < 0.05$). Inner fences determined by the Tukey method; black dots indicate outliers; CFU, colony-forming units.

Figure 5.

Squid colonized with wild-type *V. fischeri* for 24 h were either left untreated (open boxes), or treated with 4-iodo PHL (diagonal boxes) for an additional 24 h (end-point = 48 h), or 48 h (end-point = 72 h) prior to enumeration of colony-forming units (CFU). Populations of squid were also colonized by *luxI*⁻ in the presence of a DMSO solvent control (dotted boxes) as a comparison for 4-iodo-PHL inhibitory activity. The mean CFU value was compared to that of the control wild-type populations by Kruskal-Wallis test ($H=47.9$, 6 d.f., $P<0.0001$), with Post-hoc Dunn's multiple comparisons ('a' and 'b' indicate distinct groups with equivalent means, $P<0.05$).

Figure 6.

Induction of bioluminescence by LuxIR quorum sensing is required for establishment of a robust symbiont population. The bioluminescence of an established symbiont population was either inhibited or induced by exogenous exposure to 5 μ M of a HL-analog for a period of time (exposure indicated as a range of hours post-colonization: 0-18 h or 18-48 h; No ad. = no analog addition). (a) Animal bioluminescence was measured at 18 h (open boxes), and 36 h (diagonal boxes) post-inoculation for each treatment group. Differences in mean luminescence (compared to wild type without additions) were determined by the Kruskal-

Wallis test ($H = 147.3$, 12 d.f., $P < 0.0001$) with Post-Hoc Dunn's multiple comparisons ('a' and 'b' indicate distinct groups with equivalent means), dashed line indicates threshold of bioluminescence detection. **(b)** Mean CFU at 48 h post-inoculation was compared to the wild type without additions by the Kruskal-Wallis test ($H = 41.3$, 6 d.f., $P < 0.0001$), with Post-Hoc Dunn's multiple comparisons ('a', 'b' and 'c' indicate distinct groups with equivalent means, $P < 0.05$). Inner fences determined by the Tukey method; black dots indicate outliers; CFU, colony-forming units.