CORRESPONDENCE

The Management of Thyroid Nodules: A Retrospective Analysis of Health Insurance Data

by Romy Wienhold, Prof. Dr. rer. nat. Markus Scholz, Jürgen-Bernhard Adler, Christian Günster, Prof Dr. med. Ralf Paschke in volume 49/2013

Too Much Diagnostic Evaluation and Therapy

Wienhold and colleagues' key message in the article (1) on the management of thyroid nodules in Germany—that scintigraphy and surgery are used too often in Germany, whereas fine-needle aspiration cytology (FNAC) is used too rarely—is true and correct.

It is also correct that the upper billing limit for sonography and the low reimbursement for FNAC are contributing factors in this setting. The assumption that the indication for surgery is often defined by using scintigraphy is (in many cases) reality in Germany. As long as thyroid problems indicate scintigraphy almost as a reflex, and as long as the scintigraphic finding of a “cold nodule” triggers the association of “suspected malignancy” in doctors as well as patients, this kind of over-treatment is not going to change. Attractive reimbursement for thyroid nodules, irritating scintigraphic findings, and the “German Angst” ascribed to us by general opinion worldwide, provoke unnecessary surgical procedures.

In conclusion: since most thyroid disorders and functional impairments can be diagnosed conclusively by the patient's medical history, a small number of laboratory parameters, qualified color Doppler ultrasound, and, if required, elastography and FNAC, this will have to become the diagnostic standard (2). Revised guidelines should clearly state that scintigraphy (which entails all the risks associated with radiation) should not be used as the primary investigative option, and not for diagnosing malignancy, in hyperthyroidism, Graves' disease, and in order to determine autonomy in nodules with a volume (measured by ultrasound) of <1 mL and avascular nodules of <2 mL. Applying all these standards will result in scintigraphy rates below 10%, as was recently shown in an occupational thyroid screening conducted by myself, which included some 1000 participants.

The authors' comments regarding calcitonin measurement in non-surgically treated patients—not performed as often as it should—are problematic (1). For reasons of space, I refer readers to an excellent critical overview (3). Calcitonin is highly sensitive for the purpose of detecting C cell carcinomas (how many were detected in the 25 600 patients insured with the statutory sickness funds?), but its specificity is irritatingly low (2, 3). Pre-analytic laboratory problems, many false-positive findings, and cost (one calcitonin measurement accounts for about half the quarterly budget for one patient) may explain doctors' reticence in this context. Rather than demanding improved dissemination of calcitonin screening, the guideline should, conversely, be adapted to the reality of clinical practice and the development of sonography.

DOI: 10.3238/arztebl.2014.0287a

REFERENCES

1. Wienhold R, Scholz M, Adler JB, Günster C, Paschke R: The management of thyroid nodules—a retrospective analysis of health insurance data. *Dtsch Arztebl Int* 2013; 110(49): 827–34.
2. Braun B. Schilddrüse und Nebenschilddrüsen. In: Braun B, Günther R, Schwerk WB (eds.): *Ultraschalldiagnostik – Lehrbuch und Atlas*. Heidelberg, München, Landsberg: ecomed MEDIZIN 2010; III-3.1: 1–238.
3. Ross DS, Cooper DS, Mulder JE: Diagnostic approach to and treatment of thyroid nodules. Serum calcitonin concentration. www.uptodate.com/store (Last accessed on 25 September 2013).

Prof. Dr. med. Bernd Braun
Reutlingen, Prof.B.Braun@gmx.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Benefit of Follow-up not Confirmed

I disagree with the key message that sonography is used too rarely to monitor thyroid nodules not managed surgically (1). As the authors correctly state, about 20% of the population have at least one thyroid nodule which most are not aware of and live well without control examinations. Attempting to detect and control all nodules would overstretch the healthcare system.

The positive predictive value of thyroid ultrasound for malignancy is low (8–20%) given the extremely low pretest probability (2). This is certainly one reason for the number of diagnostic surgical procedures, which were criticized as excessive in the article. There is no evidence for the benefit of follow-up examinations in nodules that were initially assessed as non-malignant. We do not know which follow-up intervals are appropriate and when to stop screening stop altogether.

Every year, about 5500 persons in Germany develop thyroid cancer, which has an excellent prognosis for the most common subtypes (3). Morphological thyroid investigations should be restricted to symptomatic patients or patients with impaired thyroid function, until evidence from high quality studies with long term follow-up is available to show that asymptomatic nodules require medical attention. Most of the 16 000 000 persons with asymptomatic thyroid nodules will be grateful if a problem that is not one does not turn them into chronically ill patients, and if the limited resources are used for more relevant health problems.

DOI: 10.3238/arztebl.2014.0287b

REFERENCES

1. Wienhold R, Scholz M, Adler JB, Günster C, Paschke R: The management of thyroid nodules—a retrospective analysis of health insurance data. *Dtsch Arztebl Int* 2013; 110(49): 827–34.
2. Bonavita JA, Mayo J, Babb J, Bennett G, Oweity T, Macari M, Yee J: Pattern recognition of benign nodules at ultrasound of the thyroid: which nodules can be left alone? *AJR Am J Roentgenol* 2009; 193: 207–13.
3. Robert Koch Institut. Gesundheitsberichterstattung des Bundes. Krebs in Deutschland 2009/2010, 9th edition. 2013. www.krebsdaten.de/Krebs/DE/Content/Publikationen/Krebs_in_Deutschland/krebs_in_deutschland_node.html (last accessed on 5.3.2014)

Prof. Dr. med. Jean-François Chenot, MPH
Abteilung Allgemeinmedizin, Institut für Community
Universitätsmedizin Greifswald, jchenot@uni-greifswald.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Inconsistencies

On the basis of erroneous and incomplete data, the authors (1) suggest algorithms for the diagnostic evaluation of the thyroid, which cannot be applied in their current form—at least not in Germany, (formerly) a region with iodine deficiency.

The specificity of scintigraphy was given as 5%; this is not correct—the three citations are not conclusive: a 20-year-old study from Ireland and two review articles without concrete data.

Furthermore, the article contains some technical inconsistencies.

Firstly, the cited European guideline was introduced in 2006 and the others even later, which means they had not been published at the time of data collection. The procedural recommendations of the German Society of Nuclear Medicine of 2003 (2), which were coordinated with the German Society of Endocrinology and the German Society for Surgery and deposited with the Association of the Scientific Medical Societies in Germany (Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften, AWMF) were not cited.

Secondly, the authors combined their data with a figure from the AACE/AME/ETA guideline, which was published only in 2010, two years after the collection of data.

Thirdly, how would it have been common knowledge that 94% of patients who underwent scintigraphy also had a low concentration of thyroid stimulating hormone (TSH) or multinodular goiter? Scintigraphy in (euthyroid) uninodular goiter was “done in accordance with the guideline.” According to the valid German guideline (2), indications for thyroid scintigraphy are amongst others:

- Palpable and/or sonographically distinct focal findings (nodules of 1 cm or larger).
- Suspected focal/diffuse autonomy in manifest/latent hyperthyroidism.
- After definitive therapy in order to document therapeutic success.
- If required during the course of untreated functional autonomy.

The depicted algorithm can therefore not be applied to the situation in Germany. A single nodule ≥ 1 cm would directly undergo fine-needle aspiration cytology (FNAC) without scintigraphic investigation. For a hyperfunctional and therefore benign nodule, the cytopathological finding often is “follicular neoplasia,” and the patient would probably have unnecessary surgery because scintigraphy was not done. A recent German study found no decreased TSH concentration in 70% of all cases of focal autonomy (3).

Lastly, the cited article (Fast et al.) on thyroid hormone administration in the treatment of goiter studied the long abandoned concept of TSH suppressive administration of levothyroxine; but a recent, methodologically superior German study remains unmentioned (4).

DOI: 10.3238/arztebl.2014.0288a

REFERENCES

1. Wienhold R, Scholz M, Adler JB, Günster C, Paschke R. The management of thyroid nodules—a retrospective analysis of health insurance data. *Dtsch Arztebl Int* 2013; 110(49): 827–34.
2. Dietlein M, Dressler J, Eschner W, Leisner B, Reiners C, Schicha H; Deutsche Gesellschaft für Nuklearmedizin; Deutsche Gesellschaft für Medizinische Physik: Leitlinie zur Schilddrüsendiagnostik (Version 2). *Nuklearmedizin* 2003; 42: 120–2.
3. Graf D, Helmich-Kapp B, Graf S, Veit F, Lehmann N, Mann K: Funktionelle Aktivität fokaler Schilddrüsenautonomien in Deutschland. *Dtsch Med Wochenschr* 2012; 137: 2089–92.

4. Grussendorf M, Reiners C, Paschke R, Wegscheider K; LISA Investigators: Reduction of thyroid nodule volume by levothyroxine and iodine alone and in combination: a randomized, placebo-controlled trial. *J Clin Endocrinol Metab* 2011; 96: 2786–95.

On behalf of the working group for thyroid diseases of the German Society of Nuclear Medicine:

Dr. med. Michael C. Kreißl

Klinik für Nuklearmedizin, Klinikum Augsburg/Klinik für Nuklearmedizin, Universitätsklinikum Würzburg
Michael.Kreissl@klinikum-augsburg.de

Prof. Dr. med. Dr. rer. nat. Andreas Bockisch

Klinik für Nuklearmedizin, Universitätsklinikum Essen

Prof. Dr. med. Markus Dietlein

Klinik für Nuklearmedizin, Universitätsklinikum Köln

Prof. Dr. med. Frank Grünwald

Klinik für Nuklearmedizin, Universitätsklinikum Frankfurt am Main

Prof. Dr. med. Markus Luster

Klinik für Nuklearmedizin, Universitätsklinikum Gießen und Marburg GmbH, Marburg

Conflict of interest statement

Prof. Grünwald has received speaker honoraria from Sanofi/Henning, Merck, Thermofischer, GE. The other authors declare that no conflict of interest exists.

Better Diagnostics and Treatment

In contrast to what is shown in Figure 2 (1), thyroid scintigraphy is relevant in nodules larger than 1 cm when the TSH concentration is normal. If a hot nodule is present or the scintigram shows an autonomous nodule, then malignancy can be excluded with a probability bordering on certainty (2). In such cases, fine-needle aspiration cytology (FNAC) is not required. As multicenter study showed that most autonomous nodules are accompanied by a normal TSH concentration, although a focal raised concentration is shown by the scintigram (3). Before performing FNAC in a sonographically suspect finding, technetium-99m pertechnetate scintigraphy should be undertaken.

Unfortunately, a negligible proportion of thyroid cancers present as sonographically normal. Especially multinodular goiter (where not all nodules can be aspirated) therefore leaves us all a desire for a procedure with a high predictive value. FNAC is important, but its usefulness in excluding malignancy is limited (10–20% false-negatives). Technetium-99m-MIBI tumor scintigraphy of the thyroid can rule out malignancy with a probability of 97% if the findings are negative (4).

On the basis of the data analyzed by Wienhold et al., the unsatisfactory preoperative risk stratification of thyroid nodules can be assumed, if at all, only for uninodular goiter (it is not obvious from the coding whether local symptoms/mechanical relevance is present). And in uninodular goiter too, a significant displacement of the trachea or swallowing difficulties can be reason enough for performing surgical resection. Furthermore it should not remain unmentioned that in the article, out of 9000 patients with uninodular goiter, only 4.5% had surgery. A large number of diagnostic thyroid operations can therefore not be deduced from the data presented.

DOI: 10.3238/arztebl.2014.0288b

REFERENCES

1. Wienhold R, Scholz M, Adler JB, Günster C, Paschke R: The management of thyroid nodules—a retrospective analysis of health insurance data. *Dtsch Arztebl Int* 2013; 110(49): 827–34.
2. Gharib H, Papini E, Paschke R, et al.: American Association of Clinical Endocrinologists, AME and European Thyroid Association medical guidelines for clinical

practice for the diagnosis and management of thyroid nodules. *J Endocrinol Invest* 2010; 33: 287–91.

3. Görges R, Kandror T, Kuschner S, et al.: Scintigraphically hot thyroid nodules mainly go hand in hand with a normal TSH. *Nuklearmedizin* 2011; 50: 179–88.
4. Wale A, Miles KA, Young B, et al.: Combined 99mTc-methoxyisobutylisonitrile scintigraphy and fine-needle aspiration cytology offers an accurate and potentially cost-effective investigative strategy for the assessment of solitary or dominant thyroid nodules. *Eur J Nucl Med Mol Imaging* 2014; 41: 105–15.

Prof. Dr. med. Holger Palmedo

Kompetenznetz Schilddrüse Bonn Rhein-Sieg,
Überörtliche Gemeinschaftspraxis für Radiologie und Nuklearmedizin, PET-CT-Zentrum
Johanniter Krankenhaus Bonn
holger.palmedo@gmx.de

Prof. Dr. med. Andreas Türler

Evangelische Kliniken Bonn gGmbH, Johanniter-Krankenhaus; Kompetenzzentrum für
Schilddrüsen- & Nebenschilddrüsenchirurgie (DGAV), Bonn

Conflict of interest statement

The author declares that no conflict of interest exists.

Discrepancy Between Clinical Reality and Guidelines

The authors deserve thanks for pointing out the substantial discrepancy between clinical reality in Germany and older US guidelines (1). I have some additional comments.

If reality and guidelines differ that widely, then it is worth checking the guidelines as well as the clinical reality.

Recent evidence for therapeutic options was not included. The LISA study (2) was published in 2011—after the publication of the two US guidelines (2009 and 2011) (3, 4). This study reported a significant reduction in the size of thyroid nodules for L-thyroxine alone (7.3%) versus L-thyroxine plus iodide (17.3%) compared with placebo. (However, whether these effects are relevant or cost effective has not been the subject of discussion thus far.)

The authors assume that a 1 cm diameter of an echo-less nodule is a useful intervention threshold—but this is still under discussion.

The value of fine-needle aspiration cytology is also still under discussion, in view of the very low mortality due to thyroid cancer in an assumed incidence of micro-cancers in up to 35% of the population. This raises the question of the cost-benefit-effectiveness of an approach similar to screening—as in prostate cancer.

The authors furthermore adopt from the cited guidelines without any discussion the recommendation to perform scintigraphy in multinodular goiter for the purposes of further differentiation. In view of the fact that almost all nodules are cold, and that a malignancy may also be concealed by a hot nodule, this seems rather ineffective. In view of the radiation load, an analysis of cost:benefit and benefit:harm is urgently indicated.

Postoperative follow-up examinations, which the authors lamented as being undertaken too rarely, should also be studied regarding their cost:benefit ratio. Postoperative hypocalcemia and hyperthyroidism or hypothyroidism certainly require follow-up and treatment, whereas recurrent asymptomatic increase in size after reduction surgery rather does not.

In conclusion: the time has come for evidence based German recommendations. These should evaluate critically especially the indications for scintigraphy, fine-needle aspiration cytology, and surgery.

DOI: 10.3238/arztebl.2014.0289a

REFERENCES

1. Wienhold R, Scholz M, Adler JB, Günster C, Paschke R: The management of thyroid nodules—a retrospective analysis of health insurance data. *Dtsch Arztebl Int* 2013; 110(49): 827–34.
2. Grussendorf M, Reiners C, Paschke R, Wegscheider K; LISA Investigators. J Clin Endocrinol Metab. Reduction of thyroid nodule volume by levothyroxine and iodine alone and in combination: a randomized, placebo-controlled trial. 2011; 96: 2786–95.
3. Gharib H, Papini E, Paschke R, et al.: American Association of Clinical Endocrinologists, Italian Association of Clinical Endocrinologists and European Thyroid Association: Medical guidelines for clinical practice for the diagnosis and management of thyroid Nodules. *Endocr Pract, Hot Thyroidology, J Endocrin Invest* 2010; 33 Suppl 5: 1–56.
4. Cooper DS, Doherty GM, Haugen BR, Kloos, et al.: Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid* 2009; 19: 1167–214.

Dr. med. Uwe Popert

Abteilung Allgemeinmedizin der Universität Göttingen
Sprecher der DEGAM-Sektion Versorgung
uwe.popert@t-online.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We thank Prof Braun for his involved and succinct confirmation of our key messages and agree with him that further discussion of the guideline recommendation regarding calcitonin screening should be continued elsewhere. Prof. Chenot rightly points out the scarce evidence described in the AACE/AME/ETA guideline (1) with regard to follow-up examinations. However, a misunderstanding concerns the facts that the guidelines do not recommend thyroid ultrasonography as a screening examination and that for a patient with thyroid nodules a (low) risk for a false-negative risk stratification is well known, whereas this obviously does not apply to the general population.

With regard to the letter by Kreissl and colleagues, we wish to point out that it was not us but the AACE/AME/ETA guideline that suggested an algorithm (Figure 2). The evidence levels of the publications available for guideline recommendations are known to be different. For this reason, the current AACE/AME/ETA S3 guideline classifies the evidence levels of the consulted publications and classifies the strength of each recommendation according to the evidence—in contrast to, for example, the 10-year-old consensus statement from the German Society of Nuclear Medicine (2), which did not do this.

As no further literature for the sensitivity of scintigraphy was mentioned, we may continue to assume 5% (with a low level of evidence for the respective publications).

We wish to respond to the four further issues raised by Kreissl and coauthors as follows:

First, in order to clarify that initial statements had already been composed a long time before our data collection period, we cited the German Society of Nuclear Medicine's consensus statement from 1999 and not the, similar, 2003 version (2).

Second, we thought it made more sense to compare the results of our study with the current algorithm from the AACE/AME/ETA guideline than to generate outdated algorithms from an old consensus statement for the purposes of comparison.

Third, the claim that the algorithm suggested in the AACE/AME/ETA S3 guideline (3) (Figure 2), cannot be applied to the

situation in Germany can not be substantiated by a 10 year old, unrevised—and therefore according to AMWF rules invalid—consensus statement with 7 references (2). The AACE/AME/ETA guideline (1) is a good example of how the (internationally) published evidence can be interpreted by several scientific societies jointly and under consideration of regionally different epidemiologies. According to Figure 2 and the AACE/AME/ETA guideline (1), an individual nodule >1 cm would not be fine-needle aspirated directly but only after risk stratification using clinical presentation, TSH, calcitonin, and sonographic criteria for malignancy. By applying this strategy, up to 53% of nodules can be spared from FNAC by sonography alone, as shown in Figure 2 (3).

Fourth, because of the publication date, the combination treatment with iodide and thyroxine can only be evaluated in the current guideline update. The discussion of the relevant study was therefore not the topic of our article (4).

Since euthyroid autonomy is common, especially in multinodular goiter (5), the AACE/AME/ETA guideline (1) and our Figure 2, which was taken from the guideline, recommends scintigraphy for all cases of multinodular goiter, among other reasons in order to rule out autonomy from fine-needle aspiration cytology (FNAC), and, so far, for uninodular goiter depending on TSH and clinical presentation. The articles by Görges et al. and Graf et al.—cited by Kreissl and colleagues as well as by Prof. Palmedo and Prof. Türler—only state for 57 of 514 (42% multinodular) or for none of the patients whether a single nodule or several nodules were present.

Evaluations of FNA results do not mention hot nodules as a relevant explanation for follicular proliferations. Furthermore, only 5–10% (4) of thyroid nodules are hot nodules. The use of scintigraphy to reduce the frequency of follicular proliferations in cytology as suggested by Kreissl et al. has therefore not been mentioned in the guidelines.

The false-negative rate of FNA, as mentioned by Prof. Palmedo and Prof. Türler, is reported extremely divergently in the different publications; the guideline reports it to be 1–2% (1). This means that for FNA, as for any investigator-dependent method, quality and quality control of the cytologist and further measures by the aspirator (1) are crucial to reduce false-negative results. The guidelines do not mention technetium-99m-MIBI scintigraphy. We can conclude from our data that, while paying attention to the limitations explained in our article (for example,

a lack of clinical symptoms), that in a scenario of guideline conform diagnostic evaluation, diagnostic surgery could have been avoided in most of the patients with operated uninodular thyroids. Very similar results have in the meantime also been reported from Belgium (6).

Most of the topics raised by Popert overlap with those explained above. We did not refer to older US guidelines but to a current and international guideline. Our study objective was to compare healthcare reality with guideline recommendations, and not to discuss guideline recommendations. If reality and guidelines diverge then health services research is required in order to change the evidence base for guideline revisions. Not almost all nodules are cold, but about 45%, and about as many nodules are scintigraphically normal or poorly characterized (3,7).

DOI: 10.3238/arztebl.2014.0289b

REFERENCES

1. Gharib H, Papini E, Paschke R, Duick DS, Valcavi R, Hegedus L et al.: American Association of Clinical Endocrinologists, Italian Association of Clinical Endocrinologists and European Thyroid Association Medical Guidelines For Clinical Practice For the Diagnosis and Management of Thyroid Nodules. *Endocrine Practice* 2010; 16: Suppl 1: 1–43.
2. Dietlein M, Dressler J, Eschner W, Leisner B, Reinert C, Schicha H; Deutsche Gesellschaft für Nuklearmedizin; Deutsche Gesellschaft für Medizinische Physik: Leitlinie zur Schilddrüsendiagnostik (Version 2). *Nuklearmedizin* 2003; 42: 120–2.
3. Paschke R: Schilddrüsenknoten: Gutartig oder bösartig? *MMW* 2013; 155: 50–4.
4. Wienhold R, Scholz M, Adler JB, Günster C, Paschke R: The management of thyroid nodules—a retrospective analysis of health insurance data. *Dtsch Arztebl Int* 2013; 110(49): 827–34.
5. Berghout A, Wiersinga WM, Smits NJ, Touber JL: Interrelationships between age, thyroid volume, thyroid nodularity, and thyroid function in patients with sporadic nontoxic goiter *Am J Med* 1990; 89: 602–8.
6. Van den Bruel A, Francart J, Dubois C, Adam M, Vlayen J, De Schutter H, Stordeur S, Dellaconne B: Regional variation in thyroid cancer incidence in Belgium is associated with variation in thyroid imaging and thyroid disease management. *J Clin Endocrinol Metab* 2013; 98: 4063–71.
7. Knudsen N, Bülow I, Jørgensen T, Laurberg P, Ovesen L, Perrild H: Comparative study of thyroid function and types of thyroid dysfunction in two areas in Denmark with slightly different iodine status. *Eur J Endocrinol*. 2000; 143: 485–91.

Prof. Dr. med. Ralf Paschke
Klinik für Endokrinologie und Nephrologie
Universität Leipzig
Ralf.Paschke@medizin.uni-leipzig.de

Conflict of interest statement

The author declares that no conflict of interest exists.