Published in final edited form as:

Environ Pollut. 2014 February ; 185: 365–368. doi:10.1016/j.envpol.2013.10.018.

Temporal and spatial variation in residential soil metal concentrations: implications for exposure assessments

C. Marjorie Aelion^{1,2,*}, Harley T. Davis^{2,3}, Andrew B. Lawson⁴, Bo Cai³, and Suzanne McDermott⁵

¹School of Public Health and Health Sciences, University of Massachusetts Amherst, 715 No. Pleasant Street, Amherst, MA 01003

²Department of Environmental Health Sciences, University of South Carolina, 921 Assembly Street, Columbia, SC 29208

³Department of Epidemiology and Biostatistics, University of South Carolina, 800 Sumter Street, Columbia, SC 29208

⁴Division of Biostatistics and Epidemiology, Medical University of South Carolina, 135 Cannon Street, Suite 303, Charleston, SC 29425

⁵Department of Family and Preventive Medicine, University of South Carolina, 3209 Colonial Drive, Columbia, SC 29203

Abstract

Understanding temporal and spatial variation in soil chemicals is critical in exposure assessments. We measured eight metals in subsamples, duplicates (~0.3 m), and repeat soil samples taken 1–6 years after initial sampling (~5 m). We estimated variance components (VCs) of metal concentrations using nested analyses accounting for sampling area, land use and soil type, and calculated coefficients of variation (CVs) for repeat sample pairs. Total variance for all metals was similar, but VCs were proportioned differently by metal and sample type. Spatial variation explained the majority of variance in duplicate samples. Metal concentrations CVs were not significantly different over the long time interval, but repeat samples had larger VCs for unexplained errors. Sampling area and land use were important for Ba and Mn, and Pb and Hg, respectively. Results suggest metals are stable over long times suitable for exposure assessments, but that individual metal behavior should be considered.

Keywords

Neurotoxic metals; spatial scale; Pb, temporal variation; exposure assessments

© 2013 Elsevier Ltd. All rights reserved.

*Corresponding author: (413) 545-2525 (office); (413) 545-0501 (fax); maelion@schoolph.umass.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

Human exposure assessments rely on analytically measured concentration of contaminants from specific sample locations (Paustenbach, 2000), and understanding the spatial and temporal variation of soil chemical concentrations can provide greater confidence when using these data in exposure assessments. Chronic exposure to low concentrations of neurotoxic metals and metal mixtures in soils is a current concern, and evidence suggests fetal exposures to these metals during development may impact birth outcomes (Grandjean and Landrigan, 2006). In previous studies, we measured soil metal concentrations at several residential locations in the Southeastern United States (US), and examined their potential associations with intellectual disabilities in children (Kim et al., 2009; Kim et al., 2010; Liu et al., 2010; McDermott et al., 2011; Zhen et al., 2008; Zhen et al., 2009), and their associations with social and economic factors of the study population (Aelion et al., 2012, 2013). Metal concentrations were spatially interpolated at maternal residences, but no validation was carried out to determine spatial and temporal variation of soil metal concentrations.

Identification of spatial and temporal variation in soil metal concentrations is important. Small-scale spatial variation in soil characteristics may be important for chemical distribution. Soil organic carbon affects chemical degradation (Smalling and Aelion, 2004) and clay and sand content impacts chemical concentrations over distances of centimeters (Aelion, 1996; Einax and Kraft, 2002). Wilcke (2000) reported that the bioavailability of metals in soils can vary greatly over scales of $< 1 \text{ m}^2$. Temporal variation in soil metal concentrations is important because environmental data and health data may be collected at disparate times and assumptions are made about temporal changes in metal concentrations.

The objectives of the current study were to examine small-scale spatial variation and large-scale temporal variation in concentrations of eight soil metals. We calculated differences in metal concentrations for subsamples (multiple samples from the same grab sample) and duplicate samples (independent grab samples collected $< 0.3 \text{ m}$ apart at the same time) at 11 residential sampling sites spread over a large geographic area, and repeat samples (grab samples collected within 5 m or less of the original sample 1–6 years after the initial sampling event) at three of the 11 sites, and estimated variance components for each metal. We further examined temporal variation by comparing coefficient of variations (CV) among repeat sample pairs for the three repeat areas individually.

We hypothesize that the spatial and temporal variation will be small for all metals due to the small spatial scale, no significant anthropogenic sources of metals in these areas, and limited biotransformation of metals. Despite limited variation, we hypothesize that error variance components will be lowest in subsamples compared to duplicate and repeat samples due to lack of spatial and temporal components in subsamples. We also hypothesize that the spatial variation of metals will be greater than the temporal variation because soils are inherently geologically variable, and metals are relatively chemically stable.

Methods

From 2006–2011, grab topsoil samples (upper 5 cm; ~50 g) were collected from 11 individual residential sampling areas throughout an area in the Southeastern US approximately 320 km wide and 420 km long. Ten of the sampling areas ranged from 60–120 km², and one area was 490 km². Sampling areas included five urban, five rural and one mixed urban plus rural location. Samples were collected on a uniform 120-node grid at each sampling area as close to the nodes as possible. Distance between sampling nodes ranged from 0.8–1 km for the 10 smaller areas, and 4 km for the one large area (Area 1). Approximately 120 samples were collected at each site, with 60 samples collected at Area 1. Figure 1 shows the entire sampling area (not to scale) with Area 4 grid nodes identified.

Two duplicate grab samples were collected at approximately 10% of sampling locations selected randomly within a sampling area, in close proximity (within 0.3 m). A total of 57 duplicate soil sample pairs (114 total samples) were analyzed from the 11 sampling areas. In addition, a subset of samples was homogenized using a tumbling method (Schumacher et al., 1991) and two ~15 g subsamples were taken from the one original grab soil sample. In total, 37 subsample pairs (74 total samples) were analyzed from the 11 sampling areas. The analytical lab was blinded to the status of both duplicate and subsample soils.

All soils were acid digested and analyzed for arsenic (As), barium (Ba), chromium (Cr), copper (Cu), lead (Pb), manganese (Mn), and nickel (Ni) using inductively coupled plasma atomic emission spectroscopy (ICP-AES) using EPA Method 6010 and for total mercury (Hg) with the cold vapor method using EPA Method 7471 (Aelion and Davis, 2007; Aelion et al., 2008, 2009a, 2009b; Davis et al., 2009). All results were reported in mg/kg, and reported values below the minimum detection limit (MDL) were set to ½ the MDL for that metal for all subsequent data analyses.

In 2012, soil sample locations from three of the original sampling areas were randomly chosen for collection of repeat samples (59 sample pairs, or 118 total samples). The temporal differences in repeat samples were 1.5 (Area 22), 4.7 (Area 4) and 5.6 (Area 1) years after the original sample collection and analysis. The original latitude and longitude coordinates were entered into a GPS device (Garmin eTrex, Garmin, Olathe, KS, USA) in order to sample as close as possible to the original sampling location, recognizing the GPS device accuracy was ~5 m. Most repeat sample spatial ranges were smaller than 5 m because site photographs and sampling field notes containing information on vegetation cover and physical landmarks helped to increase the certainty of sampling at a spatial scale similar to that of the duplicate samples. Samples were collected and analyzed in the same manner as previously described.

A nested procedure was implemented in SAS version 9.4 software (SAS Institute, Cary, NC). The absolute values of differences between subsample, duplicate, and repeat soil pair metal concentrations were calculated and these and metal concentrations were log transformed due to positive skewness. Each metal was modeled as a separate dependent variable nested within sampling area, Anderson land use type (Anderson et al., 1976), and soil type (USDA, 2013). This type of procedure allows for an unbalanced sampling design,

and estimates the variance components for errors and other model covariates to soil metal concentration. The coefficient of variation (CV) for repeat soil pairs was calculated for each metal for each area, and compared using the non-parametric Kruskal-Wallis test.

Results and Discussion

In general total variance of subsample, duplicate, and repeat samples was similar (Table 1A–C). This suggests that the spatial (in repeat and duplicate sample pairs) and temporal (in repeat samples only) components contributed a similar amount of variance as the analytical +mixing component. Although total variance of the different samples was similar, there were differences in allocation of the proportion of total variance for different samples and for individual metals.

The subsampling error (analytical+mixing) variance component explained a small percentage of total variance with the exception of As and Hg (Table 1A), for which subsampling error explained 68% and 61% of the total variance, respectively. In contrast, duplicate error variance components explained >50% of the variance for five of eight metals (As, Cr, Cu, Pb, and Ni; Table 1B), suggesting that the small spatial scale component explained the greatest proportion of the total variance in these metal concentrations. Rawlins et al. (2009) compared the sampling variances from duplicate samples 20 m apart and subsamples for soils in the UK. They observed that the duplicate sample error variance components were an order of magnitude greater than subsample error variance components for the three metals they investigated, Cu, Ni and zinc (Zn). Our subsample error variance component for Ni was an order of magnitude less than the duplicate error variance component; however no consistent pattern was observed for most metals. Einax and Kraft (2002) examined sampling variance of As, Cr, Cu, Pb, Mn and Ni collected at 25-cm intervals in top soil in Germany, measured at similar ranges and concentrations as our study. Similar to our findings, they found that the sampling uncertainty, due to the large range of metal concentrations, was higher than the uncertainty of sample treatment and analytical measurements.

In contrast to our hypothesis, repeat error variance components explained a smaller proportion of the total variance than either subsample or duplicate error variance components, suggesting the spatial+temporal component intrinsic in our repeat error did not sufficiently explain the variation in metal concentrations. The repeat sample models had high proportions of unexplained variance (Table 1C), particularly for Hg (72%) and As (61%). This was not observed in the subsample and duplicate sample models. We attribute some of the unexplained variance to the fact that Hg was analyzed using a different analytical method than the other metals. This was corroborated by the subsample error variance component for Hg, which was 61% of the total variance (Table 1C), suggesting that the analytical+mixing error accounted for the majority of variance in Hg concentrations. In addition, the number of repeat samples below the MDL was higher for As (12) and Hg (14); all the other metals except Ni had no samples below the MDL. In a validation study of kriged estimated metal concentration estimates, Aelion et al. (2009a) concluded that kriged concentrations for Hg were greater than measured values because Hg had more samples below the MDL than other metals measured.

Sampling area, land use and soil type explained differing amounts of variance depending on the individual metal. The sampling area variance component explained a substantial amount of the variance for all metals in all models, except Hg in the repeat model. This was particularly consistent for Ba, Mn and Ni (Tables 1A–C). In past research at these sites Ba and Mn consistently clustered together based on principal component analysis (PCA) followed by cluster analysis (Davis et al., 2009; Aelion et al. 2012), and were identified as naturally occurring (Aelion et al., 2008, 2009b). Soil type did not explain variance to a great extent, except in the subsamples, and again primarily for Ba, Mn and Ni suggesting similar to sampling area, soil type was more important for naturally-occurring metals than for anthropogenic metals.

Like soil type, land use did not explain a large proportion of the variance in metal concentrations for most models, but did consistently for Pb and Hg. Soil type variance component estimates ranged from 3–29% for Pb and 7–15% for Hg (Tables 1A–C). In contrast to sampling area and soil type, the variance components for land use appeared to be important for anthropogenic metals regardless of sample type. In our previous PCA and cluster analysis, Hg and Pb consistently clustered together (Davis et al., 2009; Aelion et al. 2012). In addition, Pb concentrations were associated with percentage of road coverage at the US Census 2000 block group level (Aelion et al., 2012), which also is indicative of urban/built up land use.

We further investigated temporal variation by comparing CVs for repeat sample pairs separately by area (Area 1, 5.6 yr; Area 4, 4.7 yr; Area 22, 1.5 yr temporal sampling interval) (Table 2). Only Ni between Areas 1 and 22 was significantly different among mean repeat sample pair CVs. Therefore temporal variability in repeat sample metal concentrations was not evident, as repeat sample CVs were similar regardless of the time interval between sampling periods.

There are limitations to this study due to sample collection. It is not possible to exactly site the repeat sample to the original sample location, and collecting duplicate and repeat samples at the exact same spatial scale would facilitate comparisons between those samples. Although we cannot verify spatial differences between duplicate and repeat samples, photographs increased the accuracy of the relocation, and we are confident that distances between repeat samples were better than the 5 m accuracy of the GPS. Repeat samples were collected in only a subset of sampling areas and over time intervals that were not evenly spaced. Thus we cannot confirm that the findings would be consistent for all our sampling areas, or if evenly spaced and a greater number of temporal intervals had been chosen. Also, the numbers of subsample and duplicate samples were not the same for all sampling locations; while a nested analysis is acceptable for this unbalanced sample set, a balanced design would be more desirable.

Regardless of the limitations of this study, variances were small for all metals regardless of sample type, suggesting that neither the small spatial nor large temporal scales in our study contributed greatly to the variance estimates in metal concentrations. The variance contribution to metal concentrations from spatial variation was greater in the duplicate samples than for other factors in these models. Comparison of CVs indicated no differences

for the long time intervals examined but repeat samples had substantial unexplained error which should be further investigated. Sampling area explained a large proportion of variance in naturally occurring metals like Ba and Mn and land use explained some of the variance in the anthropogenic metals Pb and Hg. Metal data collected on these small spatial and long temporal scales appear to be appropriate for human exposure studies, however differences in variances and effects from other model covariates for individual metals should be considered.

Acknowledgments

Funding provided by the National Institute of Environmental Health Sciences (NIEHS), National Institutes of Health (NIH) (2R01ES012895-04A2). We thank B. Bey, J. Davis, M. Engle, S. Jayasinghe, and F. Nemeth for their help with sample collection.

References

- Aelion CM. Impact of aquifer sediment grain size on petroleum hydrocarbon distribution and biodegradation. *J. Contam. Hydrol.* 1996; 22:109–121.
- Aelion CM, Davis HT. Use of a general toxicity test to predict heavy metal concentrations in residential soils. *Chemosphere.* 2007; 67:1043–1049. [PubMed: 17140621]
- Aelion CM, Davis HT, McDermott S, Lawson AB. Metal concentrations in rural topsoil in South Carolina: potential for human health impact. *Sci. Total Environ.* 2008; 402:149–156. [PubMed: 18538375]
- Aelion CM, Davis HT, Liu Y, Lawson AB, McDermott S. Validation of Bayesian kriging of arsenic, chromium, lead and mercury in surface soils based on internode sampling. *Environ. Sci. and Technol.* 2009a; 43:4432–4438. [PubMed: 19603658]
- Aelion CM, Davis HT, McDermott S, Lawson AB. Soil metal concentrations and toxicity: associations with distances to industrial facilities and implications for human health. *Sci. Total Environ.* 2009b; 407:2216–2223. [PubMed: 19155049]
- Aelion CM, Davis HT, Lawson AB, Cai B, McDermott S. Associations of estimated residential soil arsenic and lead concentrations and community-level environmental measures with mother-child health conditions in South Carolina. *Health Place.* 2012; 18:774–781. [PubMed: 22579118]
- Aelion CM, Davis HT, Lawson AB, Cai B, McDermott S. Associations between soil lead concentrations and populations by race/ethnicity and income-to-poverty ratio in urban and rural areas. *Environ. Geochem. Health.* 2013; 35:1–12. [PubMed: 22752852]
- Anderson JR, Hardy EE, Roach JT, Witmer RE. A land use and land cover classification system for use with remote sensor data. *Geological Survey Professional Paper.* 1976:964.
- Davis HT, Aelion CM, McDermott S, Lawson AB. Identifying natural and anthropogenic sources of metals in urban and rural soils using GIS-based data, PCA, and spatial interpolation. *Environ. Pollut.* 2009; 157:2378–2386. [PubMed: 19361902]
- Einax JW, Kraft J. Small-scale variability of metals in soil and composite sampling. *Environ. Sci. Pollut. Res.* 2002; 9:257–261.
- Grandjean P, Landrigan PJ. Developmental neurotoxicity of industrial chemicals. *Lancet.* 2006; 368:2167–2178. [PubMed: 17174709]
- Kim J-I, Lawson AB, McDermott S, Aelion CM. Variable selection for spatial random field predictors under a Bayesian mixed hierarchical spatial model. *Spat. and Spatio temporal Epidemiol.* 2009; 1:95–102.
- Kim J-I, Lawson A, McDermott S, Aelion CM. Bayesian spatial modeling of disease risk in relation to multivariate environmental risk fields. *Stat. Med.* 2010; 29:142–157. [PubMed: 19904772]
- Liu Y, McDermott S, Lawson AB, Aelion CM. The relationship between mental retardation and developmental delays in children and the levels of arsenic, mercury, and lead in soil samples taken near their mother's residence during pregnancy. *Int. J. Hyg. Environ. Health.* 2010; 213:116–123. [PubMed: 20045663]

- McDermott S, Wu J, Cai B, Lawson AB, Aelion CM. Probability of intellectual disability is associated with soil concentrations of arsenic and lead. *Chemosphere*. 2011; 84:31–38. [PubMed: 21450328]
- Paustenbach DJ. The practice of exposure assessment: a state-of-the-art review. *J. Toxicol. Environ. Health, Part B*. 2000; 3:179–291.
- Rawlins BG, Scheib AJ, Lark RM, Lister TR. Sampling and analytical plus subsampling variance components for five soil indicators observed at regional scale. *Eur. J. Soil Sci.* 2009; 60:740–747.
- SAS Version 9.3 Software. Copyright 2012 of SAS Institute Inc. Cary, NC, USA:
- Schumacher, BA.; Shines, KC.; Burton, JV.; Papp, ML. A comparison of soil sample homogenization techniques. In: Simmons, MS., editor. *Hazardous Waste Measurements*. Boca Raton, FL: Lewis Publishers, Inc.; 1991. p. 53-68.
- Smalling K, Aelion CM. Distribution of atrazine into three chemical fractions: Impact of sediment depth and organic carbon content. *Environ. Toxicol. Chem.* 2004; 23:1164–1171. [PubMed: 15180367]
- United States Department of Agriculture (USDA). State Soil Geographic (STATSGO) data base for South Carolina. 2013. Available at: <http://artsandsciences.sc.edu/gis/dataindex.html>
- Wilcke W. Small-scale variability of metal concentrations in soil leachates. *Soil Sci. Soc. Am. J.* 2000; 64:138–143.
- Zhen H, Lawson AB, McDermott S, Pande Lamichhane A, Aelion CM. A spatial analysis of mental retardation of unknown cause and maternal residence during pregnancy. *Geospatial Health*. 2008; 2:173–182. [PubMed: 18686266]
- Zhen H, McDermott S, Lawson AB, Aelion CM. Are clusters of mental retardation correlated with clusters of developmental delay? *Geospatial Health*. 2009; 4:17–26. [PubMed: 19908187]

Figure 1.

Relative sizes of sampling areas (distances between areas not to scale) and close up of grid nodes for tentative sampling locations in Area 22.

Table 1

A–C. Variance component estimates (proportion of total variance) for error measurements, model covariates, and unexplained error for subsample error (A), duplicate error (B), and repeat error (C) by metal from nested analyses.

A Metal	Sampling Area	Land Use	Soil Type	Subsample Error	Unexplained Error	Total Variance
As	0.08 (23)	0	0	0.23 (68)	0.03 (9)	0.33
Ba	0.12 (50)	0	0.08 (33)	0.02 (6)	0.03 (11)	0.24
Cr	0.13 (68)	0.01 (7)	0	0.03 (18)	0.01 (7)	0.19
Cu	0.15 (39)	0.08 (19)	0	0.14 (37)	0.02 (5)	0.39
Pb	0.12 (35)	0.009 (3)	0.10 (29)	0.08 (23)	0.03 (10)	0.36
Mn	0.2 (38)	0	0.19 (36)	0.12 (23)	0.01 (3)	0.52
Ni	0.14 (59)	0	0.06 (25)	0.01 (5)	0.03 (11)	0.24
Hg	0.03 (17)	0.03 (15)	0	0.11 (61)	0.01 (7)	0.19

B Metal	Sampling Area	Land Use	Soil Type	Duplicate Error	Unexplained Error	Total Variance
	0.04 (19)	0	0	0.14 (64)	0.04 (17)	0.21
Ba	0.08 (42)	0	0	0.09 (46)	0.03 (12)	0.2
Cr	0.07 (32)	0	0	0.13 (56)	0.03 (12)	0.22
Cu	0.07 (29)	0	0	0.14 (53)	0.04 (18)	0.25
Pb	0.04 (14)	0.04 (15)	0	0.18 (61)	0.03 (10)	0.29
Mn	0.12 (50)	0.005 (2)	0.01 (5)	0.08 (32)	0.03 (11)	0.24
Ni	0.07 (25)	0.02 (9)	0	0.16 (56)	0.03 (10)	0.28
Hg	0.02 (19)	0.007 (7)	0.007 (7)	0.02 (14)	0.06 (53)	0.11

C Metal	Sampling Area	Land Use	Soil Type	Repeat Error	Unexplained Error	Total Variance
As	0.04 (18)	0	0.02 (11)	0.02 (9)	0.12 (61)	0.20
Ba	0.08 (44)	0.02 (9)	0	0.02 (10)	0.07 (37)	0.18
Cr	0.12 (42)	0	0	0.08 (28)	0.09 (30)	0.29
Cu	0.12 (36)	0.05 (16)	0.03 (9)	0.05 (15)	0.08 (24)	0.33
Pb	0.07 (25)	0.09 (29)	0	0.05 (17)	0.09 (29)	0.3

C Metal	Sampling Area	Land Use	Soil Type	Repeat Error	Unexplained Error	Total Variance
Mn	0.13 (44)	0	0	0.03 (11)	0.14 (45)	0.3
Ni	0.14 (46)	0.03 (10)	0.02 (11)	0.02 (7)	0.12 (36)	0.32
Hg	0	0.08 (14)	0.09 (14)	0	0.43 (72)	0.59

Table 2

Mean (standard deviation) of coefficient of variation (CV) percentages (%) by metal for repeat (by sampling area) soil pair metal concentrations.

	Repeat (n=59 pairs)		
	Area 1	Area 4	Area 22
As	59.2 (39.0)	46.6 (32.9)	55.2 (37.9)
Ba	46.0 (19.6)	50.7 (34.1)	31.0 (29.3)
Cr	52.4 (32.6)	49.2 (27.1)	45.3 (35.9)
Cu	53.2 (29.6)	43.6 (21.4)	49.6 (29.9)
Pb	54.9 (39.9)	40.7 (36.9)	30.6 (23.5)
Mn	71.7 (42.0)	48.2 (33.1)	46.2 (36.5)
Ni	68.4 (37.1)	52.1 (39.8)	32.8 (24.2)
Hg	67.5 (39.5)	74.3 (42.3)	53.6 (38.9)