Published in final edited form as:

Mol Cell Endocrinol. 2014 April 25; 387(0): 35–43. doi:10.1016/j.mce.2014.02.010.

Development of Adrenal Cortical Zonation and Expression of Key Elements of Adrenal Androgen Production in the Chimpanzee (*Pan troglodytes*) from Birth to Adulthood

C. R. Parker Jr^a, W. E. Grizzle^b, J. K. Blevins^c, and K. Hawkes^c

C. R. Parker: crparker@uab.edu; W. E. Grizzle: wgrizzle@uab.edu; K. Hawkes: hawkes@anthro.utah.edu

^aDepartment of Obstetrics and Gynecology, University of Alabama at Birmingham, Birmingham, AL, 35294

^bDepartment of Pathology, University of Alabama at Birmingham, Birmingham, AL, 35294

^cDepartment of Anthropology, 270S 140E., University of Utah, Salt Lake City, Utah

Abstract

The basis for the pattern of adrenal androgen production in the chimpanzee, which resembles that of humans, is poorly defined. We characterized the developmental zonation and expression of elements of the androgen biosynthetic pathway in the chimpanzee adrenal. The newborn adrenal contained a broad fetal zone (FZ) expressing CYP17, SULT2A1, and Cytochrome b5 (CB5) but not HSD3B; the outer cortex expressed HSD3B but not SULT2A1 or CB5. During infancy, the FZ involuted and the HSD3B-expressing outer cortex broadened. By 3 years of age, a thin layer of cells that expressed CB5, SULT2A1, and CYP17 adjoined the medulla and likely represented the zona reticularis; the outer cortex consisted of distinct zonae fasciculata and glomerulosa. Thereafter, the zona reticularis broadened as also occurs in the human. The adult chimpanzee adrenal displayed other human-like characteristics: intramedullary clusters of reticularis-like cells and also a cortical cuff of zona fasciculata-like cells adjoining the central vein.

Keywords

Adrenal Gland; Adrenal Androgens; Chimpanzee; Adrenal Zonation; Adrenarche; Adrenal Development

© 2014 Elsevier Ireland Ltd. All rights reserved.

Corresponding author: C.R. Parker, Jr., Department of Obstetrics and Gynecology, University of Alabama at Birmingham, Birmingham, AL, 35294. crparker@uab.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

1.1

During human fetal development, a specialized zone of the adrenal cortex, known as the fetal zone, comprises about 80% of the mass of the adrenal and produces the large quantities of DHEA and DHEA-sulfate that serve as the principal precursors for placental estrogen formation. The fetal zone of the human adrenal contains large quantities of the enzyme DHEA-sulfotransferase (SULT2A1) and little if any 3 β -hydroxysteroid dehydrogenase (HSD3B2) that could direct the adrenal steroid pathway toward delta-4 steroids such as aldosterone and cortisol rather than DHEA/DHEA-sulfate (Barker et al, 1995; Dupont et al, 1990; Mesiano et al, 1993; Parker et al, 1994, 1995). In fact, there appears to be little HSD3B2 in the fetal adrenal until the latter stages of gestation where it is primarily localized to the outer cortex known as the neocortex (Dupont et al, 1990; Mesiano et al, 1993; Parker et al, 1995). The enzyme 17 α -hydroxylase/17,20-lyase (CYP17), which can catalyze both the 17 α -hydroxylation of pregnenolone and progesterone as well as conversion of such 17-hydroxylated steroids to DHEA and androstenedione, respectively, is characteristic of the fetal and adult human adrenal (Mesiano et al, 1993; Dharia et al, 2005). Cytochrome B5 (CB5), which acts as an accessory protein to CYP17 and promotes its 17,20-lyase activity (Katagiri et al, 1995; Lee-Robichaud et al, 1995), is co-localized with CYP17 in both the fetal zone of the fetal adrenal and the zona reticularis (Dharia et al, 2005), which develops during adrenarche and persists throughout adulthood, in the human adrenal (Dhom, 1973).

1.2

Although adrenal androgen production during fetal development and/or during adult life is not common among other mammals, many primate species have been found to bear similarities in the potential for adrenal androgen synthesis to that of the human. Those species that have been found to possess a fetal zone like that of the human also appear to rely on the fetal adrenal to a variable degree to supply precursors for placental estrogen formation during pregnancy (Albrecht et al, 1980; Lanman, 1961; Walsh et al, 1979a, 1979b). Interestingly, however, there is limited information about the morphologic and functional zonation of the fetal and postnatal adrenal of our genetically closest primate relative, the chimpanzee (*Pan troglodytes*).

1.3

The available data are suggestive that in the chimpanzee, the fetal adrenal is composed of a morphologically distinct fetal zone and a neocortex (Czekela et al, 1983) and estrogen production during pregnancy follows a similar time course to that of humans (Czekela et al, 1983; Reyes et al, 1975, Smith et al, 1999). However, we are not aware of any studies demonstrating the capacity for androgen production by the fetal adrenal or a role for the fetal adrenal in estrogen formation in chimpanzee pregnancy. There are developmental increases in circulating levels of DHEA and DHEA sulfate during adolescence that resemble that seen in human adrenarche (Bernstein et al, 2012; Collins et al, 1981, Copeland et al, 1985; Smail et al, 1982), and there is maintenance in serum levels of DHEA and DHEA sulfate during young adulthood that are within the general range of circulating concentrations noted for humans (Bernstein et al, 2012). Since it is known that DHEA and DHEA sulfate are

produced in castrated chimpanzees and are responsive to ACTH (Albertson et al, 1984), these steroids must arise from the adrenal gland. Despite such findings, there are no data concerning the functional phenotype of the adrenal cortical zones of the fetal or postnatal chimpanzee. Based on these limited findings, we sought to better characterize the steroidogenic potential of the adrenal of the fetal, infant, adolescent, and adult chimpanzee by evaluating the zonation and expression of several key elements of the steroidogenic pathway.

2. Methods

2.1

Archival adrenal samples from 27 chimpanzees (13 male, 14 female) ranging in age from term newborn to 31 years of age were kindly provided to us by the Yerkes National Primate Research Center and the Southwest National Primate Research Center. At the time of routine autopsy of the chimpanzees, tissues, including the adrenal glands, were removed and were fixed in 10% neutral buffered formalin. Tissues were processed using standard techniques and embedded in paraffin. For our study, five micron sections were mounted on Plus slides (Corning) and were either stained with hematoxylin and eosin for general morphologic characterization or were utilized for immunohistochemistry using standard techniques in our laboratory.

2.2

The anti-human SULT2A1 antiserum was developed in the rabbit against purified human liver DHEA sulfotransferase by Dr. Charles Falany (Department of Pharmacology, University of Alabama at Birmingham). The anti-human CB5 antiserum was developed in the rabbit by Dr. Alan J Conley (School of Veterinary Medicine, Univ. California at Davis), utilizing recombinant human CB5 donated by Ron Estabrook and Manju Shet, (Univ Texas Southwestern Medical School, Dallas TX). The anti-human HSD3B antiserum (which we find to react with both isozymes of the human, HSD3B1 and HSD3B2, which share 95% sequence homology and are both coded for on Chromosome 1 in the human) was developed in the rabbit by Dr. Richard Parker against purified human placental 3 β -hydroxysteroid dehydrogenase/delta 5-4 isomerase kindly donated by Dr. James Thomas (Mercer University School of Medicine, Macon GA). The mouse monoclonal anti-human CYP17 antibody was developed by Dr Richard Parker against recombinant human CYP17 that was generated by an expression plasmid kindly donated by Michael Waterman (Vanderbilt University, Nashville TN). Chimpanzee CYP17 differs by only 2 amino acids from that in the human, and there is 95% homology between these and the enzyme in rhesus and baboon (Arlt et al, 2002). The antisera used in this study of the chimpanzee adrenal have been previously utilized in several published studies on human and rhesus steroidogenic tissues (Dharia et al, 2004; Mapes et al, 2002; Parker et al, 1995; Parker et al 2000, for example).

2.3

Briefly described, after the tissue sections were subjected to deparaffinization in xylene and progressive hydration in ethanol, 95% ethanol, 75% ethanol, and distilled water, the sections were incubated with 3% hydrogen peroxide for 10 min at 25°C to block endogenous

peroxidase. Then, sections were rinsed in distilled water and Tris buffer, pH 7.6, and incubated for 20 min at 25°C in 5% goat serum to block non-specific antibody binding. The sections were then drained and incubated for 1 hr at 25°C with antisera diluted in PBE buffer, pH 7.6 (0.1M Phosphate buffer that contained 1% bovine serum albumin, 1 mM EDTA, 1.5mM sodium azide). The dilutions of antisera used for immunohistochemistry were as follow: a 1:12,000 dilution of rabbit anti-human CB5, a 1:40 dilution of mouse anti-human CYP17, a 1:500 dilution of rabbit anti-human HSD3B1, and a 1:500 dilution of rabbit anti-human SULT2A1. These dilutions were based on our prior studies of human and rhesus steroidogenic tissues. We also immunostained some sections for a neuroendocrine marker of the adrenal medulla by use of rabbit anti-human Chromogranin A antiserum (DAKO, Cat # N1535), diluted 1:500 in PBE buffer and incubated as above. After being rinsed in Tris buffer, all sections were incubated with linking and labeling reagents contained in the Multi-Species Ultra Streptavidin Horseradish peroxidase detection kit with DAB as chromogen (Covance). Sections were then counterstained with hematoxylin and coverslipped after dehydration using graded ehanols to xylene.

2.4

Digital photography of the slides was accomplished through the use of a Zeiss Axio Imager M2. Images were captured as Tiff files and composite figures were prepared using Power Point software. Magnifications cited in this paper are the product of the eyepiece lens magnification (10X) and those of the objective lenses (2.5, 5, and 10X).

3. Results

3.1

Although adrenal sections from all specimens were usually found to contain regions that were immunopositive for HSD3B, CYP17, SULT2A1, and/or CB5, the distribution and abundance of these elements of the steroidogenic pathway varied with morphologic zonation and developmental stage. There were no apparent differences in the patterns or intensity of immunostaining between the adrenals of male and female chimpanzees at similar ages.

3.2 Newborn

In the newborn chimpanzee (n = 3), the cortex was found to contain a narrow outer cortical zone (approximately 25% of the total cortical thickness) resembling that of the human that has been termed the neocortex and a broad inner area previously described as the fetal zone (Fig 1A). The outer portion of the neocortical area in the chimpanzee resembles the zona glomerulosa whereas the inner part of the neocortex appears to be similar to the cell groupings in the human fetal adrenal characterized as the transitional zone (likely precursor to the zona fasciculata). HSD3B staining was mainly restricted to cells in the outer 2/3rds of the neocortex; minimal, if any, immunostaining for this steroidogenic enzyme was noted in the fetal zone (Fig 1B). CYP17 staining was noted throughout the fetal zone and the inner neocortex but was absent from the outer neocortex (Fig 1C). The immunostaining for CB5 (Fig 1D) was found throughout the fetal zone but not in the neocortex. The immunohistochemical staining pattern for SULT2A1 (not shown) was similar to that of CB5 in the adrenals of newborn chimpanzees.

3.3 Early Infancy

3.3.1—In the adrenal of a 1 month old chimpanzee, the outer cortex area was broadened compared to that of the newborns and was immunopositive for HSD3B but not the other steroidogenic markers. Immunostaining for SULT2A1 and CB5 persisted in the broad fetal zone, as did that for CYP17, but not in the neocortex.

3.3.2—In the adrenal of a 3 month old chimpanzee, however, there were marked changes compared to that of the newborn. There was evidence for marked regression of the fetal zone and broadening of the outer cortex (Fig 2A). Coincident with these morphologic changes, there was a widening of the HSD3B positive portion of the cortex (Fig 2B) and a relative shrinkage in the cortical region expressing SULT2A1 (Fig 2C) and CB5 (Fig 2D). Staining for CYP17 (not shown) extended from the innermost cells expressing HSD3B through the innermost area of the cells expressing CB5 and SULT2A1. There also was an amorphous area between the cortex and the medulla (as defined by expression of Chromogranin-A and morphologic appearance by H & E staining), that contained a few scattered clusters of cells that immunostained for CB5, SULT2A1 and CYP17. It was in this perimedullary region in which involution of the fetal zone appeared to be occurring.

3.3.3—The histologic appearance of the area immediately surrounding the medulla at 1 month, 3 months and 13 months is shown in Figure 3. Compared to that seen in the term fetal adrenal, in the 1 month adrenal (Panels A-1 and A-2), we noted a loss of cortical cells and an increase of acellular material and scattered macrophages containing brown refractile pigment that would be consistent with hemosiderin. We did not notice any such pigment containing cells in the outer cortex. There also was evidence of hemorrhage in the inner aspects of the fetal zone. Many of the cortical cells in this perimedullary region were swollen and vacuolated, consistent with that seen in autophagic cells (Edinger and Thompson, 2004; Gurpinar et al, 2013; Rodrigues et al, 2009). Some of these large cells contained nuclei and in others there were no nuclei visible; however, some nuclei of these cells appeared to be degenerated, but not apoptotic (Gurpinar et al, 2013). Also, in this area were numerous cells with spindle-shaped nuclei, consistent with fibroblast-like cells. Additionally, there were cortical cells that appeared to be apoptotic (Edinger and Thompson, 2004) in the perimedullary region.

In the adrenal of the 3 month chimpanzee, we noted (Figure 3, Panels B-1 and B-2) large numbers of cortical cells near the medulla that appeared to be autophagic (with and without nuclei visible) as well as cells appearing to be undergoing apoptosis. Autophagic and apoptotic cells were not prominent elsewhere in the adrenal. However, by 13 months of age, there were only a few cells in the perimedullary region that appeared to be autophagic or apoptotic; zona fasciculata-like cells predominated at the cortical-medullary border, which was defined by a thin capsule (Figure 3, Panels C-1 and C-2).

3.3.4—When we closely examined the immunohistochemical nature of the cells that appeared to be autophagic at 3 months of age, (Figure 4), we noted that many such cells were immunopositive for CB5 (Panels A and B) and SULT2A1 (Panels C and D). These cells did not express HSD3B. Some such cells had a prominent nucleus and in others the

nucleus was not visible upon focusing through the tissue section. The cells with the least intense immunostaining were usually those in which the nucleus was not seen.

3.4 Later Infancy

Further changes were noted in immunohistochemical characteristics of the chimpanzee adrenal at 13 months of age. Virtually the entire cortex was immunopositive for HSD3B and there was striking immunostaining for CYP17 that extended from the medulla to the junction with the cell layers resembling the zona glomerulosa. On the other hand, there were only a few scattered groups of CB5-positive cells in the innermost cortical area adjacent to the medulla (Fig 5A); the staining profile for SULT2A1 (not shown) was similar to that of CB5 but SULT2A1-positive cells were less widespread. The abundance of CB5 and SULT2A1-expressing cells was strikingly diminished in relation to that seen at 3 months.

3.5 Early Juvenile

In the adrenal of a 3 yr old chimpanzee, as shown in Figure 5B there was evidence for a distinct zona glomerulosa and zona fasciculata and an inner cortical zone that adjoined an unorganized group of cells surrounding the medulla. HSD3B staining (not shown) was present throughout the zona glomerulosa and zona fasciculata whereas CYP17 was localized to the zona fasciculata and the inner cortical zone (Fig 5C). CB5 staining (Fig 5D) was noted in the thin innermost cortex (presumably the zona reticularis and/or residual fetal zone) adjoining the medulla and the abundance of such CB5-positive cells was increased compared to that noted above at 13 months (Fig 5A). SULT2A1 staining (not shown) also was noted within this inner cortical zone but was detectable in strikingly fewer cells than was CB5.

3.6 Young Adult

The adrenal of a young adult chimpanzee (12 yr old) was found to have 3 distinct cortical zones. CYP17 staining (Fig 6A) was noted in the zona fasciculata and zona reticularis but not the outer cortical zone (zona glomerulosa). On the other hand, both the zona glomerulosa and fasciculata were positive for HSD3B (Fig 6B). In this adrenal, the zona reticularis was considerably broader than in the 3yr old and was immunopositive for CB5 (Fig 6C) as it was for CYP17 (Fig 6A); the zona reticularis was not positive for HSD3B (not shown). As had been noted in the 3 yr old, SULT2A1 (Fig 6D) staining of the young adult chimpanzee adrenal was restricted to the zona reticularis and was less widespread in this zone than that for CB5 (Fig 6C).

3.7 Mature Adult

3.7.1—We also noted that cortical cell clusters within the medulla that co-expressed both CYP17 (Fig 7A) and CB5 (Fig 7B) were fairly common in the adult chimpanzee adrenal. Many such clusters of cortical cells in the medulla also expressed SULT2A1 but not HSD3B. Note also that the zona reticularis as defined by CB5 staining (Fig 7B), although somewhat irregular in its border with the fasciculata, comprised about 1/3 of the cortical width in this 21 yr old's adrenal.

3.7.2—The adult chimpanzee adrenal also was noted to have a cortical cuff around the central adrenal vein (Fig 8A) (similar to that noted previously in the human) with cells resembling zona fasciculata that expressed HSD3B (Fig 8B) and CYP17 (Fig 8C) next to the vein and medulla, and then cells that expressed CYP17 and CB5 but not HSD3B (Fig 8D) furthest from the vein. Many of the cells that co-expressed CYP17 and CB5 also were immunopositive for SULT2A1 (not shown).

4. Discussion

4.1

The development of the human adrenal cortex involves several changes in structure and functional capacity during fetal life and after birth (Lanman, 1961; Mesiano and Jaffe, 1997). During fetal development, the adrenal cortex is composed of a large inner cortical mass known as the fetal zone and a thin outer rim of cells termed the neocortex or definitive zone. At mid-gestation, the fetal adrenal is as large as the fetal kidney and by the end of gestation, the fetal adrenal gland is as large as that of the adult human, with the fetal zone comprising over 70% of the total cortical mass. The human fetal adrenal serves an integral physiologic partner in pregnancy maintenance by virtue of the massive production in the fetal zone of estrogenic precursors, especially DHEA and DHEA sulfate, which are converted into estrogens by the placenta. Soon after birth, the fetal zone of the human adrenal involutes by a process that likely involves apoptosis (Bech et al, 1969; Bocian-Sobkowska et al, 1998; Spencer et al, 1999) and this regression is accompanied by reductions in circulating levels of DHEA and DHEA sulfate (Forest et al, 1978; Garagorri et al, 2008).

4.2.1

Development of a structurally and functionally similar adrenocortical zone during fetal life has been noted in only a few other mammalian species. While there has been considerable progress in characterizing the development and function of the rhesus and baboon fetal adrenal (Ducsay et al, 1991; Leavitt et al, 1999; McNulty et al, 1981; Nguyen et al, 2008), there is no information concerning the functional zonation of the chimpanzee adrenal cortex. In our study, we found that the adrenal gland of the newborn chimpanzee bears considerable similarity to that of the human adrenal at term in that both have a large inner cortical zone and a thin outer cortex that resembles the zona glomerulosa and a compressed zona fasciculata of the adult. Although we did not perform any quantitative measurements in this study, we believe that the relative thickness of the outer cortex of the newborn chimpanzee appears to be greater than that of the human, as also was shown by Czekala et al, 1983. By virtue of the presence of CYP17, CB5, and SULT2A1 and the absence of HSD3B staining in the fetal zone, we suggest that the fetal zone of the chimpanzee has the same steroidogenic capacity as the human fetal zone to produce DHEA and DHEA sulfate, which could serve as precursors for placental estrogen formation in pregnancy of the chimpanzee. It is known that estrogen production in chimpanzee pregnancy follows a temporal pattern that is similar to that of the human (Reyes et al, 1975; Smith et al, 1999). Fetal adrenal androgen production also is evident in other primates such as the rhesus and baboon and likely arises from the fetal zone as the result of the prominence of CYP17 and CB5 and a paucity of HSD3B

(Mesiano et al, 1993; Leavitt et al, 1999; Mapes et al, 2002) and, at least in the rhesus, the presence also of SULT2A1 (Parker et al, 2000).

4.2.2

The outer cortical zone of the newborn chimpanzee adrenal was found to express HSD3B but not CYP17, CB5 or SULT2A1, which is the same profile that we and others have found in the human neocortex and which is similar to the expected enzymology of the zona glomerulosa (Barker et al, 1995; Dharia et al, 2004; Dupont et al, 1990; Mesiano et al, 1993; Parker et al, 1994; Parker et al, 1995; Suzuki et al, 2000). In the chimpanzee, we also noted the presence of cortical cells located between the glomerulosa-like outer cortex and the fetal zone that were arranged in columns and contained HSD3B and CYP17 but little, if any, CB5 or SULT2A1. Such a cortical arrangement in the newborn chimpanzee is similar to the pattern that characterizes the 'transitional' zone of the human fetal adrenal and the zona fasciculata of the adult human adrenal. Although we are unable to document the fetal developmental changes in the chimpanzee adrenal, it is likely that the fetal zone requires trophic support of the hypothalamic-pituitary axis for its growth and steroidogenic output, as has been shown to be the case for the fetal adrenal of the human and several other primates (Lanman, 1961; Mesiano and Jaffe, 1997).

4.3

During infancy in the chimpanzee, we found evidence for adrenocortical reorganization that qualitatively resembles that noted for the human, with involution of the fetal zone along with development of a medullary capsule and expansion of the outer cortex into zona glomerulosa and zona fasciculata (Bech et al, 1969; Bocian-Sobkowska et al, 1998; Dhom, 1973; Sucheston and Cannon, 1968). There was persistence of a broad inner cortex that was immunopositive for CB5 and SULT2A1 in a 1 month old chimpanzee, but by 3 months of age, there was clear evidence for involution of this zone. Subsequently, there were only a few scattered cells remaining in the inner cortical area that were immunopositive for CB5 and SULT2A1 in the adrenal of a 13 month old chimpanzee; the rest of the cortex in this specimen displayed the characteristic staining pattern of the zona fasciculata (HSD3B and CYP17 positive) and zona glomerulosa (HSD3B positive, CYP17 negative).

4.4

The majority of the loss of the human fetal zone occurs within the first few weeks of life, followed by a much slower process occurring over the first 1–2 years of life (Bech et al, 1969; Bocian-Sobkowska et al, 1998; Sucheston and Cannon, 1968). The process of regression of the human adrenal has been described as an initial hemorrhagic process (Bech et al, 1969) that is associated with striking levels of apoptosis (Bocian-Sobkowska et al, 1998; Spencer et al, 1999). Our results are suggestive that apoptosis and autophagy (Edinger and Thompson, 2004; Gulpinar et al, 2013; Rodrigues et al, 2009;) may play a role in regression of the fetal zone in the chimpanzee. There also is some indication of hemorrhage in the chimpanzee fetal zone. Our sample size is too limited to provide more than an approximation of the timeline involved in reorganization of the adrenal cortex during infancy. Nevertheless, it appears that fetal zone regression is largely complete within the first year of life and development of the zona reticularis occurs later.

4.5

There also is regression of the fetal zone during early infancy in other related species such as the rhesus monkey, baboon, and marmoset monkey (Levine et al, 1982; McNulty et al, 1981; Nguyen et al, 2008). In the baboon and rhesus, however, the regression of the fetal zone appears to coincide with the formation of the zona reticularis during infancy. In both species, a group of small densely staining cells have been noted between the outer cortex and the fetal zone that expands as the fetal zone shrinks (Ducsay et al, 1991; McNulty et al, 1981). Ducsay et al considered the dense zone of the baboon to represent an area of degeneration of fetal zone cells; this aspect of baboon adrenal development does not appear to have been explored further. On the other hand, the dense band in the neonatal rhesus adrenal was believed to be the site of transition into zona fasciculata cells in the outer dense band and into reticularis cells in the inner portion of the dense band (McNulty et al, 1981), and immunohistochemical studies of the infant rhesus adrenal (Nguyen et al, 2008) may be consistent with such an interpretation. We did not notice the presence of a dense band of cortical cells in the newborn or infant adrenals of the chimpanzee and have not seen any reference to this type of morphologic feature in the human adrenal during the perinatal period.

4.6

By 3 years of age, the chimpanzee adrenal was found to have a thin, but continuous inner cortical zone that was positive for both CB5 and CYP17 and to a lesser extent, SULT2A1; the cells of this zone were negative for HSD3B. This staining pattern is characteristic of that seen in the zona reticularis of the human adrenal cortex that emerges during adrenarche and persists thereafter (Endoh et al, 1996; Dharia et al, 2004; Kennerson et al, 1983; Suzuki et al, 2000).

4.7

The presence of a zona reticularis that expresses a phenotype consistent with the capacity for DHEA and DHEA sulfate synthesis in the young chimpanzee and in older specimens regardless of gender, in which the zone became broader though young adulthood provides the functional basis for the developmental changes in circulating levels of adrenal androgens that has been documented during adrenarche and thereafter in the chimpanzee (Bernstein et al, 2012; Collins et al, 1981; Copeland et al, 1985; Cutler et al, 1978). The development of the zona reticularis in the chimpanzee differs somewhat from that in the rhesus and marmoset monkey and the baboon. In the rhesus and the baboon, the zona reticularis appears to develop as the fetal zone is regressing (Ducsay et al, 1991; McNulty et al, 1981; Nguyen et al, 2008), thus masking developmental changes in adrenal androgen production that would be indices of the process of adrenarche (Castracane et al, 1981; Nguyen et al, 2008). On the other hand, the zona reticularis of the marmoset monkey is only apparent in females and the enzymology and androgen production is subject to regulation by social status and ovarian function (Pattison et al, 2005; 2007).

4.8

With aging in the human, there is maintenance of corticosteroid secretion but impairment in DHEA and DHEA sulfate secretion by the adrenal (Dharia and Parker, 2004, Parker et al, 2000). There also is evidence for decreased adrenal androgen production during aging in the rhesus and baboon (Lane et al, 1997; Sapolsky et al, 1993). We have noted that there appears to be narrowing of the zona reticularis but not the zona fasciculata/glomerulosa in the human adrenal during aging (Dharia et al, 2005; Parker et al, 1997) that may contribute to the striking adrenal androgen deficiency seen in humans during aging. Recently, Blevins et al, 2013 have provided evidence for gradual reductions in circulating levels of DHEA sulfate in the aging chimpanzee; these changes, however, are not as striking as that noted in the human. Whether such reductions in the chimpanzee, (or in the rhesus and baboon) during aging can be attributed to morphologic changes in the adrenal will require further study.

Acknowledgments

These studies were supported by NSF grant BCS-0717886 (K. Hawkes), NIH grant P51 RR000165 (Yerkes National Primate Research Center), and NIH grants P51 IRR013986 and OD P51 OD011133 (Southwest National Primate Research Center). The authors wish to thank the staffs of the Yerkes National Primate Research Center and the Southwest National Primate Research Center, who provided the adrenal sections from their archival autopsy specimens for our studies. We also wish to acknowledge the important contributions of Rob Stockard to these studies.

References

- Albertson BD, Hobson WC, Burnett BS, Turner PT, Clark RV, Schiebinger RJ, Loriaux DL, Cutler GB Jr. Dissociation of cortisol and adrenal androgen secretion in the hypophysectomized adrenocorticotropin-replaced chimpanzee. *J Clin Endocrinol, Metab.* 1984; 59:13–18. [PubMed: 6327755]
- Albrecht ED, Haskins AL, Pepe GL. The influence of fetectomy at midgestation upon the serum concentrations of progesterone, estrone and estradiol in baboons. *Endocrinology.* 1980; 107:766–770. [PubMed: 6772429]
- Arlt W, Martens JWM, Song M, Wang JT, Auchus RJ, Miller WL. Adrenarche: Structural and functional analysis of P450c17 from four primate species. *Endocrinology.* 2002; 143:4665–4672. [PubMed: 12446594]
- Barker EV, Hume R, Hallas A, Coughtrie MWH. Dehydroepiandrosterone sulfotransferase in the developing human fetus: Quantitative biochemical and immunological characterization of the hepatic, renal and adrenal enzymes. *Endocrinology.* 1995; 134:982–989. [PubMed: 8299591]
- Bech K, Tygstrup I, Nerup J. The involution of the foetal adrenal cortex. *Acta Path. Microbiol. Scand.* 1969; 76:391–400. [PubMed: 5823359]
- Bernstein RM, Sterner KN, Wildman DE. Adrenal androgen production in catarrhine primates and the evolution of adrenarche. *Am. J. Phys. Anthropol.* 2012; 147:389–400. [PubMed: 22271526]
- Blevins JK, Coxworth JE, Herndon JG, Hawkes K. Brief Communication: Adrenal androgens and aging: female chimpanzees (*Pan troglodytes*) compared with women. *Am. J. Phys. Anthropol.* 2013; 151:643–648. [PubMed: 23818143]
- Bocian-Sobkowska J, Wozniak W, Malendowicz LK. Postnatal involution of the human adrenal fetal zone: Stereologic description and apoptosis. *Endocrine Res.* 1998; 24:969–983. [PubMed: 9888608]
- Bornstein SR, Gonzalez-Hernandez JA, Ehrhart-Bornstein M, Adler G, Scherbaum WA. Intimate contact of chromaffin and cortical cells within the human adrenal gland forms the cellular basis for important intraadrenal interactions. *J Clin. Endocrinol. Metab.* 1994; 78:225–232.
- Castracane VD, Cutler GB Jr, Loriaux DL. Pubertal endocrinology of the baboon: adrenarche. *Am. J. Physiol.* 1981; 241:E305–E309. [PubMed: 6459032]

- Collins DC, Nadler RD, Preedy JRK. Adrenarche in great apes. *Am. J. Primatol.* 1981; 1:344. (abstract).
- Copeland KC, Eichberg J, Parker CR Jr, Bartke A. Puberty in the chimpanzee: Somatomedin-C and its relationship to somatic growth and steroid hormone concentrations. *J. Clin. Endocrinol. Metab.* 1985; 60:1154–1159. [PubMed: 3158669]
- Cutler GB Jr, Glenn M, Bush M, Hodgen GD, Graham CE, Loriaux DL. Adrenarche: A survey of rodents, domestic animals, and primates. *Endocrinology.* 1978; 103:2112–2118. [PubMed: 155005]
- Czekala NM, Benirschke K, McClure H, Lasley BL. Urinary estrogen excretion during pregnancy in the gorilla (*Gorilla gorilla*), orangutan (*Pongo pygmaeus*), and the human (*Homo sapiens*). *Biol. Reprod.* 1983; 28:289–294. [PubMed: 6404315]
- Dharia S, Parker CR Jr. Adrenal androgens and aging. *Sem. Repro. Med.* 2004; 22:361–368.
- Dharia S, Slane A, Jian M, Conner M, Conley AJ, Brissie RM, Parker CR Jr. Effects of aging on cytochrome b5 expression in the human adrenal gland. *J. Clin. Endocrinol. Metab.* 2005; 90:4357–4361. [PubMed: 15870126]
- Dharia S, Slane A, Jian M, Conner M, Conley AJ, Parker CR Jr. Colocalization of P450C17 and cytochrome b5 in androgen synthesizing tissues of the human. *Biol. Reprod.* 2004; 71:83–88. [PubMed: 14985252]
- Dhom G. The prepuberal and puberal growth of the adrenal (Adrenarche). *Beitr. Path. Bd.* 1973; 150:357–377.
- Ducsay CA, Hess DL, McClellan MC, Novy MJ. Endocrine and morphological maturation of the fetal and neonatal adrenal cortex in baboons. *J. Clin. Endocrinol. Metab.* 1991; 73:385–395. [PubMed: 1649841]
- Dupont E, Luu-The V, Labrie F, Pelletier G. Ontogeny of 3 β -hydroxysteroid dehydrogenase / 5-4 isomerase (3 β -HSD) in human adrenal gland performed by immunohistochemistry. *Mol. Cell. Endocrinol.* 1990; 74:R7–R10.
- Edinger AL, Thompson CB. Death by design: Apoptosis, necrosis, and autophagy. *Cur. Opin. in Cell Biol.* 2004; 16:663–669.
- Endoh A, Kristiansen SB, Casson PR, Buster JE, Hornsby PJ. The zona reticularis is the site of biosynthesis of dehydroepiandrosterone and dehydroepiandrosterone sulfate in the adult human adrenal cortex resulting from its low expression of 3 β -hydroxysteroid dehydrogenase. *J. Clin. Endocrinol. Metab.* 1996; 81:3558–3565. [PubMed: 8855801]
- Forest, M.; de Peretti, E.; Bertrand, J. Developmental patterns of the plasma level of testosterone, androstenedione, 17-hydroxyprogesterone, dehydroepiandrosterone and its sulfate in normal infants and prepubertal children. In: James, VHT.; Serio, M.; Guisti, G.; Martini, L., editors. *The endocrine function of the human adrenal cortex.* New York, NY: Academic Press; 1978. p. 561-582.
- Garagorri JM, Rodriguez G, Lario-Elboj AJ, Olivares JI, Lario-Manoz A, Orden I. Reference Levels for 17-hydroxyprogesterone, 11-desoxycortisol, cortisol, testosterone, dehydroepiandrosterone sulfate and androstenedione in infants from birth to 6 months of age. *Eur. J. Pediatr.* 2008; 167:647–653. [PubMed: 17694323]
- Gurpinar E, Grizzle WE, Shacka JJ, Made, r BJ, Li N, Piazza NA, Russo S, Keeton AB, Piazza GA. A novel sulindac derivative inhibits lung adenocarcinoma cell growth through suppression of Akt/mTOR signaling and induction of autophagy. *Mol. Cancer Ther.* 2013; 12:663–674. [PubMed: 23443799]
- Katagiri M, Kagawa N, Waterman MR. The role of cytochrome b5 in the biosynthesis of androgens by human P450c17. *Arch. Biochem. Biophys.* 1995; 317:343–347. [PubMed: 7893148]
- Kennerson AR, McDonald DA, Adams JB. Dehydroepiandrosterone sulfotransferase localization in human adrenal glands: A light and electron microscopic study. *J. Clin. Endocrinol. Metab.* 1983; 56:786–790.
- Lane MA, Ingram DK, Ball SS, Roth GS. Dehydroepiandrosterone sulfate: a biomarker of primate aging slowed by calorie restriction. *J. Clin. Endocrinol. Metab.* 1997; 82:2093–2096. [PubMed: 9215277]

- Lanman JT. The adrenal gland in the human fetus. An interpretation of its physiology and unusual developmental pattern. *Pediatrics*. 1961; 27:140–158. [PubMed: 13758926]
- Leavitt MG, Albrecht ED, Pepe GJ. Development of the baboon fetal adrenal gland: Regulation of the ontogenesis of the definitive and transitional zones by adrenocorticotropin. *J. Clin. Endocrinol. Metab.* 1999; 84:3831–3835. [PubMed: 10523038]
- Lee-Robichaud P, Wright JN, Akhtar ME, Akhtar M. Modulation of the activity of human 17 α -hydroxylase, 17, 20-lyase CYP17 by cytochrome b5: endocrinological and mechanistic implications. *Biochem. J.* 1995; 302:901–908. [PubMed: 8948449]
- Levine J, Wolfe LG, Schiebinger RJ, Loriaux DL, Cutler GB Jr. Rapid regression of fetal adrenal zone and absence of adrenal reticular zone in the marmoset. *Endocrinology*. 1982; 111:1797–1802. [PubMed: 6216097]
- Mapes S, Corbin CJ, Tarantal A, Conley A. The primate adrenal zona reticularis is defined by expression of cytochrome b5, 17 α -hydroxylase/17, 20-lyase cytochrome P450 (P450c17) and NADPH-cytochrome P450 reductase (reductase) but not 3 β -hydroxysteroid dehydrogenase/delta 5-4 isomerase (3 β -HSD). *J. Clin. Endocrinol. Metab.* 1999; 84:3382–3385. [PubMed: 10487714]
- Mapes S, Tarantal AF, Parker CR, Moran FM, Bahr JM, Pyter L, Conley AJ. Adrenocortical cytochrome b5 expression during fetal development of the rhesus Macaque. *Endocrinology*. 2002; 143:1451–1458. [PubMed: 11897703]
- McNulty WP, Novy MJ, Walsh SW. Fetal and postnatal development of the adrenal glands in *Macaca mulatta*. *Biol. Reprod.* 1981; 25:1079–1089. [PubMed: 6275916]
- Mesiano S, Jaffe RB. Developmental and functional biology of the primate fetal adrenal cortex. *Endocrine Res.* 1997; 18:378–403.
- Mesiano S, Coulter CL, Jaffe RB. Localization of cytochrome P450 cholesterol-side chain cleavage, cytochrome P450 17 α -hydroxylase/17, 20-lyase, and 3 β -hydroxysteroid dehydrogenase isomerase steroidogenic enzymes in human and rhesus monkey fetal adrenal glands: Reappraisal of functional zonation. *J. Clin. Endocrinol. Metab.* 1993; 77:1184–1189.
- Nguyen AD, Mapes SM, Corbin CJ, Conley AJ. Morphological adrenarche in rhesus macaques: development of the zona reticularis is concurrent with fetal zone regression in the early neonatal period. *J. Endocrinol.* 2008; 199:367–378. [PubMed: 18787057]
- Parker CR Jr, Falany CN, Stockard CR, Stankovic AK, Grizzle WE. Immunohistochemical localization of dehydroepiandrosterone sulfotransferase in human fetal tissues. *J. Clin. Endocrinol. Metab.* 1994; 78:234–236. [PubMed: 8288708]
- Parker CR Jr, Faye-Petersen O, Stankovic AK, Mason JI, Grizzle WE. Immunohistochemical evaluation of the cellular localization and ontogeny of 3 β -hydroxysteroid dehydrogenase/delta 5-4 isomerase in the human fetal adrenal gland. *Endocrine Res.* 1995; 21:69–80. [PubMed: 7588420]
- Parker CR Jr, Jian M, Conley AJ. The localization of DHEA sulfotransferase in steroidogenic and steroid metabolizing tissues of the adult rhesus macaque monkey. *Endocrine Res.* 2000; 26:517–522. [PubMed: 11196423]
- Parker CR Jr, Mixon RL, Brissie RM, Grizzle WE. Aging alters zonation in the adrenal cortex of men. *J. Clin. Endocrinol. Metab.* 1997; 82:3898–3901. [PubMed: 9360559]
- Parker CR Jr, Slayden SM, Azziz R, Crabbe SL, Hines GA, Boots LR, Bae S. Effects of aging on adrenal function in the human: Responsiveness and sensitivity of adrenal androgens and cortisol to ACTH in premenopausal and postmenopausal women. *J. Clin. Endocrinol. Metab.* 2000; 85:48–54. [PubMed: 10634362]
- Pattison JC, Abbott DH, Saltzman W, Nguyen AD, Henderson G, Jing H, Price CR, Allen AJ, Conley AJ, Bird IM. Male marmoset monkeys express an adrenal fetal zone at birth, but not a zona reticularis in adulthood. *Endocrinology*. 2005; 146:365–374. [PubMed: 15459122]
- Pattison J, Saltzman W, Abbott DH, Hogan BK, Nguyen AD, Husen B, Einspanier A, Conley AJ, Bird IM. Gender and gonadal status difference in zona reticularis expression in marmoset monkey adrenals: Cytochrome b5 localization with respect to cytochrome P450 17, 20-lyase activity. *Mol. Cell. Endo.* 2007; 265–266:93–100.
- Pepe GJ, Albrecht ED. Regulation of the primate fetal adrenal cortex. *Endocrine Rev.* 1990; 11:151–176. [PubMed: 2180686]

- Reyes FI, Winter JSD, Faiman C, Hobson WC. Serial serum levels of gonadotropins, prolactin and sex steroids in the nonpregnant and pregnant chimpanzee. *Endocrinology*. 1975; 96:1447–1445. [PubMed: 1126314]
- Rodrigues P, Limback D, McGinnis LK, Plancha CE, Albertini DF. Multiple mechanisms of germ cell loss in the perinatal mouse ovary. *Reproduction*. 2009; 137:709–720. [PubMed: 19176312]
- Sapolsky RM, Vogelmann JH, Orentreich N, Alt-Mann J. Senescent decline in serum dehydroepiandrosterone sulfate concentrations in a population of wild baboons. *J. Gerontol*. 1993; 48:B196–B200. [PubMed: 8366259]
- Smail PJ, Faiman C, Hobson WE, Fuller GB, Winter JSD. Further studies on adrenarche in nonhuman primates. *Endocrinology*. 1982; 111:844–848. [PubMed: 6213402]
- Smith R, Wickings EJ, Bowman ME, Bellefoud A, Dubreuil G, Davies JJ, Madsen G. Corticotropin-releasing hormone in chimpanzee and gorilla pregnancies. *J. Clin. Endocrinol. Metab*. 1999; 84:2820–2825. [PubMed: 10443686]
- Spencer SJ, Mesiano S, Lee JY, Jaffe RB. Proliferation and apoptosis in the human adrenal cortex during the fetal and perinatal periods: implications for growth and remodeling. *J. Clin. Endocrinol. Metab*. 1999; 84:1110–1115. [PubMed: 10084603]
- Sucheston ME, Cannon MS. Development of the zonular patterns in the human adrenal gland. *J. Morph*. 1968; 126:477–492. [PubMed: 5716437]
- Suzuki T, Sasano H, Takeyana J, Kaneko C, Freije WA, Carr BR, Rainey WE. Developmental changes in steroidogenic enzymes in human postnatal adrenal cortex: immunohistochemical studies. *Clin. Endocrinol*. 2000; 53:739–747.
- Walsh SW, Kittinger GW, Novy MJ. Maternal peripheral concentrations of estradiol, estrone, cortisol, and progesterone during late pregnancy in rhesus monkeys (*Macaca mulatta*) and after experimental fetal anencephaly and fetal death. *Am. J. Obstet. Gynecol*. 1979; 135:37–42. [PubMed: 112865]
- Walsh SW, Norman RL, Novy MJ. In Utero regulation of rhesus monkey fetal adrenals: effects of dexamethasone, adrenocorticotropin, thyrotropin-releasing hormone, prolactin, human chorionic gonadotropin, and α -melanocyte-stimulating hormone on fetal and maternal plasma steroids. *Endocrinology*. 1979; 104:1805–1813. [PubMed: 221193]

Highlights

1. Adrenal sections from 27 autopsied chimpanzees (birth -31yrs) were immunostained
2. Chimpanzee fetal zone (FZ) expressed CYP17, SULT2A1, and Cytochrome b5 but not HSD3B
3. FZ involution was paralleled by reduced expression of SULT2A1 and Cytochrome B5
4. Z reticularis development was evident by 3 yrs and had similar expression to the FZ
5. Adult chimpanzee and human adrenals share many similarities

Figure 1.
Adrenal sections from a newborn chimpanzee. The dark lines approximate the borders between the adrenal capsule, the neocortex (NC), the fetal zone (FZ), and the medulla (M).
Panel A: Hematoxylin and eosin stained section, 50X. Panel B: HSD3B, 100X. Panel C: CYP17, 100X. Panel D: CB5, 100X.

Figure 2.

Adrenal sections from a 3 month old chimpanzee. The dark lines approximate the borders of the regressing fetal zone (FZ); the medulla is indicated by M. Panel A: Hematoxylin and eosin stained section, 100X. Panel B: HSD3B, 50X. Panel C: SULT2A1, 100X. Panel D: CB5, 50X.

Figure 3.

Morphologic characteristics of chimpanzee fetal zone involution. The medulla is indicated by M, capillaries are indicated by C and the zona fasciculata is indicated by ZF. Hematoxylin and eosin stained sections showing the region of the fetal zone near the medulla are shown for the 1 month (Panels A-1 and A-2, both 400X), 3 month (Panels B-1 and B-2, both 400X) and 13 month (Panels C-1, 200x, and C-2, 400X) infant chimpanzees. Autophagic appearing cortical cells with a nucleus are indicated by solid black arrows and those without nuclei are shown by brown filled arrows; of note, multiple nuclei of the autophagic appearing cells are degenerated, but not apoptotic. Apoptotic appearing cortical cells are indicated by unfilled arrows. Macrophages containing brown pigment, presumably hemosiderin, are indicated by blue arrows.

Figure 4.

Immunohistochemical characteristics of cortical cells in the fetal zone of the 3 month old chimpanzee. Panels A and B: CB5 at 200X and 400X, respectively. Panels C and D: SULT2A1 at 200X and 400X, respectively. Autophagic appearing cortical cells with a nucleus are indicated by black arrows and those with no visible nucleus are indicated by brown filled arrows; again, degenerative changes are present in the nuclei of some of these cells.

Figure 5. Adrenal sections from a 13 month old chimpanzee (A) and a 3 year old chimpanzee (B, C, D). Panel A: CB5, 50 X. The few scattered immunopositive cells are shown within the circle near the medulla (M). Panel B: Hematoxylin and eosin stained section, 50X. Panel C: CYP17, 50X. Panel D: CB5, 50X. The junction between the zona fasciculata and zona glomerulosa is indicated by the arrows.

Figure 6. Adrenal sections from a 12 year old chimpanzee. Panel A: CYP17, 25X. Panel B: HSD3B, 100X. Panel C: CB5, 25X. Panel D: SULT2A1, 50X. The arrows point out the zona glomerulosa. The V indicates the central medullary vein.

Figure 7.
Adrenal sections from a 21 year old chimpanzee showing intramedullary cortical cells. Panel A: CYP17, 25X. The medulla (M), central vein (V) and adrenal capsule (arrow) are pointed out. Panel B: CB5, 25X.

Figure 8.

Adrenal sections from a 31 year old chimpanzee demonstrating the cortical cuff. Panel A: hematoxylin and eosin stained section, 100X. The dashed line in each panel indicates the border between the cortex and the medulla (M) and the central vein (V). Panel B: HSD3B, 100X. Panel C: CYP17, 100X. Panel D: CB5, 100X.