Anesthesiology and Pain Medicine

Journal home page: www.AnesthPain.com

Quality of Life Evaluation

Laxmaiah Manchikanti^{1*}

¹ Pain Management Center of Paducah, Paducah, USA

ARTICLE INFO

Article type:
Letter to Editor

Article history:
Received: 16 Nov 2011
Revised: 17 Nov 2011
Accepted: 19 Nov 2011

Keywords:
Quality of Life
Discectomy
Surgery

► Please cite this paper as:

Manchikanti L. Quality of Life Evaluation. *Anesth Pain*. 2012;1(3):203-4. DOI: 10.5812/kowsar.22287523.3548

Copyright © 2012, ISRAPM, Published by Kowsar Corp.

Dear Editor,

Farzanegan *et al.* described quality of life evaluations of patients undergoing lumbar discectomy using the short-form 36 (SF-36) (1). They provided physical and mental health scores of patients which improved significantly 6 and 12 months after lumbar discectomy, leading to the conclusion that lumbar discectomy improves both the physical and mental health subscale of the quality of life in patients with chronic disc herniation.

Quality of life and functional status improvement is evaluated by many available tests; one of them SF-36 as utilized in this study. While all these instruments are considered as objective evaluations, they all depend on subjective information. SF-36 is a generic measure, as opposed to one that targets a specific age, disease, or treatment group. Thus, for low back pain, either the Oswestry Disability Index (ODI) (2) or Roland Morris Disability Questionnaire (RDQ) (3) are considered as more specific measures. Even so, the studies of validity of SF-36 from many types of research have yielded content, concurrent, criterion, construct, and predictive evidence of validity (4). Consequently, the SF-36 has been shown to be sensitive to change (5, 6) and able to differentiate be-

tween treatment responders and non-responders (7, 8). Further, SF-36 has been used as a validation tool in the development of new disease-specific instruments (9, 10), including a pain-specific tool (10, 11).

However, there has not been any significant description of a clinically significant change in any condition for SF-36. In contrast, clinically significant change has been described as a 15-point change in patients who undergo spinal fusion before surgery and at follow-up for ODI (4). Others have described a change of 4 points as the minimum difference in mean scores between the group that carried clinical significance. On the same grounds, it has been suggested that the smallest change likely to be clinically significant for RDQ lies between 2.5 and 5 points (4). In recent years, it has been stated that a 40% or 50% change from pre-treatment level as the appropriate change (12, 13).

Consequently, when these issues are taken into consideration, do the changes described in Farzanegan *et al.*'s manuscript (1), which looked rather dramatic with significantly high *P* values, really indicate clinical significance? Essentially the changes described in this manuscript for physical health scores, which are the best, show only 18% improvement, whereas for mental health, it appears that they may be less than 15% from 38.16 to 43.48 (value for 12 months appears to be wrong in Figure 2).

The clinical implications of the study include not only its usefulness in evaluating lumbar discectomy, but also the questions about meaningful change, which is clini-

* Corresponding author: Laxmaiah Manchikanti, Pain Management Center of Paducah, Paducah, KY, USA. Tel: +1-2705548373 ext: 101, Fax: +1-2705545394, E-mail: drm@asipp.org

cally significant. In the future, studies such as this one should be compared with other disease specific tests such as RDQ. In this case, most patients had subacute pain, whereas ODI is for patients with chronic pain.

Financial Disclosure

None declared.

References

1. Farzanegan G, Alghasi M, Safari S. Quality-of-Life Evaluation of Patients Undergoing Lumbar Discectomy Using Short Form 36. *Anesth Pain*. 2011;**1**(2):73-6.
2. Fairbank JC, Pynsent PB. The Oswestry Disability Index. *Spine (Phila Pa 1976)*. 2000;**25**(22):2940-52. discussion 52.
3. Roland M, Morris R. A study of the natural history of back pain. Part I: development of a reliable and sensitive measure of disability in low-back pain. *Spine (Phila Pa 1976)*. 1983;**8**(2):141-4.
4. Manchikanti L, Hirsch JA, Smith HS. Evidence-based medicine, systematic reviews, and guidelines in interventional pain management: Part 2: Randomized controlled trials. *Pain Physician*. 2008;**11**(6):717-73.
5. Beaton DE, Hogg-Johnson S, Bombardier C. Evaluating changes in health status: reliability and responsiveness of five generic health status measures in workers with musculoskeletal disorders. *J Clin Epidemiol*. 1997;**50**(1):79-93.
6. Essink-Bot ML, Krabbe PF, Bonsel GJ, Aaronson NK. An empirical comparison of four generic health status measures. The Nottingham Health Profile, the Medical Outcomes Study 36-item Short-Form Health Survey, the COOP/WONCA charts, and the EuroQol instrument. *Med Care*. 1997;**35**(5):522-37.
7. Bronfort G, Bouter LM. Responsiveness of general health status in chronic low back pain: a comparison of the COOP charts and the SF-36. *Pain*. 1999;**83**(2):201-9.
8. Kosinski M, Keller SD, Ware JE, Jr., Hatoum HT, Kong SX. The SF-36 Health Survey as a generic outcome measure in clinical trials of patients with osteoarthritis and rheumatoid arthritis: relative validity of scales in relation to clinical measures of arthritis severity. *Med Care*. 1999;**37**(5 Suppl):MS23-39.
9. Roos EM, Klassbo M, Lohmander LS. WOMAC osteoarthritis index. Reliability, validity, and responsiveness in patients with arthroscopically assessed osteoarthritis. Western Ontario and MacMaster Universities. *Scand J Rheumatol*. 1999;**28**(4):210-5.
10. Smith BH, Penny KI, Purves AM, Munro C, Wilson B, Grimshaw J, et al. The Chronic Pain Grade questionnaire: validation and reliability in postal research. *Pain*. 1997;**71**(2):141-7.
11. Elliott AM, Smith BH, Smith WC, Chambers WA. Changes in chronic pain severity over time: the Chronic Pain Grade as a valid measure. *Pain*. 2000;**88**(3):303-8.
12. Manchikanti L, Singh V, Cash KA, Pampati V, Damron KS, Boswell MV. A randomized, controlled, double-blind trial of fluoroscopic caudal epidural injections in the treatment of lumbar disc herniation and radiculitis. *Spine (Phila Pa 1976)*. 2011;**36**(23):1897-905.
13. Manchikanti L, Singh V, Cash KA, Pampati V, Datta S. A comparative effectiveness evaluation of percutaneous adhesiolysis and epidural steroid injections in managing lumbar post surgery syndrome: a randomized, equivalence controlled trial. *Pain Physician*. 2009;**12**(6):E355-68.