Trigeminal Neuralgia: A Condition to Be Better Understood

Felipe Porto ^{1*}

¹ University of Connecticut, School of Dental Medicine, Division of Behavioral Sciences and Community Health, Farmington, USA

ARTICLE INFO

Article type:
Letter to Editor

Article history:
Received: 11 Nov 2011
Revised: 12 Nov 2011
Accepted: 13 Nov 2011

Keywords:
Trigeminal Neuralgia
Patients
Multiple Sclerosis

► Please cite this paper as:

Porto F. Trigeminal Neuralgia: A Condition to Be Better Understood. *Anesth Pain*. 2012;1(3):201-2.

DOI: 10.5812/kowsar.22287523.3492

Copyright © 2012, ISRAPM, Published by Kowsar Corp.

Dear Editor,

I read the paper entitled "Trigeminal Neuralgia: Frequency of Occurrence in Different Nerve Branches", published in your journal (1). The data presented confirms the findings of previous papers (2-4), as was also mentioned by the author. Such findings help the clinicians to diagnose the disease. However, some other points need to be mentioned; even though this study reports no case of bilateral trigeminal neuralgia, previous publications reported that it may affect 1-5% of the patients (5, 6), and another author mentioned that bilateral cases should suggest a systemic involvement, such as multiple sclerosis (MS) or cranial tumor (7). We can conclude that bilateral cases are rare, but not impossible; and a clinician should be suspicious of a systemic condition when examining a patient with bilateral trigeminal neuralgia (TN). Considering that TN is a clinical diagnosis, the clinician should not only rely on the epidemiologic data, even though it might help on the differential diagnosis. Clinical characteristics should be considered during the history taking and clinical examination, and those characteristics can be found on the diagnostic criteria published by the International Headache Society (IHS). The

IHS divides TN in two subsets:

A) Classical TN, which is not related to any pathology, but vascular compression.

B) Symptomatic TN, which is caused by an identifiable lesion or pathology.

It is known that over 85% of the cases with TN have the classical TN. All clinicians involved on the management of pain patients should be familiar with the diagnostic criteria published by the IHS, which helps to correctly identify patients with TN, as well as differentiate between classical TN and symptomatic TN (8).

Although diagnosing TN appears to be easy, many patients have undergone unnecessary dental treatment (9), and about 90% of patients diagnosed with TN experienced pain for more than 1 year before receiving the correct diagnosis (10).

Before establishing a long-term treatment, the health care provider should first try to identify the etiology of the TN. As the Bangash's paper well mentioned, the clinician should closely follow the patients up, in order to select the best treatment, watch for side effects and often evaluate the patient for any change on the medical history (including new intra and extra oral exam). Clinicians should become familiar not only with treatment options, indications and contraindications for each treatment modality, side effects and adverse reactions associated with the treatment, but also with new studies about diagnosis and treatment of TN.

As all the other neuropathic pain conditions, TN is not

* Corresponding author: Felipe Porto, University of Connecticut, School of Dental Medicine, Division of Behavioral Sciences and Community Health, Farmington, USA. Tel: +1-8606792778, E-mail: porto@uchc.edu

completely understood yet. Finally, further investigations should focus not only on the treatment of TN, but also on the pathogenesis of this condition. Epidemiologic studies as the article published by Bangash on your journal are the first step to develop more specific studies regarding etiology and new treatment options in order to decrease/eliminate the patient's suffering and clinician's frustration.

Financial Disclosure

None declared.

References

1. Bangash TH. Trigeminal Neuralgia: Frequency of Occurrence in Different Nerve Branches. *Anesth Pain*. 2011;**1**(2):70-2.
2. Hall GC, Carroll D, Parry D, McQuay HJ. Epidemiology and treatment of neuropathic pain: the UK primary care perspective. *Pain*. 2006;**122**(1-2):156-62.
3. Neto HS, Camilli JA, Marques MJ. Trigeminal neuralgia is caused by maxillary and mandibular nerve entrapment: greater incidence of right-sided facial symptoms is due to the foramen rotundum and foramen ovale being narrower on the right side of the cranium. *Med Hypotheses*. 2005;**65**(6):1179-82.
4. Hunt K, Patwardhan R. Trigeminal neuralgia: a modern-day review. *Int Rev Neurobiol*. 2007;**79**:621-31.
5. Kuncz A, Voros E, Barzo P, Tajti J, Milassin P, Mucsi Z, et al. Comparison of clinical symptoms and magnetic resonance angiographic (MRA) results in patients with trigeminal neuralgia and persistent idiopathic facial pain. Medium-term outcome after microvascular decompression of cases with positive MRA findings. *Cephalalgia*. 2006;**26**(3):266-76.
6. Leeuw R., American Academy of Orofacial Pain. *Orofacial pain: guidelines for assessment, diagnosis, and management*. 4th ed. Chicago: Quintessence; 2008.
7. Okeson JP, Bell WE. *Bell's orofacial pains: the clinical management of orofacial pain*. 4th ed. Chicago: Quintessence Pub. Co.; 2005.
8. Olesen J, Boussier MG, Diener HC. The International Classification of Headache Disorders 2nd Edition: Part One: The Primary Headaches. *Cephalalgia*. 2004;**24**:23-136.
9. de Siqueira SR, Nobrega JC, Valle LB, Teixeira MJ, de Siqueira JT. Idiopathic trigeminal neuralgia: clinical aspects and dental procedures. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2004;**98**(3):311-5.
10. Casey KF. Role of patient history and physical examination in the diagnosis of trigeminal neuralgia. *Neurosurg Focus*. 2005;**18**(5):E1.