Published in final edited form as:

Biochemistry. 2010 February 9; 49(5): 1014–1023. doi:10.1021/bi9015549.

Enhancing Resistance to Cephalosporins in class C β -lactamases: impact of Glu214Gly in CMY-2

Andrea Endimiani^{1,2}, Yohei Doi³, Christopher R. Bethel¹, Magdalena Taracila^{1,2}, Jennifer M. Adams-Haduch³, Alexandra O'Keefe³, Andrea M. Hujer^{1,2}, David L. Paterson^{3,4}, Marion J. Skalweit^{1,2}, Malcolm G. P. Page⁵, Sarah M. Drawz⁶, and Robert A. Bonomo^{1,2,7,8,*}

¹Research Service, Louis Stokes Cleveland Department of Veterans Affairs Medical Center, Cleveland, OH ²Departments of Medicine, Case Western Reserve University School of Medicine, Cleveland, OH ³University of Pittsburgh, Medical Center, Pittsburgh, PA ⁴Centre for Clinical Research, University of Queensland, Royal Brisbane and Women's Hospital, Brisbane, Australia ⁵Basilea Pharmaceutica International Ltd, Basel, Switzerland ⁶Department of Pathology Case Western Reserve University School of Medicine, Cleveland, OH ⁷Department of Pharmacology Case Western Reserve University School of Medicine, Cleveland, OH ⁸Departments of Molecular Biology and Microbiology Case Western Reserve University School of Medicine, Cleveland, OH

Abstract

The biochemical properties of CMY-32, a class C enzyme possessing a single amino acid substitution in the Ω loop (Gly214Glu) were compared to the parent enzyme, CMY-2, a widespread class C β -lactamase. In parallel with our microbiological characterization, the Gly214Glu substitution in CMY-32 reduced catalytic efficiency (k_{cat}/K_m) by 50-70% against “good” substrates (i.e., cephalothin) while increasing k_{cat}/K_m against “poor” substrates (i.e., cefotaxime). Additionally, CMY-32 was more susceptible to inactivation by sulfone β -lactamase inhibitors (i.e., sulbactam and tazobactam) than CMY-2. Timed electrospray ionization mass spectrometry (ESI-MS) analysis of the reaction of CMY-2 and CMY-32 with different substrates and inhibitors suggested that both β -lactamases formed similar intermediates during catalysis and inactivation. We next showed that the carbapenems (imipenem, meropenem, and doripenem) form long-lived acyl-enzyme intermediates and present evidence that there is β -lactamase-catalyzed elimination of the C₆ hydroxyethyl substituent. Furthermore, we discovered that the monobactam aztreonam and BAL29880, a new β -lactamase inhibitor of the monobactam class, inactivate CMY-2 and CMY-32 by forming an acyl-enzyme intermediate that undergoes elimination of -SO₃²⁻. Molecular modeling and dynamics simulations suggest that the Ω loop is more constrained in CMY-32 than CMY-2. Our model also proposes that Gln120 adopts a novel conformation in the active site while new interactions form between Glu214 and Tyr221, thus explaining increased cefotaxime hydrolysis. When docked in the active site, we observe that BAL29880 exploits contacts with highly conserved residues Lys67 and Asn152 in CMY-2 and CMY-32. These

*Corresponding author: Robert A. Bonomo, MD. Louis Stokes Cleveland Department of Veterans Affairs Medical Center, 10701 East Blvd., Cleveland, Ohio, 44106 Ph: 216-791-3800 x 4399; Fax: 216-231-3482; robert.bonomo@med.va.gov.

Supporting Information: Supporting Information is available free of charge and may be accessed at the following internet address (<http://pubs.acs.org>).

findings highlight: *i*) the impact of single amino acid substitutions on protein evolution in clinically important AmpC enzymes; and *ii*) the novel insights into the mechanisms by which carbapenems and monobactams interact with CMY-2 and CMY-32 β -lactamases.

The most common mechanism responsible for β -lactam resistance in Gram-negative bacteria is the production of β -lactamases (EC 3.5.2.6) (1). At present, there are more than 850 different β -lactamases described in nature (<http://www.lahey.org/Studies/>). As a family of proteins, β -lactamases are grouped into four main classes (i.e., A to D) based upon amino acid sequence homology (2). Classes A, C, and D employ serine in the active site as the reactive nucleophile, while class B β -lactamases use a single or pair of metal ions (Zn^{2+}) to catalyze the opening of the β -lactam ring. Both serine and metallo- β -lactamases make use of a strategically positioned water molecule to hydrolyze the lactam bond (3-5).

In general, class A, C, and D β -lactamases follow a three-step reaction mechanism that is represented as follows:

Here, E is the β -lactamase enzyme, S is the β -lactam substrate, E:S is the Henri-Michaelis complex, E-S is the acyl enzyme, and P is the inactivated β -lactam. The rate constants for each step are designated by k_1 , k_{-1} , k_2 , and k_3 .

Usually, class A and C β -lactamases are found in the *Enterobacteriaceae*. Acquired class A enzymes, often carried on mobile plasmids, are common in *Escherichia coli* and *Klebsiella pneumoniae*, are constitutively expressed and confer resistance to penicillins and narrow-spectrum cephalosporins (1). Class C, or AmpC-type β -lactamases (AmpCs), are encoded by chromosomal genes in many Gram-negative pathogens (e.g., *Citrobacter freundii*, *Enterobacter* spp., and *Pseudomonas aeruginosa*), are inducible, and confer resistance to narrow and extended-spectrum cephalosporins. Most recently, an increasing number of AmpC β -lactamase genes are being discovered on plasmids that also spread by horizontal transfer (6-8). Unlike class A enzymes, AmpCs are poorly inhibited by the commercially available β -lactamase inhibitors (clavulanate, sulbactam, and tazobactam) (7).

Among class C enzymes, CMY-type β -lactamases represent the most frequently detected plasmid encoded AmpCs (pAmpCs) (8-13). Unfortunately, we know little regarding the ability of clinically important CMY enzymes to hydrolyze different classes of β -lactam antibiotics (14-16). Furthermore, the activity of novel β -lactamase inhibitors against the CMY type pAmpCs has not yet been fully studied (17).

In class A β -lactamases, single amino acid substitutions (primarily, but not exclusively, at Ambler positions Gly238 on the b3 β -strand and Arg164 or Asp179 in the Ω loop) remodel a "broad-spectrum" β -lactamase into an extended-spectrum β -lactamase (ESBL) (18-20). The bacterium possessing a class A ESBL becomes resistant to ceftazidime and/or cefotaxime. However, these changes come at a "price" (18, 21). Among class A ESBLs, one uniformly

detects a decrease in penicillin resistance (lower MICs) and an increase in the susceptibility to β -lactamase inhibitors and carbapenems (18, 21).

Herein, we analyzed the biochemical properties of a novel CMY-2-like pAmpC (i.e., CMY-32) cloned from an *E. coli* isolate found in the clinic. In comparison with CMY-2 (wild type, WT), we determined that CMY-32 possesses a single amino acid substitution in the Ω loop (i.e., Gly214Glu) resulting in significant changes in the resistance phenotype and hydrolytic profile (making this β -lactamase an “extended-spectrum type”). We also evaluated the mechanisms by which carbapenems, a monobactam antibiotic aztreonam, and the novel monobactam derivative, BAL29880, react and inhibit CMY-2 and CMY-32 enzymes. Our data reveal the impact of single amino acid substitutions on protein evolution in this emerging family of resistance enzymes and highlight the manner in which carbapenems and monobactams inactivate class C enzymes.

Materials and Methods

Detection and cloning of *bla*_{CMY} genes and expression in *E. coli* DH10B cells

We used two clinical *E. coli* isolates to obtain the constructs employed in this study. The first strain carried the *bla*_{CMY-2} gene and was detected in 2002 in Cleveland (OH), whereas the second strain possessed the *bla*_{CMY-32} gene and was discovered in 2006 in Pittsburgh (PA) (22, 23).

Specific primers were used to amplify both *bla*_{CMY} genes (For: 5'-CCGGACACCTTTTGTCTTT-3'; Rev: 5'-TATCCTGGGCCTCATCGTCAGTTA-3'). As previously reported, sequence analysis demonstrates that CMY-32 (GeneBank accession number: EU496815) possesses a single amino acid substitution (i.e., Gly214Glu) compared to CMY-2 (GeneBank accession number: EF406116) (23). Each *bla*_{CMY} gene was cloned into the pBC SK(-) phagemid vector (Stratagene, La Jolla, CA) as previously described (24). Commercially purchased electrocompetent *E. coli* DH10B cells (Invitrogen Corporation, Carlsbad, CA) were electroporated with the ligation mixture. We selected strains expressing the *bla*_{CMY} genes on Luria-Bertani (LB) agar plates containing chloramphenicol (20 μ g/mL) and ampicillin (100 μ g/mL).

Phenotypic characterization

Minimum inhibitory concentrations (MICs) for the following β -lactams were obtained using the Etest method (AB Biodisk, Solna, Sweden) on cation-adjusted Mueller-Hinton agar (BBL, Becton Dickinson, Sparks, MD): ampicillin; ampicillin-sulbactam; amoxicillin-clavulanate; piperacillin; piperacillin-tazobactam; cefoxitin; cefotaxime (CTX); ceftazidime; cefepime; aztreonam (ATM); imipenem; and meropenem (see Figure 1 for the chemical structures of several of these compounds).

The inhibitory activity of BAL29880 (Basilea Pharmaceutica International Ltd, Basel, Switzerland) was evaluated by microdilution in cation-adjusted Mueller-Hinton broth, according to the Clinical Laboratory Standard Institute criteria (25). This inhibitor was also combined with ampicillin and piperacillin (Sigma Chemical Co., St. Louis, MO) at a constant concentration of 4 μ g/mL. The following American Type Culture Collection

(ATCC) control strains were used as controls: *E. coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 27853, and *K. pneumoniae* ATCC 700603.

β -Lactamase expression

Polyclonal anti-CMY-2 antibodies were produced by Genosys Biotech, Inc. (Woodlands, TX) and purified using a HiTrapTM Protein G HP column (GE Healthcare Bio-Sciences AB, Uppsala, Sweden) as previously reported (26). *E. coli* DH10B isolates containing *bla*_{CMY-2} and *bla*_{CMY-32} were grown in LB broth to an OD₆₀₀ of 0.8. Preparation of samples, immunoblotting, and assessment of β -lactamase expression were performed using a method previously described (26).

β -lactamase purification

Each *E. coli* DH10B strain containing the *bla*_{CMY} gene expressed in the pBC SK(-) vector was grown overnight at 37°C in 4 liters of SOB broth supplemented with ampicillin (100 μ g/mL) and chloramphenicol (20 μ g/mL). Cells were then pelleted and resuspended in 50 ml of 50 mM Tris buffer (pH 7.4). β -lactamases were released using lysozyme, EDTA (Sigma Chemical Co.), and DNase (Promega, Madison, WI) according to a previously established protocol (27).

We performed preparative isoelectric focusing (pIEF) using a Sephadex G100 gel matrix and commercially prepared ampholines (pH 3.5 to 10.0; Amersham Biosciences, Piscataway, NJ). The parameters used for pIEF were described previously (28). After running pIEF overnight, areas of the gel demonstrating β -lactamase activity by nitrocefin (NCF) overlay were cut from the gel, eluted with PBS (pH 7.4) on polyethylene glycol columns (Amersham Biosciences), and concentrated using an Amicon Ultra-4 concentrator (MW cutoff = 10,000; Millipore, Carrigtwohill, Co Cork, Ireland). Protein concentrations were determined by spectrophotometric assay (Bio-Rad Laboratories, Hercules, CA) using bovine serum albumin as a standard.

We further purified CMY-2 and CMY-32 β -lactamases to homogeneity using size exclusion chromatography with a Pharmacia ÄKTA purifier system (GE Healthcare Bio-Sciences AB). We employed a HiLoad 16/60 Superdex 75 column (GE Healthcare Bio-Sciences AB) and eluted with PBS (pH 7.4). We assessed the purity of β -lactamases by running the samples in sodium dodecyl-sulfate-polyacrylamide gels. Gels were stained using Coomassie blue R-250 (Fisher, Pittsburgh, PA), and protein molecular weight was estimated using pre-stained low-molecular-weight standards (Bio-Rad Laboratories).

Kinetic experiments

Based upon the reaction scheme outlined in Eq. 1, we measured steady-state kinetic parameters (i.e., K_m , k_{cat} , and k_{cat}/K_m) by continuous assays at room temperature. The Michaelis constants were determined using nonlinear least squares fit of the data (Henri-Michaelis equation, Eq.2) employing the program Enzfitter (Biosoft Corp., Ferguson, MO) (29, 30) where:

$$v = V_{\max}[S]/(K_m + [S]) \quad (\text{Eq. 2})$$

We used an Agilent™ 8453 diode array spectrophotometer with a 1-cm path length and determined each parameter in triplicate in PBS buffer (pH 7.4). Initial velocity measurements were obtained for the following “good substrates” (as defined by $k_{\text{cat}} \geq 2 \text{ s}^{-1}$) (31): NCF ($\epsilon_{482} = 17,400 \text{ M}^{-1} \text{ cm}^{-1}$), cephaloridine ($\epsilon_{260} = 10,200 \text{ M}^{-1} \text{ cm}^{-1}$), cephalothin ($\epsilon_{262} = 7,660 \text{ M}^{-1} \text{ cm}^{-1}$), and cefoxitin ($\epsilon_{260} = 7,700 \text{ M}^{-1} \text{ cm}^{-1}$). For “poor substrates” ($k_{\text{cat}} < 2 \text{ s}^{-1}$ or non-measurable; i.e., cefotaxime, $\epsilon_{260} = 7,500 \text{ M}^{-1} \text{ cm}^{-1}$), the apparent K_m values were obtained as competitive inhibition constants (K_i) in the presence of NCF.

To characterize the β -lactamase inhibitors (clavulanic acid, sulbactam, tazobactam and BAL29880) we measured the first-order rate constant for enzyme and inhibitor complex inactivation (k_{inact}) by monitoring the reaction time courses in the presence of inhibitor and NCF under steady state conditions. A fixed concentration of enzyme (i.e., $E = 1.6 \text{ nM}$ for CMY-2 or 3.1 nM for CMY-32), $100 \mu\text{M}$ NCF, and increasing concentrations of inhibitor (I) were used in each assay. The k_{obs} values were determined using a non-linear least squares fit of the data employing Origin 7.5® (OriginLab Co., Northampton, MA):

$$A = A_0 + v_f t + (v_0 - v_f)[1 - \exp(-k_{\text{obs}} t)]/k_{\text{obs}} \quad (\text{Eq. 3})$$

Here, A is absorbance, v_0 (expressed in variation of absorbance per unit time) is initial velocity, v_f is final velocity, and t is time (32). Each k_{obs} was plotted vs I and fit to determine k_{inact}

$$k_{\text{obs}} = (k_{\text{inact}} I)/(k_{\text{inact}} + I) \quad (\text{Eq. 4})$$

In a similar manner to our determination of K_i for poor substrates, we determined K_i of each inhibitor. We corrected the K_i values for the presence of NCF using Eq. 5.

$$K_i \text{ corrected} = K_i \text{ observed} / (1 + [S]/K_{m \text{ NCF}}) \quad (\text{Eq. 5})$$

We used Eq. 6 to calculate the percentage of the standard error (SE):

$$\text{SE of } k_{\text{cat}}/K_m \text{ or } k_{\text{inact}}/K_i = \sqrt{[(\text{SE of } k_{\text{cat}} \text{ or } k_{\text{inact}}/k_{\text{cat}} \text{ or } k_{\text{inact}})^2 + (\text{SE of } K_m \text{ or } K_i/K_m \text{ or } K_i)^2]} \quad (\text{Eq. 6})$$

Ultraviolet difference (UVD) spectroscopy

UVD spectra of absorption were obtained for CMY-2 and CMY-32 (both 8 nM) reacted with CTX ($8 \mu\text{M}$) in 10 mM PBS (pH of 7.4). Wavelengths 200 to 500 nm were analyzed using the Agilent™ 8453 spectrophotometer for 1 hour at room temperature.

Electrospray Ionization (ESI) Mass Spectrometry (MS)

We performed timed ESI-MS of each CMY β -lactamase alone or in combination with β -lactam antibiotics or β -lactamase inhibitors on a Q-STAR XL quadrupole time-of-flight mass spectrometer (Applied Biosystems, Framingham, MA) equipped with a nanospray source as described (33). Two incubation times (i.e., 1 and 15 min) were analyzed for each β -lactamase inhibitor tested, whereas for the carbapenems only a 15 min incubation was performed. We desalted the reaction mixtures using C₁₈ ZipTips (Millipore, Billerica, MA). The CMY-2 or CMY-32 β -lactamase sample was then diluted with 50% acetonitrile, 0.1% trifluoroacetic acid to a concentration of 10 μ M. We next infused this protein solution at a rate of 0.5 μ l/min, and spectral data were collected for 2 min. Spectra were deconvoluted using the Analyst program (Applied Biosystems). Notably, all data reported in the Results section should be considered with a $\pm$ 3 Da standard error.

Molecular modeling

To generate a model of the CMY-32 enzyme we used the crystal structure of CMY-2 (Protein Data Bank, 1ZC2) and Discovery Studio (DS) 2.1 software (Accelrys, San Diego, CA). The protein was prepared for molecular dynamics simulations (MDS) by adding hydrogen atoms and setting the pH at 7.4. The CMY-32 model was created using the Build Module of DS by inserting a substitution at position 214 (i.e., Gly to Glu).

First, the crystallographic waters were removed and the two β -lactamase models were immersed in a water box (i.e., 7 Å from any face of the box) using explicit periodic boundary conditions (PBCs). Then, the complexes were minimized in several steps, using Steepest Descent and Conjugate Gradient algorithms to reach the minimum convergence (i.e., 0.02 kcal/mol $\times$ Å) after 10,000 iterations). All energy minimizations and MDSs were performed using CHARMM force-field parameters. We employed the Particle Mesh Ewald (PME) method to treat long-range electrostatic interactions. Bonds that involved hydrogen atoms were constrained with SHAKE algorithm. Following equilibration, two separate (2fs) MDS (Heating/Cooling and Production), at constant pressure and temperature (300°K), were carried out for the CMY-2 and CMY-32 models. The trajectories collected after the production steps were analyzed and the conformations with the minimum energies were chosen. The CMY-2 and CMY-32 models were superimposed.

The model of the pre-acylation complex of the CMY-32 and CTX was developed using the Flexible Docking Module of DS 2.1. The side-chains of the active site amino acids (i.e., Ser64, Glu120, Asn152, Tyr221, Tyr150, Lys67, and Ser318) were allowed to move during docking. The low energy CTX conformations were docked into the active site of the CMY-32 model using LibDock (34). In the presence of CTX, the active site side-chain residues were refined using the ChiRotor algorithm (35). A final simulated annealing and energy minimization of each CTX conformation was performed using CDOCKER (34). The 10 resulting conformations of CTX were manually analyzed for hydrogen bonds, steric clashes and distances between atoms. The energies were evaluated and the most favorable conformation was chosen. The complex that was created was further energy minimized.

A model of the acyl-enzyme complex of the CMY-2 enzyme with BAL29880 was also constructed. The chemical structure of BAL29880, built using the Fragment Builder tools of DS 2.1, was minimized using a Standard Dynamics Cascade protocol. CDOCKER algorithm was employed to dock BAL29880 in the active site of CMY-2 β -lactamase (34). The BAL29880 placement in the active site was specified using a binding site sphere. Random ligand conformations were generated from the initial structure through high temperature MDS. Grid-based simulated annealing followed by force field minimization refined the generated conformations. Starting with the most energetically favorable conformation, we created and energy minimized the acyl-enzyme complex.

Results

Phenotype characterization

In order to measure the impact of a single amino acid substitution on the hydrolytic activity of the CMY-2 and CMY-32 β -lactamases in *E. coli*, we performed MICs in a uniform genetic background. As shown in Table 1, both CMY β -lactamases expressed in *E. coli* DH10B confer high-level resistance to ampicillin, amoxicillin-clavulanate, piperacillin and ceftazidime (MICs > 256 μ g/mL), whereas piperacillin-tazobactam, cefepime, and carbapenems demonstrated MICs in the susceptible ranges (< 6 μ g/mL). Noticeably, the strain expressing CMY-32 was more susceptible to ampicillin-sulbactam and ceftazidime than strain harboring CMY-2 (36). In contrast, the MIC for CTX was significantly higher for CMY-32 than CMY-2 when tested in *E. coli* (64 vs. 16 μ g/mL). We observed the same trend when testing ATM (MIC of 12 vs. 6 μ g/mL). Although a slight increase of MIC was observed for both *E. coli* clones containing CMY-2 or CMY-32 when compared to the *E. coli* DH10B, full resistance to cefepime was not established (Table 1).

To further characterize the phenotype of CMY-32 in *E. coli* DH10B, we also tested the activity of the novel monobactam inhibitor and compared it to ATM. BAL29880 alone did not show activity against *E. coli* DH10B, including those expressing the *bla*_{CMY} genes (MICs > 32 μ g/mL). However, in combination with ampicillin or piperacillin, BAL29880 lowered MICs considerably (i.e., < 4 μ g/mL) for strains expressing CMY-2 or CMY-32 (Table 1).

β -Lactamase expression

In order to ensure that the differences in MICs between the *E. coli* DH10B producing *bla*_{CMY-2} and that producing *bla*_{CMY-32} were not due to changes in expression levels of the proteins, we performed immunoblots using a specific anti-CMY-2 polyclonal antibody. Steady-state expression levels of CMY-2 and CMY-32 in *E. coli* DH10B showed that the strain expressing CMY-2 produced similar quantities of β -lactamase as the strain harboring CMY-32 pAmpC (data not shown).

Kinetics and UVD spectroscopy

Steady-state kinetic parameters revealed that CMY-32 possessed significantly less catalytic efficiency (k_{cat}/K_m) for hydrolyzing the narrow-spectrum cephalosporins (NCF, cephaloridine, and cephalothin) compared to CMY-2 (Table 2). These results were mainly

due to reductions in k_{cat} rather than K_{m} . A decrease in turnover was also seen with cefoxitin, a 7- α -methoxy cephalosporin (14, 37), (Table 2).

The turnover of CTX, an oxyimino amino-thiazole cephalosporin that is regarded as a “poor substrate”, is often difficult to measure due to its low k_{cat} (14, 31). Here, we followed the hydrolysis of CTX by CMY-2 and -32 using UVD absorption spectra. In the Supporting Information (Figure 1 supplement) we show the increase in hydrolysis of CTX by CMY-32 vs. CMY-2 (see $\lambda_{\text{max}} = 228 \text{ nm}$). Consistent with MICs, cefepime, an extended-spectrum cephalosporin that is also “poor substrate” for class C enzymes, was poorly hydrolyzed by CMY-2 and CMY-32 ($K_{\text{m}} > 100 \mu\text{M}$, k_{cat} not measurable; Table 2) (14, 37).

For ATM and imipenem, we did not observe hydrolysis by either CMY-2 or CMY-32 under saturating conditions (Table 2). To understand why this occurred, we employed ESI-MS (see below) and observed that ATM and imipenem formed long-lived covalent complexes with both β -lactamases. In competition experiments with NCF, we observed that the affinity for CMY-2 and CMY-32 was greater than for any substrate tested ($K_{\text{i}} \sim 600 \text{ nM}$)

The commercially available β -lactamases inhibitors (tazobactam, sulbactam, and clavulanate) were not very effective at inhibiting the two CMY enzymes (Table 3). Consistent with our MIC determinations showing increase in susceptibility to ampicillin-sulbactam and piperacillin-tazobactam, these inhibitors possessed lower K_{i} (by at least 40%) against CMY-32 β -lactamase than CMY-2 (Table 3). The K_{i} of BAL29880, the novel monobactam, for CMY-2 and CMY-32 was lower than the three commercially available inhibitors. Furthermore, our MIC data indicated that BAL29880 requires a β -lactam partner for full efficacy (MIC $\sim 2 \mu\text{g/mL}$).

Additionally, we observed a notable finding regarding the kinetics of inactivation of CMY-2 and CMY-32 with tazobactam. Compared to the other inhibitors, tazobactam demonstrates a k_{inact} value at least 3-fold greater than sulbactam, clavulanate or BAL29880. This more rapid inactivation constant coupled with greater affinity (lower K_{i}) is also consistent with the MIC results.

ESI-MS

We employed ESI-MS to gain insight into the nature of the intermediates formed in the inactivation of CMY-2 and CMY-32 β -lactamases. To begin, we established that the molecular weight (MW) of CMY-2 is 39,854 Da, whereas that of CMY-32 is 39,926 Da, consistent with the predicted molecular weights of the proteins (Figure 2).

Since carbapenems are known to be poorly hydrolyzed by both class A and C β -lactamases, we first sought to determine if imipenem, meropenem and doripenem would form a stable acyl-enzyme adduct with CMY-2 and CMY-32 β -lactamases (38). At 15 minutes, the inactivation of CMY-2 and CMY-32 with imipenem in a ratio of 1:1,000 produced adducts of +299 Da, corresponding to the molecular weight of imipenem (Figure 2). However, when the ratio of E:I was reduced to 1:20, an additional minor adduct with a mass defect of 44 Da was seen (data not shown). We observed the same phenomenon with meropenem and doripenem (Figure 2).

We next inactivated CMY-2 and CMY-32 with tazobactam, sulbactam, ATM, and BAL29880. In the course of our analysis, it became evident that the ESI-MS determinations obtained after 1 and 15 min incubations were equivalent. Therefore, in Figure 3 we present only one of the timed reactions.

As shown in Figure 3, inactivation of both β -lactamases with the sulfones yielded products of +72 and +87 Da. The fragmentation of sulbactam to yield similar adducts as tazobactam supports a common mechanism of inactivation by sulfone inhibitors. As stated above, we also demonstrated that both ATM and BAL29880 form acyl-enzyme species by ESI-MS. As with the carbapenems, we detected two species (Figure 3), one 80 Da less than the other. This suggested to us that there is fragmentation of ATM and BAL29880 as is seen with the carbapenems.

Discussion

In the present work, we studied the biochemical characteristics of CMY-2 and its variant, CMY-32. Our analysis points to the distinctive properties that define the “extended-spectrum” phenotype of CMY-32. To emphasize the singular features of this “class C ESBL”, we draw comparisons to the well-studied class A ESBLs. Our work highlights the consequence of “natural protein engineering” (i.e. previous antibiotic selection pressure) on substrate specificity (i.e., emergence of CTX-resistant phenotype). Next, we describe the reaction intermediates observed when these carbapenems, sulfones, and monobactams react with CMY-2 and CMY-32 β -lactamases.

Impact of Gly214Glu substitution on phenotype

The phenotype we observed because of the Gly214Glu substitution is very reminiscent of class A ESBLs in *E. coli* and illustrates that in class C β -lactamases, a rather specific substitution located in the Ω loop that alters susceptibility and resistance to many β -lactams. Most strikingly, resistance to CTX and ATM is increased for *E. coli* bearing CMY-32 compared to CMY-2. In contrast, the class C CMY-32 β -lactamase expressed in *E. coli* DH10B resulted in a lower level of resistance to cefoxitin, as compared to CMY-2. Moreover, there is increased susceptibility to β -lactam/ β -lactamase inhibitor combinations and carbapenems. These properties are the “clinical hallmarks” that alert microbiology laboratories to the presence of ESBLs in *E. coli*, *K. pneumoniae*, and *Proteus* spp. Interestingly, there are many class A ESBLs of the TEM, SHV families that possess substitutions in the Ω loop that behave similarly (18, 21).

Impact of Gly214Glu on turnover of good and poor substrates

The steady-state kinetic parameters of both CMY-2 and CMY-32 pAmpCs were consistent with the antimicrobial susceptibility profiles. First, we saw an overall reduction in k_{cat}/K_m for first-generation cephalosporins (i.e., cephaloridine and cephalothin), substrates that are generally “preferred” by class C enzymes (7). Likewise, cefoxitin, a 7- α -methoxy cephalosporin, which is usually hydrolyzed by AmpCs, has a 45% reduction in k_{cat}/K_m when hydrolyzed by CMY-32. Most strikingly, CMY-32 exhibits the ability to hydrolyze CTX

($k_{\text{cat}}/K_{\text{m}} = 0.22 \mu\text{M}^{-1} \text{s}^{-1}$ for CMY-32 compared to not measurable for CMY-2, see Table 2 and Supporting Information, Figure 1).

Consistent with the kinetic behavior of other class C enzymes, CMY-2 and CMY-32 hydrolyzes imipenem poorly (k_{cat} not measurable). Following the reaction with timed ESI-MS, we see that imipenem accumulates in the unfragmented acyl-enzyme form for both CMY pAmpCs (Figure 2). We observed this phenomenon only when the ratio of imipenem to CMYs was 1,000:1. Reducing the ratio to 20:1, we noted identical patterns with all three carbapenems. After the covalent adduct is formed, there is fragmentation to an acyl-enzyme species that is 44 Da less (Figure 2). Similar results were observed with meropenem and doripenem. As previously observed for a chromosomal class A β -lactamase expressed by *Mycobacterium tuberculosis* and the ADC-7 P-lactamase of *Acinetobacter baumannii*, these results suggest that, after β -lactam ring opening, the covalently bound carbapenem inactivation pathway splits between: i) direct hydrolysis with formation of unfragmented acyl-enzyme; ii) cleavage of the hydroxyethyl substituent (39, 40). Investigations are underway to define the impact of this finding and identify of the residues that may facilitate this process.

Impact of Gly214Glu on β -lactamase inhibitors

For CMY-32, the Gly214Glu substitution lowers the K_{i} for all clinically used inhibitors (the most dramatic change was for tazobactam and sulbactam), and significantly increases the k_{inact} , a property that is also very common among class A ESBLs (1). Timed ESI-MS showed that both sulfone inhibitors follow a pattern of inhibition that generates the same postulated intermediates for both AmpCs (i.e., $m/z +88$, the hydrated aldehyde and $m/z +72$, the aldehyde). Originally, this was described for class A β -lactamases and CMY-2 (17, 41). As sulbactam is regarded as a very poor inhibitor of class C enzymes ($K_{\text{i}} > 100 \mu\text{M}$), we were surprised to see adducts similar to those formed by tazobactam, a more effective inhibitor. Therefore, we maintain that even these class C serine β -lactamases pass through the imine and enamine transitional stages and that CMY-32 may form certain intermediates more readily than CMY-2. We stress that these findings also shed significant insight into why tazobactam serves as a good inhibitor of CMY β -lactamases, albeit unrecognized as such. Our data show that tazobactam possesses a first order rate constant of inactivation (k_{inact}) for CMY-32 that was three fold greater than commercially available inhibitors. Together with the lower K_{i} , the more rapid inactivation adds to the potency of the β -lactam/ β -lactamase inhibitor combinations.

Aztreonam and BAL29880, a monobactam substrate and a monobactam inactivator

The most intriguing observations were made with ATM and BAL29880. We generally regard ATM as a “poor substrate” and BAL29880 is an inhibitor. As result of the Gly214Glu substitution, CMY-32 expressed in *E. coli* has increased MICs against aztreonam. In contrast, BAL29880 does not appear to possess antibacterial activity (MICs $> 32 \mu\text{g/mL}$ for *E. coli* with CMY-2 and CMY-32 β -lactamases). However, when combined with ampicillin or piperacillin, BAL29880 lowers MICs at least 6-twofold dilutions (256 to $4 \mu\text{g/mL}$), a range similar to ATM alone against CMY-2 and CMY-32 (Table 1).

To understand why two monobactams behave so differently, we performed direct competition experiments using NCF as an indicator substrate and studied the reactions using timed ESI-MS. Here, ATM showed 30- to 55-fold greater affinity for CMY-2 and CMY-32 when compared to BAL29880 (Table 2 and 3). This compelled us to explore possible mechanisms by which ATM and BAL29880 interact with CMY-2 and CMY-32. Timed ESI-MS analysis demonstrated that BAL29880 and ATM follow similar pathways of inhibition against both CMY-2 and CMY-32 pAmpCs (Figure 3). Both monobactams acylate the CMY β -lactamases (i.e., $+411$ Da for BAL29880 or $+435$ Da for ATM) and form long-lived acyl enzymes. These findings were reminiscent of Oefner et al. who demonstrated that ATM, a poor substrate, acylates the active site Ser64 of the *C. freundii* AmpC β -lactamase (PDB, 1FR1), an enzyme with 97% amino acid homology to CMY-2 and CMY-32 (42). Therefore, we anticipated the presence of the full adduct added to CMY-2 and CMY-32 when ATM and BAL29880 were reacted with these class C cephalosporinases (42, 43). Unexpectedly, we discovered that a second adduct was also formed. The MW of the second adduct is approximately 70 Da less than the first, suggesting that both monobactams undergo fragmentation after acylation of the CMY β -lactamase (Figure 3). Mechanistically, the addition of water to the acyl-enzyme species would allow hydrolysis of the sulfonamide to yield the free sulfate (Figure 4a and 4b).

Molecular models

In order to explain the basis of affinity and specificity of the Gly214Glu substitution in the Ω loop of CMY-2, we constructed three molecular models.

In the first model, we superimposed the energy minimized structures of CMY-32 and CMY-2 (rmsd = 1.0 Å) using backbone atoms and compared the conformation of the active sites paying particular attention to the Ω loop region (Val211 to Val223) (Figure 5a and 5b). The analysis of the trajectories generated after MDS showed an average potential movement of 1-3 Å in the region of Gly214Glu substitution. Our model shows that there is also significant change in the position of the Tyr221 side chain in the CMY-32 β -lactamase (Figure 5b). As result of the Gly to Glu substitution, Tyr221 likely moves within hydrogen bond distance to Glu214 (1.9 Å, Figure 5b). Our interpretation is that this newly positioned hydrogen bond limits the overall flexibility and mobility of the Ω loop. Additionally, we observed that because of the Glu214-Tyr221 hydrogen bond, Gln120 adopts a new conformation in which the side chain is flipped more than 3.5 Å (see Figure 6).

In our second model, we present an explanation of why CTX is hydrolyzed by CMY-32 β -lactamase. Using the energy-minimized structure of CMY-32, we docked the carbonyl of the 4-member ring of CTX in the oxyanion hole (Figure 6). Because of the repositioning of Gln120 in CMY-32, we see a new hydrogen bond contact between Glu120:O and the terminal amide of the R1 side chain of CTX. In this model, the C4 carboxylate is in hydrogen bond distance with Arg349 and the carbonyl oxygen of the R2 side chain makes favorable interactions with Asn346 (Figure 1 and Figure 6).

Lastly, we modeled the acyl-enzyme complex of BAL29880 without the $-\text{SO}_3^{-2}$ group (Figure 7). In this energy-minimized model, we docked the carbonyl of BAL29880 into the oxyanion hole of CMY-2 (see Figure 4b). In this conformation, the oxygen is

advantageously positioned to form hydrogen bonds with the protonated Lys67. Rotating the acyl-amide bond, we observe a stabilizing hydrogen bond interaction with Asn152 (Figure 9) (44). Thus, the inhibitor BAL29880, like ATM, adopts an arrangement in the acyl-enzyme that primarily exploits contacts with conserved residues (Lys67 and Asn152) of the active site. Despite these similarities, ATM and BAL29880 behave very differently (one a poor substrate and the other an inhibitor).

Conclusions

We emphasize four important findings arising from this work. Firstly, we demonstrate how a single amino acid change in the Ω loop of CMY-2 can considerably alter MICs, modify the spectrum of substrate hydrolysis, and vary the efficacy of β -lactamase inhibitors. In this context, we argue that in “class C ESBLs”, movement of the Ω loop and repositioning of key amino acids may be the structural explanation of the phenotypic and kinetic properties. In contrast to the findings in class A ESBLs, the major impact of this substitution is to constrain rather than expand or disorder the Ω loop and active site cavity (20, 45, 46). Here, the inherently larger space in the WT CMY-2 enzyme active site, that is optimally configured to hydrolyze narrow-spectrum cephalosporin substrates, appears to be tailored in CMY-32 to also accommodate CTX and ATM.

Secondly, comparing our analysis to other class C β -lactamases that are also considered “extended-spectrum” types, we see that amino acid substitutions at multiple positions, additions, and deletions all give rise to this ESBL phenotype in class C β -lactamases (15, 47, 48). Among class A enzymes, the number and types of changes able to confer ESBL phenotype are limited (18). Hence, we predict that there will be even more opportunities in class C enzymes to evolve novel catalytic abilities by natural protein engineering.

Thirdly, we propose a novel mechanism for the interaction of class C β -lactamases with monobactam substrates and inhibitors. One wonders if the elimination of the $-\text{SO}_3^-$ group from both ATM and the BAL29880 facilitates hydrolysis or serves to further stabilize the acyl-enzyme complex and result in a long lived terminal species. We also do not know if this phenomenon is unique to CMY type β -lactamases (49). In NMC-A, a similar pattern was seen by Mourey et al studying monobactam inhibitors (49).

Lastly, we observed in this class C serine β -lactamase that the formation of the acyl-enzyme intermediates of carbapenems is accompanied by fragmentation of the substrate with loss of the hydroxyethyl substituent at the C6 position. This seems to be a property of CMY- and ADC-type class C β -lactamases.

As we contemplate these major conclusions, we are certain that the complexity of β -lactamases and the mechanisms responsible for β -lactamase mediated resistance will continue to pose significant challenges to our understanding of the nature and the evolution of protein structure-function relationships in the years ahead. Our study into the mechanism of inactivation by monobactams and carbapenems form templates for the future rationale design of new inhibitors of this class of enzymes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding. The National Institutes of Health (grant RO1-AI063517 to RAB), the Veterans Affairs Merit Review Program (RAB and MJS), the Geriatric Research Education and Clinical Care VISN 10 (RAB), and NIH training grant (T32 AI007333 to YD) supported this work. A Joint Fellowship Grant from the National Foundation of Infectious Diseases and the Infectious Diseases Society of America supported YD.

References

1. Paterson DL, Bonomo RA. Extended-spectrum β -lactamases: a clinical update. Clin Microbiol Rev. 2005; 18:657–686. [PubMed: 16223952]
2. Ambler RP. The structure of β -lactamases. Philos Trans R Soc Lond B Biol Sci. 1980; 289:321–331. [PubMed: 6109327]
3. Bebrone C. Metallo- β -lactamases (classification, activity, genetic organization, structure, zinc coordination) and their superfamily. Biochem Pharmacol. 2007; 74:1686–1701. [PubMed: 17597585]
4. Leonard DA, Hujer AM, Smith BA, Schneider KD, Bethel CR, Hujer KM, Bonomo RA. The role of OXA-1 β -lactamase Asp(66) in the stabilization of the active-site carbamate group and in substrate turnover. Biochem J. 2008; 410:455–462. [PubMed: 18031291]
5. Kuzin AP, Nukaga M, Nukaga Y, Hujer AM, Bonomo RA, Knox JR. Structure of the SHV-1 β -lactamase. Biochemistry. 1999; 38:5720–5727. [PubMed: 10231522]
6. Pai H, Kang CI, Byeon JH, Lee KD, Park WB, Kim HB, Kim EC, Oh MD, Choe KW. Epidemiology and clinical features of bloodstream infections caused by AmpC-type- β -lactamase-producing *Klebsiella pneumoniae*. Antimicrob Agents Chemother. 2004; 48:3720–3728. [PubMed: 15388426]
7. Jacoby GA. AmpC β -lactamases. Clin Microbiol Rev. 2009; 22:161–182. [PubMed: 19136439]
8. Sidjabat HE, Paterson DL, Qureshi ZA, Adams-Haduch JM, O'Keefe A, Pascual A, Rodriguez-Bano J, Doi Y. Clinical Features and Molecular Epidemiology of CMY-Type β -Lactamase-Producing *Escherichia coli*. Clin Infect Dis. 2009; 48:739–744. [PubMed: 19187027]
9. Deshpande LM, Jones RN, Fritsche TR, Sader HS. Occurrence of plasmidic AmpC type β -lactamase-mediated resistance in *Escherichia coli*: report from the SENTRY Antimicrobial Surveillance Program (North America, 2004). Int J Antimicrob Agents. 2006; 28:578–581. [PubMed: 17112706]
10. Li Y, Li Q, Du Y, Jiang X, Tang J, Wang J, Li G, Jiang Y. Prevalence of plasmid-mediated AmpC β -lactamases in a Chinese university hospital from 2003 to 2005: first report of CMY-2-Type AmpC β -lactamase resistance in China. J Clin Microbiol. 2008; 46:1317–1321. [PubMed: 18305137]
11. Empel J, Baraniak A, Literacka E, Mrowka A, Fiett J, Sadowy E, Hryniewicz W, Gniadkowski M. Molecular survey of β -lactamases conferring resistance to newer beta-lactams in *Enterobacteriaceae* isolates from Polish hospitals. Antimicrob Agents Chemother. 2008; 52:2449–2454. [PubMed: 18458126]
12. Luzzaro F, Brigante G, D'Andrea MM, Pini B, Giani T, Mantengoli E, Rossolini GM, Toniolo A. Spread of multidrug-resistant *Proteus mirabilis* isolates producing an AmpC-type β -lactamase: epidemiology and clinical management. Int J Antimicrob Agents. 2009; 33:328–333. [PubMed: 19095415]
13. Hanson ND, Moland ES, Hong SG, Propst K, Novak DJ, Cavalieri SJ. Surveillance of community-based reservoirs reveals the presence of CTX-M, imported AmpC, and OXA-30 β -lactamases in urine isolates of *Klebsiella pneumoniae* and *Escherichia coli* in a U.S. community. Antimicrob Agents Chemother. 2008; 52:3814–3816. [PubMed: 18663030]

14. Bauvois C, Ibuka AS, Celso A, Alba J, Ishii Y, Frere JM, Galleni M. Kinetic properties of four plasmid-mediated AmpC β -lactamases. *Antimicrob Agents Chemother*. 2005; 49:4240–4246. [PubMed: 16189104]
15. Kim JY, Jung HI, An YJ, Lee JH, Kim SJ, Jeong SH, Lee KJ, Suh PG, Lee HS, Lee SH, Cha SS. Structural basis for the extended substrate spectrum of CMY-10, a plasmid-encoded class C β -lactamase. *Mol Microbiol*. 2006; 60:907–916. [PubMed: 16677302]
16. Doi Y, Shibata N, Shibayama K, Kamachi K, Kurokawa H, Yokoyama K, Yagi T, Arakawa Y. Characterization of a novel plasmid-mediated cephalosporinase (CMY-9) and its genetic environment in an *Escherichia coli* clinical isolate. *Antimicrob Agents Chemother*. 2002; 46:2427–2434. [PubMed: 12121914]
17. Bonomo RA, Liu J, Chen Y, Ng L, Hujer AM, Anderson VE. Inactivation of CMY-2 β -lactamase by tazobactam: initial mass spectroscopic characterization. *Biochim Biophys Acta*. 2001; 1547:196–205. [PubMed: 11410275]
18. Gniadkowski M. Evolution of extended-spectrum β -lactamases by mutation. *Clin Microbiol Infect*. 2008; 14(1):11–32. [PubMed: 18154525]
19. Orenica MC, Yoon JS, Ness JE, Stemmer WP, Stevens RC. Predicting the emergence of antibiotic resistance by directed evolution and structural analysis. *Nat Struct Biol*. 2001; 8:238–242. [PubMed: 11224569]
20. Ibuka A, Taguchi A, Ishiguro M, Fushinobu S, Ishii Y, Kamitori S, Okuyama K, Yamaguchi K, Konno M, Matsuzawa H. Crystal structure of the E166A mutant of extended-spectrum β -lactamase Toho-I at 1.8 Å resolution. *J Mol Biol*. 1999; 285:2079–2087. [PubMed: 9925786]
21. Page MG. Extended-spectrum β -lactamases: structure and kinetic mechanism. *Clin Microbiol Infect*. 2008; 14(Suppl 1):63–74. [PubMed: 18154529]
22. Huyen CM, Hujer AM, Hujer KM, Marshall SH, Carias L, Toltzis P, Rice LB, Bonomo RA. A clinical strain of *Escherichia coli* possessing CMY-2 plasmid-mediated amp C β -lactamase: an emerging concern in pediatrics? *Microb Drug Resist*. 2002; 8:329–333. [PubMed: 12523630]
23. Endimiani A, Doi Y, Bethel CR, Taracila M, Hujer AM, Adams-Haduch JD, O'Keefe A, Yamane K, Helfand MS, Paterson DL, Bonomo RA. CMY-32, a novel β -lactamase of the CMY-2 family with enhanced susceptibility to sulfone inhibitors. *Abstr 48th Intersci Conf Antimicrob Agents Chemother*, abstr C1-888. 2008
24. D'Andrea MM, Nucleo E, Luzzaro F, Giani T, Migliavacca R, Vailati F, Kroumova V, Pagani L, Rossolini GM. CMY-16, a novel acquired AmpC-type β -lactamase of the CMY/LAT lineage in multifocal monophyletic isolates of *Proteus mirabilis* from northern Italy. *Antimicrob Agents Chemother*. 2006; 50:618–624. [PubMed: 16436718]
25. CLSI. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard-Seventh edition. Wayne, PA: 2006. Clinical and Laboratory Standard Institute. CLSI document M7-A7
26. Hujer AM, Page MG, Helfand MS, Yeiser B, Bonomo RA. Development of a sensitive and specific enzyme-linked immunosorbent assay for detecting and quantifying CMY-2 and SHV β -lactamases. *J Clin Microbiol*. 2002; 40:1947–1957. [PubMed: 12037047]
27. Paterson DL, Rice LB, Bonomo RA. Rapid method of extraction and analysis of extended-spectrum β -lactamases from clinical strains of *Klebsiella pneumoniae*. *Clin Microbiol Infect*. 2001; 7:709–711. [PubMed: 11843918]
28. Hujer AM, Hujer KM, Bonomo RA. Mutagenesis of amino acid residues in the SHV-1 beta-lactamase: the premier role of Gly238Ser in penicillin and cephalosporin resistance. *Biochim Biophys Acta*. 2001; 1547:37–50. [PubMed: 11343789]
29. Raquet X, Lamotte-Brasseur J, Fonce E, Goussard S, Courvalin P, Frere JM. TEM β -lactamase mutants hydrolysing third-generation cephalosporins. A kinetic and molecular modelling analysis. *J Mol Biol*. 1994; 244:625–639. [PubMed: 7990143]
30. Bethel CR, Hujer AM, Hujer KM, Thomson JM, Ruzsyczky MW, Anderson VE, Pusztai-Carey M, Taracila M, Helfand MS, Bonomo RA. Role of Asp104 in the SHV β -lactamase. *Antimicrob Agents Chemother*. 2006; 50:4124–4131. [PubMed: 16982784]

31. Galleni M, Amicosante G, Frere JM. A survey of the kinetic parameters of class C β -lactamases. Cephalosporins and other β -lactam compounds. *Biochem J.* 1988; 255:123–129. [PubMed: 3264155]
32. Copeland, RA. Evaluation of enzymes inhibitors in drug discovery: a guide for medicinal chemists and pharmacologists. Wiley-Intersciences; Hoboken, New Jersey: 2005.
33. Thomson JM, Distler AM, Prati F, Bonomo RA. Probing active site chemistry in SHV β -lactamase variants at Ambler position 244. Understanding unique properties of inhibitor resistance. *J Biol Chem.* 2006; 281:26734–26744. [PubMed: 16803899]
34. Wu G, Robertson DH, Brooks CL 3rd, Vieth M. Detailed analysis of grid-based molecular docking: A case study of CDOCKER-A CHARMM-based MD docking algorithm. *J Comput Chem.* 2003; 24:1549–1562. [PubMed: 12925999]
35. Spassov VZ, Yan L, Flook PK. The dominant role of side-chain backbone interactions in structural realization of amino acid code. ChiRotor: a side-chain prediction algorithm based on side-chain backbone interactions. *Protein Sci.* 2007; 16:494–506. [PubMed: 17242380]
36. CLSI. Performance standards for antimicrobial susceptibility testing: 17th informational supplement. Wayne, PA: 2008. Clinical and Laboratory Standard Institute. CLSI document M100-S18
37. Galleni M, Amicosante G, Frere JM. A survey of the kinetic parameters of class C beta-lactamases. Cephalosporins and other beta-lactam compounds. *Biochem J.* 1988; 255:123–129. [PubMed: 3264155]
38. Nukaga M, Bethel CR, Thomson JM, Hujer AM, Distler A, Anderson VE, Knox JR, Bonomo RA. Inhibition of class A β -lactamases by carbapenems: crystallographic observation of two conformations of meropenem in SHV-1. *J Am Chem Soc.* 2008; 130:12656–12662. [PubMed: 18761444]
39. Hugonnet JE, Tremblay LW, Boshoff HI, Barry CE 3rd, Blanchard JS. Meropenem-clavulanate is effective against extensively drug-resistant *Mycobacterium tuberculosis*. *Science.* 2009; 323:1215–1218. [PubMed: 19251630]
40. Drawz SM, Babic M, Bethel CR, Taracila M, Distler AM, Ori C, Caselli E, Prati F, Bonomo RA. Inhibition of the Class C β -Lactamase from *Acinetobacter* spp.: Insights into Effective Inhibitor Design. *Biochemistry.* 2009
41. Sulton D, Pagan-Rodriguez D, Zhou X, Liu Y, Hujer AM, Bethel CR, Helfand MS, Thomson JM, Anderson VE, Buynak JD, Ng LM, Bonomo RA. Clavulanic acid inactivation of SHV-1 and the inhibitor-resistant S130G SHV-1 β -lactamase. Insights into the mechanism of inhibition. *J Biol Chem.* 2005; 280:35528–35536. [PubMed: 15987690]
42. Oefner C, D'Arcy A, Daly JJ, Gubernator K, Charnas RL, Heinze I, Hubschwerlen C, Winkler FK. Refined crystal structure of β -lactamase from *Citrobacter freundii* indicates a mechanism for β -lactam hydrolysis. *Nature.* 1990; 343:284–288. [PubMed: 2300174]
43. Heinze-Krauss I, Angehrn P, Charnas RL, Gubernator K, Gutknecht EM, Hubschwerlen C, Kania M, Oefner C, Page MG, Sogabe S, Specklin JL, Winkler F. Structure-based design of β -lactamase inhibitors. I. Synthesis and evaluation of bridged monobactams. *J Med Chem.* 1998; 41:3961–3971. [PubMed: 9767633]
44. Nukaga M, Kumar S, Nukaga K, Pratt RF, Knox JR. Hydrolysis of third-generation cephalosporins by class C beta-lactamases. Structures of a transition state analog of cefotaxime in wild-type and extended spectrum enzymes. *J Biol Chem.* 2004; 279:9344–9352. [PubMed: 14660590]
45. Nukaga M, Mayama K, Hujer AM, Bonomo RA, Knox JR. Ultrahigh resolution structure of a class A β -lactamase: on the mechanism and specificity of the extended-spectrum SHV-2 enzyme. *J Mol Biol.* 2003; 328:289–301. [PubMed: 12684014]
46. Wang X, Minasov G, Blazquez J, Caselli E, Prati F, Shoichet BK. Recognition and resistance in TEM β -lactamase. *Biochemistry.* 2003; 42:8434–8444. [PubMed: 12859188]
47. Doi Y, Wachino J, Ishiguro M, Kurokawa H, Yamane K, Shibata N, Shibayama K, Yokoyama K, Kato H, Yagi T, Arakawa Y. Inhibitor-sensitive AmpC β -lactamase variant produced by an *Escherichia coli* clinical isolate resistant to oxyminocephalosporins and cephamycins. *Antimicrob Agents Chemother.* 2004; 48:2652–2658. [PubMed: 15215122]

48. Nukaga M, Haruta S, Tanimoto K, Kogure K, Taniguchi K, Tamaki M, Sawai T. Molecular evolution of a class C β -lactamase extending its substrate specificity. *J Biol Chem.* 1995; 270:5729–5735. [PubMed: 7890700]
49. Mourey L, Kotra LP, Bellettini J, Bulychiev A, O'Brien M, Miller MJ, Mobashery S, Samama JP. Inhibition of the broad spectrum nonmetallo-carbapenemase of class A (NMC-A) beta-lactamase from *Enterobacter cloacae* by monocyclic beta-lactams. *J Biol Chem.* 1999; 274:25260–25265. [PubMed: 10464248]

Abbreviations

CMY	Active on <u>c</u> eph <u>m</u> ycins
TEM	Class A β -lactamase named after the patient (<u>T</u> emoneira) providing the first sample
SHV	<u>S</u> ulfhydryl reagent <u>v</u> ariable β -lactamase
CTX-M	Active on <u>c</u> efot <u>a</u> xime, first isolated at <u>M</u> unich
ESBL	Extended spectrum β -lactamase
ESI MS	Electrospray ionization mass spectroscopy
MIC	Minimum inhibitory concentration
ATCC	American type culture collection
CTX	Cefotaxime
ATM	Aztreonam

Figure 1.
Chemical structures of several β -lactams.

Figure 2.

Mass spectrometry of the two purified CMY pAmpC β -lactamases alone and in combination with different carbapenems. Left column: CMY-2 β -lactamase. Right column: CMY-32 β -lactamase. Molecular weight (MW), time of incubation and ratio between CMY protein and carbapenem are indicated within the squares. All results should be considered with a ± 3 Da standard error.

Figure 3.

Mass spectrometry of the two purified CMY pAmpC β -lactamases alone and in combination with different β -lactamase inhibitors and ATM. Left column: CMY-2 β -lactamase. Right column: CMY-32 β -lactamase. Molecular weight (MW), time of incubation and ratio between CMY protein and inhibitor are indicated within the squares. All results should be considered with a ± 3 Da standard error.

Figure 4.

(a) Chemical structures of ATM and its alternative mechanism of inactivation by CMY β -lactamases responsible for the adduct of +353 Da seen with ESI-MS (see Figure 4); (b) Chemical structures of BAL29880 and its alternative mechanism of inactivation by CMY β -lactamases responsible for the adduct of +333 Da seen with ESI-MS (see Figure 4). In the case of ATM and BAL29880, the N-sulfonate monobactam binds in the oxyanion hole and the Ser64 hydroxyl attacks the activated carbonyl (C=O), forming the acyl-enzyme and a sulfonamide ($-\text{NH}-\text{SO}_3^{-1}$). The sulfonamide is subsequently hydrolyzed (by H_2O in the active site) to generate the amine ($-\text{NH}_3^{+}$) and sulfate (SO_4^{-2}).

Figure 5.

(a) Molecular modeling of CMY-32 (purple) superimposed onto CMY-2 (yellow) enzyme (r.m.s.d. = 1Å) showing the conformational changes introduced by the mutation at position 214. The most significant difference between the conformations of the two enzymes were observed in the Ω -loop region (Val211-Val223) with a displacement up to 3Å between the

backbone of the amino acids at position 214; **(b)** The change in the position of Tyr221 side chain (more than 90° rotation of the aromatic ring) and Gly214Glu (approximately 3Å change in the position of the backbone) allows the formation of the hydrogen bond between Tyr221:H and Glu214:OE2 ($a = 1.93\text{\AA}$). The new conformation in the CMY-32 enzyme limits the mobility of the Ω -loop. CMY-2 residues are represented in green and CMY-32 in yellow with the oxygen's in red. The hydrogen atoms are not shown.

Figure 6.

CTX modeled in the active site of CMY-32 β -lactamase as a pre-acylation complex. CTX (carbon atoms colored in blue) was docked with the carbonyl of the β -lactam ring in the oxyanion hole of CMY-32 in alignment with the backbone amides of Ser64 and Ser318 ($a = 2.1\text{\AA}$; $b = 2.2\text{\AA}$). The hydrogen bonds of CTX with CMY-32 (colored by element) are represented as dashed lines ($c = 2.0\text{\AA}$; $d = 2.1\text{\AA}$; $e = 2.3\text{\AA}$). In CMY-32, Gln120 and Tyr221 (grey) adapt a different conformation than in CMY-2 (green).

Figure 7. BAL29880 (carbon atoms colored in orange) modeled in the active site of CMY-2 β -lactamase as a acyl-enzyme complex with the carbonyl group in the oxyanion hole formed by Ser318 and Ser64 ($a = 1.9 \text{ \AA}$). Dashed lines represent the hydrogen bonds between BAL29880 and CMY-2 (colored by element; $b = 1.8 \text{ \AA}$; $c = 2.1 \text{ \AA}$; $d = 2 \text{ \AA}$). BAL29880 was modeled without the SO_3^{-2} group.

Table 1

Antimicrobial susceptibility tests for the *E. coli* DH10B clones expressing the CMY-2 or CMY-32 β -lactamases.

Antibiotic	MICs ($\mu\text{g/mL}$) ^a		
	<i>E. coli</i> DH10B pBC SK(-)	<i>E. coli</i> DH10B pBC SK(-)/CMY-2	<i>E. coli</i> DH10B pBC SK(-)/CMY-32
Ampicillin	4	>256	>256
Ampicillin-sulbactam	4	128	24
Ampicillin-BAL29880	2	2	2
Amoxicillin-clavulanate	4	>256	>256
Piperacillin	2	>256	>256
Piperacillin-tazobactam	2	6	4
Piperacillin-BAL29880	1	4	2
Cefoxitin	4	>256	32
Cefotaxime (CTX)	<0.032	16	64
Ceftazidime	<0.032	>256	>256
Cefepime	<0.032	0.25	0.19
Aztreonam (ATM)	<0.032	6	12
BAL29880	>32	>32	>32
Imipenem	<0.032	0.5	0.38
Meropenem	<0.032	0.047	<0.032

^a MICs were obtained using the Etest method. MICs for BAL29880 alone or in combination with ampicillin or piperacillin were obtained by microdilution according to CLSI criteria (25).

Table 2

Steady-state kinetic parameters of CMY-2 and CMY-32.^{a,b}

β -lactam	K_m [μ M] (or K_i corrected)		k_{cat} [s^{-1}]		k_{cat} / K_m [$\mu M^{-1} s^{-1}$]	
	CMY-2	CMY-32	CMY-2	CMY-32	CMY-2	CMY-32
<i>“Good substrates”</i>						
Nitrocefin (NCF)	11.2 (± 1.2)	10.7 (± 0.7)	534.8 (± 16.0)	225.9 (± 3.2)	47.6 (± 5.23)	21.1 (± 1.4)
Cephaloridine	42.7 (± 7.2)	53.8 (± 6.6)	408.5 (± 24.1)	139.6 (± 6.7)	9.6 (± 1.7)	2.6 (± 0.3)
Cephalothin	7.8 (± 0.6)	9.8 (± 1.2)	140.0 (± 22.4)	74.5 ($\pm 2.2\%$)	17.9 (± 3.2)	7.6 (± 1.0)
Cefoxitin	17.9 (± 5.5)	22.9 (± 4.3)	6.8 (± 0.7)	4.7 (± 0.3)	0.38 (± 0.04)	0.21 (± 0.04)
<i>“Poor substrates”</i>						
Cefotaxime (CTX)	1.8 (± 0.2)	4.05 (± 0.26)	NM	0.90 (± 0.06)	NM	0.22 (± 0.02)
Cefepime ^{b,c}	108.1 (± 7.0)	988.9 (± 0.3)	NM	NM	NM	NM
Aztreonam (ATM) ^{b,c}	0.12 (± 0.008)	0.10 (± 0.003)	NM	NM	NM	NM
Imipenem ^{b,c}	0.57 (± 0.059)	0.40 (± 0.036)	NM	NM	NM	NM

^a Standard errors were reported in parentheses;^b k_{cat} not measurable;^c K_i measured; NM, not measurable

Table 3

Determination of K_i , first-order rate constant (k_{inact}), and k_{inact} / K_i ; ^{a,b}

β -lactam	K_i [μM] corrected			k_{inact} [s^{-1}]			k_{inact} / K_i [$\mu\text{M}^{-1} \text{s}^{-1}$]		
	CMY-2	CMY-32		CMY-2	CMY-32		CMY-2	CMY-32	
Tazobactam	50.0 (± 9.5)	13.9 (± 1.2)		0.045 (± 0.0018)	0.129 (± 0.0024)		0.0009 (± 0.00005)	0.009 (± 0.0002)	
Sulbactam	101.3 (± 7.6)	87.9 (± 3.1)		0.025 (± 0.0007)	0.038 (± 0.001)		0.0002 (± 0.00001)	0.0004 (± 0.00002)	
Clavulanate	4,365 (± 471)	500 (± 24.5)		NM	0.031 (± 0.0007)		NM	0.00006 (± 0.0000004)	
BAL29880	4.1 (± 0.40)	5.5 (± 0.60)		0.04 (± 0.001)	0.06 (± 0.005)		0.0098 (± 0.0025)	0.011 (± 0.0023)	

^a Standard errors were reported in parentheses; NM not measurable