Published in final edited form as:

Environ Res. 2014 May ; 131: 174–180. doi:10.1016/j.envres.2014.03.011.

Plasma Polychlorinated Biphenyl Concentrations and Immune Function in Postmenopausal Women*

June T. Spector^{a,b,*,l}, Anneclaire J. De Roos^{c,d,l}, Cornelia M. Ulrich^{d,e,f,l}, Lianne Sheppard^{a,g,l}, Andreas Sjodin^{h,l}, Mark H. Wener^{b,l}, Brent Wood^{b,l}, and Anne McTiernan^{b,c,d,l}

^aDepartment of Environmental & Occupational Health Sciences, School of Public Health, University of Washington, Seattle, WA, USA

^bDepartment of Medicine, School of Medicine, University of Washington, Seattle, WA, USA

^cEpidemiology Program, Public Health Sciences Division, Fred Hutchinson Cancer Research Center, Seattle, WA, USA

^dDepartment of Epidemiology, School of Public Health, University of Washington, Seattle, WA, USA

^eCancer Prevention Program, Fred Hutchinson Cancer Research Center, Seattle, WA, USA

^fNational Center for Tumor Diseases and German Cancer Research Center, Heidelberg, Germany

^gDepartment of Biostatistics, School of Public Health, University of Washington, Seattle, WA, USA

^hNational Center for Environmental Health, CDC, Atlanta, GA, USA

Abstract

Background—Polychlorinated biphenyl (PCB) exposure has been associated with non-Hodgkin lymphoma in several studies, and the immune system is a potential mediator.

Objectives—We analyzed associations of plasma PCBs with immune function measures. We hypothesized that higher plasma PCB concentrations are associated with lower immune function cross-sectionally, and that increases in PCB concentrations over a one year period are associated with decreases in immune function.

*Funding Sources: This research was funded by grants from the National Institutes of Health National Institute of Environmental Health Sciences (NIEHS) (R03 ES015787) and National Cancer Institute (R01CA/AG9334). Support for JTS was provided by the NIEHS (T32 ES015459).

© 2014 Elsevier Inc. All rights reserved.

*Corresponding author at: University of Washington, Departments of Environmental & Occupational Health Sciences and Medicine, 4225 Roosevelt Way NE, Suite 100, Seattle, WA 98105, USA. Tel.: +00 1 206 897 1979..

^lspectj@uw.edu ajd335@drexel.edu neli.ulrich@nct-heidelberg.de sheppard@uw.edu asjodin@cdc.gov wener@u.washington.edu woodbl@u.washington.edu amctiern@fhcrc.org

All authors declare they have no actual or potential competing financial interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Methods—Plasma PCB concentrations and immune function [natural killer (NK) cell cytotoxicity and PHA-induced T-lymphocyte proliferation (PHA-TLP)] were measured at baseline and one year in 109 postmenopausal overweight women participating in an exercise intervention study in the Seattle, Washington (USA) area. Mixed models, with adjustment for body mass index and other potential confounders, were used to estimate associations of PCBs with immune function cross-sectionally and longitudinally.

Results—Associations of PCBs with immune function measures differed across groups of PCBs (e.g., medium- and high-chlorinated and dioxin-like [mono-ortho substituted]) and by the time frame for the comparison (cross-sectional vs. longitudinal). Higher concentrations of medium- and high-chlorinated PCBs were associated with higher PHA-TLP cross-sectionally but not longitudinally. The mean decrease in 0.5 µg/mL PHA-TLP per 50.0 pmol/g-lipid increase in dioxin-like PCBs over one year was 51.6 (95% confidence interval 2.7, 100.5; $P=0.039$). There was no association between plasma PCBs and NK cytotoxicity.

Conclusions—These results do not provide strong evidence of impaired cellular immunity from PCB exposure. Larger longitudinal studies with greater variability in PCB exposures are needed to further examine temporal associations of PCBs with immune function.

Keywords

Cytotoxicity; Immune function; Lymphocyte proliferation; Persistent organic pollutants; Polychlorinated biphenyls

1. INTRODUCTION

Polychlorinated biphenyls (PCBs) continue to be found in human tissues despite the introduction of manufacturing bans in the United States and other countries decades ago (CDC 2009; Dallaire et al. 2002). Organochlorine pollutants, which include PCBs and organochlorine pesticides/pesticide metabolites, are lipid soluble compounds that do not break down easily in the environment or in humans. The main route of human exposure to organochlorines is through the diet, and these compounds concentrate as they move up the food chain. Circulating blood organochlorine concentrations are regulated by homeostasis between adipose tissue storage and blood lipid levels (Brown and Lawton 1984).

Exposure to organochlorines has been implicated as a risk factor for non-Hodgkin lymphoma (NHL) (Ballschmiter and Zell 1980; Hardell et al. 1996; Rothman et al. 1997; Colt et al. 2005; DeRoos et al. 2005; Engel et al. 2007; Spinelli et al. 2007; Colt et al. 2009; Bertrand et al. 2010; Ng et al. 2010), and immune suppression is among the proposed mechanisms for this increased risk. An association between organochlorines and immunosuppressive effects, including decreases in natural killer (NK) cell cytotoxicity and T-lymphocyte proliferation (TLP), has been observed *in vitro* (Daniel et al. 2001; Hammond et al. 2005), in animal studies (Exon et al. 1985; Talcott et al. 1985; Ross et al. 1996; Beckman et al. 2003; Beineke et al. 2005; Mori et al. 2006; Sormo et al. 2009), and in human studies (Lü and Wu 1985; Svensson et al. 1994; Leijls et al. 2009), although associations are not entirely consistent. Toxic effects and potential mechanisms of action of PCBs and related organochlorine pesticides/pesticide by-products appear to vary by

structural characteristics. PCB congeners with mono-*ortho*-substituted (dioxin-like) structures exert toxicity through binding to the aryl hydrocarbon receptor, while different mechanisms may exist for other organochlorines (CDC 2009; Duffy and Zelikoff 2006; Lyche et al. 2006; Ng et al. 2010). Several studies of NHL found the strongest associations with PCB 180, a moderately chlorinated PCB, compared to other PCB congeners (Bertrand et al. 2010; DeRoos et al. 2005; Spinelli et al. 2007), indicating possible differences in biologic effects between individual, non-dioxin-like congeners that may vary by degree of chlorination. Few studies have examined the association of organochlorines with immune effects in humans (Lü and Wu 1985; Svensson et al. 1994; Leijds et al. 2009), and no study that we are aware of has reported on the effect of within-person changes in organochlorines with within-person changes in immune function.

We investigated the association between plasma PCB concentrations and immune function in a study of postmenopausal, overweight and obese women living in the Seattle, Washington metropolitan area. We focused on three summed measures of plasma PCBs: dioxin-like (monoortho substituted), high-chlorinated (8 or more chlorines), and medium-chlorinated (5-8 chlorines) PCBs (CDC 2009; Ng et al. 2010). Cellular immune function was evaluated by TLP (lower levels are believed to reflect less effective immune responses) and NK cell cytotoxicity (lower levels may predict risk of future adverse health events) (Levy et al. 1991; Mizutani et al. 1996; Vedhara et al. 1999; Imai et al. 2000; Albers et al. 2005). We hypothesized that: 1) higher plasma PCB concentrations are associated with lower baseline immune function cross-sectionally; and 2) increases in PCB concentrations over time are associated with decreases in immune parameters. We based our hypotheses that higher-chlorinated and dioxin-like PCBs, in particular, have immune effects on existing studies of the relationship between PCBs and immune measures (Daniel et al. 2001; Leijds et al. 2009) and between PCBs and NHL (Bertrand et al. 2010; DeRoos et al. 2005; Spinelli et al. 2007).

2. MATERIALS AND METHODS

2.1. Study Population

The study included participants in a previously conducted exercise intervention trial and ancillary study of immune function in sedentary and overweight/obese postmenopausal women. The Physical Activity for Total Health study, conducted at the Fred Hutchinson Cancer Research Center (FHCRC), was a randomized controlled trial comparing the effects of a one year moderate-intensity aerobic exercise intervention, with goal of 225 minutes/week in a combined facility and home program, versus a stretching control program on sex hormone concentrations (as biomarkers of breast cancer risk) (McTiernan et al. 1999). As previously described, participants were selected to maximize the possible effects of exercise on endogenous sex hormones and to avoid other factors known to affect sex hormones (Irwin et al. 2003). Women were enrolled into the Physical Activity for Total Health study between 1998 and 2000. One hundred and seventy three women from the Seattle, Washington area participated in the study and were evaluated at baseline, three months, and one year. The ancillary study of Immune Function and Exercise (IMEX) was conducted

among 115 participants of the Physical Activity for Total Health study that met additional IMEX selection criteria (Shade et al. 2004).

The present study included the IMEX subset of the Physical Activity for Total Health study participants. We excluded participants at baseline or one year who were missing all primary outcome or primary exposure values or total lipid measures necessary to compute lipid-adjusted plasma organochlorine concentrations (Phillips et al. 1989; Schisterman et al. 2005). Six of the 115 IMEX participants were missing these values at baseline, and seventeen participants (including three that were lost to follow-up) were missing these values at one year. A sensitivity analysis excluding observations with any missing covariate data in any of the analyses yielded similar results.

2.2. Questionnaires

Information on demographics and smoking history was collected via self-administered questionnaire. Questionnaire data also included body weight history, medical history, health habits, medication use, and dietary intake over the past three months. Dietary intake and alcohol consumption over the past three months were assessed using a 120-item food frequency questionnaire designed and validated at the FHCRC (Patterson et al. 1999).

2.3. Body Composition Measures

Body weight and height were measured to the nearest 0.1 kg and 0.1 cm, respectively, using a balance-beam scale and stadiometer at corresponding clinic visits. Both measurements were taken in duplicate and averaged. Body mass index (BMI) was calculated as $(\text{weight}[\text{kg}]/\text{height}[\text{m}]^2)$.

2.4. Blood Draw

Blood samples were collected for previously described studies conducted in the same population as our study population (Shade et al. 2004; Troen et al. 2006; Campbell et al. 2008). Blood draws took place at the UW Department of Laboratory Medicine. Fasting blood samples were taken between 7:30 and 8:30 AM. Samples were processed within one hour of collection.

2.5. Immune Function Measures

Assays of cellular immunity were conducted by the UW Department of Laboratory Medicine on blood samples from the baseline and one year IMEX visits. NK cytotoxicity was measured in fresh NK cells using a flow-cytometric assay. NK cells were isolated from peripheral blood mononuclear cells (PBMCs) by Ficoll-Hypaque separation. Cells were washed and diluted to a mononuclear cell concentration of 7.7×10^6 cell/mL. K562 target cells were washed in the log phase of growth twice and incubated with label 3,3'-diiododecylcarboxycyanine perchlorate (DiO; Live/Dead cytotoxicity kit no. L7010; Molecular Probes, Eugene, OR). Cells were incubated, washed, and re-suspended to a concentration of 1×10^6 cells/mL and then filtered through a 35- μm strainer. Culture-suspended NK cells were diluted to four effector-to-target cell (E:T) ratios of 50:1, 25:1, 12.5:1, and 6.25:1, pelleted, and incubated. Propidium iodide was added to a final concentration of 0.03 mg/mL, and cells were transferred to a polypropylene tube for flow

cytometric analysis to identify dead cells. The percentage of dead target cells among a total DiO-identified target cells was used as the measure of NK cytotoxicity.

T-lymphocyte proliferation was assessed using cryopreserved PBMCs with ^3H -thymidine incorporation in response to the mitogen phytohemagglutinin (PHA), as previously described (Boynton et al. 2007; Campbell et al. 2008). Cells were prepared by Ficoll-Hypaque separation and frozen in 30% fetal calf serum, 60% RPMI medium, and 10% dimethyl sulfoxide (Gibco, Gaithersburg, MD). For the ^3H -thymidine incorporation, PBMCs were incubated with PHA of 0.1 and 0.5 $\mu\text{g/mL}$ in five replicates each. After incubation for 3 days at 37°C , cells were pulsed for 24 hr with ^3H -thymidine, harvested, and counted with a β -counter. The PHA-stimulated TLP (PHA-TLP) index was expressed as counts per minute of stimulated cells divided by counts per minute of unstimulated cells.

2.6. Polychlorinated Biphenyl Measures

Plasma samples available from 111 IMEX participants were analyzed for PCBs at the Persistent Organic Pollutants Biomonitoring Laboratory of the Centers for Disease Control and Prevention (CDC) in Atlanta, GA. Measurements of 36 PCB congeners [International Union of Pure and Applied Chemistry [IUPAC] scheme (Ballschmiter and Zell 1980) numbers: 18, 28, 44, 49, 52, 66, 74, 87, 99, 101, 105, 110, 118, 128, 138/158, 146, 149, 151, 153, 156, 157, 167, 170, 172, 177, 178, 180, 183, 187, 189, 194, 195, 196/203, 199, 206, 209] were conducted by high-resolution gas chromatography/isotope-dilution high-resolution mass spectrometry (Sjödén et al. 2004). Measurements were conducted in samples from the baseline and one year IMEX visits. The plasma sample amount available for the assays ranged from 0.23 g to 0.84 g with a median of 0.38 g. The analytic results were reported as wet-weight concentrations (pg/g plasma) and lipid-standardized concentrations (using previously measured triglyceride and total cholesterol values from the same blood draw) in ng/g-lipid (Phillips et al. 1989; Mohanka et al. 2006).

We limited our study to the 15 PCB congeners detected in >80% of the samples. Repeatability of measurements was excellent, with coefficients of variation (CVs) between blinded quality control replicate samples ranging from 2.4 to 11.2 for these analytes. The number of samples with measured values above the limit of detection (LOD) and the average detection limits for these PCBs have been previously described (DeRoos et al. 2012) and are shown for the PCBs assessed in our primary analyses in Appendix A, Table 1. We imputed values below the LOD with a value equal to the LOD divided by two (DeRoos et al. 2012).

2.7. Statistical Analysis

Statistical analyses were performed using Stata 10 (StataCorp, College Station, TX). We used mixed models to investigate the association of PCB concentrations with immune function measures, adjusting for potential confounding factors. We chose primary exposures *a priori* to be lipid-standardized summed dioxin-like PCBs (PCB 105, 118, and 156), high-chlorinated PCBs (PCB 194, 196/203, 199, 206, 209), and medium-chlorinated PCBs (PCB 138/158, 146, 153, 156, 170, 180, 183, 187). In a secondary analysis, we examined lipid-

standardized summed low-chlorinated PCBs (PCB 74, 99, 105, 118). The summed values were obtained by adding the molar concentrations of individual PCBs in each group.

Our primary outcomes were NK cytotoxicity and PHA-TLP, chosen based on associations with PCBs reported in previous studies. We considered these to be independent outcomes. Only NK cytotoxicity % based on effector-target ratios of 12.5:1 and 25:1 were analyzed in this study, as these dilutions were considered to have the greatest reproducibility and were in the linear range (Shade et al. 2004).

To investigate associations between each summed PCB exposure and immune function outcome both cross-sectionally at baseline and longitudinally from baseline to one year of follow-up, we used random-intercept mixed models of the form:

$$Y_{ij} = \beta_{0i} + \beta_1 PCB_{i0} + \beta_2 TIME_{ij} (PCB_{ij} - PCB_{i0}) + \sum_1^s \alpha_p COV_{i0p} + \sum_1^s \gamma_p TIME_{ij} (COV_{ijp} - COV_{i0p}) + \epsilon_{ij},$$

where $\beta_{0i} = \beta_0 + \delta_{0i}$,

and Y_{ij} = immune function for subject i at time j ; PCB_{ij} = PCB level for subject i at time j ; COV_{ijp} = the p^{th} covariate for subject i at time j ; $TIME_{ij}$ = time (years) from baseline visit for subject i at time j ($TIME_{i0}=0$ at baseline; $TIME_{i1}=1$ at one year); δ_{0i} = residual variation in initial status; and ϵ_{ij} = measurement error associated with Y_{ij} . Mixed models were fit via restricted maximum likelihood using STATA *xtmixed*. At baseline ($TIME=0$), $Y_{i0} = \beta_{0i} + \beta_1 PCB_{i0} + \sum_1^s \alpha_p COV_{i0p} + \epsilon_{i0}$, and β_1 represents the adjusted baseline cross-sectional estimate of the mean change in immune function measure per unit difference in PCB concentration. This cross-sectional effect estimate was scaled to reflect the magnitude of variation in PCB concentrations and is reported as the mean change in immune function measure per 50 pmol/g-lipid difference in PCB concentration. Between baseline ($TIME=0$) and one year of follow-up ($TIME=1$), $Y_{i1} - Y_{i0} = \beta_2 (PCB_{i1} - PCB_{i0}) + \sum_1^s \gamma_p (COV_{i1p} - COV_{i0p}) + \epsilon_{i1} - \epsilon_{i0}$, and β_2 represents the adjusted estimate of the mean change in immune function measure difference from baseline to one year per unit change in PCB concentration difference from baseline to one year. This longitudinal effect estimate was also scaled to reflect the magnitude of variation in change in PCB concentrations and is reported as the mean change in immune function measure difference from baseline to one year per 50 pmol/g-lipid change in PCB concentration difference from baseline to one year. To evaluate the consistency of our results, we examined in secondary analyses the association between exposures and outcomes cross-sectionally using generalized estimating equations (GEE) with exchangeable working correlations and robust variance estimators.

We selected variables *a priori* for adjustment based on previously described confounders (Johnson et al. 1990; Nieman et al. 1999; Deutch et al. 2003; Deutch et al. 2007; Romeo et al. 2007) and biological plausibility. Variables included baseline age (years), race (white non-Hispanic, other), baseline education status (high school or less; vocational school, some college, or associate degree; bachelor degree; master degree or doctorate), baseline smoking status (never, former), blood draw season (winter, spring, summer, fall), alcohol use (<1, 1-5, > 5 g/day), BMI, and fish intake (<1.5, 1.5-2.4, > 2.5 servings/week). We adjusted for BMI because relationships of excess adiposity with impaired immune function (Marti et al.

2001) and lower plasma concentrations of most PCBs (De Roos et al. 2012) have been described. We also adjusted for the Physical Activity for Total Health study intervention group (one year moderate intensity aerobic exercise vs. stretching control).

3. RESULTS

3.1. Study population

Baseline characteristics of the 109 women in our study are shown in Table 1. Participants had a mean age of 60.6 years and were mostly highly educated (41% with a bachelor's degree or higher) and white (89%). All participants were non-smokers, per inclusion criteria of the parent study, and approximately half were former smokers. Alcohol intake was generally low (<1g/day for 53% of participants). Forty five percent of participants reported eating less than 1.5 servings of fish per week. Forty seven percent were obese (BMI ≥ 30). There was a very modest change in BMI over one year (mean change −0.2 kg/m² in the exercise group, 0.2 kg/m² in the stretching group).

Median (inter-quartile range) dioxin-like, high-, and medium-chlorinated PCB concentrations were 87.4 (60.4, 134.9), 83.9 (58.1, 109.0), and 407.3 (324.9, 536.9) pmol/g-lipid, respectively, at baseline and 93.9 (69.1, 135.9), 86.4 (66.5, 108.8), and 440.6 (360.9, 545.7) pmol/g-lipid at one year. Older participants were more likely than younger participants to have higher high- and medium-chlorinated plasma PCB concentrations. Changes in PCB concentrations between baseline and one year are described in Table 1. There was little variability in PCB concentrations over time; the median (interquartile range) change in PCB concentrations between baseline and one year was 1.4 (−8.6, 10.1), −0.3 (−8.9, 9.9), and 7.5 (−31.4, 47.9) pmol/g-lipid for dioxin-like, high-, and medium-chlorinated PCBs, respectively. Immune function measure distributions in the study population at baseline and one year are shown in Appendix A, Table 2. There was no significant difference in the mean (standard deviation) absolute number of T cells (CD3+, CD45+) between baseline [1302 (363)] and one year [1344 (412)] or in the percent of T cells between baseline [71 (6)] and one year [71(7)].

3.2. Cross-sectional associations

Baseline cross-sectional associations of plasma PCB concentrations with immune function obtained via mixed models are shown in Table 2. There were positive cross-sectional associations of baseline PCB concentrations with PHA-TLP indices, although confidence intervals were very wide. The mean change in PHA-TLP measure per 50.0 pmol/g-lipid increase in PCB concentration was 4.0 (95% CI 0.6, 7.4) and 26.3 (95% CI 3.4, 49.2) for medium-chlorinated and high-chlorinated PCBs, respectively, with 0.5 µg/mL PHA-TLP, and 1.7 (95% CI 0.1, 3.3) for medium-chlorinated PCBs with 0.1 µg/mL PHA-TLP. There was no significant association of PCB levels with NK cytotoxicity. Cross-sectional GEE models, which included observations from baseline and one year, yielded similar results (not shown).

3.3. Longitudinal associations

Longitudinal associations of plasma PCB concentrations with immune function obtained via mixed models are shown in Table 2. In general, increases in PCB levels from baseline to one year were associated with decreases in PHA-TLP indices. Although very imprecise, the mean decrease in 0.5 µg/mL PHA-TLP per 50.0 pmol/g-lipid increase in dioxin-like PCBs over one year was 51.6 (95% confidence interval 2.7, 100.5; $P=0.039$). The mean decrease in 0.5 µg/mL PHA-TLP per 50.0 pmol/g-lipid increase in low-chlorinated PCBs over one year was 29.0 (95% confidence interval 1.9, 56.1; $P=0.036$). There was no significant association of PCB level differences over time with NK cytotoxicity % differences (Table 2).

4. DISCUSSION

This is the first study, to our knowledge, to report on associations of plasma PCB concentrations and functional assays of cellular immunity over time. Cross-sectionally, we observed positive associations of PCBs with PHA-TLP, particularly for medium- and high-chlorinated PCBs. We observed a negative association of differences in 0.5 µg/mL PHA-TLP index with differences in dioxin-like and low-chlorinated PCBs over one year. These results must be interpreted with caution given the modest size of the study, the small variability in PCB concentrations over time, and the relatively short duration of the study.

Although counter to our hypothesis, our findings of cross-sectional associations of higher concentrations of medium- and high-chlorinated PCBs with higher PHA-TLP are consistent with previous findings in patients exposed to high levels of PCBs at Yu-Cheng (Lü and Wu 1985). Phytohemagglutinin-TLP in these patients was noted to be significantly elevated one and three years after cessation of high-level exposure, compared to control subjects, and it was hypothesized that these findings reflected a rebound phenomenon resulting from sub-lethal toxic doses of PCBs (Lü and Wu 1985). However, compared to levels in our study, PCB levels were several orders of magnitude higher in the Lü et al. (1985) study.

Our study is not directly comparable to other studies that have used different methods to assess immunity in individuals with lower-level PCB exposures. Leijds et al. (2009) reported a significant negative correlation between current serum levels of dioxin-like PCBs and concentrations of segmental neutrophils in the blood/serum, but immune function was not assessed. Svensson et al. (1994) reported lower absolute numbers of NK cells, but no significant difference in Concanavalin A (Con A)-induced LP, in high- versus non-consumers of fish. High and non-consumers had median wet-weight levels of PCB 118 of 500 and 200 pg/g, respectively (136 pg/g in our study).

The generally smaller cross-sectional effect estimates of dioxin-like, compared to non-dioxin-like, PCBs with PHA-TLP observed in our study are consistent with previous studies. Mori et al. (2006) reported that changes in Con A-induced LP were mostly explained by non-dioxin-like PCBs, particularly medium-chlorinated PCBs 138 and 180, rather than dioxin-like (specifically, coplanar) PCBs in certain marine mammals. Differential effects of dioxin-like and non-dioxin-like PCBs on immune function may reflect different mechanisms of toxicity.

Although variability in exposures was extremely small and effect estimates imprecise in longitudinal analyses, increases in PCB levels from baseline to one year were generally associated with decreases in PHA-TLP. The higher longitudinal effect sizes for dioxin-like compared to non-dioxin-like (medium and high-chlorinated) PCBs may again reflect differences in immunotoxic mechanisms. A similar pattern in effect estimates was observed for dioxin-like and low-chlorinated PCBs. This may be explained by the overlap in PCBs in these two groups; PCBs 105 and 118 are both dioxin-like and low-chlorinated congeners. Although these findings need to first be confirmed in larger longitudinal studies with greater variability in PCB exposures, declines in LP may have public health implications, for example in the elderly, who are also more likely to have had higher PCB exposures. Lymphocyte proliferation has been reported to correlate inversely with mortality in elderly patients and patients with HIV (Albers et al. 2005). In addition, a reduction in LP may result in impaired tumor immunity by reducing the accumulation of tumor-specific T cells (Datta and Sarvetnick 2009).

Effect sizes of NK cytotoxicity with PCB levels were generally small. NK cells are involved in the innate immune response and can destroy virally infected cells and tumor cells if activated (Friberg et al. 1996), but the relationship between *in vivo* NK function and NK cytotoxicity assays is unclear. Further, findings of the relationship between PCBs and NK function have not been entirely consistent. Omara et al. (1998) found no evidence of decreased NK cytotoxicity with *in vitro* exposure of rat leukocytes to low levels of PCB mixtures, but dosing of rats with higher levels of PCBs has been reported to cause decreases in NK cytotoxicity (Exon et al. 1985; Talcott et al. 1985).

Strengths of our study include the use of functional assays for characterizing immune status, the strict criteria for blood draws, and the quantitative exposure assessment using plasma PCB concentrations. Polychlorinated biphenyl measures were consistent with previous reports. Baseline plasma PCB concentrations in our study, and unadjusted trends of increasing PCB levels with age, were comparable to those reported in the National Health and Nutrition Examination Survey (NHANES) (CDC 2009; Patterson et al. 2009).

Our study also has important limitations. First, there was very little variability in PCB concentrations over the one-year time frame of the study. PCB levels are expected to be relatively stable over short time periods in the absence of significant exogenous or endogenous exposures (Carpenter 2006), and the modest weight change in the Physical Activity for Total Health trial participants contributed to the lack of substantial variability seen in PCB levels. Observation and interpretation of longitudinal changes in immune function is difficult with such small variability in PCB concentrations over time, as the possibility of chance findings exists. We limited the number of comparisons (e.g. exposure and immune function outcomes) by identifying *a priori* primary exposure and outcome measures in order to minimize the possibility of chance findings. However, longitudinal studies with greater changes in PCB concentrations are needed to confirm these findings. Second, we used summed PCB measures rather than isolated congener measures. An argument for this approach is that organochlorine mixtures are more commonly present in humans, and our findings may therefore be more biologically relevant than for individual congeners. Third, we adjusted for fish intake over the past three months but not over the

longer term. A previous study in the same population reported an association between fish intake over the past three months and a summed PCB measure (De Roos et al. 2012). However, it is possible that residual confounding by dietary factors or dietary intake over time periods other than those addressed in our study is present.

The extent to which our findings may generalize to populations other than postmenopausal, overweight/obese women, and in other settings, is unknown. It is also unknown whether the NK cytotoxicity assay used in our study is sufficiently sensitive to detect mildly impaired NK function, and PHA-TLP may have been affected by cryopreservation of cells. Future studies should incorporate additional methods, such as assays involving CD107a degranulation and stimulation with anti-CD3/CD28/CD49, to assess NK cell and T-lymphocyte function, respectively.

5. CONCLUSIONS

In this study, higher concentrations of medium- and high-chlorinated PCBs were associated with higher PHA-TLP cross-sectionally, and an increase in dioxin-like PCB concentrations was associated with a decrease in PHA-TLP over one year. Larger longitudinal studies with greater variability in PCB exposures are needed to confirm these findings and to further examine temporal associations of PCBs with immune function. Such studies may help elucidate whether negative associations of differences in LP with differences in dioxin-like PCBs over time are relevant to the development of cancers such as NHL.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

All authors certify that their freedom to design, conduct, interpret, and publish research is not compromised by any controlling sponsor. The findings and conclusions in this report are those of the authors and do not necessarily represent the Fred Hutchinson Cancer Research Center, the University of Washington, or the Centers for Disease Control and Prevention.

Abbreviations

CDC	Centers for Disease Control and Prevention
Con A	Concanavalin A
CV	Coefficient of variation
FHCRC	Fred Hutchinson Cancer Research Center
IMEX	Immune Function and Exercise
LOD	limit of detection
LP	lymphocyte proliferation
NK	natural killer

NHL	non-Hodgkin lymphoma
PBMCs	peripheral blood mononuclear cells
PCB	polychlorinated biphenyl
PHA	phytohemagglutinin
PHATLP	PHA-induced T-lymphocyte proliferation
UW	University of Washington

REFERENCES

- Albers R, Antoine JM, Bourdet-Sicard R, Calder PC, Gleeson M, Lesourd B, Samartin S, Sanderson IR, Van Loo J, Vas Dias FW, Watzl B. Markers to measure immunomodulation in human nutrition intervention studies. *Br. J. Nutr.* 2005; 94:452–481. [PubMed: 16176618]
- Ballschmiter K, Zell M. Baseline studies of the global pollution. I. Occurrence of organohalogens in pristine European and antarctic aquatic environments. *Int. J. Environ. Anal. Chem.* 1980; 8:15–35. [PubMed: 7399798]
- Beckman KB, Blake JE, Ylitalo GM, Stott JL, O'Hara TM. Organochlorine contaminant exposure and associations with hematological and humoral immune functional assays with dam age as a factor in free-ranging northern fur seal pups (*Callorhinus ursinus*). *Mar. Pollut. Bull.* 2003; 46:594–606. [PubMed: 12735957]
- Beineke A, Siebert U, McLachlan M, Bruhn R, Thron K, Failing K, Muller G, Baumgartner W. Investigations of the potential influence of environmental contaminants on the thymus and spleen of harbor porpoises (*Phocoena phocoena*). *Environ. Sci. Technol.* 2005; 39:3933–3938. [PubMed: 15984767]
- Bertrand KA, Spiegelman D, Aster JC, Altshul LM, Korrick SA, Rodig SJ, Zhang SM, Kurth T, Laden F. Plasma organochlorine levels and risk of non-Hodgkin lymphoma in a cohort of men. *Epidemiology.* 2010; 21:172–180. [PubMed: 20087190]
- Boynton A, Neuhauser ML, Wener MH, Wood B, Sorensen B, Chen-Levy Z, Kirk EA, Yasui Y, McTiernan A, Ulrich CM. Associations between healthy eating patterns and immune function or inflammation in overweight or obese postmenopausal women. *Am. J. Clin. Nutr.* 2007; 86:1445–55. [PubMed: 17991658]
- Brown JF, Lawton RW. Polychlorinated biphenyl (PCB) partitioning between adipose tissue and serum. *Bull. Environ. Contam. Toxicol.* 1984; 33:277–80. [PubMed: 6434008]
- Campbell PT, Wener MH, Sorensen B, Wood B, Potter JD, McTiernan A, Ulrich CM. Effect of exercise on immune function: A 12-month randomized controlled trial among postmenopausal women. *J. Appl. Physiol.* 2008; 104:1648–55. [PubMed: 18403448]
- Carpenter DO. Polychlorinated biphenyls (PCBs): routes of exposure and effects on human health. *Rev. Environ. Health.* 2006; 21:1–23. [PubMed: 16700427]
- Centers for Disease Control and Prevention (CDC). Fourth National Report on Human Exposure to Environmental Chemicals. Atlanta, GA: 2009.
- Colt JS, Rothman N, Severson RK, Hartge P, Cerhan JR, Chatterjee N, Cozen W, Morton LM, DeRoos AJ, Davis S, Chanock S, Wang SS. Organochlorine exposure, immune gene variation, and risk of non-Hodgkin lymphoma. *Blood.* 2009; 113:1899–1905. [PubMed: 19066394]
- Colt JS, Severson RK, Lubin J, Rothman N, Camann D, Davis S, Cerhan JR, Cozen W, Hartge P. Organochlorines in carpet dust and non-Hodgkin lymphoma. *Epidemiology.* 2005; 16:516–525. [PubMed: 15951670]
- Dallaire F, Dewailly E, Laliberte C, Muckle G, Ayotte P. Temporal trends of organochlorine concentrations in umbilical cord blood of newborns from the lower north shore of the St. Lawrence River. *Environ. Health. Perspect.* 2002; 110:835–838. [PubMed: 12153768]

- Daniel V, Huber W, Bauer K, Suesal C, Conradt C, Opelz G. Associations of blood levels of PCB, HCHS, and HCB with numbers of lymphocyte subpopulations, in vitro lymphocyte response, plasma cytokine levels, and immunoglobulin autoantibodies. *Environ. Health Perspect.* 2001; 109:173–178. [PubMed: 11266329]
- Datta S, Sarvetnick N. Lymphocyte proliferation in immune-mediated diseases. *Trends Immunol.* 2009; 30:430–438. [PubMed: 19699149]
- DeRoos AJ, Hartge P, Lubin JH, Colt JS, Davis S, Cerhan JR, Severson RK, Cozen W, Patterson DG, Needham LL, Rothman N. Persistent organochlorine chemicals in plasma and risk of non-Hodgkin's lymphoma. *Cancer Res.* 2005; 65:11214–11226. [PubMed: 16322272]
- DeRoos AJ, Ulrich CM, Sjodin A, McTiernan A. Adiposity, body composition, and weight change in relation to organochlorine pollutant plasma concentrations. *J. Expo. Sci. Environ. Epidemiol.* 2012; 22:617–24. [PubMed: 22588213]
- Deutch B, Pedersen HS, Jorgensen EC, Hansen JC. Smoking as a determinant of high organochlorine levels in Greenland. *Arch. Environ. Health.* 2003; 58:30–36. [PubMed: 12747516]
- Deutch B, Petersen HS, Asmund G, Hansen JC. Contaminants, diet, plasma fatty acids and smoking in Greenland 1999–2005. *Sci. Total Environ.* 2007; 372:486–496. [PubMed: 17157358]
- Duffy JE, Zelikoff JT. The relationship between noncoplanar PCB-induced immunotoxicity and hepatic CYP1A induction in a fish model. *J. Immunotoxicol.* 2006; 3:39–47. [PubMed: 18958684]
- Engel LS, Laden F, Andersen A, Strickland PT, Blair A, Needham LL, Barr DB, Wolff MS, Helzlsouer K, Hunger DJ, Lan Q, Cantor KP, Comstock GW, Brock JW, Bush D, Hoover RN, Rothman N. Polychlorinated biphenyl levels in peripheral blood and non-Hodgkin's lymphoma: A report from three cohorts. *Cancer. Res.* 2007; 67:5545–5552. [PubMed: 17545638]
- Exon JH, Talcott PA, Koller LD. Effect of lead, polychlorinated biphenyls, and cyclophosphamide on rat natural killer cells, interleukin 2, and antibody synthesis. *Fund. Appl. Toxicol.* 1985; 5:158–64.
- Friberg DD, Bryant JL, Whiteside TL. Measurements of natural killer (NK) activity and NK-cell quantification. *Methods.* 1996; 9:316–326. [PubMed: 8812685]
- Hammond JA, Hall AJ, Dyrnya EA. Comparison of polychlorinated biphenyl (PCB) induced effects on innate immune functions in harbour and grey seals. *Aquat. Toxicol.* 2005; 74:126–138. [PubMed: 15982755]
- Hardell L, Vanbavel B, Lindstrom G, Fredrikson M, Hagberg H, Liljegren G, Nordstrom M, Johansson B. Higher concentrations of specific polychlorinated biphenyl congeners in adipose tissue from non-Hodgkin's lymphoma patients compared with controls without a malignant disease. *Int. J. Oncol.* 1996; 9:603–608. [PubMed: 21541557]
- Imai K, Matsuyama S, Miyake S, Suga K, Nakachi K. Natural cytotoxic activity of peripheral-blood lymphocytes and cancer incidence: An 11-year follow-up study of a general population. *Lancet.* 2000; 356:1795–1799. [PubMed: 11117911]
- Irwin ML, Yasui Y, Ulrich CM, Bowen D, Rudolph RE, Schwartz RS, Yukawa M, Aiello E, Potter JD, McTiernan A. Effect of exercise on total and intra-abdominal body fat in postmenopausal women: A randomized controlled trial. *JAMA.* 2003; 289:323–330. [PubMed: 12525233]
- Johnson JD, Houchens DP, Kluwe WM, Craig DK, Fisher GL. Effects of mainstream and environmental tobacco smoke on the immune system in animals and humans: A review. *Crit. Rev. Toxicol.* 1990; 20:369–395. [PubMed: 2202327]
- Leijds MM, Koppe JG, Olie K, van Aalderen WM, de Voogt P, ten Tusscher GW. Effects of dioxins, PCBs, and PBDEs on immunology and hematology in adolescents. *Environ. Sci. Technol.* 2009; 43:7946–7951. [PubMed: 19921918]
- Levy SM, Herberman RB, Lee J, Whiteside T, Beadle M, Heiden L, Simons A. Persistently low natural killer cell activity, age, and environmental stress as predictors of infectious morbidity. *Nat. Immun. Cell Growth Regul.* 1991; 10:289–307. [PubMed: 1787835]
- Lü YC, Wu YC. Clinical findings and immunological abnormalities in Yu-Cheng patients. *Environ. Health Perspect.* 1985; 59:17–29. [PubMed: 3921359]
- Lyche JL, Larsen HJ, Skaare JU, Tverdal A, Johansen GM, Ropstad E. Perinatal exposure to low doses of PCB 153 and PCB 126 affects maternal and neonatal immunity in goat kids. *J. Toxicol. Environ. Health.* 2006; 69:139–158.

- Marti A, Marcos A, Martinez JA. Obesity and immune function relationships. *Obes. Rev.* 2001; 2:131–140. [PubMed: 12119664]
- McTiernan A, Ulrich CM, Yancey D, Slate S, Nakamura H, Oestreich N, Bowen D, Yasui Y, Potter J, Schwartz R. The Physical Activity for Total Health (PATH) Study: Rationale and design. *Med. Sci. Sports Exerc.* 1999; 31:1307–12. [PubMed: 10487373]
- Mizutani Y, Okada Y, Terachi T, Yoshida O. Prognostic significance of circulating cytotoxic lymphocytes against autologous tumors in patients with bladder cancer. *J. Urol.* 1996; 155:888–892. [PubMed: 8583600]
- Mohanka M, Irwin M, Heckbert SR, Yasui Y, Sorensen B, Chubak J, Tworoger SS, Ulrich CM, McTiernan A. Serum lipoproteins in overweight/obese postmenopausal women: a one-year exercise trial. *Med. Sci. Sports Exerc.* 2006; 38:231–239.
- Mori C, Morsey B, Levin M, Nambiar PR, De Guise S. Immunomodulatory effects of in vitro exposure to organochlorines on T-cell proliferation in marine mammals and mice. *J. Toxicol. Environ. Health.* 2006; 69:283–302.
- Ng CH, Janoo-Gilani R, Sipahimalani P, Gallagher RP, Gascoyne RD, Connors JM, Weber JP, Lai AS, Leach S, Le ND, Brooks-Wilson AR, Spinelli JJ. Interaction between organochlorines and the AHR gene, and risk of non-Hodgkin lymphoma. *Cancer Causes Control.* 2010; 21:11–22. [PubMed: 19821039]
- Nieman DC, Henson DA, Nehlsen-Cannarella SL, Ekkens M, Utter AC, Butterworth DE, Fagoaga OR. Influence of obesity on immune function. *J. Am. Diet. Assoc.* 1999; 99:294–299. [PubMed: 10076580]
- Omara FO, Flipo D, Brochu C, Denizeau F, Brousseau P, Potworowski EF, Fournier M. Lack of suppressive effects of mixtures containing low levels of methylmercury (MeHg), polychlorinated dibenzo-p-dioxins (PCDDs), polychlorinated dibenzofurans (PCDFS), and aroclor biphenyls (PCBS) on mixed lymphocyte reaction, phagocytic, and natural killer cell activities of rat leukocytes in vitro. *J. Toxicol. Environ. Health.* 1998; 54:561–77.
- Patterson DG, Wong LY, Turner WE, Caudill SP, Dipietro ES, McClure PC, Cash TP, Osterloh JD, Pirkle JL, Sampson EJ, Needham LL. Levels in the U.S. population of those persistent organic pollutants (2003-2004) included in the Stockholm Convention or in other long range transboundary air pollution agreements. *Environ. Sci. Technol.* 2009; 43:1211–8. [PubMed: 19320182]
- Patterson RE, Kristal AR, Tinker LF, Carter RA, Bolton MP, Agurs-Collins T. Measurement characteristics of the Women's Health Initiative food frequency questionnaire. *Ann. Epidemiol.* 1999; 9:178–87. [PubMed: 10192650]
- Phillips DL, Pirkle JL, Burse VW, Bernert JT, Henderson LO, Needham LL. Chlorinated hydrocarbon levels in human serum: Effects of fasting and feeding. *Arch. Environ. Contam. Toxicol.* 1989; 18:495–500. [PubMed: 2505694]
- Romeo J, Warnberg J, Nova E, Diaz LE, Gomez-Martinez S, Marcos A. Moderate alcohol consumption and the immune system: A review. *Br. J. Nutr.* 2007; 98:S111–5. [PubMed: 17922947]
- Ross P, De Swart R, Addison R, Van Loveren H, Vos J, Osterhaus A. Contaminant-induced immunotoxicity in harbour seals: Wildlife at risk? *Toxicology.* 1996; 112:157–169. [PubMed: 8814345]
- Rothman N, Cantor KP, Blair A, Bush D, Brock JW, Helzlsouer K, Zahm SH, Needham LL, Pearson GR, Hoover RN, Comstock GW, Strickland PT. A nested case-control study of non-Hodgkin lymphoma and serum organochlorine residues. *Lancet.* 1997; 350:240–244. [PubMed: 9242800]
- Schisterman EF, Whitcomb BW, Louis GM, Louis TA. Lipid adjustment in the analysis of environmental contaminants and human health risks. *Environ. Health Perspect.* 2005; 113:853–857. [PubMed: 16002372]
- Shade ED, Ulrich CM, Wener MH, Wood B, Yasui Y, Lacroix K, Potter JD, McTiernan A. Frequent intentional weight loss is associated with lower natural killer cell cytotoxicity in postmenopausal women: Possible long-term immune effects. *J. Am. Diet. Assoc.* 2004; 104:903–12. [PubMed: 15175588]

- Sjödin A, Jones RS, Lapeza CR, Focant JF, McGahee EE, Patterson DG. Semiautomated high-throughput extraction and cleanup method for the measurement of polybrominated diphenyl ethers, polybrominated biphenyls, and polychlorinated biphenyls in human serum. *Anal Chem.* 2004; 76:1921–7. [PubMed: 15053652]
- Sormo EG, Larsen HJ, Johansen GM, Skaare JU, Jenssen BM. Immunotoxicity of polychlorinated biphenyls (PCB) in free-ranging gray seal pups with special emphasis on dioxin-like congeners. *J. Toxicol. Environ. Health.* 2009; 72:266–276.
- Spinelli JJ, Ng CH, Weber JP, Connors JM, Gascoyne RD, Lai AS, Brooks-Wilson AR, Le ND, Berry BR, Gallagher RP. Organochlorines and risk of non-Hodgkin lymphoma. *Int. J. Cancer.* 2007; 121:2767–2775. [PubMed: 17722095]
- Svensson BG, Hallberg T, Nilsson A, Schutz A, Hagmar L. Parameters of immunological competence in subjects with high consumption of fish contaminated with persistent organochlorine compounds. *Int. Arch. Occup. Environ. Health.* 1994; 65:351–358. [PubMed: 8034358]
- Talcott PA, Koller LD, Exon JH. The effect of lead and polychlorinated biphenyl exposure on rat natural killer cell cytotoxicity. *Int. J. Immunopathol. Pharmacol.* 1985; 7:255–61.
- Troen AM, Mitchell B, Sorensen B, Wener MH, Johnston A, Wood B, Selhub J, McTiernan A, Yasui Y, Potter JD, Ulrich CM. Unmetabolized folic acid in plasma is associated with reduced natural killer cell cytotoxicity among postmenopausal women. *J. Nutr.* 2006; 136:189–94. [PubMed: 16365081]
- Vedhara K, Fox JD, Wang EC. The measurement of stress-related immune dysfunction in psychoneuroimmunology. *Neurosci. Biobehav. Rev.* 1999; 23:699–715. [PubMed: 10392661]

Highlights

- Plasma PCBs and immune function were measured in 109 women at baseline and one year.
- Immune measures included T lymphocyte proliferation (TLP) and NK cell cytotoxicity.
- Higher-chlorinated PCBs were positively associated with TLP in cross section.
- An increase in dioxin-like PCBs was associated with a decrease in TLP over one year.
- We did not find strong evidence of impaired cellular immunity from PCB exposure.

Table 1

Baseline and change (baseline to one year) in plasma polychlorinated biphenyl concentrations (pmol/g-lipid) by participant characteristics.

Characteristic		Baseline			Baseline to one year			
	Baseline n (%)	Dioxin-like PCBs ^a , median (IQR)	High-chlorinated PCBs ^b , median (IQR)	Medium-chlorinated PCBs ^c , median (IQR)	One year n (%)	Change in dioxin-like PCBs ^a , median (IQR)	Change in high-chlorinated PCBs ^b , median (IQR)	Change in medium-chlorinated PCBs ^c , median (IQR)
Total	109 (100%)				98 (100%)			
Group								
Stretching	60 (55.0)	82.6 (57.8,134.9)	84.2 (59.8,107.9)	403.5 (313.4,529.5)	55 (56.1)	2.0 (–8.7,8.4)	–2.2 (–9.1,7.7)	10.6 (–40.0,48.7)
Exercise	49 (45.0)	91.1 (68.4,140.3)	82.8 (57.8,114.7)	408.7 (345.9,561.7)	43 (43.9)	0.0 (–6.1,13.1)	3.7 (–8.2,12.0)	4.0 (–24.1,42.6)
Baseline age (years)								
50 to 55	32 (29.4)	79.9 (52.8,99.3)	56.9 (45.3,91.7)	369.0 (270.8,466.9)	27 (27.6)	2.0 (–9.8,7.3)	–1.4 (–7.6,7.7)	4.0 (–37.7,42.6)
56 to 60	31 (28.4)	76.4 (53.9,127.9)	69.2 (49.4,99.7)	378.4 (294.7,535.3)	25 (25.5)	4.4 (–7.5,17.0)	1.8 (–7.4,11.8)	13.0 (–30.5,63.2)
61 to 65	16 (14.7)	96.6 (71.7,131.9)	84.2 (73.4,106.5)	438.2 (372.7,574.8)	15 (15.3)	–4.9 (–10.5,8.0)	–5.1 (–8.2,3.2)	–11.4 (–40.6,40.2)
66 to 70	14 (12.8)	92.6 (60.4,164.6)	105.4 (88.6,125.1)	465.7 (395.6,660.9)	15 (15.3)	–2.2 (–6.1,8.0)	–10.4 (–17.2,5.0)	–12.6 (–33.9,15.5)
71 to 75	16 (14.7)	105.0 (84.5,169.5)	134.4 (94.5,162.9)	551.8 (402.2,772.6)	16 (16.3)	4.9 (–5.2,10.4)	6.5 (–2.2,10.3)	40.3 (–24.1,80.2)
Race								
White non-Hispanic	96 (88.9)	88.4 (60.0,134.9)	84.7 (60.3,111.3)	404.2 (322.7,536.6)	86 (88.7)	0.4 (–8.6,10.1)	–0.3 (–9.0,8.9)	5.4 (–30.9,42.2)
Other	12 (11.1)	84.8 (58.0,117.4)	61.7 (44.3,88.3)	427.7 (328.0,546.5)	11 (11.3)	5.2 (–3.7,12.2)	–0.1 (–6.2,23.0)	48.7 (–16.7,91.6)
Education								
High school or less	16 (14.7)	113.7 (62.5,169.5)	91.9 (70.4,128.0)	438.2 (375.8,632.8)	17 (17.3)	–4.0 (–9.6,6.6)	2.1 (–5.6,10.4)	–12.3 (–30.8,53.1)

Characteristic		Baseline			Baseline to one year			
	Baseline n (%)	Dioxin-like PCBs ^a , median (IQR)	High-chlorinated PCBs ^b , median (IQR)	Medium-chlorinated PCBs ^c , median (IQR)	One year n (%)	Change in dioxin-like PCBs ^a , median (IQR)	Change in high-chlorinated PCBs ^b , median (IQR)	Change in medium-chlorinated PCBs ^c , median (IQR)
Vocational school, some college, or associate's degree	48 (44.0)	90.1 (63.0,134.9)	75.4 (51.3,102..2)	388.6 (290.7,531.4)	43 (43.9)	2.0 (–7.3,10.1)	–0.6 (–8.7,8.8)	16.9 (–29.3,42.2)
Bachelor's degree	25 (22.9)	86.8 (62.3,139.3)	97.7 (65.0,116.7)	466.8 (361.2, 605.4)	22 (22.4)	3.2 (–8.6,11.3)	–4.2 (–15.3,2.8)	–12.6 (–37.5,31.3)
Master's degree or doctorate	20 (18.3)	80.0 (55.7,97.7)	78.0 (58.1,109.5)	389.1 (302.0, 417.5)	16 (16.3)	3.8 (–9.4,17.1)	7.5 (–0.2,12.4)	17.8 (–24.9,58.6)
Season of blood draw								
Winter	23 (21.1)	99.4 (60.4,152.5)	80.3 (51.5,119.7)	397.1 (304.4, 527.6)	22 (22.9)	–2.3 (–9.4,6.1)	–2.0 (–10.3,8.2)	–11.6 (–32.0,15.5)
Spring	41 (37.6)	82.9 (53.9,114.2)	70.6 (59.3,102.5)	389.1 (306.3, 479.7)	37 (38.5)	5.4 (–7.3,12.6)	4.1 (–7.2,10.2)	29.0 (–32.7,63.3)
Summer	21 (19.3)	89.2 (60.4,110.8)	84.4 (55.7,116.7)	420.3 (320.4, 605.4)	17 (17.7)	0.5 (–7.5,9.3)	–3.3 (–12.3,5.0)	10.2 (–34.7,54.0)
Fall	24 (22.0)	102.1 (61.3,165.8)	93.8 (73.7,111.3)	478.1 (357.3, 566.3)	20 (20.8)	–2.1 (–10.5,3.5)	0.7 (–7.6,10.4)	–4.0 (–30.6,31.3)
Smoking status								
Never smoker	54 (49.5)	86.4 (63.9,143.8)	81.5 (56.0,109.0)	400.0 (320.4,580.3)	51 (52.0)	–0.6 (–9.9,13.1)	–1.8 (–8.9,7.6)	4.9 (–43.3,52.0)
Former smoker	55 (50.5)	89.3 (58.1,129.0)	84.4 (61.4,109.5)	420.3 (324.9,527.6)	47 (48.0)	1.7 (–6.1,8.0)	–0.2 (–8.2,11.0)	9.4 (–29.3,42.2)
Alcohol intake								

Characteristic		Baseline			Baseline to one year			
	Baseline n (%)	Dioxin-like PCBs ^a , median (IQR)	High-chlorinated PCBs ^b , median (IQR)	Medium-chlorinated PCBs ^c , median (IQR)	One year n (%)	Change in dioxin-like PCBs ^a , median (IQR)	Change in high-chlorinated PCBs ^b , median (IQR)	Change in medium-chlorinated PCBs ^c , median (IQR)
<1g/day	58 (53.2)	89.7 (58.1,149.0)	82.7 (56.0,113.1)	407.6 (294.7,567.5)	46 (49.5)	0.9 (–5.3,10.1)	–0.3 (–8.7,11.0)	13.2 (–29.9,45.5)
1–5g/day	24 (22.0)	96.2 (64.2,127.3)	76.0 (56.9,98.7)	392.3 (303.7,504.9)	28 (30.1)	1.8 (–7.5,8.6)	1.3 (–11.3,10.1)	5.1 (–30.9,55.8)
> 5g/day	27 (24.8)	82.3 (62.3,114.2)	92.2 (60.3,113.1)	426.8 (355.9,536.9)	19 (20.4)	3.6 (–9.8,13.1)	3.2 (–7.7, 8.2)	8.1 (–33.9,52.0)
Fish intake								
< 1.5/week	49 (45.4)	93.0 (58.1,134.9)	70.6 (59.3,100.4)	395.6 (320.4,498.2)	33 (37.5)	2.9 (–5.7,12.6)	–1.3 (–8.8,10.2)	2.9 (–31.8,42.6)
1.5–2.4/week	26 (24.1)	87.1 (59.1,109.7)	100.6 (56.8,119.7)	398.2 (294.7,536.3)	23 (26.1)	1.4 (–3.9,15.0)	2.8 (–3.5,11.0)	19.8 (–6.9,59.2)
> 2.5/week	33 (30.6)	89.3 (68.4,139.3)	86.5 (69.2,132.6)	449.3 (375.6,618.1)	32 (36.4)	–4.4 (–11.4,9.7)	0.7 (–10.3,8.8)	10.6 (–64.4,47.0)
Body mass index (kg/m ²)								
24 to <30	58 (53.2)	84.5 (59.1,126.8)	96.5 (65.0,124.0)	440.9 (352.8,567.5)	53 (54.6)	2.3 (–5.6,13.9)	3.9 (–8.2,11.2)	16.3 (–27.5,58.6)
30 to 34	36 (33.0)	102.5 (65.2,147.5)	77.9 (48.9,102.2)	398.0 (291.5,535.8)	33 (34.0)	–2.2 (–9.8,7.3)	–1.7 (–10.0,8.9)	–1.9 (–43.3,40.2)
35	15 (13.8)	82.9 (57.4,143.8)	49.8 (45.0,75.0)	369.7 (273.3,406.0)	11 (11.3)	0.0 (–19.5,6.1)	–3.1 (–8.2,1.7)	–20.0 (–45.0,42.2)

Interquartile range (IQR); polychlorinated biphenyl (PCB).

^aPCB 105, 118, 156.

^bPCB 194, 196/203, 199, 206, 209.

^cPCB 138/158, 146, 153, 156, 170, 180, 183, 187.

Table 2

Multivariable adjusted^a effect estimates^{b,c} (95% confidence intervals) from mixed models of associations of plasma polychlorinated biphenyl concentrations with immune function.

			NK cytotoxicity %		Immune Function	PHA-stimulated lymphocyte proliferation index ^d			
PCB Exposure	Effect Estimate	n	25:1 ^e	n	12.5:1 ^e	n	0.1µg/mL	n	0.5µg/mL
Dioxin- like ^f		186		186		182		182	
	Cross- sectional (β_1)		1.0 (– 0.1,2.1)		1.0 (0.0,1.9)		1.8 (– 1.3,4.8)		3.2 (– 3.3,9.7)
	Longitudinal (β_2)		0.7 (– 5.3,6.7)		3.2 (– 2.2,8.5)		–19.2 (– 41.8,3.5)		–51.6 (– 100.5,– 2.7)
Medium- chlorinated g		185		185		181		181	
	Cross- sectional (β_1)		0.2 (– 0.3,0.8)		0.2 (– 0.3,0.8)		1.7 (0.1,3.3)		4.0 (0.6,7.4)
	Longitudinal (β_2)		–1.0 (– 2.8,0.8)		–0.4 (– 2.0,1.2)		–2.6 (– 9.2,3.9)		–4.6 (– 18.9,9.7)
High- chlorinated h		185		185		181		181	
	Cross- sectional (β_1)		–0.3 (– 4.1,3.6)		0.2 (– 3.3,3.7)		7.3 (– 3.7,18.2)		26.3 (3.4,49.2)
	Longitudinal (β_2)		–4.9 (– 11.4,1.6)		–3.1 (– 9.1,2.9)		–5.1 (– 29.2,18.9)		–3.0 (– 55.1,49.1)
Low- chlorinated i		186		186		182		182	
	Cross- sectional (β_1)		0.7 (0.0,1.4)		0.7 (0.0,1.3)		1.1 (– 0.9,3.2)		2.0 (– 2.4,6.4)
	Longitudinal (β_2)		0.5 (– 2.8,3.8)		1.9 (– 1.0,4.8)		–9.5 (– 22.1,3.1)		–29.0 (– 56.1,– 1.9)

Natural killer (NK); polychlorinated biphenyl (PCB); phytohemagglutinin (PHA).

^a Adjusted for age, race, education, intervention group, smoking status, blood draw season, alcohol intake, fish intake, and body mass index

^b Cross-sectional effect estimate (β_1) represents the mean change in immune function measure per 50 pmol/g-lipid difference in PCB concentration

^c Longitudinal effect estimate (β_2) represents the mean change in immune function measure difference per 50 pmol/g-lipid change in PCB concentration difference from baseline to one year

^d PHA=Counts per minute of simulated cells/counts per minute of unstimulated cells

^e Effector-to-target ratio

f PCB 105, 118, 156

g PCB 138/158, 146, 153, 156, 170, 180, 183, 187

h PCB 194, 196/203, 199, 206, 209

i PCB 74, 99, 105, 118