Cloned human snRNP proteins B and B' differ only in their carboxy-terminal part

Alje van Dam, Irene Winkel,
Josee Zijlstra-Baalbergen, Ruud Smeenk and
H.Theo Cuypers

Central Laboratory of the Red Cross Blood Transfusion Service,
Plesmanlaan 125, 1066 CX Amsterdam, The Netherlands

Communicated by H.L.Ploegh

Antibodies to the snRNP proteins B' and B are frequently observed in autoimmune diseases. We analyzed different types of cDNAs encoding these proteins. One type of cDNA encoded a protein whose predicted mol. wt is 24.6 kd, whereas another type encoded a protein with a predicted mol wt of 23.7 kd. When translated *in vitro* from cDNA transcripts, the apparent mol. wts of these proteins on SDS–polyacrylamide gel were 29.5 and 28.5 kd respectively. The main difference between these two types of cDNAs proved to be the presence of an additional sequence of 146 nucleotides in the 3' part of the open reading frame of the clone encoding the shorter protein. This insert contains a termination codon in frame, at 9 nucleotides downstream from the 5' end of the insert. The additional sequence revealed at the 3' end a consensus sequence of vertebrates for intron–exon junctions. We demonstrated the presence of mRNAs corresponding with both types of cDNA in human cells. We hypothesize that the B' and B protein are derived from one pre-mRNA by alternative splicing, and show that they differ only at the carboxy terminus, where a proline rich motive is repeated once more in B'. A comparison with amino acid sequences of other cloned snRNP proteins is included.

Key words: alternative splicing/anti-Sm antibodies/snRNP/systemic lupus erythematosus

Introduction

Eukaryotic cells synthesize a group of abundant small nuclear ribonucleoprotein (snRNP) particles, each containing an uridine (U)-rich RNA molecule and a number of proteins (Reddy and Busch, 1988). Substantial evidence now exists that the U1, U2, U4, U5 and U6 snRNPs all play an essential role in pre-mRNA splicing in the nucleus of the cell (reviewed by Steitz *et al.*, 1988). The snRNP particles are contained in large 50–60S structures, normally referred to as 'spliceosomes', in which pre-mRNA splicing occurs (Frendewey *et al.*, 1987; Konarska and Sharp, 1987; Zillmann *et al.*, 1988). The 50–60S structures contain the products of the first step of splicing, 5'-exon and lariat exon, indicating that these spliceosomes are functional structures (Grabowski *et al.*, 1985; Agris *et al.*, 1989).

In addition to the U-RNA molecules, the snRNPs contain a number of proteins, some of which are shared by all snRNPs (B, B', D, D', E, F and G), whereas others are

specific to the type of U-RNA (70k, A and C for U1-RNP, and A' and BB' for U2-RNPs) (Hinterberger *et al.*, 1983; Billings and Hoch, 1984; Pettersson *et al.*, 1984; Bringmann and Lührmann, 1986). The exact role of the proteins in the splicing process remains to be established, but U1 snRNP assembly studies *in vivo* and *in vitro* made it possible to determine the protein distribution on the U1-RNA (Fresco *et al.*, 1987; Hamm *et al.*, 1987, 1988).

Antibodies against the different snRNP proteins are a characteristic feature of patients with autoimmune diseases. So-called anti-Sm antibodies, defined in a counter-immunoelectrophoresis method with the use of a standard reference serum, have been shown to react with the snRNP proteins B, B', D and to a lesser extent with E (Tan, 1982). Since anti-Sm antibodies are only rarely produced by patients with other connective tissue diseases, the presence of this specificity serves as a useful marker for systemic lupus erythematosus (Tan, 1982). Molecular characterization of the snRNP-associated proteins and their immunoreactive epitopes can be helpful in understanding their role in the splicing process, as well as in unravelling the mechanisms responsible for the autoantibody response in autoimmune patients.

A number of snRNP-associated proteins have recently been cloned, including the U1-RNP proteins 70K (Theissen *et al.*, 1986; Spritz *et al.*, 1987), A (Sillekens *et al.*, 1987), and C (Sillekens *et al.*, 1988; Yamamoto *et al.*, 1988), the U2 snRNP proteins A' (Sillekens *et al.*, 1989) and B' (Habets *et al.*, 1987), and the Sm snRNP proteins D, E and N (Stanford *et al.*, 1987; Rokeach *et al.*, 1988; Schmauss *et al.*, 1989). Furthermore it has been reported that B/B' and D share antigenic determinants, as was demonstrated by cross-reactivity of affinity-purified human sera (Habets *et al.*, 1985) and monoclonal antibodies (Williams *et al.*, 1986). However, Reuter *et al.* (1987) concluded from peptide mapping and cross-reactivity studies with polyclonal rabbit antisera that the D protein is structurally different from B/B'. Their data support a striking homology between the proteins B and B' and they suggest that B' differs from B only by an additional amino- or carboxy-terminal amino acid sequence. Another snRNP associated polypeptide called 'N', containing an anti-Sm epitope, has been cloned from rat and human cDNA libraries and is expressed in a tissue-specific manner in brain and to a lesser extent in heart (McAllister *et al.*, 1988). Rokeach *et al.* (1989) have determined the amino acid sequence of the first 19 amino-terminal-residues of Sm-B and the first 20 of Sm-B' from the polypeptides purified from HeLa cells. Both polypeptides share an identical amino-terminal sequence, which is also present as an amino-terminal sequence in the proposed rat and human N-protein sequences (McAllister *et al.*, 1989; Schmauss *et al.*, 1989).

In this paper we report the characterization of different cDNA clones, encoding the proteins B and B', and show that both mRNAs most likely are derived from a common

Fig. 1. Characterization of β -galactosidase B/B' fusion proteins by immunoblotting. Proteins of the induced lysogenic strains, encoding β -galactosidase (lane 1), fusion protein of clone 5 (lane 2), of clone 9 (lane 3), and of clone 16 (lane 4) were separated on a 6% SDS-polyacrylamide gel and Western blotted. The blot on the left was subsequently stained for total protein with Ponceau-S. The blot in the centre was incubated with the monoclonal antibody CLB-ANA-125. The blot on the right was incubated with a human anti-Sm serum. The centre and right blot were subsequently incubated with enzyme-labeled anti-mouse and anti-human conjugates and with substrates as described in Materials and methods. At the left, the mol. wt markers are indicated.

pre-mRNA by differential processing events. The determined nucleotide sequences are related to the other published B/B' clones (Ohosone *et al.*, 1989; Rokeach *et al.*, 1989), and the predicted amino acid sequences are compared to other snRNP proteins that have been cloned.

Results

Selection and identification of cDNA clones

A mouse monoclonal antibody (CLB-ANA-125) was shown to react with affinity-purified Sm protein from rabbit thymus. On HeLa nuclear and cytoplasmic blots CLB-ANA-125 reacted specifically with B and B'. With this monoclonal antibody, 700 000 independent clones of a λ gt11 expression library, made from poly(A)⁺ RNA of HL60 cells, were screened. We obtained three clones (nos. 5, 9 and 16), which also showed a positive reaction with a second monoclonal antibody (CLB-ANA-98) reactive with B/B', and with a human serum containing antibodies to B and B'. No reaction was observed with control sera and with a panel of control monoclonal antibodies. Screening of $\sim 1.5 \times 10^6$ independent recombinant phages from a human placenta cDNA library with a random-primed labeled cDNA insert of clone 16, yielded another 19 independent cross-hybridizing recombinants. Rescreening of these recombinants with a synthetic oligonucleotide complementary to the 5' end of clone 16 revealed that eight of these clones also hybridized with this oligonucleotide.

To characterize β -galactosidase fusion proteins, clones 5, 9 and 16 were introduced into a temperature-sensitive lysogenic host. After induction, cell lysate of these strains was separated on an SDS-polyacrylamide gel and fusion

Fig. 2. Immunoblotting with antibodies affinity-purified from Sm-B' fusion protein. A mixture of a serum containing anti-Sm antibodies and a serum containing antibodies against ribosomal P-proteins was absorbed on two nitrocellulose strips, containing clone 16 fusion protein and β -galactosidase respectively. Bound antibodies were eluted and reused to probe an immunoblot with HeLa cytoplasmic cell extract. Lanes 1, the antibodies, eluted from the control β -galactosidase strip; 2, affinity-purified antibodies eluted from clone 16 fusion protein; 3, the mixture of anti-Sm and anti-P serum, used as starting material for the affinity purification; 4, anti-Sm serum; 5, anti P-serum. Bands were visualized with horseradish peroxidase conjugated anti-human immunoglobulin and substrate, as described.

proteins were analyzed by immunoblotting. The fusion proteins of clones 5, 9 and 16 were reactive with CLB-ANA-125 and with a human anti-Sm serum (Figure 1). The fusion proteins had apparent mol. wts of ~ 140 kd. The fusion protein encoded by clone 16 was reactive on immunoblots with eight different anti-Sm sera, whereas no reactivity was found with sera containing antibodies specific for 70k, A, centromere, SS-B or rRNP (results not shown).

To prove that human antibodies reactive with fusion protein encoded by clone 16 recognized B/B', we separated extracts from lysogenic strains containing either fusion protein or control β -galactosidase protein on SDS-polyacrylamide gels. The proteins were subsequently blotted onto nitrocellulose paper and identified by Ponceau-S protein staining. The areas of each blot containing the fusion protein or the β -galactosidase were excised. To prove that remaining anti-B/B' activity of affinity-purified antibodies was not due to aspecific binding of IgG to fusion protein, we used a 1:1 mixture of two sera for incubation with these strips. One serum contained anti-Sm/RNP antibodies; the other serum contained antibodies directed to the ribosomal proteins P0, P1 and P2 (Francoeur *et al.*, 1985). Bound antibodies were eluted from the strips and subsequently incubated with a HeLa cytoplasmic blot (Figure 2). Antibodies eluted from the clone 16 fusion protein strongly reacted with both B and B', whereas no reactivity with P0, P1 and the Sm-D protein was observed. In addition, the antibodies eluted from the fusion protein also recognized the 33 kd RNP A protein.

Fig. 3. Partial restriction map and sequence strategy for the cDNA inserts of clone 16 (A) and clone 5 (B). Restriction enzyme sites: B, *Bam*HI; E, *Eco*RI; P, *Pst*I; S, *Sph*I; Sm, *Sma*I; T, *Taq*I. The thick bars represent the protein coding regions. The open part of the bar in (B) marks the site of the additional sequence of 146 nucleotides in clone 5. The determined nucleotide sequences (horizontal arrows) were obtained from subclones and in part from the use of synthetic oligonucleotide primers.

The eluate from the control β -galactosidase strip was not reactive.

Characterization of cDNA inserts in reactive clones

Determination of the insert size revealed a 1.1 kb cDNA insert for clone 5, whereas clone 16 contained an 0.9 kb insert. From partial restriction enzyme analysis and DNA sequencing we concluded that we had isolated almost completely identical cDNAs, which differed only in the 3'-situated restriction fragment *Bam*HI–*Eco*RI (Figure 3). Sequencing of the cDNA inserts was performed on both strands by employing the dideoxy method. Partial restriction maps of the clones and the sequencing strategy are shown in Figure 3, and the DNA sequences are presented in Figure 4. The cDNA insert of clone 5 comprises 1049 nucleotides and contains an open reading frame, encoding 249 amino acids in frame with the β -galactosidase coding region. The first ATG within this fusion reading frame is at position 55. This codon is situated within a sequence propitious for translation-initiation, the sequence 50–58 CCACCATGA only diverges in the last position from the optimal context for initiation (Kozak, 1986). If the methionine residue encoded by this codon is considered as the initiation site for translation, the total length of the encoded protein is 231 amino acids. The TGA translation stop signal at position 748 is followed by a non-coding region of 298 nucleotides. At the 3' end a stretch of five adenines is found, which is preceded by the polyadenylation signal AATAAA at nucleotide 1015 (Leff *et al.*, 1986). Clone 9 starts at exactly the same nucleotide as clone 5 and is identical but for nucleotide 359, where a G/T substitution leads to an amino acid mutation from glycine to valine.

The cDNA insert from clone 16 covers 914 nucleotides

(Figure 4). This clone starts 15 nucleotides more downstream compared to clone 5. Subsequently, 719 nucleotides are identical in the two clones, up to position 740. At this position, an extra insert of 146 nucleotides is found in clone 5. From position 886 onwards, up to the 3' end, nucleotide sequences of clones 5 and 16 are again identical. Absence of the 146 nucleotide fragment in clone 16, which contains the translation stop codon in clone 5, results in a longer coding region up to the TAG stop codon at position 921. The deduced amino acid sequence encoded by clone 16 is, therefore, nine amino acids longer at the carboxy terminus.

Restriction enzyme analyses of four B/B' cDNA clones, obtained from the human placenta cDNA library, which were homologous up to the 5' end of clone 16 (see above), revealed that two of these clones comprised the insert sequence of 146 nucleotides (identical to clone 5), whereas the other two lacked the insert. Sequence analyses of one representative of each group confirmed these results (not shown).

A striking feature of the 3' end of the insert of 146 nucleotides in clone 5 is the excellent homology with the consensus sequence for the 3' intron acceptor junction for pre-mRNA splicing of vertebrates (Padgett *et al.*, 1986) (Figure 5). The homology to the consensus site includes 12 nucleotides at the 5' site of the junction and the G residue at the 3' site. No homology with the consensus sequence for the 5' intron donor junction was found at the 5' end of the insert. These results indicate, that clone 5 and 16 are representative for two mRNAs, which probably are derived from one pre-mRNA by alternative splicing from one donor site to two different acceptor sites.

In vitro translation and immunoprecipitation

The identity of the cDNAs was confirmed by *in vitro* transcription, followed by *in vitro* translation of the resulting RNAs in a rabbit reticulocyte lysate. The synthesized products were run on an SDS–polyacrylamide gel and compared to [35 S]methionine labeled HeLa snRNP proteins, which were immunoprecipitated with a human anti-Sm serum. The major translation product of clone 16 was a protein with a mol. wt of 29.5 kd on SDS gels, whereas a number of smaller bands could be seen (Figure 6, lane 2). *In vitro* translation of clone 5 resulted in a major band of 28.5 kd (Figure 6, lane 3). We observed a slight difference in mol. wt between the translation products of clones 16 and 5, and *in vivo* labeled proteins B' and B (29 and 28 kd respectively) (Figure 6, lane 1). All translation products from clones 5 and 16 could be immunoprecipitated with an anti-Sm serum (Figure 6, lanes 5 and 6), in contrast to the translation products of the control TMV RNA (lane 8). If normal human serum was used for immunoprecipitation, a faint band was seen at the site of the major translation product after prolonged exposure (lane 7). From these experiments we conclude that the protein encoded by clone 16 is the snRNP B' protein, whereas clone 5 encodes B.

Characterization of mRNA

Northern blot analysis of total RNAs from RC2A and Jurkat cells with cDNAs from clones 5 and 16 resulted in a single broad band of ~ 1.3 kb (not shown). Since we presumed that the two mRNA species, differing by only 146 nucleotides in length, could not be separated on an agarose gel, we performed an RNase mapping experiment, using the

5	1	GAATTTCCTGAGGAGGCGAATCCGGCGGGTATCAGAGCCATCAGAACC <u>GCACCC</u>																				ATG	ACG	GTG	GGC	AAG	AGC														
16		GAATTC.....CCGGCGGGTATCAGAGCCATCAGAACC <u>GCACCC</u>																				ATG	ACG	GTG	GGC	AAG	AGC														
B/B'																						Met	Thr	Val	Gly	Lys	Ser														
																						1				5															
5	73	AGC	AAG	ATG	CTG	CAG	CAT	ATT	GAT	TAC	AGG	ATG	AGG	TGC	ATC	CTG	CAG	GAC	GGC	CGG	ATC	TTC	ATT	GGC	ACC																
16		AGC	AAG	ATG	CTG	CAG	CAT	ATT	GAT	TAC	AGG	ATG	AGG	TGC	ATC	CTG	CAG	GAC	GGC	CGG	ATC	TTC	ATT	GGC	ACC																
B/B'		Ser	Lys	Met	Leu	Gln	His	Ile	Asp	Tyr	Arg	Met	Arg	Cys	Ile	Leu	Gln	Asp	Gly	Arg	Ile	Phe	Ile	Gly	Thr																
																						10	15	20	25	30															
5	145	TTC	AAG	GCT	TTT	GAC	AAG	CAC	ATG	AAT	TTG	ATC	CTC	TGT	GAC	TGT	GAT	GAG	TTC	AGA	AAG	ATC	AAG	CCA	AAG																
16		TTC	AAG	GCT	TTT	GAC	AAG	CAC	ATG	AAT	TTG	ATC	CTC	TGT	GAC	TGT	GAT	GAG	TTC	AGA	AAG	ATC	AAG	CCA	AAG																
B/B'		Phe	Lys	Ala	Phe	Asp	Lys	His	Met	Asn	Leu	Ile	Leu	Cys	Asp	Cys	Asp	Glu	Phe	Arg	Lys	Ile	Lys	Pro	Lys																
																						35	40	45	50																
5	217	AAC	TCC	AAA	CAA	GCA	GAA	AGG	GAA	GAG	AAG	CGA	GTC	CTC	GGT	CTG	GTG	CTG	CTG	CGA	GGG	GAG	AAT	CTG	GTC																
16		AAC	TCC	AAA	CAA	GCA	GAA	AGG	GAA	GAG	AAG	CGA	GTC	CTC	GGT	CTG	GTG	CTG	CTG	CGA	GGG	GAG	AAT	CTG	GTC																
B/B'		Asn	Ser	Lys	Gln	Ala	Glu	Arg	Glu	Glu	Lys	Val	Leu	Gly	Leu	Val	Leu	Leu	Arg	Gly	Glu	Asn	Leu	Val																	
																						55	60	65	70	75															
5	289	TCA	ATG	ACA	GTA	GAG	GGA	CCT	CCT	CCC	AAA	GAT	ACT	GGT	ATT	GCT	CGA	GTT	CCA	CTT	GCT	GGA	GCT	GCC	GGG																
16		TCA	ATG	ACA	GTA	GAG	GGA	CCT	CCT	CCC	AAA	GAT	ACT	GGT	ATT	GCT	CGA	GTT	CCA	CTT	GCT	GGA	GCT	GCC	GGG																
B/B'		Ser	Met	Thr	Val	Glu	Gly	Pro	Pro	Pro	Lys	Asp	Thr	Gly	Ile	Ala	Arg	Val	Pro	Leu	Ala	Gly	Ala	Gly																	
																						80	85	90	95	100															
5	361	GGC	CCA	GGG	ATC	GGC	AGG	GCT	GCT	GGC	AGA	GGA	ATC	CCA	GCT	GGG	GTT	CCC	ATG	CCC	CAG	GCT	CCT	GCA	GGA																
16		GGC	CCA	GGG	ATC	GGC	AGG	GCT	GCT	GGC	AGA	GGA	ATC	CCA	GCT	GGG	GTT	CCC	ATG	CCC	CAG	GCT	CCT	GCA	GGA																
B/B'		Gly	Pro	Gly	Ile	Gly	Arg	Ala	Ala	Gly	Arg	Gly	Ile	Pro	Ala	Gly	Val	Pro	Met	Pro	Gln	Ala	Pro	Ala	Gly																
																						105	110	115	120	125															
5	433	CTT	GCT	GGG	CCA	GTC	CGT	GGG	GTT	GGC	GGG	CCA	TCC	CAA	CAG	GTG	ATG	ACC	CCA	CAA	GGA	AGA	GGT	ACT	GTT																
16		CTT	GCT	GGG	CCA	GTC	CGT	GGG	GTT	GGC	GGG	CCA	TCC	CAA	CAG	GTG	ATG	ACC	CCA	CAA	GGA	AGA	GGT	ACT	GTT																
B/B'		Leu	Ala	Gly	Pro	Val	Arg	Gly	Val	Gly	Gly	Pro	Ser	Gln	Gln	Val	Met	Thr	Pro	Gln	Gly	Arg	Gly	Thr	Val																
																						130	135	140	145	150															
5	505	GCA	GCC	GCT	GCA	GCT	GCT	GCC	ACA	GCC	AGT	ATT	GCC	GGG	GCT	CCA	ACC	CAG	TAC	CCA	CCT	GGC	CGT	GGG	GGT																
16		GCA	GCC	GCT	GCA	GCT	GCT	GCC	ACA	GCC	AGT	ATT	GCC	GGG	GCT	CCA	ACC	CAG	TAC	CCA	CCT	GGC	CGT	GGG	GGT																
B/B'		Ala	Ala	Ala	Ala	Ala	Ala	Thr	Ala	Thr	Ala	Ser	Ile	Ala	Gly	Ala	Pro	Thr	Gln	Tyr	Pro	Gly	Arg	Gly	Gly																
																						155	160	165	170																
5	577	CCT	CCC	CCA	CCT	ATG	GGC	CGA	GGG	GCA	CCC	CCT	CCA	GGC	ATG	ATG	GGC	CCA	CCT	CCT	GGT	ATG	AGA	CCT	CCT																
16		CCT	CCC	CCA	CCT	ATG	GGC	CGA	GGG	GCA	CCC	CCT	CCA	GGC	ATG	ATG	GGC	CCA	CCT	CCT	GGT	ATG	AGA	CCT	CCT																
B/B'		Pro	Pro	Pro	Pro	Met	Gly	Arg	Gly	Ala	Pro	Pro	Pro	Gly	Met	Met	Gly	Pro	Pro	Pro	Gly	Met	Arg	Pro	Pro																
																						175	180	185	190	195															
5	649	ATG	GGT	CCC	CCA	ATG	GGG	ATC	CCC	CCT	GGA	AGA	GGG	ACT	CCA	ATG	GGC	ATG	CCC	CCT	CCG	GGA	ATG	CGG	CCT																
16		ATG	GGT	CCC	CCA	ATG	GGG	ATC	CCC	CCT	GGA	AGA	GGG	ACT	CCA	ATG	GGC	ATG	CCC	CCT	CCG	GGA	ATG	CGG	CCT																
B/B'		Met	Gly	Pro	Pro	Met	Gly	Ile	Pro	Pro	Gly	Arg	Gly	Thr	Pro	Met	Gly	Met	Pro	Pro	Pro	Gly	Met	Arg	Pro																
																						200	205	210	215	220															
5	721	CCT	CCC	CCT	GGG	ATG	CGA	GGC	CTT	CTT	TGA	CCCTTGGCCACAGAGTATGGAAGTAGCTCCGCAGAGGCGTGCGCTCGATTCTCTCA																													
16		CCT	CCC	CCT	GGG	ATG	CGA	G.....																																	
B		Pro	Pro	Pro	Gly	Met	Arg	Gly	Leu	Leu	TER																														
B'		Pro	Pro	Pro	Gly	Met	Arg	Gly																																	
																						225	230																		
5	806	GGGCCACGTTACACAGACCTGTTTGTTCCTTATGCTGTTGTTCGTGGAGTCTCATGGGATTGTCTGGTTTCCCTTACAG																																							
16															*****																										
B'																								GG	GG	CCC	CCT	CCC													
																											Pro	Pro	Pro												
5	897	CCG	GGA	ATG	CGC	CCA	CCA	AGG	CCC	TAG	ACTCATCTTGGCCCTCCTCAGCTCCCTGCCTGTTTCCCGTAAGGCTGTACATAGTCCTT																														
16		CCG	GGA	ATG	CGC	CCA	CCA	AGG	CCC	TAG	ACTCATCTTGGCCCTCCTCAGCTCCCTGCCTGTTTCCCGTAAGGCTGTACATAGTCCTT																														
B'		Pro	Gly	Met	Arg	Pro	Pro	Arg	Pro	TER																															
																						235	240																		
5	983	TTATCTCCTTGTGGCCTATGAAACTGGTTTATAATAAACTCTTAAGAGAACATTATAATTGCAAAAA																																							
16		TTATCTCCTTGTGGCCTATGAAACTGGTTTATAATAAACTCTTAAGAGAACATTATAATTGCAAAAA																							1049																
																												914													

Fig. 4. Nucleotide sequence of clones 5 and 16, and deduced amino acid sequence of B' and B. The predicted amino acids are shown below their triplets. Nucleotides are numbered starting at the first nucleotide of clone 5. Nucleotides are numbered on the left, amino acids below the sequence. The consensus sequences for initiation as proposed by Kozak (1986), and the polyadenylation signal (Leff *et al.*, 1986) are underlined. The consensus sequence for the intron-exon junction is indicated with asterisks. Boxes indicate repetitive motives, consisting of several proline residues, followed by the sequence Gly-Met-Arg, or Met-Gly-Arg.

anti-sense *Bam*HI–*Sma*I fragments of clones 5 and 16 as probe. The fragment of clone 5 spans the additional insert sequence in this clone (see Figure 3). The uniformly labeled fragments of clones 16 and 5 comprise, including the transcribed pSP65 fragment, 109 and 255 nucleotides respectively (Figure 7, lanes 1 and 2). The most abundant RNase-resistant product produced, if HeLa cell RNA and HL60 RNA [total cytoplasmic and poly(A)⁺] were hybridized to the clone 5 *Bam*HI–*Sma*I RNA probe, was

233 ± 3 nucleotides in length (Figure 7, lanes 4, 8, 13 and 15). This value corresponds with the fragment we could expect to be protected by mRNA corresponding to clone 5. If the fragment *Bam*HI–*Sma*I of clone 16 was used as hybridization probe, the expected fragment of 87 ± 2 nucleotides was protected (Figure 7, lanes 3, 7, 12 and 14).

From these results we conclude that HL60 and HeLa cells contain the two cytoplasmic mRNA species, which correspond with the cDNA clones 5 and 16 respectively.

Fig. 5. Comparison of the nucleotide sequence of the insert in clone 5 to the consensus sequence of vertebrates for the intron-exon junctions (Padgett *et al.*, 1986). Sequences were compared both at the 5' and 3' ends of the insert. 'Y' represents a pyrimidine residue.

Fig. 6. *In vitro* translation of clone 5 and 16 encoded RNAs. In lanes 1–3 the size of *in vivo* labeled B' and B was compared to the size of the translated products. **Lanes 1**, an immunoprecipitation of a [35 S]-methionine-labeled HeLa extract; **2 and 3**, [35 S]-methionine-labeled products translated *in vitro*, from synthetic mRNAs obtained from clones 16 and 5 respectively; **4**, the result of a control *in vitro* translation of TMV RNA in the presence of [35 S]-methionine; **5 and 6**, immunoprecipitation of the translation products of clones 16 and 5 respectively, with a human anti-Sm serum; **7**, the result of a control immunoprecipitation of *in vitro* translation product from clone 5 with a normal human serum; **8**, control immunoprecipitation of the translation product of TMV RNA with the anti-Sm serum.

Discussion

We have isolated cDNA clones, encoding the human Sm-B' and B proteins. The characteristic difference between the clones encoding B' and B is the existence of an additional sequence of 146 nucleotides at the 3' end of the coding region of the clone encoding B, which is absent in the clone encoding B'. This results in the formation of two proteins, which are identical but for their carboxy-terminal end. The clone containing the 146 bp insert encodes a protein of 231 amino acids, as the result of a termination codon in the insert at 9 nucleotides downstream from the start of the insert. The clone without insert encodes a protein comprising 240 amino acids, including one more repeat of a proline-rich motif at the carboxy-terminal end. RNase-mapping experiments showed that mRNAs corresponding to both types of cDNAs were present in HeLa cells.

In vitro transcription and translation of the clone without

Fig. 7. RNase mapping of B/B' mRNAs. Uniformly labeled anti-sense RNA probes, corresponding to the *Bam*HI–*Sma*I fragments of clone 5 and 16 were hybridized to either HeLa or HL60 RNA (total cytoplasmic and poly(A)⁺ RNA), or as control to tRNA or no RNA. The samples were subsequently digested with RNases T1 and A. The protected fragments were run on a 6% polyacrylamide, 6 M urea gel. Sizes of protected fragments were determined by comparison to M13 sequencing ladders and to end-labeled *Hae*III fragments of pBR322 (lane indicated with M), run on the same gel. **Lanes 1 and 2**, uniformly labeled probes of clone 16 (**lane 1**) and 5 (**lane 2**), used as input in the protection experiments; **3 and 4**, 20 μ g total HeLa RNA + the probe of clone 16 (**lane 3**) and 5 (**lane 4**); **5 and 6**, control experiment without RNA of the probe of clone 16 (**lane 5**) and of clone 5 (**lane 6**); **7 and 8**, 2 μ g HL60 poly(A)⁺ RNA + the probe of clone 16 (**lane 7**) and of clone 5 (**lane 8**); **9**, empty lane; **10**, 20 μ g yeast tRNA + the probe of clone 16; **11**, 10 μ g total HL60 RNA + the probe of clone 16; **12**, 20 μ g yeast tRNA + the probe of clone 5; **13**, 10 μ g HL60 RNA + the probe of clone 5; **14 and 15**, 2 μ g HeLa poly(A)⁺ RNA + the probe of clone 16 (**lane 14**) and of clone 5 (**lane 15**). The sizes of the uniformly labeled input fragments and of the protected fragments are indicated on the right. The protected band in lane 11 is of low intensity and hardly visible in the figure.

insert resulted in a protein with a mol. wt on SDS-polyacrylamide gel of 29.5 kd, whereas the cDNA clone with the insert encoded a protein with an apparent mol. wt of 28.5 kd. These *in vitro* translation products had a slightly higher mol. wt than the *in vivo* synthesized B' and B. Such a difference of ~ 1.0 kd is also observed for the U1-snRNP-specific C protein (Sillekens *et al.*, 1988), and there is experimental evidence that this C protein is post-translationally modified *in vivo* (Fisher *et al.*, 1985).

Both *in vitro* translated proteins could be immunoprecipitated with a human anti-Sm serum. The observed difference in mol. wt between the two *in vitro* synthesized

	1	20	40	60	80	100
B' (1)	MTVCKSSKMLQHIDYRMCTIQDGRIFGTGKAFDKHMLILCCDEFKTIKPKNSKQAEREKRVLGLVLLRGENLVSMTVEGPPPKDTGIARVPLAGA					
B (1)	-----					
<hr/>						
N (2)	-----A--P-----					
B/B' (3)	-----A--P-----					
<hr/>						
	120	140	160	180	200	
B' (1)	ACGPGIGRAAGRGIPAGVMPQAPAGLAGPVRGVGGPSQVMTPQGRGTVAATAAATASIAGAPTQYPPGRGGPPPMGRGAPPPGMGPPPGMRPPMG					
B (1)	-#-----					
<hr/>						
N (2)	----V-----V-----I-----V-----T--V--AT--I-A-----					
B/B' (3)	----V-----V-----I-----V-----T--V--AT--I-A-----					
<hr/>						
	220	240				
B' (1)	PPMGIPPGRGTHGMPPPGMRPPPPGMRGPPPPGMRPPRP					
B (1)	-----LL					
<hr/>						
N (2)	--I-L--A--I-----I-----					
B/B' (3)	--I-L--A--I-----I-----					

Fig. 8. Comparison of different published amino acid sequences of B, B' and N. (1) This report; (2) Schmauss *et al.* (1989), McAllister *et al.* (1989); (3) Rokeach *et al.* (1989). Only amino acids differing from B' are shown. On position 102 in B, indicated with #, a polymorphism was found in clone 9, resulting in a triplet encoding a valine instead of a glycine. After correction of the data, the cDNA clone encoding the B protein described by Ohosone *et al.* (1989) was shown to be identical to the cDNA clone encoding B described in this article (Y.Ohosone *et al.*, correction submitted for publication).

products is in accordance with the difference in mol. wt between B' and B, which is generally estimated to be 1 kd. The difference in calculated mol. wt (B', 24 613 daltons; B, 23 659 daltons), and mol. wt observed on SDS gels may be explained by aberrant electrophoretic properties of the proteins. Such an anomalous migration of polypeptides on SDS-polyacrylamide gels is more often observed for nuclear proteins, the reason for this being unknown (Earnshaw *et al.*, 1987; Spritz *et al.*, 1987).

In our cDNA clones, the AUG at position 55 is preceded by a nearly perfect sequence for initiation, and is therefore probably used as a translation-initiation site *in vivo*. Minor bands, as seen in Figure 6, can be explained by the use of more downstream AUG codons as initiation sites for translation. Arguments for the use of AUG position 55 as initiation site for translation *in vivo* can be obtained from the protein sequence experiments of Rokeach *et al.* (1989). Amino acid sequencing of the first 20 amino-terminal residues of Sm-B' and the first 19 of Sm-B showed that both polypeptides share an identical amino-terminal sequence. These sequences correspond with our proposed B' and B sequences from position 2. Probably the amino-terminal methionine is removed during post-translational processing. Because Rokeach *et al.* (1989) purified their B' and B polypeptides from HeLa cells, which do not express the N-protein (McAllister *et al.*, 1989), mistaking the B/B' proteins for the N-protein is not very likely. From this fact and from the immunological data we conclude that the clones we obtained encode B and B' respectively.

The high homology between B, B' and N has caused a lot of confusion. Although the amino-terminal amino acid sequences of B/B' and N are identical (Schmauss *et al.*, 1989), obvious amino acid differences are observed further on in the sequences, resulting in 92.5% homology on amino acid sequence level for N and B' (Figure 8). The non-coding regions of the B/B' cDNA sequence we present and the published human N-protein cDNA (Schmauss *et al.*, 1989) are completely non-homologous, indicating that N and B'

are encoded by different genes. Since in our view both B' and N contain 240 amino acids, their co-migration on SDS gels is by no means surprising. Differences between N and B' are found in 17 amino acids (Figure 8). All charged amino acids are identical in both proteins. The most remarkable difference, occurring six times, is a substitution of a valine or isoleucine in the N-protein to a methionine residue in B'. It is still unclear whether the minor differences in protein structure between B/B' and N give rise to differences in protein function. Using oligonucleotides derived from the amino-terminal amino acid sequence, Rokeach *et al.* (1989) characterized cDNA clones from a human lymphocyte (Raji cells) cDNA library, which also included the B/B' amino-terminal amino acid sequence. Their cDNAs are, however, completely homologous, also in the non-coding regions, to the human N-protein cDNA as published by Schmauss *et al.* (1989). Therefore, it must be concluded that they isolated cDNA clones encoding N. This is in agreement with the fact that Raji cells were shown to contain both mRNA species—a 1.3 kb message encoding B/B' and a 1.6 kb mRNA encoding the N-protein (McAllister *et al.*, 1989).

During revision of this manuscript, another Sm B'/B cDNA was published (Ohosone *et al.*, 1989; Schmauss *et al.*, 1989). This cDNA resembles closely our B clone, but the authors propose an amino acid sequence that differs in several places from our proposed B-polypeptide, among others at the carboxy terminus, where they propose an additional stretch of 57 residues. Resequencing of the clone described by Ohosone *et al.* (1989) has shown that their clone is completely homologous, also in the non-coding regions, to our clone 5 (Figure 4) (Y.Ohosone *et al.*, correction submitted). A comparison of the complete data concerning the published amino acid sequences of B'/B and N is shown in Figure 8.

In our view, B' and B result from alternative splicing of one pre-mRNA. Arguments for this hypothesis are: (i) the existence of a sequence representing a consensus sequence for the 3' intron acceptor site for splicing at the 3' end of

the additional sequence of 146 nucleotides in clone 5; (ii) the completely identical nucleotide sequences also in the non-coding regions of the cDNAs 5 and 16, with the exception of the insert sequence; (iii) the demonstration of two different types of mRNA in cells; and (iv) the indication that genomic DNA contains only one copy of the B/B' gene, as concluded from Southern blot analysis of genomic DNA with the cDNA insert (result not shown). The existence of different mRNAs for B and B' has already been suggested by Fisher *et al.* (1983). Moreover, they already suggested that after size fractionation of poly(A)⁺ RNA by sucrose gradient centrifugation, the peak fractions of mRNA encoding B and B' were in reverse order relative to the size of the proteins. Our results are also in agreement with the protein mapping findings of Reuter *et al.* (1987), who found a strong structural homology between B' and B, and suggested that B' differs from B by an additional amino- or carboxy-terminal amino acid sequence. Alternative splicing of pre-mRNA has been suggested earlier for the U1-RNP-specific 70K protein (Spritz *et al.*, 1987).

The main characteristic of the proteins B/B' is the extreme proline-richness at their carboxy-terminal end. From amino acid 165 onwards, 45% of the B' protein consists of proline residues. The sequence PPPGMR is repeated four times in B' and three times in B. An RNP consensus sequence (Swanson *et al.*, 1987) is absent. The nucleic acid binding property of the proteins B and B' can in part be accounted for by the relatively high content of positively charged amino acids (12%), in comparison to negatively charged residues (5%). Also, proline-rich domains have been described in nucleic acid binding proteins (Garoff *et al.*, 1980; Kruijer *et al.*, 1981).

Since shorter polypeptides than the full-length protein, synthesized by the use of alternative AUG codons during *in vitro* translation, were also immunoprecipitated by the anti-Sm serum (Figure 6, lanes 5 and 6), we conclude that at least one epitope is present on the carboxy-terminal part of the protein.

Homologies are found between the deduced amino acid sequence for B/B' and the described sequences of the U1-snRNP-specific proteins A (Sillekens *et al.*, 1987) and C (Sillekens *et al.*, 1988). These include all proline-rich motifs. The sequence PGMPPMG is shared by B/B' (193–200) and C (120–127). A sequence GPPPPGM is also found in both B' (229–235) and C (107–113), whereas the motif MPPPPGM is common to A (164–169) and B/B' (215–220). These homologies with U1-snRNP-specific A and C proteins could account for the cross-reactive epitopes described for B/B' and C (Yamamoto *et al.*, 1988) and observed for B/B' and A (Figure 2). Although a cross-reaction between B/B' and D has been observed with affinity-purified human antibodies (Habets *et al.*, 1985), as well with monoclonal antibodies (Williams *et al.*, 1986), no homology was found between B/B' and D. Because most of the additional amino acid sequence present in B', is a repeat already present four times in the B sequence, one can easily understand why no immunological differences between B' and B have been found.

The amino acid sequence of the proteins B and B' may present a clue to their function as autoantigens. In murine graft-versus-host disease, an autoimmune disease can be elicited by immunization of F1 mice with parental T cells (Gleichmann *et al.*, 1984). This autoimmune disease is

similar to the human disease systemic lupus erythematosus. In this model, parental T cells can present aspecific help to autoreactive B cells, resulting in the formation of various specificities of antinuclear autoantibodies, including anti-DNA, and a range of antibodies, directed against membrane antigens. Recently, anti-B/B' antibodies have also been found in these animals (Gelpi *et al.*, 1988; A. van Dam *et al.*, in preparation). It is also noticed that only antigens able to cross link B cell membrane receptors, either by being membrane-bound or by having repetitive epitopes, elicited an auto-immune response in this model (Gleichmann *et al.*, 1984). The deduced amino acid sequence of the proteins B and B' shows that these proteins indeed contain highly repetitive amino acid sequences. Therefore, they are likely to give rise to autoimmune responses.

Materials and methods

Monoclonal and human antibodies

The IgG2a monoclonal antibody CLB-ANA-125 was obtained from a fusion of spleen cells from an anti-Sm producing MRL/lpr mouse with SP2/0 cells. Affinity-purified Sm (Maddison *et al.*, 1985) was coated onto an ELISA plate and culture supernatant from wells containing growing cells was tested for reactivity with this antigen. Positively reacting clones were subsequently subcloned by limiting dilution and characterized by immunoblotting on HeLa nuclear blots. CLB-ANA-125 reacted exclusively with the Sm proteins B and B'. The monoclonal antibody ANA-98, reactive against B/B', was a gift of Dr A.G. Rolink, Basel Institute for Immunology. Human sera containing anti-Sm antibodies were obtained from patients with systemic lupus erythematosus.

Isolation of human B/B' Sm protein cDNA clones

A λ gt11 cDNA library made from HL60 poly(A)⁺ RNA was purchased from Clontech Laboratories (Palo Alto, CA) and was screened with the use of the monoclonal antibody CLB-ANA-125. To detect specifically bound antibody, a polyclonal goat-anti-mouse conjugate, coupled to alkaline phosphatase (Promega, Madison, WI) was used. EcoRI cDNA fragments from immunologically detected recombinants were tested for cross-hybridization by Southern blotting (Maniatis *et al.*, 1982). Subsequently, other cDNA clones were isolated from a human placenta library by hybridization.

Preparation of lysogens

For immunoblotting, fusion protein was prepared essentially as described (Huynh *et al.*, 1985). After induction, cells were grown for another hour at 37°C, harvested, resuspended in phosphate-buffered saline (PBS; 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.4) containing 1 mM phenylmethylsulfonylfluoride (PMSF), and sonicated 4 × 45 s.

Preparation of HeLa extracts

HeLa extracts were prepared as described (Westgeest *et al.*, 1986). Radio-labeling of HeLa cells was performed by incubating 2 × 10⁷ cells for 5 h in methionine-free medium, to which 18 MBq [³⁵S]methionine (Amersham, UK) was added. A cell extract was prepared according to Mimori *et al.* (1984).

Electrophoresis and immunoblotting of proteins

SDS-PAGE was performed as described (Westgeest *et al.*, 1986), with some slight modifications. Human sera were pre-absorbed with an Y1090 *E. coli* extract before they were incubated with the blots. As conjugates, polyclonal goat-anti-mouse antibodies, coupled to alkaline phosphatase (Promega) or mouse-anti-human monoclonal antibodies, coupled to horseradish peroxidase (CLB-MH-16-E, this institute) were used.

Elution of antibodies from nitrocellulose was performed according to Smith and Fisher (1984). In order to ascertain the specificity of the reaction of the affinity-purified antibodies, a 1:1 mixture of an anti-Sm serum and a serum containing antibodies to the ribosomal proteins P0, P1 and P2 was used for incubation with the strips containing fusion protein and β -galactosidase. A titration of this mixture of anti-Sm and anti-ribosomal serum on a HeLa cytoplasmic blot showed, that both types of antibodies were present in equal amounts.

RNA transcription, in vitro translation and immunoprecipitation

RNA transcripts were synthesized *in vitro* from *Sst*I-digested pSP65 plasmids containing clone 5 and 16 inserts respectively with SP6 RNA polymerase (Melton *et al.*, 1984).

In vitro translations in a nuclease-treated reticulocyte lysate (Promega) were performed for 60 min at 30°C in 25 µl reaction mixtures. Translation products were immunoprecipitated as described (Fisher *et al.*, 1983).

RNA blot hybridization and RNase mapping

Total RNA was isolated from HeLa and HL60 cells and analyzed by electrophoresis on 1.5% agarose gel in the presence of formaldehyde as described (Maniatis *et al.*, 1982). Poly(A)⁺ RNA was prepared from cytoplasmic HeLa RNA by repeated chromatography on oligo(dT)-cellulose (Maniatis *et al.*, 1982).

RNase mapping was essentially performed as described (Melton *et al.*, 1984). pSP65 plasmids containing the *Bam*HI-*Sma*I fragments of clones 5 and 16 (nucleotides 665–896 for 5 and 665–750 for 16, Figure 4) were used to generate uniformly labeled anti-sense RNA probes. SP6 RNA polymerase (Promega) was used to synthesize the RNA probe according to the manufacturers' specifications except that 80 µCi of [³²P]CTP and 10 units of human placental ribonuclease inhibitor (Promega) were included in the reaction mixture.

DNA sequence analysis

For sequence analysis, suitable cDNA fragments were cloned into M13mp18 or mp19, and subjected to dideoxy chain termination reactions with the standard M13 primers or synthetic B/B' primers, using Sequenase (USB, Cleveland, OH). All clones have also been sequenced using dITP instead of dGTP.

Acknowledgements

We thank Dr A.J.G.Swaak, the Rheumatology Department of the Daniel den Hoed Clinic, Rotterdam, for providing the anti-Sm sera and Eric Eldering and Han Huisman for critically reading the manuscript. This work was supported by the Netherlands League against Rheumatism (grant no. 87/CR/38).

References

Agris, C.H., Nemeroff, M.E. and Krug, R.M. (1989) *Mol. Cell. Biol.*, **9**, 259–267.
 Billings, P.B. and Hoch, S.O. (1984) *J. Biol. Chem.*, **259**, 12850–12856.
 Bringmann, P. and Lührmann, R. (1986) *EMBO J.*, **5**, 3509–3516.
 Earnshaw, W.C., Sullivan, K.F., Machlin, P.S., Cooke, C.A., Kaiser, D.A., Pollard, T.D., Rothfield, N.F. and Cleveland, D.W. (1987) *J. Cell. Biol.*, **104**, 817–829.
 Fisher, D.E., Conner, G.E., Reeves, W.H., Blobel, G. and Kunkel, H.G. (1983) *Proc. Natl. Acad. Sci. USA*, **80**, 6356–6360.
 Fisher, D.E., Conner, G.E., Reeves, W.H., Wisniewski, R. and Blobel, G. (1985) *Cell*, **42**, 751–758.
 Francoeur, A., Peebles, C.L., Heckman, K.J., Lee, J.C. and Tan, E.M. (1985) *J. Immunol.*, **135**, 2378–2384.
 Frendewey, D., Kramer, A. and Keller, W. (1987) *Cold Spring Harbor Symp. Quant. Biol.*, **52**, 287–298.
 Fresco, L.D., Kurilla, M.G. and Keene, J.D. (1987) *Mol. Cell. Biol.*, **7**, 1148–1155.
 Garoff, H., Frischauf, A.M., Simons, K., Lehrach, H. and Delius, H. (1980) *Proc. Natl. Acad. Sci. USA*, **77**, 6376–6380.
 Gelpi, C., Rodriguez-Sanchez, J.L., Martinez, M.A., Craft, J. and Hardin, J.A. (1988) *J. Immunol.*, **140**, 4160–4166.
 Gleichmann, E., Pals, S.T., Rolink, A.G., Radaszkiewicz, T. and Gleichmann, H. (1984) *Immunol. Today*, **5**, 324–332.
 Grabowski, P.J., Seiler, S.R. and Sharp, P.A. (1985) *Cell*, **42**, 345–353.
 Habets, W.J., Berden, J.H.M., Hoch, S.O. and van Venrooy, W.J. (1985) *Eur. J. Immunol.*, **15**, 992–997.
 Habets, W.J., Sillekens, P.T.G., Hoet, M.H., Schalken, J.A., Roebroek, A.J.M., Leunissen, J.A.M., Van de Ven, W.J.M. and Van Venrooy, W.J. (1987) *Proc. Natl. Acad. Sci. USA*, **84**, 2421–2425.
 Hamm, J., Kazmaier, M. and Mattaj, J.W. (1987) *EMBO J.*, **6**, 3479–3485.
 Hamm, J., Van Santen, V.L., Spritz, R.A. and Mattaj, J.W. (1988) *Mol. Cell. Biol.*, **8**, 4787–4791.
 Hinterberger, M., Pettersson, I. and Steitz, J.A. (1983) *J. Biol. Chem.*, **258**, 2604–2613.
 Huynh, T.V., Young, R.A. and Davis, R.W. (1985) In Glover, D.M. (ed.), *DNA Cloning: A Practical Approach*. IRL Press, Oxford, pp. 49–78.

Konarska, M.M. and Sharp, P.A. (1987) *Cell*, **49**, 763–774.
 Kozak, M. (1986) *Cell*, **44**, 283–292.
 Kruijer, W., van Schaik, F.M.A. and Sussenbach, J.S. (1981) *Nucleic Acids Res.*, **9**, 4439–4457.
 Leff, S.E., Rosenfeld, M.G. and Evans, R.M. (1986) *Annu. Rev. Biochem.*, **55**, 1091–1117.
 Maddison, P.J., Skinner, R.P., Vlachoyiannopoulos, P., Brennand, D.M. and Hough, D. (1985) *Clin. Exp. Immunol.*, **62**, 337–345.
 Maniatis, T., Fritsch, E.F. and Sambrook, J. (1982) *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
 McAllister, G., Amara, S.G. and Lerner, M.R. (1988) *Proc. Natl. Acad. Sci. USA*, **85**, 5296–5300.
 McAllister, G., Roby-Shemkowitz, A., Amara, S.G. and Lerner, M.R. (1989) *EMBO J.*, **8**, 1177–1181.
 Melton, D.A., Krieg, P.A., Rebagliati, M.R., Maniatis, T., Zinn, K. and Green, M.R. (1984) *Nucleic Acids Res.*, **12**, 7035–7056.
 Mimori, T., Hinterberger, M., Pettersson, I. and Steitz, J.A. (1984) *J. Biol. Chem.*, **259**, 560–565.
 Ohosone, Y., Mimori, T., Griffith, A., Akizuki, M., Homma, M., Craft, J. and Hardin, J.A. (1989) *Proc. Natl. Acad. Sci. USA*, **86**, 4249–4253.
 Padgett, R.A., Grabowski, P.J., Konarska, M.M., Seiler, S. and Sharp, P.A. (1986) *Annu. Rev. Biochem.*, **55**, 1119–1150.
 Pettersson, I., Hinterberger, M., Mimori, T., Gottlieb, E. and Steitz, J.A. (1984) *J. Biol. Chem.*, **259**, 5907–5914.
 Reddy, R. and Busch, H. (1988) In Birnstiel, M.L. (ed.), *Structure and Function of Major and Minor Small Nuclear Ribonucleoprotein Particles*. Springer Verlag, Heidelberg, pp. 1–37.
 Reuter, R., Rothe, S. and Lührmann, R. (1987) *Nucleic Acids Res.*, **15**, 4021–4034.
 Rokeach, L.A., Haselby, J.A. and Hoch, S.O. (1988) *Proc. Natl. Acad. Sci. USA*, **85**, 4832–4836.
 Rokeach, L.A., Jannatipour, M., Haselby, J.A. and Hoch, S.O. (1989) *J. Biol. Chem.*, **264**, 5024–5030.
 Schmauss, C., McAllister, G., Ohosone, Y., Hardin, J.A. and Lerner, M.R. (1989) *Nucleic Acids Res.*, **17**, 1733–1743.
 Sillekens, P.T.G., Habets, W.J., Beijer, R.P. and Van Venrooy, W.J. (1987) *EMBO J.*, **6**, 3841–3848.
 Sillekens, P.T.G., Beijer, R.P., Habets, W.J. and Van Venrooy, W.J. (1988) *Nucleic Acids Res.*, **16**, 8307–8322.
 Sillekens, P.T.G., Beijer, R.P., Habets, W.J. and Van Venrooy, W.J. (1989) *Nucleic Acids Res.*, **17**, 1893–1906.
 Smith, D.E. and Fisher, P.A. (1984) *J. Cell Biol.*, **99**, 20–28.
 Spritz, R.A., Strunk, K., Surowy, C.S., Hoch, S.O., Barton, D.E. and Francke, U. (1987) *Nucleic Acids Res.*, **15**, 10373–10391.
 Stanford, D.R., Rohleder, A., Neiswanger, K. and Wieben, E.D. (1987) *J. Biol. Chem.*, **262**, 9931–9934.
 Steitz, J.A., Black, D.L., Gerke, V., Parker, K.A., Kramer, A., Frendewey, D. and Keller, W. (1988) In Birnstiel, M.L. (ed.), *Structure and Function of the Major and Minor Small Ribonucleoprotein Particles*. Springer Verlag, Heidelberg, pp. 115–154.
 Swanson, M.S., Nakagawa, T.Y., LeVan, K. and Dreyfuss, G. (1987) *Mol. Cell. Biol.*, **7**, 1731–1739.
 Tan, E.M. (1982) *Adv. Immunol.*, **33**, 167–240.
 Theissen, H., Etzerodt, M., Reuter, R., Schneider, C., Lottspeich, F., Argos, P., Luehrmann, R. and Philipson, L. (1986) *EMBO J.*, **5**, 3209–3217.
 Westgeest, A.A.A., Bons, J.C., Van den Brink, H.G., Aarden, L.A. and Smeenk, R.J.T. (1986) *J. Immunol. Methods*, **95**, 283–288.
 Wieben, E.D., Madore, S.J. and Pederson, T. (1983) *Proc. Natl. Acad. Sci. USA*, **80**, 1217–1220.
 Williams, D.G., Stocks, M.R., Smith, P.R. and Maini, R.N. (1986) *Immunology*, **58**, 496–500.
 Yamamoto, K., Miura, H., Moroi, Y., Yoshinoya, S., Goto, M., Nishioka, K. and Miyamoto, T. (1988) *J. Immunol.*, **140**, 311–317.
 Zillmann, M., Zapp, M.L. and Berget, S.M. (1988) *Mol. Cell. Biol.*, **8**, 814–821.

Received on May 5, 1989; revised on August 1, 1989