Published in final edited form as:

J Agric Food Chem. 2012 August 15; 60(32): 7949–7956. doi:10.1021/jf204755g.

Effects of Estrogen and Estrus Cycle on Pharmacokinetics, Absorption and Disposition of Genistein in Female Sprague-Dawley Rats

Kaustubh H. Kulkarni, Zhen Yang, Niu Tao, and Ming Hu*

Department of Pharmacological and Pharmaceutical Sciences, College of Pharmacy, University of Houston, 1441 Moursund Street, Houston, TX, 77030, USA

Abstract

Genistein is an active soy isoflavone with anticancer activities but it is unknown why it has a higher oral bioavailability in female than in male rats. Our study determined the effects of estrus cycle on genistein's oral bioavailability. Female rats with various levels of estrogen were orally administered with genistein or used in a four-site rat intestinal perfusion experiment. Rats in "proestrus" group (with elevated estrogen) had significantly reduced (57% decrease, $p < 0.05$) oral bioavailability of total genistein (aglycone+conjugates) than those in "metoestrus" group (with basal level of estrogen). Female ovariectomized rats, due to lack of estrogen, showed oral bioavailability of total genistein similar to the "metoestrus" group but higher (155% increase, $p < 0.05$) than the "proestrus" group. Based on intestinal perfusion studies, the increased bioavailability was partially attributed to the higher ($> 100\%$ increase, $p < 0.05$) hepatic disposition via glucuronidation and possibly more efficient enterohepatic recycling of genistein in the "metoestrus" group. Furthermore, chronic exogenous supplementation of estradiol in ovariectomized rats significantly reduced (77%, $p < 0.05$) the oral bioavailability of total genistein, mostly via increased sulfation (> 10 folds) in liver, to a level comparable to those in the "proestrus" group. In conclusion, the oral bioavailability of total genistein was inversely proportional to elevated estrogen levels in female rats, which is partially mediated through the regulation of hepatic enzymes responsible disposition of genistein.

Keywords

Genistein; Oral bioavailability; pharmacokinetics; estrus cycle; estrogen

INTRODUCTION

Genistein, a soy isoflavone present in nature as genistin (genistein-7-O-glucoside) but absorbed as aglycone¹, has been labeled as a "miracle" compound with lots of claimed beneficial effects such as its anti-oxidant and anticancer activities^{2, 3}, many of which have been demonstrated in small-scale clinical trials²⁻⁴. Genistein has also displayed potential favorable impact on prevention of coronary heart disease and post-menopausal symptoms⁵.

* Author to whom correspondence should be addressed [Tel: (713) 795-8320; Fax: (713) 795-8305. mhu@uh.edu].

To this date, the major difficulty in demonstrating the efficacy of genistein in humans is its poor, highly variable and gender-dependent oral bioavailability^{6 7}.

Gender-dependent oral bioavailability was demonstrated in an earlier study, where a significantly higher (2 fold) oral bioavailability of total ¹⁴C-genistein (aglycone+metabolite) was observed in female (15%) than in male (7%) Sprague-Dawley or SD rats⁶. We observed similar results for oral PK study in male and female SD rats, in that significantly higher oral bioavailability of genistein (<2 fold) and genistein glucuronide (>4 fold) and thereby total genistein was observed in female than in male SD rats (see Supplemental Data). Until now, no study has been done to determine if female sex hormone(s) will impact oral bioavailability of pharmaceutical or dietary polyphenol in rodents, primates or humans. Furthermore, no study has been done to determine the role of estrus cycle in regulation of oral bioavailability of pharmacologically active polyphenols in female rats, which is important since expression levels of certain hepatic phase II metabolizing enzymes (e.g., UDP-glucuronosyltransferases, or UGTs) are female sex-hormone regulated⁸.

Female sex hormone appears to impact the expression levels of certain phase II enzymes in that the pregnant female rats showed significantly lower hepatic Ugt1A, Ugt1A1, Ugt1A5, Ugt1A6 and Ugt1A8 enzyme expression levels than in control female SD rats⁸. In the same study, significantly higher levels of these hepatic UGT isoforms were observed in female rats 10 to 12 days post-partum compared to those in the control female rats⁸. However, there is a lack of studies done to determine the regulation of uptake/efflux transporter proteins by female sex hormones in rodents.

Female ovariectomized rodents with a diminished level of estrogen (a major female sex hormone) have been the model of choice for testing the effectiveness of antiestrogens such as tamoxifen and phytoestrogens such as genistein against breast cancer and bone loss for more than two decades^{9, 10}. Raloxifene, tamoxifen and genistein are three extensively studied compounds for breast cancer development, prevention and treatment in female ovariectomized rats^{11–13}. No studies have been done to clearly understand the effect of diminishing and resurrected levels of estrogen on oral bioavailability of these phenolic compounds in female ovariectomized rats, which could significantly impact their efficacy.

Therefore, the purpose of this study was to determine the role of sex hormones, estrogen, on oral bioavailability of genistein by using female ovariectomized rats, and control female rats in different phases of the estrus cycle. We divided female rats into two groups (the “proestrus” and the “metoestrus”) based on the levels of female sex hormones¹⁴. At proestrus and oestrus phases (called the “proestrus” group) of the estrus cycle, female rats showed elevated levels of estrogen, whereas at metoestrus and diestrus phases (called the “metoestrus” group) of the estrus cycle, female rats showed basal levels of estrogen. Oral bioavailability of genistein was compared between female ovariectomized and control female rats, and between female rats in the “proestrus” and the “metoestrus” groups. We also determined the effect of a chronic exogenous dose of estradiol on oral bioavailability of genistein in female ovariectomized rats. Additional absorption and intestinal disposition studies were conducted using a four-site rat intestinal perfusion model with bile duct

cannulation to determine the contribution of gut and liver metabolism to the observed differences in oral bioavailability.

MATERIALS AND METHODS

Chemicals

Genistein and daidzein (internal standard) were purchased from Indofine Chemicals (Somerville, NJ). Estradiol, mineral oil and Hank's balanced salt solution (HBSS, powder form) was purchased from Sigma-Aldrich (St. Louis, MO). All other materials (analytical grade or better) used were analytical grade or better.

Animals

Female Sprague Dawley rats (6 to 8 weeks old) were purchased from Harlan Laboratories (Madison, WI). The rats were fed with an AIN 76A non-soy diet (W) purchased from Harlan Laboratories (Madison, WI). The rats were fasted for up to 15 hours before the day of the pharmacokinetic (or PK) or perfusion experiment.

Estrus cycle status

Estrus cycle status of a female SD rat was monitored daily. Vaginal smears for all the female rats were taken twice (morning and evening) daily for four consecutive days at the same time and only the female rats showing all four phases of estrus cycle were chosen for the study. The vaginal smear was taken using the same method described previously^{15, 16} with minor modifications. Briefly, the vaginal smear was taken using plastic pipette on a clean glass slide; air dried and stained using methylene blue dye to identify the presence of different cells confirming the phases of estrus cycle in female rats (See Supplemental Data). For perfusion/PK experiments, the female rats were grouped as the "proestrus" group for those in "proestrus + oestrus" phases (at elevated plasma estrogen levels) or as the "metoestrus" group for those in "metoestrus" + diestrus" phases (at basal plasma estrogen levels).

Exogenous estradiol administration

Estradiol 100 µg/ml was prepared in mineral oil and 10 µg/kg subcutaneous dose was administered to female ovariectomized rats every day at 9:00am for a period of ten days. On the eleventh day, a single dose of genistein (20 mg/kg) was administered orally. Another group of rats were dosed with vehicle (i.e., mineral oil) for seven days and on the eighth day, they were dosed with a single dose of genistein (20 mg/kg).

Pharmacokinetic (PK) experiment

The procedures for this and other animal experimentation were approved by the University of Houston's Institutional Animal Care and Use Committee. Genistein suspension (20 mg/ml) freshly prepared in oral suspending vehicle from PCCA Laboratories (Houston, TX) was administered by oral gavage at a dose of 20 mg/kg¹⁷. Blood samples were taken from the tail vein at 15, 30, 45, 60, 120, 180, 240, 360, 480, 720 and 1440 minutes after oral administration.

Sample extraction

Blood samples (20 μ l) from the pharmacokinetic studies were precipitated with 100% acetonitrile (80 μ l) containing 3 μ M of daidzein to collect >90% of genistein and its phase II metabolites. The mixture was centrifuged at 15,500 rpm for 15 minutes and 80 μ l of supernatant was collected and dried under air. The residue was reconstituted in 100 μ l of 30% acetonitrile solution, centrifuged at 15,500 rpm for 15 minutes and injected in UPLC-MS/MS.

Animal surgery

The intestinal surgical procedures were modified from our previous publications^{18, 19}, in that four segments of the intestine were cannulated simultaneously along with a bile duct cannulation^{1, 20, 21}. The blood circulation to the liver and intestine was not disrupted in this model. To maintain the temperature of the perfusate constant at 37°C, the inlet cannulae were insulated and kept warm by a 37°C circulating water bath.

Transport and metabolism experiments in perfused rat intestinal model

A single-pass perfusion method described previously was used^{21, 22}. At least four rats were used for each set of the perfusion study. Briefly, four segments (duodenum, upper jejunum, terminal ileum, and colon) of the rat intestine approximately 10 to 15 cm each were perfused simultaneously with a Hanks' balanced salt solution (HBSS, pH 7.4) containing 10 μ M genistein as the perfusate. The flow of the perfusate was driven by an infusion pump (Harvard Apparatus, Cambridge, MA) at a flow rate of 0.191 ml/min. After a 30-min washout period, which is usually sufficient to achieve steady-state absorption, intestinal perfusate samples were collected from the outlet cannulae every 30 min. Bile samples (approximately 0.5 ml) were collected before perfusion started and every 30 min afterward. After perfusion, the length of the intestine was measured as described previously^{18, 19}. The perfusate sample concentrations of genistein and its phase II metabolites were determined using UPLC-MS/MS. Bile samples were diluted (1:40) with HBSS buffer, centrifuged, and supernatant removed for UPLC-MS/MS analysis.

UPLC-MS/MS analysis of genistein and its conjugates

We implemented a published method for analysis of genistein and its conjugates in rat plasma, perfusion and microsome samples using UPLC-MS/MS¹⁷. An API 3200 QTRAP® triple quadrupole mass spectrometer (Applied Biosystems/MDS SCIEX, Foster City, CA, USA) equipped with a TurboIonSpray™ source was operated in negative ion mode to perform the analysis. The flow dependent parameters for introduction of the samples to the mass spectrometers ionization source were set as follows: ionspray voltage, -4.0 kV; ion source temperature, 700°C; the nebulizer gas (gas1), nitrogen, 40 psi; turbo gas (gas2), nitrogen, 40 psi; curtain gas, nitrogen, 10 psi. The quantification was performed using MRM method with the transitions of m/z 269 → m/z 133 for genistein, m/z 445 → m/z 269 for genistein glucuronides, m/z 349 → m/z 269 for genistein sulfate and m/z 253 → m/z 132 for daidzein (IS)– Unit mass resolution was set in both mass-resolving quadrupole Q1 and Q3. The standard curves for genistein, genistein glucuronide and genistein sulfate were prepared in blank rat blood. The standard curves were linear in the concentration range of

0.019 μM to 10 μM for genistein; 0.02 μM to 10 μM for genistein glucuronide and 0.002 μM to 1.6 μM for genistein sulfate.

Data analysis in pharmacokinetic studies

The plasma concentration-time profiles for each rat were analyzed by using WinNonlin3.3. C_{max} and the area under the concentration-time curve [$\text{AUC}_{(0-24\text{hr})}$] was calculated according to a non-compartmental model using the WinnonlinTM software.

Data analysis in rat intestinal perfusion experiments

Amounts of genistein absorbed (M_{ab}), amounts of conjugated genistein excreted into the intestinal lumen (M_{gut}), amounts of conjugated genistein excreted via the bile (M_{bile}) values were calculated as described previously^{20, 21, 23}. Briefly, M_{ab} was expressed as:

$$M_{\text{ab}} = Q \cdot \tau \cdot (CA_{\text{in}} - CA_{\text{out}}) \quad \text{Equation (1)}$$

where Q is the flow rate (ml/min), τ is the sampling interval (30min), and CA_{in} and CA_{out} are the inlet and outlet concentrations of genistein aglycone corrected for water flux, respectively. M_{gut} was expressed as

$$M_{\text{gut}} = Q \cdot \tau \cdot CM_{\text{out}} \quad \text{Equation (2)}$$

where CM_{out} was the outlet concentration (nmol/ml) of metabolites corrected for water flux. And M_{bile} was expressed as

$$M_{\text{bile}} = V \cdot CM_{\text{bile}} \quad \text{Equation (3)}$$

where CM_{bile} was the bile concentration (nmol/ml) of metabolites and V was the volume of bile collected over a 30 min time period.

Statistical analysis

The data in this paper were presented as mean $\pm$ SD, if not specified otherwise. Significant differences were assessed by using unpaired Student's t-test with "p" value of <0.05.

RESULTS

Estrus cycle-dependent oral bioavailability of genistein in female SD rats

Female rats were divided into two separate groups; one for elevated (the "proestrus") and the other for basal (the "metoestrus") levels of estrogen using vaginal smears as described previously¹⁴. A single oral dose of genistein (20 mg/kg) was given to female SD rats from both groups to identify the effects of estrus cycle on plasma profiles of genistein and its metabolites (glucuronide and sulfate) using UPLC-MS/MS analysis.

Significant estrus cycle-dependent differences were observed in total genistein (aglycone +metabolite) plasma pharmacokinetic profiles (Figure 1; Table 1). Genistein plasma concentration profiles showed significantly higher C_{max} (<2 fold, $p < 0.05$) and $\text{AUC}_{(0-24\text{hr})}$ (>2 fold, $p < 0.05$) values in the "metoestrus" group than in the "proestrus" group (Figure 1A;

Table 1). Furthermore, genistein glucuronide plasma concentration profiles also showed higher $AUC_{(0-24hr)}$ (2.4 fold, $p<0.01$) values in the “metoestrus” group than in the “proestrus” group (Figure 1B; Table 1). Genistein sulfate plasma concentrations, however, were too low to determine the effects of estrus cycle on its level in these rats.

Overall, the 2-fold higher oral bioavailability of total genistein in the “metoestrus” group as compared to the “proestrus” group was contributed predominantly by 2.4 fold ($p<0.01$) higher plasma concentrations of genistein glucuronide in “metoestrus” group than in “proestrus” group of the rats (Table 1).

Ovarian-dependent oral bioavailability of genistein in female SD rats

Single dose oral pharmacokinetics of genistein (20 mg/kg) was also determined in ovariectomized rats and plasma concentrations of genistein and its metabolites (glucuronide and sulfate) were also analyzed. Significant ovarian (sex hormone)-dependent difference was observed in total genistein (aglycone+metabolite) plasma pharmacokinetic profiles (Table 1; Figure 1, 2, and S3 in Supplemental Data). Genistein plasma concentrations showed significantly higher C_{max} (>2.5 fold, $p<0.05$) and $AUC_{(0-24hr)}$ (3 fold, $p<0.05$) values in ovariectomized rats than the “proestrus” group respectively (Table 1; Figure 1, 2, and S3). Surprisingly, genistein sulfate plasma concentrations also showed significantly higher C_{max} and $AUC_{(0-24hr)}$ (>5 fold each, $p<0.01$) values in ovariectomized rats than not just the “proestrus” but also the “metoestrus” group (Table 1; Figure 1, 2, and S3).

Overall the 2.5-fold higher oral bioavailability of total genistein in ovariectomized rats as compared to the “proestrus” group was contributed predominantly by 3 fold ($p<0.01$) higher plasma concentrations of genistein in ovariectomized rats than the “proestrus” group (Table 1). No significant difference in oral bioavailability of total genistein in ovariectomized rats was observed when compared to the female rats in the “metoestrus” group (Table 1).

Exogenous estradiol-dependent oral bioavailability of genistein in female SD

The pharmacokinetics of genistein after a single oral dose (20 mg/kg) was further determined in ovariectomized rats after chronic (10 days) subcutaneous estradiol (10 μ g/kg/day) administration, and plasma concentrations of genistein and its metabolites (glucuronide and sulfate) were again analyzed using UPLC-MS/MS analysis. Our hypothesis was that the exogenous estrogen supplement might reverse the higher oral bioavailability plasma profiles of genistein and its metabolites (glucuronide and sulfate) in female ovariectomized rats. The results indicated that a significant exogenous estradiol-dependent difference was observed in total genistein (aglycone + metabolites) plasma pharmacokinetic profiles in ovariectomized rats after chronic estradiol administration. Genistein, genistein glucuronide and genistein sulfate plasma concentrations showed significantly lower C_{max} (4, 2 and 5 fold, $p<0.05$) and $AUC_{(0-24hr)}$ (6.5, 2 and 13 fold, $p<0.05$) values after exogenous estradiol supplement in ovariectomized rats (Figure 2A/B/C; Table 1).

Overall, the 3-fold lower oral bioavailability of total genistein after chronic (10day) subcutaneous estradiol (10 μ g/kg/day) administration in ovariectomized rats as compared to the control ovariectomized rats was contributed predominantly by 6.5 fold ($p<0.01$) and 2

fold ($p<0.05$) lower plasma concentrations of genistein and genistein glucuronide respectively (Table 1).

Comparison of oral bioavailability of total genistein between chronic (10 days) estradiol (10 µg/kg) administered ovariectomized and control rats in the “proestrus” group

Oral bioavailability of total genistein (aglycone + metabolite) plasma pharmacokinetic profile comparison showed no significant difference (except one) between exogenous estradiol supplemented ovariectomized rats and the “proestrus” group. Genistein plasma concentrations showed significantly lower $AUC_{(0-24hr)}$ (<2 fold, $p<0.05$) with no difference in C_{max} values in estradiol supplemented ovariectomized rats than the “proestrus” group (Table 1; Figure 1, 2, and S3). Genistein glucuronide and total genistein (aglycone + metabolite) plasma concentrations showed no significant difference in C_{max} and $AUC_{(0-24hr)}$ values between estradiol supplemented ovariectomized rats and the “proestrus” group (Table 1; Figure 1, 2, and S3). Furthermore, genistein sulfate plasma concentrations, in both groups of rats were too low to determine the effects of exogenous estradiol administration on its level (Table 1; Figure 1, 2, and S3).

Overall, no significant difference in oral bioavailability of total genistein in ovariectomized rats after chronic estradiol administration and control rats in “proestrus” group was observed (Table 1).

Comparison of oral bioavailability of total genistein between ovariectomized rats after chronic (10 days) estrogen (10 µg/kg) dose administration and control rats in the “metoestrus” group

Oral bioavailability of total genistein (aglycone+metabolite) plasma pharmacokinetic profile comparison showed significant difference between exogenous estradiol supplemented ovariectomized rats and the “metoestrus” group. Genistein and genistein glucuronide plasma concentrations showed significantly lower C_{max} (>2 fold each, $p<0.05$) and $AUC_{(0-24hr)}$ (4 and >2.5 fold, $p<0.01$) values in estradiol supplemented ovariectomized rats than the “metoestrus” group respectively (Table 1; Figure 1, 2, and S3). Genistein sulfate plasma concentrations, however, in both groups of rats were too low to determine the effects of exogenous estradiol administration on its level (Table 1; Figure 1, 2, and S3).

Overall, the 3-fold lower oral bioavailability of total genistein after chronic estradiol (10 µg/kg) administration in ovariectomized rats as compared to the control rats in the “metoestrus” group was contributed predominantly by 2.5 fold ($p<0.01$) lower plasma concentrations of genistein glucuronide (Table 1).

Estrus cycle-dependent genistein absorption and disposition in female SD rats

Female rats were divided into two separate groups; elevated (the “proestrus”) and basal (the “metoestrus”) levels of female sex hormone, estrogen, using vaginal smears as described previously¹⁴. Four-site rat intestinal perfusion with bile duct cannulation was used to identify the estrus cycle-dependent differences in intestinal perfusate and biliary excretion profiles of genistein and its metabolites (glucuronide and sulfate). Significant estrus cycle-dependent difference was observed in hepatic disposition but not in intestinal absorption and

disposition of genistein (Figure 3). Cumulative biliary excretion profiles showed significantly higher (<2 fold each, $p<0.05$) amount of genistein (1–12 nmol) and genistein glucuronide (500–1500 nmol) in the “metoestrus” than the “proestrus” group (Figure 3D/E). Intestinal absorption and metabolite (glucuronide and sulfate) excretion of genistein showed no difference between the “metoestrus” and the “proestrus” groups (Figure 3A/B/C). Cumulative biliary excretion of genistein sulfate in both groups was too low to determine the effects of changes in the estrus cycle (Figure 3F).

Overall, significantly higher biliary excretion of genistein and conjugates in the “metoestrus” as compared to the “proestrus” group was mainly due to the higher (~2 fold, $p<0.05$) cumulative biliary efflux of genistein glucuronide (micromolar range) in the “metoestrus” than in the “proestrus” group of rats.

Ovarian-dependent genistein absorption and disposition in female SD rat

Four-site rat intestinal perfusion with bile duct cannulation was used to determine the impact of ovary on intestinal and biliary excretion of genistein conjugates. Significant ovarian-dependent difference in biliary excretion was observed in ovariectomized rats when compared to rats in either the “proestrus” or the “metoestrus” or the control (containing both proestrus and metoestrus) group, but intestinal absorption and disposition of genistein was not observed, with one minor exception (Figure 3): duodenal genistein glucuronide excretion was significantly higher (<2 fold, $p<0.05$) in female ovariectomized rats as compared to the “proestrus” group (Figure 3B). In case of biliary excretion of genistein conjugates, no detectable amount of genistein aglycone and significantly lower genistein glucuronide (150–1250 nmol range) (4 and 8 fold respectively, $p<0.01$) was observed in cumulative biliary excretion of genistein conjugates in ovariectomized rats (Figure 3E). In contrast, significantly higher amount of genistein sulfate (10–30 nmol range) was observed in biliary excretion in ovariectomized rats as compared to undetectable levels in the other groups (Figure 3F).

Overall, significantly lower biliary excretion of genistein and conjugates in the ovariectomized rats as compared to control group of rats (containing rats from both the “proestrus” and the “metoestrus” groups) was mainly due to 4 and 8 fold lower ($p<0.01$) cumulative biliary efflux of genistein glucuronide in the ovariectomized rats.

DISCUSSION

Our results showed for the first time that the estrus cycle differences and associated changes in sex hormones, especially estrogen, significantly affected oral bioavailability of total genistein in female rats. The observed higher oral bioavailability (AUC value >2 fold, $p<0.05$) of total genistein in the “metoestrus (low estrogen)” group as opposed to the “proestrus (high estrogen)” group (Table 1) supports the conclusion that higher level of female sex hormone estrogen down regulates phase II enzyme activities. The conclusion is further corroborated by the fact that ovariectomized rats (without ovarian sex hormone estrogen) also showed equally higher oral bioavailability (AUC value >2 fold, $p<0.05$) of total genistein than that in the “proestrus” group (Table 1). Interestingly, chronic (10 days) estradiol supplementation in ovariectomized rats significantly reduced the oral

bioavailability (AUC value >2 fold, $p < 0.05$) of total genistein in ovariectomized rats (Table 1) to a level that is comparable to that in “proestrus” group (Table 1), suggesting that estradiol alone may contribute to the bulk of the female hormonal regulation of phase II enzymes.

Until now, to our surprise, we did not see any publications studying the differences in oral bioavailability of pharmaceutical or dietary compounds in female rats in relation to differences in female sex hormone (especially estrogen) levels. Control female and female ovariectomized rats have been used as a model of choice for pharmacological testing of these compounds in a variety of pharmacological models including breast cancer^{12, 24, 25}. The major assumption for using ovariectomized rats as an ideal model for testing the effectiveness of these agents against breast cancer was that sex hormones did not alter their oral bioavailabilities. Considering that breast cancer is not just limited to postmenopausal women (completely diminished plasma levels of major sex hormone, estrogen), one should consider how to extrapolate the results of various translational research that test the pharmacological activities of these compounds’ efficacy using ovariectomized rats. For antiestrogenic agents such as raloxifene and genistein, which are extensively studied in female ovariectomized rats^{12, 26}, it is now imperative to determine the effect of female sex hormones on their oral bioavailability and efficacy.

While studying the effect of fluctuating levels of female sex hormone, estrogen, during the estrus cycle phases on single oral dose pharmacokinetics of genistein (20mg/kg), we observed that plasma estrogen levels played an important role in determining oral bioavailability of total genistein (>2 fold, $p < 0.05$) in female rats. Significantly higher oral bioavailability of total genistein in the “metoestrus” (basal levels of estrogen) than the “proestrus” (elevated levels of estrogen) group (Figure 1; Table 1) suggested that oral bioavailability of total genistein was inversely proportional to the elevated plasma sex hormone levels in female rats.

To further confirm our theory, we used female ovariectomized rats to identify the effect of hormone ablation on oral bioavailability of total genistein in female rats. As expected, we observed significantly higher oral bioavailability of total genistein (2.5 fold, $p < 0.05$) with predominant contribution from genistein (3 fold, $p < 0.01$) and genistein glucuronide (<2 fold, $p < 0.05$) in ovariectomized than the “proestrus” (elevated estrogen) group. Furthermore, there was no significant difference in oral bioavailability of total genistein in ovariectomized rats and the “metoestrus” (basal estrogen) group. This further confirmed our theory that oral bioavailability of total genistein is inversely proportional to elevated estrogen levels in female rats.

In the next set of studies, after chronic (10 days) exogenous estradiol (10 µg/kg/day) supplementation to ovariectomized rats, oral bioavailability of total genistein significantly dropped (Figure 2; Table 1), making it comparable to the “proestrus” group. This observed significant drop in oral bioavailability further bolstered our theory that oral bioavailability of total genistein is inversely proportional to elevated plasma estrogen levels in female rats.

Because oral bioavailability is affected by both absorption and first-pass metabolism, we determined if intestinal absorption and metabolism of genistein contributed to the differences in oral bioavailability observed above. To study the differences in absorption and disposition of genistein, we used a four-site rat intestinal perfusion model with bile duct cannulation. The results suggested that female sex hormone, estrogen, predominantly regulated the hepatic disposition (phase II metabolizing enzymes and/or efflux transporters) of genistein and its conjugates in female rats since the bile excretion patterns were drastically altered (Figure 3). The basal levels of estrogen in the “metoestrus” group of rats were associated with a significantly higher biliary efflux of genistein glucuronide (Figure 3), which in turn meant more efficient enterohepatic recycling than in the “proestrus” group (elevated levels of estrogen). Efficient enterohepatic recycling of genistein metabolites contributed to a longer apparent half-life which in turn meant higher oral bioavailability of total genistein in the “metoestrus” than in the “proestrus” group of rats, as observed in oral pharmacokinetic studies.

In case of ovariectomized rats, the rat intestinal perfusion experiments showed significantly lower biliary efflux of genistein glucuronide than in the “proestrus” as well as the “metoestrus” group of rats (Figure 3). The lowered biliary excretion was inconsistent with the oral pharmacokinetics data observed in ovariectomized rats. Currently, we do not have any explanation for these observed inconsistencies. A possible explanation is that intestinal phase II metabolism was significantly increased but the majority of metabolites were transported to the plasma via the basolateral efflux transporters that were up-regulated in absence of the ovary hormones. More work is still required to explain this phenomenon.

In conclusion, we have clearly demonstrated for the first time that oral bioavailability of total genistein was inversely proportional to elevated plasma estrogen levels in female rats, which is partially mediated via changes in hepatic phase II metabolism of genistein. Therefore, plasma estrogen levels played an important role in oral bioavailability of total genistein in female rats through regulation of hepatic disposition of genistein and efficient enterohepatic recycling of genistein conjugates.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ABBREVIATIONS USED

AUC	Area under the curve
Pro	“Proestrus” phase (or elevated plasma estrogen levels) group of estrus cycle in female rats
Meto	“Metoestrus” phase (or basal plasma estrogen levels) group of estrus cycle in female rats
Ova	“Ovariectomized” (or undetectable levels of plasma estrogen) group of female rats with no significant estrogen

Ova+E	“Ovariectomized+Estrogen” (or undetectable levels of plasma estrogen) group of female rats with exogenous dose of estradiol 10 µg/kg/day
UPLC-MS/MS	Ultra performance liquid chromatography coupled with tandem mass spectrometry
UGT	Uridine diphospho-glucuronosyl transferase enzymes
HBSS	Hank’s balanced salt solution
MSDR	Male sprague dawley rats
FSDR	Female sprague dawley rats

References

1. Liu Y, Hu M. Absorption and metabolism of flavonoids in the caco-2 cell culture model and a perused rat intestinal model. *Drug Metab Dispos.* 2002; 30(4):370–377. [PubMed: 11901089]
2. Perabo FG, Von Low EC, Ellinger J, von Rucker A, Muller SC, Bastian PJ. Soy isoflavone genistein in prevention and treatment of prostate cancer. *Prostate Cancer Prostatic Dis.* 2008; 11(1):6–12. [PubMed: 17923857]
3. Sarkar FH, Li Y. The role of isoflavones in cancer chemoprevention. *Front Biosci.* 2004; 9:2714–2724. [PubMed: 15353308]
4. Lazarevic B, Hammarstrom C, Yang J, Ramberg H, Diep LM, Karlsen SJ, Kucuk O, Saatcioglu F, Tasken KA, Svindland A. The effects of short-term genistein intervention on prostate biomarker expression in patients with localised prostate cancer before radical prostatectomy. *Br J Nutr.* 2012;1–10.
5. Pan W, Ikeda K, Takebe M, Yamori Y. Genistein, daidzein and glycitein inhibit growth and DNA synthesis of aortic smooth muscle cells from stroke-prone spontaneously hypertensive rats. *J Nutr.* 2001; 131(4):1154–1158. [PubMed: 11285318]
6. Coldham NG, Zhang AQ, Key P, Sauer MJ. Absolute bioavailability of [14C] genistein in the rat; plasma pharmacokinetics of parent compound, genistein glucuronide and total radioactivity. *Eur J Drug Metab Pharmacokinet.* 2002; 27(4):249–258. [PubMed: 12587954]
7. Sarkar FH, Li Y, Wang Z, Padhye S. Lesson learned from nature for the development of novel anti-cancer agents: implication of isoflavone, curcumin, and their synthetic analogs. *Curr Pharm Des.* 2010; 16(16):1801–1812. [PubMed: 20345353]
8. Luquita MG, Catania VA, Pozzi EJ, Veggi LM, Hoffman T, Pellegrino JM, Ikushiro S, Emi Y, Iyanagi T, Vore M, Mottino AD. Molecular basis of perinatal changes in UDP-glucuronosyltransferase activity in maternal rat liver. *J Pharmacol Exp Ther.* 2001; 298(1):49–56. [PubMed: 11408524]
9. Cohen LA, Chan PC, Wynder EL. The role of a high-fat diet in enhancing the development of mammary tumors in ovariectomized rats. *Cancer.* 1981; 47(1):66–71. [PubMed: 6780183]
10. Jordan VC. Effects of tamoxifen in relation to breast cancer. *Br Med J.* 1977; 1(6075):1534–1535. [PubMed: 871651]
11. Fudge MA, Kavaliers M, Baird JP, Ossenkopp KP. Tamoxifen produces conditioned taste avoidance in male rats: an analysis of microstructural licking patterns and taste reactivity. *Horm Behav.* 2009; 56(3):322–331. [PubMed: 19576896]
12. Kang X, Jin S, Zhang Q. Antitumor and antiangiogenic activity of soy phytoestrogen on 7,12-dimethylbenz[alpha]anthracene-induced mammary tumors following ovariectomy in Sprague-Dawley rats. *J Food Sci.* 2009; 74(7):H237–H242. [PubMed: 19895476]
13. Fudge MA, Kavaliers M, Baird JP, Ossenkopp KP. Tamoxifen and raloxifene produce conditioned taste avoidance in female rats: a microstructural analysis of licking patterns. *Life Sci.* 2009; 84(9–10):282–289. [PubMed: 19146862]

14. Ogle TF, Kitay JJ. Ovarian and adrenal steroids during pregnancy and the oestrous cycle in the rat. *J Endocrinol.* 1977; 74(1):89–98. [PubMed: 559719]
15. Marcondes FK, Bianchi FJ, Tanno AP. Determination of the estrous cycle phases of rats: some helpful considerations. *Braz J Biol.* 2002; 62(4A):609–614. [PubMed: 12659010]
16. Marcondes FK, Miguel KJ, Melo LL, Spadari-Bratfisch RC. Estrous cycle influences the response of female rats in the elevated plus-maze test. *Physiol Behav.* 2001; 74(4–5):435–440. [PubMed: 11790402]
17. Yang Z, Zhu W, Gao S, Xu H, Wu B, Kulkarni K, Singh R, Tang L, Hu M. Simultaneous determination of genistein and its four phase II metabolites in blood by a sensitive and robust UPLC-MS/MS method: Application to an oral bioavailability study of genistein in mice. *J Pharm Biomed Anal.* 2010; 53(1):81–89. [PubMed: 20378296]
18. Hu M, Roland K, Ge L, Chen J, Li Y, Tyle P, Roy S. Determination of absorption characteristics of AG337, a novel thymidylate synthase inhibitor, using a perfused rat intestinal model. *J Pharm Sci.* 1998; 87(7):886–890. [PubMed: 9649359]
19. Hu M, Sinko PJ, deMeere AL, Johnson DA, Amidon GL. Membrane permeability parameters for some amino acids and beta-lactam antibiotics: application of the boundary layer approach. *J Theor Biol.* 1988; 131(1):107–114. [PubMed: 3419188]
20. Chen J, Lin H, Hu M. Metabolism of flavonoids via enteric recycling: role of intestinal disposition. *J Pharmacol Exp Ther.* 2003; 304(3):1228–1235. [PubMed: 12604700]
21. Wang SW, Chen J, Jia X, Tam VH, Hu M. Disposition of flavonoids via enteric recycling: structural effects and lack of correlations between in vitro and in situ metabolic properties. *Drug Metab Dispos.* 2006; 34(11):1837–1848. [PubMed: 16882763]
22. Wang S, Kulkarni KH, Tang L, Wang J, Daidoji T, Yokota H, Hu M. Disposition of Flavonoids via Enteric Recycling: UGT1As Deficiency in Gunn Rats Is Compensated by Increases in UGT2Bs Activities. *J. Pharmacol. Exp. Therap.* 2009; 329:1023–1031. 2009. [PubMed: 19264971]
23. Wang SW, Kulkarni KH, Tang L, Wang JR, Yin T, Daidoji T, Yokota H, Hu M. Disposition of flavonoids via enteric recycling: UDP-glucuronosyltransferase (UGT) 1As deficiency in Gunn rats is compensated by increases in UGT2Bs activities. *J Pharmacol Exp Ther.* 2009; 329(3):1023–1031. [PubMed: 19264971]
24. Foster PA, Chander SK, Newman SP, Woo LW, Sutcliffe OB, Bubert C, Zhou D, Chen S, Potter BV, Reed MJ, Purohit A. A new therapeutic strategy against hormone-dependent breast cancer: the preclinical development of a dual aromatase and sulfatase inhibitor. *Clin Cancer Res.* 2008; 14(20):6469–6477. [PubMed: 18927286]
25. Cho K, Mabasa L, Fowler AW, Walsh DM, Park CS. Canola oil inhibits breast cancer cell growth in cultures and in vivo and acts synergistically with chemotherapeutic drugs. *Lipids.* 45(9):777–784. [PubMed: 20730604]
26. Fudge MA, Kavaliers M, Baird JP, Ossenkopp KP. Tamoxifen and raloxifene produce conditioned taste avoidance in female rats: a microstructural analysis of licking patterns. *Life Sci.* 2009; 84(9–10):282–289. [PubMed: 19146862]

Figure 1.

Plasma concentrations of genistein (A), genistein glucuronide (B) and genistein sulfate (C) in female rats in the “proestrus” (diamonds) or the “metoestrus” (triangles) phase of the estrus cycle after single oral dose of genistein 20mg/kg. The symbol “*” indicates $p < 0.05$ compared with “proestrus” group.

Figure 2.

Plasma concentrations of genistein aglycone (A), genistein glucuronide (B) and genistein sulfate (C) in female ovariectomized (circles) and female ovariectomized SD rats with chronic exogenous estradiol (@10μg/kg for 10 days) (diamonds) administration after single oral dose of genistein at 20mg/kg. The symbol “*” indicates $p < 0.05$ compared with “Ovariectomized+E” group.

Figure 3.

Absorption and disposition of genistein in a four-site rat intestinal perfusion model (n = 4). Four segments of the intestine (duodenum, upper jejunum, terminal ileum and colon) were perfused simultaneously at a flow rate of 0.191 ml/min using perfusate containing 10 μ M genistein in the “proestrus” group (columns with dots); the “metoestrus” group (columns with stripes) and ovariectomized rats (columns with broken vertical lines). Amount absorbed (A); intestinal glucuronide excreted (B) and intestinal sulfate excreted (C) were normalized to 10 cm intestinal length using equations 1 to 3 in data analysis. Biliary excretion samples

(n = 4) were collected at every 30min time interval during the entire perfusion experiment. Cumulative amount of biliary excretion of genistein aglycone (D); glucuronide (E) and sulfate (F) was depicted. The arrow indicates a statistically significant difference between two groups.

Comparison of pharmacokinetic parameters for genistein (**Aglycone**), genistein glucuronide (**Glucuronide**), genistein sulfate (**Sulfate**), and total genistein (aglycone+conjugates or **Total**) in female SD rats at different estrus cycle phases, and in female ovariectomized SD rats with or without estrogen supplementation. A single dose of genistein at 20mg/kg was given to each rat, and pharmacokinetic parameters were calculated using a non-compartmental model (Winnonlin 3.3). In this table, “Pro” represents ““proestrus” (elevated estrogen) phase,” “Meto” represents ““metoestrus” (basal estrogen) phase, “Ova” represents ovariectomized female rats without estrogen, and “Ova+E” represents ovariectomized rats supplemented with exogenous estrogen. Comparisons of the results using various statistical means were shown in the “Results” section. Results of statistical analysis of data were shown in Supplemental Materials (Table S3).

Table 1

	Aglycone				Glucuronide				Sulfate				Total			
	Pro	Meto	Ova	Ova+E	Pro	Meto	Ova	Ova+E	Pro	Meto	Ova	Ova+E	Pro	Meto	Ova	Ova+E
C _{max} (µM)	0.10 ± 0.03	0.14 ± 0.07	0.25 ± 0.09	0.06 ± 0.02	0.47 ± 0.14	0.53 ± 0.37	0.52 ± 0.12	0.22 ± 0.04	0.01 ± 0.00	0.01 ± 0.00	0.05 ± 0.01	0.00 ± 0.00	0.53 ± 0.13	0.80 ± 0.30	0.83 ± 0.21	0.26 ± 0.04
AUC (0–24hr) (µM · hr)	0.79 ± 0.10	1.70 ± 0.72	2.38 ± 1.12	0.36 ± 0.1	2.50 ± 1.11	5.97 ± 1.41	4.38 ± 1.97	2.00 ± 1.00	0.07 ± 0.02	0.07 ± 0.02	0.53 ± 0.47	0.04 ± 0.01	3.36 ± 1.18	7.74 ± 1.90	8.58 ± 3.09	2.38 ± 1.02