Variation in leaf flushing date influences autumnal senescence and next year's flushing date in two temperate tree species

Yongshuo S. H. Fu^{a,b,1}, Matteo Campioli^a, Yann Vitasse^c, Hans J. De Boeck^a, Joke Van den Berge^a, Hamada AbdElgawad^d, Han Asard^d, Shilong Piao^b, Gaby Deckmyn^a, and Ivan A. Janssens^a

^aResearch Group of Plant and Vegetation Ecology, Department of Biology, University of Antwerp, B-2610 Wilrijk, Belgium; ^bCollege of Urban and Environmental Sciences, Peking University, Beijing 100871, China; ^cInstitute of Botany, University of Basel, 4056 Basel, Switzerland; and ^dResearch Group of Molecular Plant Physiology and Biotechnology, Department of Biology, University of Antwerp, 2020 Antwerpen, Belgium

Edited by William H. Schlesinger, Cary Institute of Ecosystem Studies, Millbrook, NY, and approved April 8, 2014 (received for review November 21, 2013)

Recent temperature increases have elicited strong phenological shifts in temperate tree species, with subsequent effects on photosynthesis. Here, we assess the impact of advanced leaf flushing in a winter warming experiment on the current year's senescence and next year's leaf flushing dates in two common tree species: *Quercus robur* L. and *Fagus sylvatica* L. Results suggest that earlier leaf flushing translated into earlier senescence, thereby partially offsetting the lengthening of the growing season. Moreover, saplings that were warmed in winter–spring 2009–2010 still exhibited earlier leaf flushing in 2011, even though the saplings had been exposed to similar ambient conditions for almost 1 y. Interestingly, for both species similar trends were found in mature trees using a long-term series of phenological records gathered from various locations in Europe. We hypothesize that this long-term legacy effect is related to an advancement of the endormancy phase (chilling phase) in response to the earlier autumnal senescence. Given the importance of phenology in plant and ecosystem functioning, and the prediction of more frequent extremely warm winters, our observations and postulated underlying mechanisms should be tested in other species.

climate change | tree phenology | spring flushing | leaf senescence

Leaf phenology of temperate trees has recently received particular attention because of its sensitivity to the ongoing climate change (1–3), and because of its crucial role in the forest ecosystem, water and carbon balances, and species distribution (4–6).

A wide variety of methods, such as long-term phenological records (7), indirect measurements of ecosystem greening by remote sensing using satellites or webcam digital images (8–10), and modeling approaches (11–13), have been applied to monitor and study phenological changes. These different approaches, conducted at different spatial scales (from individual plants to biomes), have documented a clear advancement of leaf flushing in temperate climate zones and, to a lesser extent, a delay in leaf senescence (14, 15). Furthermore, various temperature manipulation experiments have simulated the impact of future winter warming on leaf phenology and confirmed an advancement in the timing of leaf flushing in response to warming (16–18). However, the response of leaf flushing to climate warming is highly nonlinear (16, 19, 20), because trees also depend on cold temperatures to break bud dormancy (21–23). This chilling requirement may not (fully) be met in a warming climate, especially at the southern edges of species distribution ranges (5, 24, 25).

Most previous phenological studies have focused on specific phenophases, but how a phenological change (e.g., advanced leaf flushing) affects subsequent phenological events is rarely investigated. Nonetheless, the annual growth cycle of boreal and temperate trees forms an integrated system, where one phenophase in the cycle can affect the subsequent phases (26, 27). Such carryover effects have already been detected in fruit and nut

trees, where winter warming resulted in insufficient chilling (28, 29), which subsequently postponed the onset of flowering, with an associated negative impact on crop yields and crop quality (30, 31). Heide (32) also found that delayed senescence in warm autumns delayed spring leaf flushing in the following year in boreal trees. To our knowledge, however, no study has explored the lagged effect of winter warming-induced earlier leaf flushing on the current year's senescence and on leaf flushing dates after one growing season.

In this study, we exposed young trees to manipulated winter temperature to assess the legacy effect of warming-induced variation in leaf flushing (spring 2010) on the timing of leaf senescence (autumn 2010) and flushing in the following year (spring 2011) in two common deciduous and late-successional temperate tree species: pedunculate oak (*Quercus robur* L.) and European beech (*Fagus sylvatica* L.). Specifically, we tested the hypothesis that the physiological impact of winter warming lasts longer than the current growing season. To confirm our experimental results on young trees, we further explored the legacy effects on mature trees of these two study species using the long-term phenological observations of the European phenology network (www.pep725.eu).

Results

Impact of Winter and Spring Warming on Current Year's Growth. Leaf flushing and leaf senescence in 2010 were both advanced by the

Significance

Leaf phenology of temperate ecosystems is shifting in response to global warming. This affects surface albedo, ecosystem carbon balance, and evapotranspiration, and the response of leaf phenology to climatic drivers has therefore received particular interest. However, despite considerable effort, models have failed to accurately reproduce phenology patterns, likely because mechanistic understanding is incomplete. Here, we show that earlier leaf flushing in response to a warm winter translated into earlier leaf senescence and even earlier leaf flushing in the following year. This legacy effect of winter warming on leaf phenology has important implications for understanding and modelling leaf phenology and its impact on ecosystem functioning, especially in relation to global warming, and is likely to open new research lines.

Author contributions: Y.S.H.F., H.J.D.B., and I.A.J. designed research; Y.S.H.F., M.C., and I.A.J. performed research; Y.S.H.F., M.C., Y.V., H.J.D.B., J.V.d.B., H. AbdElgawad, H. Asard, S.P., G.D., and I.A.J. analyzed data; and Y.S.H.F., M.C., Y.V., H.J.D.B., J.V.d.B., H. Asard, S.P., G.D., and I.A.J. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. E-mail: yongshuo.fu@uantwerpen.be.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1321727111/-DCSupplemental.

+6 °C winter–spring warming for both species ($n = 10$ and $n = 5$, respectively, $P < 0.01$, Fig. 1). Leaf senescence occurred 15 and 18 d earlier in the +6 °C winter–spring-warming treatment for oak and beech, respectively. Per day of advance in leaf flushing, leaf senescence was advanced by 0.36 and 0.47 d for oak and beech, respectively. In agreement with these experimental results on young trees, we also observed an indication for a relationship between leaf flushing and leaf senescence dates in the long-term phenological observations on mature trees of these two study species. After removing the effect of preseason temperature in the partial correlation analysis, the timing of leaf senescence was positively correlated with spring leaf flushing dates in around 70% of study sites, and more than 27% were significant, with mean partial correlation coefficient $r = 0.1$ (PEP network results, Fig. 2, red histograms). In contrast, only 30% of all sites showed negative correlations, and only 9% (oak) and 7% (beech) were significant. The reader should be aware that in our manipulation experiment all trees were exposed to the same growth conditions and only differed in winter temperature and leaf flushing date, whereas the trees in the phenology database not only differed in leaf flushing date but also in growth conditions during the growing season. Taking this variability into account, observing such a clear pattern in the dataset toward positive partial correlations between leaf flushing and senescence dates was remarkable.

Despite the earlier leaf senescence in our manipulation experiment, the length of the growing season was significantly extended ($n = 5$, $P_{\text{unadj}} < 0.01$) in the +6 °C winter–spring-warming treatment for both oak (+27 d) and beech (+20 d), which was due to the larger shift in leaf flushing dates (Table 1). The advanced leaf flushing in 2010 in the +6 °C winter–spring-warming treatment was also associated with some physiological and morphological changes, which were substantial for oak (increased leaf number, higher leaf area per tree, and higher starch accumulation, $n = 5$, all $P_{\text{unadj}} < 0.05$), but less for beech (significant increase in leaf number, $n = 5$, $P_{\text{unadj}} < 0.05$ and a tendency to increase leaf area per tree and total sugar content, $n = 5$, $P_{\text{unadj}} < 0.10$) (Table 1). No significant difference was found between the +6 °C winter–spring-warming treatment and ambient treatment for the other growth and physiological traits

Fig. 1. Leaf flushing and leaf senescence dates for both oak and beech in 2010 in ambient and +6 °C warming treatments during the winter–spring season of 2009–2010. Values are means ± 1 SE for each treatment ($n = 10$ for leaf flushing and $n = 5$ for leaf senescence). The degrees of freedom are 1, 19 (for leaf flushing) and 1, 9 (for leaf senescence). The F values are 128.6 and 228.7 (12.0 and 20.2) for leaf flushing (leaf senescence) between ambient and +6 °C winter–spring-warming treatment for oak and beech, respectively. Different letters denote a significant difference (at $P < 0.05$); standard font is used for oak and italic is used for beech.

Fig. 2. Frequency distribution of partial correlation coefficients between leaf senescence and leaf flushing (red), after controlling for preseason temperature, and between leaf senescence and preseason temperature (blue), after controlling for leaf flushing, for two study species: *F. sylvatica* L. (Upper) and *Q. robur* L. (Lower). Mean values of the partial correlation coefficients across all phenology stations (n), and percentages of the total number of positive correlations, as well as the percentages of statistically significant correlations (in parentheses), are also provided.

measured [i.e., stem diameter and height increment, specific leaf area (SLA), light-saturated net photosynthetic capacity (A_{max}), bud diameter and length, bud numbers, and stem N and C concentration].

Impact of Winter–Spring Warming on Flushing Dates After One Growing Season. A significant positive linear relation was found between leaf flushing dates observed in 2010 (directly after the 2009–2010 warming treatments) and in 2011 for both oak and beech (Fig. 3). This suggests that warming-induced advanced leaf flushing leads to earlier leaf flushing in the following year as well, even though saplings from all treatments had been exposed to the same ambient conditions for almost one full year (post-leaf flushing 2010–April 2011). For example, in 2011, leaf flushing of saplings exposed to the +6 °C winter–spring-warming treatment in 2009–2010 occurred 6 d earlier than leaf flushing of saplings in the ambient treatment for oak ($n = 5$, $P_{\text{unadj}} = 0.002$) and tended to occur also earlier for beech (5 d; $n = 5$, $P_{\text{unadj}} = 0.06$). The relatively flat slope of the linear regression was due to the different range in leaf flushing dates between the two years. In 2010, the warming treatments of winter and spring 2009–2010 enforced a wide range of leaf flushing dates, whereas the ambient conditions of winter 2010–2011, similar for all trees, resulted in a narrow range of leaf flushing dates (Fig. 3).

Interestingly, the positive correlation between two years' leaf flushing dates observed in our warming experiment was also found in mature trees in the long-term field-based phenology observations. The results of partial correlation analysis suggested that, consistent with previous studies, the timing of leaf flushing is associated negatively with preseason temperature in almost all sites (99.6%; significant relationship in 90% of the sites; Fig. 4, blue histograms). This confirms that warmer spring temperatures almost always advance the timing of leaf flushing. However, after removing this dominant impact of preseason temperature, for both oak and beech we also found that the current year's timing of leaf flushing is associated positively with the previous year's leaf flushing dates. This positive correlation was found in ~80% of the study stations, of which 40% exhibited a statistically significant positive partial correlation. In contrast, only 20% of all sites show negative correlations, and only 3% (oak) and 5% (beech) were significant (Fig. 4, red histograms).

Fig. 3. Linear regression between leaf flushing dates in 2010 and in 2011 of oak and beech. Slope indicates the slope of the linear regression. R^2 indicates the coefficient of determination. DF is the degrees of freedom. The larger leaf flushing variation in spring 2010 was caused by the prior different warming manipulation. Between leaf flushing in 2010 and in 2011 all saplings were exposed to the same climate (in the field). Different symbols or numbers correspond to different warming treatments from December 1, 2009 until leaf flushing in spring 2010. Winter–spring-warming treatments (+1 °C, +2 °C, +5 °C, and +6 °C) are shown. The main graphs show the regression through the means per treatment for both oak and beech; insets show the linear regression using the individual saplings data. WS, spring-only warming treatment (+6 °C during the spring period only, mid-February to leaf flushing); WW, winter-only warming treatment (+6 °C during the winter period only, December 1 to mid-February).

dynamics at least for late successional deciduous species, such as oak and beech, in future warmer climate conditions. The carryover effect of the previous winter temperature may be particularly beneficial for ecosystem C sequestration because it can be associated with an advanced leaf flushing and longer growing season. It is known that the C balance of terrestrial ecosystems is particularly sensitive to the changes at the edges of the growing season, such as during leaf flushing (6, 41, 42). The advanced spring flushing would likely improve the C uptake (43) if late frost does not occur (44). However, these feedbacks might not take place if the legacy effect of winter warming on leaf flushing is counterbalanced by a negative effect of winter warming during endodormancy [insufficient chilling leading to an increase of forcing requirement (22)]. Thus, the impact of the legacy effect of warming-induced earlier leaf flushing might be tightly linked to the magnitude and impact of the chilling deficit.

Relationship Between Spring Flushing and Autumn Senescence. Our ancillary measurements on leaf senescence were available for only a limited number of trees and therefore decisive conclusions cannot be drawn. Nevertheless, using the European phenology network dataset, after removing the dominant climate effect in the partial correlation analysis a similar pattern was found in mature trees over long-term series of phenological records. Our results therefore open the door for new insights in our understanding of the drivers of autumnal leaf senescence in temperate trees. The most widely accepted mechanism underlying the onset of leaf senescence in deciduous trees is represented by the environmental control hypothesis, which proposes that leaf senescence is triggered when the unfavorable autumn season comes, that is, a decrease in day length (45), in temperature (25, 46–49), or in both day length and temperature (50). However, our study shows that in both oak and beech saplings leaf senescence started significantly later for trees maintained at ambient conditions during the previous winter than trees exposed to

winter warming, despite the fact that all saplings experienced identical conditions during the growing season. This suggests that the environmental cues alone cannot fully explain the onset of leaf senescence, although they may dominate the variability in senescence date. Further research should confirm these first findings and analyze the physiological drivers of this carryover phenomenon that influences or overrules the environmental control of leaf senescence. In particular, it needs to be verified whether sink limitation partially controls leaf senescence. We observed a larger starch/total sugar content in saplings exhibiting earliest leaf senescence, which might indicate that leaf senescence occurs once the sugar content has reached the maximum carbohydrate storage capacity (51, 52).

Limits and Conclusion

In our experiment we used saplings from a single genotype of two species and grew them in soil fertilized and irrigated in the same way, and in the same ambient light regime (photoperiod). Although the elimination of these potentially confounding determinants of leaf flushing (49, 53) allowed us to detect and elucidate the legacy effects in leaf phenological processes, our study has some shortcomings that future studies need to address. For example, further investigations are necessary to check whether these legacy effects are present in more tree species, especially in indeterminate growth species, because both *F. sylvatica* and *Q. robur* exhibit determinate flushing behavior. The European phenology database included one species exhibiting this growth pattern with sufficient replication to allow a partial correlation analysis, birch (*Betula pendula*), and for this species we found similar, but weaker, patterns compared with the patterns for the two species included in this study (Fig. S1). More experimental results are therefore needed to identify how general our observations are across tree species.

In conclusion, our study suggests that the physiological impact of winter warming lasts longer than the subsequent growing season. The legacy effect of earlier leaf flushing on autumnal leaf senescence and even on leaf flushing after one growing season calls for a renewed attention on the variables responsible for the

Fig. 4. Frequency distribution of partial correlation coefficients between consecutive years' leaf flushing dates (red), after controlling for preseason temperature, and between leaf flushing and preseason temperature (blue), after controlling for leaf flushing in the previous year, for two study species: *F. sylvatica* L. (Upper) and *Q. robur* L. (Lower). Mean values of the partial correlation coefficients across all phenology stations (n) and percentages of the total number of positive correlations, as well as the percentages of statistically significant correlations (in parentheses), are also provided.

interseasonal and interannual “tree memory”, such as, for instance, C reserves. Furthermore, developers of leaf flushing models should be aware that meteorological winter/spring conditions (temperature, daylength, humidity, etc.) are not the only drivers of the leaf-flushing timing, but that the previous year’s meteorological conditions can play also a significant role by shifting the different dormancy phases.

Materials and Methods

Experiment. Seventy saplings (3–4 y old) of both single-genotype oak and beech, originating from a local nursery, were subjected to different warming treatments in climate-controlled, sunlit growth chambers (20, 54) from December 1, 2009 until leaf flushing in spring 2010 at the University of Antwerp (51°19′ N, 4°21′ E). Up to the start, as well as after the temperature manipulations, all nursery-grown saplings were placed outside and subjected to uniform conditions (i.e., equal fertilization, irrigation, and light conditions). Saplings were transplanted into plastic pots (diameter 25 cm, depth 30 cm) with sandy soil in late November 2009 and sufficient slow-release fertilizer was added with 100 g·m⁻². The composition of the slow-release fertilizer was 13–10–20 for N, P, and K, respectively (all in percentage). The saplings were watered as soon as the topsoil seemed dry, normally once or twice a week.

Treatments comprised different intensities of winter–spring warming (+1, +2, +5, or +6 °C above ambient temperatures), winter-only warming (December 1–mid-February, +6 °C), and spring-only warming (mid-February–leaf flushing, +6 °C), and an ambient treatment (+0 °C) in which saplings were kept out of the chambers (for details on warming and chambers see Table S1 and Fig. S2). The chambers provided a stable warming treatment and actual warming was within ±5% of the prescribed value (20, 54). As expected, these different warming treatments elicited different leaf flushing dates in spring 2010 (up to 40-d variation in leaf flushing date for both oak and beech; for more details see refs. 20 and 23). Each sapling was moved out of the chambers to a nearby field in spring 2010, as soon as flushing of the first leaves was complete (i.e., at different time depending on the phenology of each sapling). Within the field, saplings from all treatments were arranged randomly in rows (with 50 cm between rows), fertilized again, and irrigated as soon as the topsoil seemed dry. All saplings were kept together in the field until spring 2011, except for 15 saplings of both species that were harvested destructively in late November 2010 to measure a range of physiological and growth traits in the ambient, +2 °C, and +6 °C winter–spring-warming treatments (details are discussed below). Unfortunately, data were lost for the +2 °C treatment. Leaf flushing dates were thus recorded on 70 saplings per species in spring 2010 and on 55 saplings per species in 2011 (Table S1).

Measurements of Phenology, Growth, and Traits. Buds of oak and beech were formed in late summer, in line with previous studies (55, 56). Leaf flushing observations were conducted on the terminal bud of each individual sapling, according to the following phenology scale: (1) undeveloped bud: bud still in winter dormancy; (2) swollen bud: green or elongated bud with broken scales; (3) leaf flushing: leaf bases still hidden in bud scales but leaf tips detached from the bud axis; and (4) leaf unfolded: the entire leaf blade and the leaf stalk were visible. Monitoring started on February 1 in 2010 and March 1 in 2011 and was repeated every 3 d (between stages 1 and 2) and every 2 d (between stages 2 and 3), always at the same time (2:00–3:00 PM). In this study, we used the starting date of stage 3 to determine leaf flushing date. We obtained exactly the same results when using stage 2. This is likely due to the fact that stage 3 followed stage 2 within 2–5 d across all treatments and species. To simplify, only results using stage 3 were presented. Stem diameter (at 20 cm above the soil) and height were measured for all saplings on December 1, 2009 and November 20, 2010.

In autumn 2010, leaf senescence was recorded for five saplings in the ambient (+0 °C) and +6 °C winter–spring-warming treatments. The selected saplings from these contrasted treatments covered the extremes of observed leaf flushing dates in 2010 for both oak and beech. Leaf senescence was defined as the date at which half of the leaves were colored or dropped, following the method described in Vitasse et al. (57). Growing season length was quantified for individual trees as the difference between days of the year of senescence and leaf flushing.

We compared a wide range of physiological and morphological traits between the saplings maintained at ambient conditions and those exposed to the +6 °C winter–spring-warming treatment, which had exhibited much earlier leaf flushing. Total leaf number per tree, $A_{\max}$, SLA, and total leaf area per tree were measured on July 22, 2010. $A_{\max}$ was measured at

photosynthetically active radiation = 2,000 μmol·m⁻²·s⁻¹ with a portable open photosynthetic system (LI-6400; Li-Cor). To calculate the SLA (expressed as the ratio of leaf area to leaf dry mass, square centimeters per gram), five leaves were collected from each individual in each treatment. The area of each leaf was measured with a planimeter and then all of the leaves were dried at 70 °C for 3 d to determine the dry weight. Furthermore, we measured the length of each leaf from each sapling and calculated the total leaf area per tree using an allometric function relating leaf length to leaf area, using the data derived from the leaves used to determine SLA. In early December 2010 the total number of buds per individual tree was counted and the size (length and diameter) of the five uppermost apical buds was measured. Branches, stems, and roots were weighed separately to obtain the total fresh biomass (for beech, most senesced leaves were still attached). Subsamples thereof were then dried at 70 °C for 3 d and weighed again to obtain the dry biomass, from which the dry/fresh weight ratio was determined that was used to determine total dry weight per tree.

A 5-cm segment of stem, root, and branch was taken from each sapling and analyzed for carbohydrate content and C and N concentration. The starch and sugar contents were measured by the anthrone method (58). Sugars and carbohydrates were extracted from dried and ground plant material. First, soluble sugars were extracted with aqueous ethanol, and then starch was extracted with 80% ethanol. The concentrations of total sugars (soluble sugars + insoluble sugars) and starch were expressed as milligrams of glucose equivalents per gram of dry weight. The C and N concentrations were measured with a dynamic flush combustion method in a NC 2100 Soil Analyzer (Carlo Erba Strumentazione).

Data Analysis. The relation between the 2010 and 2011 leaf flushing dates was analyzed with a linear regression, both through the means of the different treatments ($n = 7$) and through the individual sapling data ($n = 55$). One-way ANOVA was used to evaluate the difference between +6 °C winter–spring warming and ambient treatments of leaf flushing ($n = 10$), leaf senescence ($n = 5$), and growing season length ($n = 5$), as well as the physiological and morphological traits ($n = 5$), all in 2010. Additionally, a Bonferroni-based correction for multiple comparisons was applied (59), taking into account possible correlations between the growth traits. The adjusted P value is

$$P_{adj} = 1 - \left(1 - P_{unadj}(k)^{g(k)}\right) \quad \text{where} \quad g(k) = M^{1-r(k)},$$

where $P_{unadj}(k)$ is the unadjusted P value for the k^{th} growth trait, $r(k)$ is the mean correlation among the outcomes other than outcome k , and M is the number of growth traits being tested. P values <0.05 were considered significant and values <0.10 as indicating a tendency. All statistical analyses were conducted using SPSS 16.0 (SPSS Inc.).

Testing the Legacy Effect on Mature Trees. We further explored the legacy effects of phenological events on mature trees of the two same species, *Q. robur* L. (oak) and *F. sylvatica* L. (beech), using data from the European phenology network (www.pep725.eu) at more than 1,000 locations for each species (Fig. S3). The leaf flushing and leaf senescence dates were defined according to the BBCH (Biologische Bundesanstalt, Bundesartenamt und Chemische Industrie) code (60).

Preseason temperature is known to determine the timing of spring leaf flushing (7), as well as the timing of leaf senescence (61). To test the correlation between the leaf flushing dates in two consecutive years, as well as between the dates of the current year’s leaf flushing and the current year’s leaf senescence, we applied a partial correlation analysis to remove the covariate effects of preseason temperature. This method has been successfully applied to remove the covariate effects between study factors in other ecological studies (62, 63). The preseason periods were defined as 90 d preceding the day of leaf senescence or leaf flushing. The selected sites from the European phenology database each had more than 30 y of leaf flushing observations for which also the previous year’s leaf flushing date was recorded (needed for testing the correlation between flushing dates in consecutive years) or included at least 30 y of observations on both leaf flushing and leaf senescence dates within the same year (needed for testing the correlation between the current year’s leaf flushing vs. leaf senescence) during the period 1950–2011. The daily mean air temperature of each site was derived from a gridded climate dataset of daily mean temperature at 0.25° spatial resolution (~25 km, ERAWATCH).

ACKNOWLEDGMENTS. The authors thank Prof. Christian Körner and Prof. Van Dongen Stefan for comments on data analysis and Dr. Maarten Op de Beeck and Dr. Raphael Bequet for field assistance. We thank the reviewers

and editor of this manuscript for their valuable comments and suggestions that helped us to substantially improve the paper. Long-term phenological data were provided by the members of the PEP725 project (www.pep725.eu). This research has been financially supported by the research

project Greenhouse Gas Management in European Land Use Systems (Contract 244122). Y.S.H.F. holds a research grant from the China Scholarship Council. H.J.D.B. and M.C. are Postdoctoral Fellows of the Research Foundation–Flanders.

- Walther G-R, et al. (2002) Ecological responses to recent climate change. *Nature* 416(6879):389–395.
- Peñuelas J, Rutishauser T, Filella I (2009) Ecology. Phenology feedbacks on climate change. *Science* 324(5929):887–888.
- Polgar CA, Primack RB (2011) Leaf-out phenology of temperate woody plants: From trees to ecosystems. *New Phytol* 191(4):926–941.
- Baldocchi D, Wilson K (2001) Modeling CO₂ and water vapor exchange of a temperate broadleaved forest across hourly to decadal time scales. *Ecol Modell* 142(1–2): 155–184.
- Chaine I (2010) Why does phenology drive species distribution? *Philos Trans R Soc Lond B Biol Sci* 365(1555):3149–3160.
- Piao S, P. Friedlingstein, P. Ciais, N. Viovy, Demarty J (2007) Growing season extension and its impact on terrestrial carbon cycle in the Northern Hemisphere over the past 2 decades. *Global Biogeochemical Cycles* 21(3):GB3018, 3010.1029/2006GB002888.
- Menzel A, et al. (2006) European phenological response to climate change matches the warming pattern. *Glob Change Biol* 12(10):1969–1976.
- Schwartz MD, ed (2003) *Phenology: An Integrative Environmental Science* (Kluwer, Dordrecht), p 564.
- Zhang XY, et al. (2003) Monitoring vegetation phenology using MODIS. *Remote Sens Environ* 84(3):471–475.
- Richardson AD, et al. (2007) Use of digital webcam images to track spring green-up in a deciduous broadleaf forest. *Oecologia* 152(2):323–334.
- Chaine I (2000) A unified model for budburst of trees. *J Theor Biol* 207(3):337–347.
- Hänninen H (1990) Modelling bud dormancy release in trees from cool and temperate regions. *Acta For Fenn* 213:1–47.
- Fu YSH, Campioli M, Deckmyn G, Janssens IA (2012) Bayesian comparison of six different temperature-based budburst models for four temperate tree species. *Ecol Modell* 230:92–100.
- Vitasse Y, et al. (2009) Leaf phenology sensitivity to temperature in European trees: Do within-species populations exhibit similar responses? *Agric For Meteorol* 149(5): 735–744.
- Jeong SJ, Ho CH, Gim HJ, Brown ME (2011) Phenology shifts at start vs. end of growing season in temperate vegetation over the Northern Hemisphere for the period 1982–2008. *Glob Change Biol* 17(7):2385–2399.
- Morin X, Roy J, Sonié L, Chaine I (2010) Changes in leaf phenology of three European oak species in response to experimental climate change. *New Phytol* 186(4):900–910.
- Wolkovich EM, et al. (2012) Warming experiments underpredict plant phenological responses to climate change. *Nature* 485(7399):494–497.
- Chmielewski F, Rotzer T (2001) Response of tree phenology to climate change across Europe. *Agric For Meteorol* 108(2):101–112.
- Yu HY, Luedeling E, Xu JC (2010) Winter and spring warming result in delayed spring phenology on the Tibetan Plateau. *Proc Natl Acad Sci USA* 107(51):22151–22156.
- Fu YH, Campioli M, Deckmyn G, Janssens IA (2012) The impact of winter and spring temperatures on temperate tree budburst dates: Results from an experimental climate manipulation. *PLoS ONE* 7(10):e47324.
- Chaine I, Morin X, Bugmann H (2010) Warming, photoperiods, and tree phenology. *Science* 329(5989):277–278, author reply 278.
- Murray MB, Cannell MGR, Smith RI (1989) Date of budburst of 15 tree species in Britain following climatic warming. *J Appl Ecol* 26(2):693–700.
- Fu YSH, Campioli M, Deckmyn G, Janssens IA (2013) Sensitivity of leaf unfolding to experimental warming in three temperate tree species. *Agric Meteorol* 181:125–132.
- Dantec CF, et al. (2014) Chilling and heat requirements for leaf unfolding in European beech and sessile oak populations at the southern limit of their distribution range. *Int J Biometeorol*, in press.
- Vitasse Y, et al. (2011) Assessing the effects of climate change on the phenology of European temperate trees. *Agric For Meteorol* 151(7):969–980.
- Sarvas R (1974) Investigations on the annual cycle of development of forest trees. II. Autumn dormancy and winter dormancy. *Commun Inst For Fenn* 84:1–101.
- Hänninen H, Tanino KK (2011) Tree seasonality in a warming climate. *Trends Plant Sci* 16(8):412–416.
- Baldocchi D, Wong S (2008) Accumulated winter chill is decreasing in the fruit growing regions of California. *Clim Change* 87:5153–5166.
- Luedeling E, Girvetz EH, Semenov MA, Brown PH (2011) Climate change affects winter chill for temperate fruit and nut trees. *PLoS ONE* 6(5):e20155.
- Chmielewski MD (2003) Phenology and agriculture. *Phenology: An Integrative Environmental Science*, ed Schwartz MD (Kluwer, Dordrecht), Vol 39, pp 505–522.
- Petri JL, Berenhausen LG (2004) Consequences of insufficient winter chilling on apple tree bud-break. *Acta Hort* 662:53–60.
- Heide OM (2003) High autumn temperature delays spring bud burst in boreal trees, counterbalancing the effect of climatic warming. *Tree Physiol* 23(13):931–936.
- Lang GA, Early JD, Martin GC, Darnell RL (1987) Endo-, para- and eco-dormancy, physiological terminology and classification for dormancy research. *HortScience* 22(3): 371–377.
- Vitasse Y, Basler D (2013) What role for photoperiod in the bud burst phenology of European beech. *Eur J For Res* 132(1):1–8.
- Myking T, Heide OM (1995) Dormancy release and chilling requirement of buds of latitudinal ecotypes of *Betula pendula* and *B. pubescens*. *Tree Physiol* 15(11):697–704.
- Charrier G, Ameglio T (2011) The timing of leaf fall affects cold acclimation by interactions with air temperature through water and carbohydrate contents. *Environ Exp Bot* 72(3):351–357.
- Keller F, Körner C (2003) The role of photoperiodism in alpine plant development. *Arct Antarct Alp Res* 35(3):361–368.
- Morin X, et al. (2007) Variation in cold hardiness and carbohydrate concentration from dormancy induction to bud burst among provenances of three European oak species. *Tree Physiol* 27(6):817–825.
- Dufrène E, et al. (2005) Modelling carbon and water cycles in a beech forest. Part I: Model description and uncertainty analysis on modelled NEE. *Ecol Modell* 185: 407–436.
- Campioli M, et al. (2013) Can decision rules simulate carbon allocation for years with contrasting and extreme weather conditions? A case study for three temperate beech forests. *Ecol Modell* 243(10):42–55.
- Luyssaert S, et al. (2007) CO₂ balance of boreal, temperate, and tropical forests derived from a global database. *Glob Change Biol* 13(12):2509–2537.
- Piao SL, et al. (2008) Net carbon dioxide losses of northern ecosystems in response to autumn warming. *Nature* 451(7174):49–52.
- Piao SL, Friedlingstein P, Ciais P, Viovy N, Demarty J (2007) Growing season extension and its impact on terrestrial carbon cycle in the Northern Hemisphere over the past 2 decades. *Global Biogeochem Cy* 21(3).
- Hufkens K, et al. (2012) Ecological impacts of a widespread frost event following early spring leaf-out. *Glob Change Biol* 18(7):2365–2377.
- Keskitalo J, Bergquist G, Gardeström P, Jansson S (2005) A cellular timetable of autumn senescence. *Plant Physiol* 139(4):1635–1648.
- Richardson AD, Bailey AS, Denny EG, Martin CW, O’Keefe J (2006) Phenology of a northern hardwood forest canopy. *Glob Change Biol* 12(7):1174–1188.
- White MA, Thornton PE, Running SW (1997) A continental phenology model for monitoring vegetation responses to interannual climatic variability. *Global Biogeochem Cycles* 11(2):217–234.
- Estrella N, Menzel A (2006) Response of leaf colouring of four deciduous tree species to climate and weather in Germany. *Clim Res* 32:253–267.
- Sparks TH, Menzel A (2002) Observed changes in seasons: An overview. *Int J Climatol* 22:1715–1725.
- Farnsworth E (2004) Hormones and shifting ecology throughout plant development. *Ecology* 85:5–15.
- Herold A (1980) Regulation of photosynthesis by sink activity - the missing link. *New Phytol* 86(2):131–144.
- Paul MJ, Foyer CH (2001) Sink regulation of photosynthesis. *J Exp Bot* 52(360): 1383–1400.
- Cleland EE, Chaine I, Menzel A, Mooney HA, Schwartz MD (2007) Shifting plant phenology in response to global change. *Trends Ecol Evol* 22(7):357–365.
- Naudts K, van den Berge J, Janssens AI, Nijs I, Ceulemans R (2011) Does an extreme drought event alter the response of grassland communities to a changing climate? *Environ Exp Bot* 70:151–157.
- Meier IC, Leuschner C (2008) Leaf size and leaf area index in *Fagus sylvatica* forests: Competing effects of precipitation, temperature and nitrogen availability. *Ecosystems* (NY) 11:655–669.
- Vitasse Y, et al. (2013) Elevational adaptation and plasticity in seedling phenology of temperate deciduous tree species. *Oecologia* 171(3):663–678.
- Vitasse Y, Porté AJ, Kremer A, Michalet R, Delzon S (2009) Responses of canopy duration to temperature changes in four temperate tree species: Relative contributions of spring and autumn leaf phenology. *Oecologia* 161(1):187–198.
- Marshall JD (1985) Carbohydrate status as a measure of seedling quality. *Evaluating Seedling Quality: Principles, Procedures, and Predictive Abilities of Major Tests*, ed Duryea ML (Oregon State Univ, Corvallis, OR), pp 49–58.
- Sankoh AJ, Huque MF, Dubey SD (1997) Some comments on frequently used multiple endpoint adjustment methods in clinical trials. *Stat Med* 16(22):2529–2542.
- Meier N, Rutishauser T, Pfister C, Wanner H, Luterbacher J (2007) Grape harvest dates as a proxy for Swiss April to August temperature reconstructions back to AD 1480. *Geophys Res Lett* 34(20):L20705, 10.1029/2007GL031381.
- Fracheboud Y, et al. (2009) The control of autumn senescence in European aspen. *Plant Physiol* 149(4):1982–1991.
- Beer C, et al. (2010) Terrestrial gross carbon dioxide uptake: Global distribution and covariation with climate. *Science* 329(5993):834–838.
- Peng SS, et al. (2013) Asymmetric effects of daytime and night-time warming on Northern Hemisphere vegetation. *Nature* 501(7465):88–92.