Published in final edited form as:

Eur J Pharmacol. 2014 September 15; 0: 4–18. doi:10.1016/j.ejphar.2013.11.015.

Intracellular Calcium Channels in Protozoa

Roberto Docampo^{a,*}, Silvia N.J. Moreno^a, and Helmut Plattner^{b,**}

^aCenter for Tropical and Emerging Global Diseases and Department of Cellular Biology, University of Georgia, GA 30606, USA

^bDepartment of Biology, University of Konstanz, Konstanz 78457, Germany

Abstract

Ca²⁺-signaling pathways and intracellular Ca²⁺ channels are present in protozoa. Ancient origin of inositol 1,4,5-trisphosphate receptors (IP₃Rs) and other intracellular channels predates the divergence of animals and fungi as evidenced by their presence in the choanoflagellate *Monosiga brevicollis*, the closest known relative to metazoans. The first protozoan IP₃R cloned, from the ciliate *Paramecium*, displays strong sequence similarity to the rat type 3 IP₃R. This ciliate has a large number of IP₃- and ryanodine(Ry)-like receptors in 6 subfamilies suggesting the evolutionary adaptation to local requirements for an expanding diversification of vesicle trafficking. IP₃Rs have also been functionally characterized in trypanosomatids, where they are essential for growth, differentiation, and establishment of infection. The presence of the mitochondrial calcium uniporter (MCU) in a number of protozoa indicates that mitochondrial regulation of Ca²⁺ signaling is also an early appearance in evolution, and contributed to the discovery of the molecular nature of this channel in mammalian cells. There is only sequence evidence for the occurrence of two-pore channels (TPCs), transient receptor potential Ca²⁺ channels (TRPCs) and intracellular mechanosensitive Ca²⁺-channels in *Paramecium* and in parasitic protozoa.

Keywords

Calcium channels; Protozoa; *Paramecium*; *Toxoplasma*; *Trypanosoma*; mitochondrial calcium uniporter

1. Introduction

Calcium ion (Ca²⁺) controls a variety of cellular functions in protozoa. As occurs with mammalian cells, the cytosolic Ca²⁺ concentration [Ca²⁺]_i of protozoa is maintained at very low levels (of the order of 10⁻⁷ M). The cytosolic Ca²⁺ level is responsible for the regulation of Ca²⁺-dependent and Ca²⁺-controlled proteins. Although the total calcium

© 2013 Elsevier B.V. All rights reserved.

*Corresponding author. Tel.: +706 542 8104; fax: 706 542 9493. rdocampo@uga.edu. **Corresponding author. Tel.: +7531 882 228. helmut.plattner@uni-konstanz.de.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

inside protozoan cells is much higher than 10^{-7} M, the bulk of this calcium is either bound to proteins, polyphosphate, membranes or other cellular constituents, or is sequestered inside intracellular organelles through the activity of pumps, channels, and exchangers, and released when needed by a variety of intracellular Ca^{2+} channels.

Recent genomic studies (King *et al.*, 2008) have revealed that many ion channels including Ca^{2+} channels previously thought to be restricted to animals, can be traced back to one of the unicellular ancestor of animals, *Monosiga brevicollis*, a choanoflagellate protozoan belonging the supergroup Opisthokonta, which also includes animals, and fungi. Genes encoding homologues to various types of plasma membrane Ca^{2+} channels are present: store-operated channel (Orai) and the endoplasmic reticulum sensor protein stromal interaction molecule (Stim); voltage-operated channel (similar to dihydropyridine-sensitive L-type Ca^{2+} channel); ligand-operated channels (nicotinic acetylcholine receptor and P2X purinergic receptor); transient receptor potential (TRP) channels; and second messenger-operated channel (cyclic nucleotide-gated channel) (Cai, 2008). This protozoan appears to possess all 5 modes of regulated Ca^{2+} entry across the plasma membrane identified in animals (Parekh and Putney, 2005), although their physiological validation is needed (Cai, 2008). *Monosiga brevicollis* has also 4 homologues of the inositol 1,4,5-trisphosphate receptor (IP_3R), and a homologue to the mitochondrial calcium uniporter (XP_001749044), but no homologues to ryanodine receptors (RyR) (Cai, 2008). However, no functional studies have been reported with any of these channels.

Evidently the evolution of eukaryotic cells is characterized by increasing genomic information that allows for increasing complexity of intracellular structure, dynamics and signaling mechanisms. Target-oriented vesicle trafficking requires not only an inventory of membrane-specific proteins, such as SNAREs (Malsam *et al.*, 2008) and small GTPases (Zerial and McBride, 2001), but also provisions for Ca^{2+} signaling in a very local area where membranes have to interact (Neher, 1998). Ca^{2+} may come from the external medium or be locally released from stores via Ca^{2+} -release channels (CRC) so that Ca^{2+} can locally drive docking, priming and eventual fusion of membranes (Rizo *et al.*, 2006). Cell contraction is another example. Ca^{2+} is most appropriate for such functions because of its specific, reversible binding to Ca^{2+} -binding proteins, CaBP, which in the end transmit the signal by a conformational change in effector protein molecules (Klee *et al.*, 1980; Rizo *et al.*, 2006). On the one hand global regulation of intracellular Ca^{2+} concentration, $[\text{Ca}^{2+}]_i$, is mandatory to avoid the overall toxic effect of Ca^{2+} (Case *et al.*, 2007). On the other hand, local $[\text{Ca}^{2+}]_i$ regulation also has to account for diffusional spread by a square function, whereas most molecular effects of Ca^{2+} depend on a higher power-function of $[\text{Ca}^{2+}]_i$ (Neher, 1998). Binding to CaBPs, sequestration into organelles and extrusion from the cell antagonize the occurrence of too high and diffuse $[\text{Ca}^{2+}]_i$ values after stimulation. Remarkably, the phenomena described in this review, as well as the CRC types mentioned, are all found already in protozoa. Nevertheless, with these cells stringent analyses of Ca^{2+} signaling and the subsiding intracellular CRCs have remained elusive until quite recently.

The protozoan organisms whose Ca^{2+} signaling and subsiding CRCs are currently investigated in our labs include ciliates (*Paramecium*), their close relatives, Apicomplexa (including pathogenic species of *Plasmodium* [malaria causing agent] and *Toxoplasma*) as

well as some pathogenic flagellates (trypanosomatids). With these organisms, CRCs have been characterized at a molecular level, in conjunction with functional studies. There is a wide gap between evolutionary levels: ciliates close to recent forms have emerged ~800 to 850 million years ago, non-parasitic Apicomplexa ~500 million years (Douzery *et al.*, 2004) and mammalian apicomplexan parasites ~13 million years ago (Ricklefs and Outlaw, 2010). There is also some information available on the Ca^{2+} dynamics in social amoeba of the myxomycete *Dictyostelium*, which clearly possesses Ca^{2+} signaling pathways (Allan and Fisher, 2009), but information about CRCs in these cells is scant.

A *Paramecium tetraurelia* cell is up to ~100 μm in size and exhibits distinct intracellular vesicle trafficking pathways (Allen and Fok, 2000), essentially including all those known from metazoan cells. The pathogenic forms discussed are ~10 times smaller, but also contain specific vesicle-trafficking pathways, such as endocytosis vesicles and organelles for intracellular digestion (trypanosomatids, Apicomplexa). Apicomplexa also possess secretory organelles for exocytosis. Due to their small size and their complicated lifestyle the parasites are much more difficult to study than their free-living relatives. Using fluorescent dyes in both ciliates and Apicomplexa, a considerable Ca^{2+} signal could be recorded during exocytosis of secretory organelles, such as trichocysts (Klauke and Plattner, 1997) and during motility (Lovett and Sibley, 2003), respectively.

Values for steady state $[\text{Ca}^{2+}]_i$ in widely different cells, from protozoa to mammals, are of the order of 50 to 100 nM at rest and stimulation generally causes an increase by a factor of 10 to 100 (Bootman and Berridge, 1995). This frame also applies to ciliates (Klauke and Plattner, 1997) and to parasitic protozoa (Vieira and Moreno, 2000; Moreno *et al.*, 1994). $[\text{Ca}^{2+}]_i$ determined in *Paramecium* under steady state conditions yields values between 60 and 100 nM. It has to be stressed that measurements performed with fluorescent dyes, even when calibrated, systematically underestimate the real local $[\text{Ca}^{2+}]_i$ increase during activation because of its considerable local restriction. More realistic local, functionally relevant values are obtained by probing the threshold inhibitory effect of Ca^{2+} chelators with appropriate binding properties (Neher, 1995). For instance, during exocytosis stimulation $[\text{Ca}^{2+}]_i$ in the cell cortex peaked at ~400 nM with fluorescent dyes measurements, whereas chelator application during stimulation indicated the increase in $[\text{Ca}^{2+}]_i$ to the micromolar range (Klauke and Plattner, 1997).

2. Calcium stores

The paradigm of a Ca^{2+} store in all eukaryotic cells is the endoplasmic reticulum (ER), together with the sarcoplasmic reticulum (SR) in muscle cells (Berridge *et al.*, 2000; Berridge *et al.*, 2003; Clapham, 2007; Cai, 2008). Since Ca^{2+} is stored in many more organelles such stores and their CRCs deserve special attention also in protozoa, including ciliates and parasitic protozoa (Plattner *et al.*, 2012).

Subsequent to stimulation Ca^{2+} is sequestered into different organelles and then may be available later on for release via CRCs in a constitutive manner or in the context of signaling processes. Ca^{2+} can, thus, regulate exocytosis, endocytosis, phagocytosis, fusion of endosomes of different stages with phagosomes, phagosome formation, membrane

recycling, phago-lysosome fusion etc. (Hay, 2007; Zampese and Pizzo, 2012). In mammalian cells, many of these organelles, specifically early endosomes (Luzio *et al.*, 2010) and lysosomes (Christensen *et al.*, 2002), are known to store Ca^{2+} (Hay, 2007; Sherwood *et al.*, 2007) and the membranes of many of them contain CRCs (Zampese and Pizzo, 2012). The main types of CRCs found in metazoans up to mammalian cells are IP_3R (Taylor *et al.*, 2004; Bezprozvanny, 2005), RyR (Hamilton, 2005; Mackrill, 2012), transient receptor potential Ca^{2+} channels, TRPC (Patel and Docampo, 2009), and two pore channels, TPC, occurring mainly in acidic compartments (Galione *et al.*, 2009; Galione *et al.*, 2010; Patel and Docampo, 2010). All these channel types also occur in protozoa.

Work with *P. tetraurelia* was started with *Paramecium* database (DB) analysis and further evaluation by expression, localization and functional studies. Thus, a plethora of CRCs related to RyRs and to IP_3Rs , or to both, were identified (Ladenburger *et al.*, 2006; Ladenburger *et al.*, 2009; Ladenburger and Plattner, 2011). The different CRC types are scattered over the many sites of specific membrane interactions. The functions of IP_3Rs (Ladenburger *et al.*, 2006) and of RyRs (more safely to be addressed as RyR-like proteins, RyR-LP) (Ladenburger *et al.*, 2009) were investigated in more detail. Further on the analysis concentrated on cortical stores (alveolar sacs) and dense core-secretory organelle (trichocyst) exocytosis as well as on the contractile vacuole complex that serves for osmoregulation and maintaining the internal ionic balance, particularly of Ca^{2+} .

In Apicomplexa secretory organelles include rhoptries, micronemes, and dense granules whose exocytosis is mandatory for attachment to a host cell, invasion and establishment of the parasitophorous vacuole. Their secretion requires Ca^{2+} signals, presumably based on IP_3 (Lovett *et al.*, 2002; Lovett and Sibley, 2003) or cADP ribose (cADPR) (Chini *et al.*, 2005; Nagamune *et al.*, 2008) signaling. Database search by different groups did not allow for the identification of either IP_3Rs or of RyRs (or RyR-LPs) in Apicomplexa (Nagamune and Sibley, 2006; Plattner *et al.*, 2012) although intracellular application of IP_3 has facilitated host cell infection (Lovett *et al.*, 2002). In trypanosomatids IP_3Rs have recently been identified on a molecular level, and probed functionally, in two species, *Trypanosoma brucei* (Huang *et al.*, 2013) and *T. cruzi* (Hashimoto *et al.*, 2013). In *Paramecium*, the dense core-secretory organelles called trichocysts can explosively be released by exocytosis within fractions of a second, thus making this system amenable to sub-second analysis (Plattner and Hentschel, 2006). The reaction serves for warding off predators very efficiently (Harumoto and Miyake, 1991).

In summary, CRCs must have evolved early in evolution, i.e. already at the level of protozoa. These CRCs include not only IP_3Rs and RyR-LPs (Plattner and Verkhatsky, 2013) but also TRPCs and TPCs (Patel and Docampo, 2010; Plattner *et al.*, 2012) as well as the mitochondrial calcium uniporter (Docampo and Lukes, 2012) that will be discussed later. On a speculative basis one may envisage also intracellular mechano-sensitive Ca^{2+} channels as part of an ancient CRC inventory. This expectation is based on the finding of stomatin in the membranes of the contractile vacuole complex and of food vacuoles (Reuter *et al.*, 2013), considering that generally the scaffolding protein stomatin is structurally and functionally associated with mechano-sensitive Ca^{2+} -channels in metazoans (Lapatsina *et al.*, 2012). By sensing the internal tension in these organelles such channels may initiate a

Ca^{2+} signal for the release of contractile vacuole contents by exocytosis or for fusion processes along the food vacuole pathway (Reuter *et al.*, 2013).

Scrutiny at a molecular level, including domain analysis, intracellular localization and functional analysis, including also gene silencing, will provide us with important new insight into Ca^{2+} signaling mechanisms not only in free-living, but also in pathogenic protozoan species. Also the precise function of many of the CRCs recently identified remains to be elucidated.

3. Intracellular calcium channels in ciliates

3.1 IP_3R - and RyR-type CRCs in *Paramecium* – identification and localization

Only quite recently could such channels be identified, based on genomic sequences indicative of characteristic domain structures. Six subfamilies of putative CRCs, *PtCRC*-I to *PtCRC*-VI, which all encompass several paralogs, were found and cloned. These are identified by addition of arabic numbers, e.g. *PtCRC*-IV-2 (some with further subforms, such as *PtCRC*-III-1a and -1b). For detailed terminology, see (Ladenburger and Plattner, 2011). In part they show characteristics of IP_3Rs and in part of RyRs and they were the first of these types unambiguously identified in protozoa at a molecular level. Some other CRCs of this family show one or the other, but not all of the characteristic features of either IP_3Rs or RyRs. Domains under consideration include the IP_3 -binding domain (IP_3 -BD), the RyR/ IP_3R -homology (RIH) domain, the pore domain with its transmembrane domains and the selectivity filter as well as regions with variable similarity to mammalian IP_3Rs or RyRs. It was thus identified an unexpected total of 34 IP_3R - and RyR-like channels in the *P. tetraurelia* cell (Ladenburger and Plattner, 2011). Generally only a selected paralog of one subfamily has been analyzed in more detail. This high number of *PtCRC*s clearly is the result of several whole genome duplications (Aury *et al.*, 2006) as one can derive from the high similarity of many – though not all - paralogs contained in each subfamily. Remarkably all sequences are expressed, except one from subfamily VI, which may be on the way to pseudogene formation (Ladenburger and Plattner, 2011).

These attempts of identification were complemented by localization of *PtCRC*s at the light and electron microscope level and by functional analyses: IP_3 binding and activation of injected caged IP_3 on the one hand (Ladenburger *et al.*, 2006) and on the other hand by activation by RyR agonists (Ladenburger *et al.*, 2009) whose secretagogue effect had been probed before (Klauke and Plattner, 1998; Klauke *et al.*, 2000; Plattner and Klauke, 2001). Also discussed in more detail below are fluorescent imaging experiments.

The members of the 6 subfamilies are all distinctly placed in the cell (Fig. 1) and, according to gene silencing experiments, they can account for the regulation of widely different functions in different regions of the highly complex *Paramecium* cell. In detail, subfamily I channels (in our designation *PtCRC*-I) are associated with the ER (Ladenburger and Plattner, 2011). *PtCRC*-II/ IP_3Rs are restricted to the contractile vacuole complex (serving in fresh water organisms for the expulsion of water and of some ions, including an excess of Ca^{2+}) and, therefore, may fine-tune Ca^{2+} homeostasis by partial reflux of Ca^{2+} (Ladenburger *et al.*, 2006). Spontaneous Ca^{2+} puffs are seen along the tubular extensions of

this organelle, indicating constitutively active IP₃R type channels - a phenomenon reported later on also for a chicken lymphocyte cell line (Cardenas *et al.*, 2010) and mammalian atrial myocytes (Horn *et al.*, 2013). *PtCRC*-III molecules are associated with recycling vesicles engaged in phagosome formation (Ladenburger and Plattner, 2011). A sequence indicating an IP₃-BD occurs in the members of these three subfamilies, I to III, but has been experimentally verified only with *PtCRC*-II type channels. *PtCRC*-IV channels display structural and functional characteristics of RyRs (Ladenburger *et al.*, 2009). They are localized to the established subplasmalemmal Ca²⁺-stores, the alveolar sacs. Note that in apicomplexan parasites, the structural equivalent is the “inner membrane complex”, whose relevance for Ca²⁺ signaling is not known (Plattner *et al.*, 2012). Silencing reduces stimulated exocytosis in response to RyR agonists. The remaining *PtCRC* subfamily types, *PtCRC*-V and *PtCRC*-VI, have a more complex distribution (Ladenburger and Plattner, 2011), as shown in Fig. 1.

Altogether, immunolocalization studies with antibodies against subfamily *PtCRC* members revealed widely different, but distinct localization and functional engagement of the different Ca²⁺-release channel types also turned out to be different. Some *PtCRC*s occur in different organelles and some organelles possess different CRC types, as summarized in Fig. 1, with some selected examples presented in Figs. 2 and 3. In alveolar sacs *PtCRC*-IV molecules are scattered over the entire peripheral part where they face the cell membrane (Ladenburger *et al.*, 2009), thus allowing spilling of Ca²⁺ over exocytosis sites. An additional type of CRCs is found in this organelle, i.e. *PtCRC*-V-4, with characteristics of an IP₃R; it is positioned laterally, i.e. where adjacent sacs approach each other (Ladenburger and Plattner, 2011). Other *PtCRC*s are generally associated with vesicles participating in trafficking, as known from *in vivo* and ultrastructural analyses, as well as from the topology of specific SNARE proteins (Plattner, 2010) that mediate specific membrane interactions.

3.2 Functional aspects of IP₃R- and RyR-type CRCs in Paramecium

Only *PtCRC*-II (IP₃Rs) and *PtCRC*-IV (RyR-LP) channels have been analyzed in some more detail by post-transcriptional gene silencing, i.e. by applying the method described by (Galvani and Sperling, 2002). Genes diverging by >15 percent can thus be differentially silenced at a post-transcriptional level. This means that not all paralogs of a subfamily can be safely addressed and a similar uncertainty holds for antibodies used for immunolocalization.

3.2.1 IP₃ receptors—The *PtCRC*-II/IP₃R localized selectively to the contractile vacuole system, from the contractile vacuole bladder, over emanating radial (connecting) arms to the entire “smooth spongione”, a network of anastomosing membrane-bounded tubules. Here, a H⁺ gradient generated by a vacuolar H⁺-ATPase pump in the adjacent “decorated spongione” drives chemiosmotically the uptake of Ca²⁺ and water (Fok *et al.*, 1995; Wassmer *et al.*, 2009) for collection by the smooth spongione, delivery to the vacuole bladder and periodic extrusion by exocytosis (Stock *et al.*, 2002). What may the function of the organelle-specific CRCs then be? In fact, using a Ca²⁺-sensitive fluorescent dye, irregular, spontaneous Ca²⁺ puffs along the radial arms/smooth spongione part of the organelle were observed (Ladenburger *et al.*, 2006). Ca²⁺ signals have been altered by UV-

activation of caged Ca^{2+} and Ca^{2+} -dependent biosynthetic pathways (organelle biogenesis) have been inhibited by silencing the *PtCRC-II/IP₃R*; this, together with the unambiguous identification of an *IP₃-BD* (with characteristics of a low affinity type of mammalian *IP₃R*), shown in Fig. 3A, led the authors (Ladenburger *et al.*, 2006) to conclude that these channels allow for some Ca^{2+} reflux from the organelle. However, the function may be dual. First, they may serve for fine-tuning of Ca^{2+} secretion (Ladenburger *et al.*, 2006) similar to systemic functions in kidney. Second, they may provide Ca^{2+} for local restructuring of the spongione by reversible fusion and fission processes by the numerous organelle-specific SNARE proteins (Schönemann *et al.*, 2013). This may enable the adjustment of organelle structure and function to the actual physiological requirements (Plattner, 2013).

3.2.2 Ryanodine receptor-like proteins—The *PtCRC-IV/RyR-LP* localized to the alveolar sacs (Ladenburger *et al.*, 2009) that had been previously identified (Stelly *et al.*, 1991) and further characterized (Lange *et al.*, 1995; Hardt and Plattner, 2000) as cortical Ca^{2+} -stores. Alveolar sacs are tightly attached to the cell membrane. By immunogold EM analysis these channels have been localized to the outer part of alveolar sacs facing the cell membrane (Ladenburger *et al.*, 2009). Aspects of functional characterization are contained in Fig. 3B, C. These were identified by activation with the secretagogue aminoethyl dextran (AED), caffeine, or with the ryanodine substitute 4-chloro-*meta*-cresol (4CmC) in conjunction with Ca^{2+} imaging, paralleled by gene silencing (Fig. 3C). Since gene silencing inhibited secretagogue-induced Ca^{2+} signals (Fig. 3C), as well as trichocyst exocytosis (Fig. 3B), these channels were concluded to transport Ca^{2+} . This function has been identified as the first step of signal transduction, which induces, as a second step, a superimposed Ca^{2+} -influx from the medium (store-operated Ca^{2+} -entry, SOCE). The activity of both components in concert had been envisioned previously by whole cell-patch electrophysiology (Erxleben and Plattner, 1994; Erxleben *et al.*, 1997), by fluorochrome analysis under selective conditions (Klauke *et al.*, 2000) and by elemental analysis at the EM level using energy-dispersive x-ray microanalysis (Hardt and Plattner, 2000). The situation clearly recalls that of the junction between the SR and the plasma membrane in striated muscle, although SOCE is much more widely distributed.

The simultaneous occurrence of *PtCRC-IV*- channels (RyR type) and additionally of *PtCRC-V-4* (*IP₃R* type) in alveolar sacs is also not without precedent. It also occurs in the ER of rat sensory neurons (Solovyova and Verkhatsky, 2003) and in mammalian skeletal muscle RyRs and *IP₃R*s cooperate to activate Ca^{2+} -signaling via the SR (Tjondrokoesoemo *et al.*, 2013). A similar cooperativity has been detected in atrial cells of the heart (Horn *et al.*, 2013).

There are several unexplored properties of *PtCRC*s including the occurrence of mixed features in some subfamilies. For instance, the molecular size of *PtCRC*s of the RyR type (e.g. *PtCRC-IV*) is unusually small. According to their amino acid sequence all *PtCRC*s are around 300 kDa in size (Ladenburger and Plattner, 2011) and thus resemble *IP₃R*s in metazoans. Functional implications of this peculiarity remain to be explored. This may be one of the criteria indicating the occurrence of a common ancestral form of *IP₃R*s and RyR-LPs in protozoa (Plattner and Verkhatsky, 2013), whereas in metazoa RyRs are much larger

than IP₃Rs (Taylor et al., 2009). Also unusual is the occurrence of two large loops in the *PtCRC-II-1* (IP₃R) molecule (Ladenburger *et al.*, 2006) (Fig. 3A).

Evaluation by very recently developed data based algorithms for the determination of transmembrane domains as specified (Ladenburger and Plattner, 2011), has indicated that *PtCRCs*, including RyR-LPs, possess 6 transmembrane domains (TMD) – previously a matter of debate. Thus, *PtRyR-LPs* differ from the most widely maintained assumption of only four TMDs. A more recent computational evaluation of the mammalian RyR came to the same conclusion (Ramachandran *et al.*, 2013). Domains for tetramerization are available also in the *PtCRC* molecules (Ladenburger and Plattner, 2011).

From their occurrence in protozoa, together with the occurrence of mixed type molecules with overlapping characteristics of IP₃Rs and RyRs, we assume that some of the *PtCRCs* may represent ancestral CRC types, close to such primeval forms. Also the amino acids composing the selectivity filter, i.e. Gly-Ile-Gly-Asp, are identical in both types of *PtCRCs* (Ladenburger and Plattner, 2011). This also occurs in other lower eukaryotes (Plattner and Verkhratsky, 2013) and, thus, is in contrast to the sequence Gly-Val-Gly-Asp in the IP₃R of metazoan cells (Boehning, 2010). In *Paramecium*, the latter sequence occurs in *PtCRC-I-1a*, -1b, and -1c, all with and IP₃-BD (Ladenburger and Plattner, 2011), though its IP₃-binding capacity has not been probed experimentally as yet. Otherwise the diversification of these channels in *Paramecium* during evolution is quite similar to the diversification of other molecules pertinent to vesicle trafficking, such as SNAREs (Plattner, 2010). This also includes more or less diversification as well as partial elimination by pseudogenization.

3.3 Pharmacology of IP₃R- and RyR-type CRCs and effect of injected cADPR and NAADP on cell function in *Paramecium*

Although the mechanism of AED as a secretagogue (Plattner and Hentschel, 2006) is not known in detail, some exogenous polyamines are known to induce a SOCE mechanism and exocytosis in various metazoan cells (Williams, 1997; Gamberucci *et al.*, 1998; Plattner and Klauke, 2001; Plattner and Hentschel, 2006). By contrast, caffeine and 4CmC are established activators of RyRs up to mammalian cells (Cheek and Barry, 1993; Westerblad *et al.*, 1998). The amino acids necessary for 4CmC binding (Fessenden *et al.*, 2006) are found in the *PtCRC-IV* molecule (Ladenburger *et al.*, 2009). Assays for Ca²⁺-dependent ryanodine binding to isolated alveolar sacs gave no results. This is not surprising considering the largely aberrant pharmacology of ciliates (Plattner *et al.*, 2009) and the absence of any evolutionary pressure for this plant toxin. Also the inhibitory effect of Li⁺ on phenomena related to IP₃ in *Paramecium* (Ladenburger *et al.*, 2006) are no stringent argument considering its pleiotropic effects, whereas the usefulness of compound U73122 as an inhibitor of phosphoinositide-specific phospholipase C in ciliates (Leondaritis *et al.*, 2011) has not yet been known at the time of those analyses.

It is well known that Ca²⁺ effects are strictly locally confined and rapidly counteracted by different mechanisms (see “Introduction”). This explains two observations: microinjection of an excess of Ca²⁺ into a *Paramecium* cell does not cause any exocytosis (Klauke and Plattner, 1997). Similarly a diffuse Ca²⁺-influx, as achieved by sudden increase of [Ca²⁺]_e, does not result in exocytosis (Erxleben *et al.*, 1997). Also injection of a likely activator of

RyRs in mammalian cells (Zalk *et al.*, 2007), cADPR, causes no trichocyst exocytosis, as is the case with injected NAADP, the putative activator of TPCs (Galione *et al.*, 2009; Galione *et al.*, 2010). The localization of such channels is not known for *Paramecium*. Since either compound changes contractile vacuole pulsations (Plattner *et al.*, 2012), organelles regulating Ca^{2+} homeostasis, one may assume the occurrence of the respective target molecules. However, the proper target may not be reached by the injected compounds, as described above for Ca^{2+} , and the effects achieved with the contractile vacuole may be due to Ca^{2+} activation from remote organelles. With *Paramecium* homogenates a $K_d \sim 3.5$ nM for NAADP binding was determined (Plattner *et al.*, 2012). Although a variety of acidic organelles, which could harbor TPCs, are known from *Paramecium* (Wassmer *et al.*, 2009) acidocalcisomes have not yet been identified. Also unknown is whether vacuoles containing Ca/Mg-phosphate crystals (Grover *et al.*, 1997) would be dynamic Ca^{2+} -stores or just waste disposal containers.

4. Intracellular calcium channels in parasitic protozoa

4.1 IP₃R in trypanosomatids

Ca^{2+} homeostasis in trypanosomes differs significantly from that in mammalian cells. There are no orthologs for receptor-operated or store-operated Ca^{2+} channels although the parasites possess orthologs for a putative voltage-sensitive Ca^{2+} channel that, in the case of *T. brucei*, localizes to the flagellum (Oberholzer *et al.*, 2011). No orthologs for $\text{Na}^+/\text{Ca}^{2+}$ exchangers are present in trypanosomes. In contrast to the plasma membrane Ca^{2+} -ATPase (PMCA) of higher eukaryotes, trypanosome PMCA apparently lacks a calmodulin-BD and is also localized intracellularly (Lu *et al.*, 1998; Luo *et al.*, 2004). The sarcoplasmic-endoplasmic reticulum Ca^{2+} -ATPase (SERCA) of trypanosomatids is insensitive to inhibitors of mammalian SERCA ATPases, such as thapsigargin (Docampo *et al.*, 1993; Vercesi *et al.*, 1993; Furuya *et al.*, 2001). In addition, trypanosomes possess an important acidic calcium store, the acidocalcisomes, which is rich in polyphosphate (Docampo *et al.*, 2005).

Recent results (Huang *et al.*, 2013) have indicated that the IP₃R of *T. brucei* is localized to acidocalcisomes rather than to the ER (Fig. 4A). The demonstration of the localization of the IP₃R in acidocalcisomes was obtained by tagging the C-terminus of *TbIP₃R* of procyclic trypomastigotes with an hemagglutinin tag using homologous recombination with the endogenous gene locus (Huang *et al.*, 2013). The *TbIP₃R* partially co-localized with antibodies against *T. brucei* acidocalcisome marker vacuolar proton pyrophosphatase (V-H⁺-PPase, or *TbVPP1*) (Fig. 4A). An additional punctate staining of *TbVPP1* that did not co-localize with *TbIP₃R* was detected and could correspond to trafficking vesicles. It was described before that adaptor protein-3 (AP-3) complex is involved in sorting proteins, like *TbVPP1*, to acidocalcisomes from the Golgi or from endosomes in both *L. major* (Besteiro *et al.*, 2008) and *T. brucei* (Huang *et al.*, 2013). No co-localization with *TbBiP*, an ER marker (Bangs *et al.*, 1993) with a clear reticular labeling (Fig. 4B), was detected, thus ruling out ER localization of the *TbIP₃R*. The acidocalcisome localization was confirmed using specific antibodies against *TbIP₃R* (unpublished results).

Proteomic analysis of contractile vacuole complex (Ulrich *et al.*, 2011) and acidocalcisome fractions (unpublished) of *T. cruzi* provided evidence of the presence of the *TcIP₃R* ortholog in these organelles. These results coincided with the punctate and vacuolar localization reported for *TcIP₃R* by other authors (Hashimoto *et al.*, 2013). These authors suggested an ER localization of *TcIP₃R* although no clear co-localization with *TbBiP* antibodies was observed.

The acidocalcisome localization of *TbIP₃R* led to test for Ca^{2+} release by IP_3 in permeabilized cells under conditions of optimal acidocalcisome function, i.e., in the presence of pyrophosphate (PP_i). Addition of PP_i is necessary to acidify acidocalcisomes by the action of *TbVP1*. This acidification allows Ca^{2+} uptake by the acidocalcisome Ca^{2+} -ATPase, which is $\text{Ca}^{2+}/\text{H}^+$ countertransporting (it transports Ca^{2+} in exchange for H^+). Under such conditions, IP_3 addition results in significant Ca^{2+} release (Huang *et al.*, 2013). Similar experiments were done with isolated acidocalcisomes resulting in significant Ca^{2+} release by IP_3 (Huang *et al.*, 2013). Previous attempts to show Ca^{2+} release by IP_3 had been unsuccessful (Moreno *et al.*, 1992a; Moreno *et al.*, 1992b), which was difficult to explain considering that the parasites do have a phosphoinositide phospholipase C (PI-PLC) (Furuya *et al.*, 2000; Okura *et al.*, 2005; Martins *et al.*, 2010), the enzyme that cleaves phosphatidylinositol 4,5-bisphosphate to generate diacylglycerol and IP_3 , and that IP_3 was detected in both *T. cruzi* (Moreno *et al.*, 1992b) and *T. brucei* (Moreno *et al.*, 1992a). When those experiments were done (1991–2) the presence of acidocalcisomes (Vercesi *et al.*, 1994; Docampo *et al.*, 1995) and an acidocalcisome vacuolar H^+ -PPase in trypanosomes were not known, as they were discovered much latter (Scott *et al.*, 1998; Rodrigues *et al.*, 1999). The reason for the lack of Ca^{2+} release in those experiments was that permeabilization results in dilution of substrates (ATP, PP_i) and alkalization of acidocalcisomes, as a result of lack of function of the proton pumps in the absence of ATP and PP_i .

The gene coding for *TbIP₃R* (Tb927.8.2770) shares 41% amino acid identity with *T. cruzi* IP_3R (TcCLB.509461.90), and orthologs are also present in several *Leishmania* spp. (Prole and Taylor, 2011). Structural analysis (ELM and TMHMM servers) predicted 5 transmembrane domains in the C-terminal region of these receptors. The ORFs of *T. brucei* and *T. cruzi* IP_3Rs predict 3099, and 3011 amino acid proteins, with apparent molecular weights of 343, and 337 kDa, respectively. Trypanosome IP_3Rs possess a series of conserved domains including putative suppressor domain-like (SD), ryanodine receptor IP_3R homology (RIH), and RIH-associated (RIAD) domains (Prole and Taylor, 2011). A motif for a Ca^{2+} -specific selectivity filter (GVGD) (Boehning *et al.*, 2001; Boehning, 2010) is present in the putative intraluminal loop between transmembrane domains at the C-terminal region (Huang *et al.*, 2013). This sequence resembles that of IP_3Rs in higher eukaryotes (Boehning, 2010) and also occurs in *PtCRC-I* type channels (with a putative IP_3 -BD), opposite to *PtCRC-II* channels (with an established IP_3 -BD (Ladenburger and Plattner, 2011; Plattner and Verkhratsky, 2013). Of the ten residues that have been proposed to form a basic pocket that binds IP_3 (Yoshikawa *et al.*, 1996; Bosanac *et al.*, 2002), 4 are conserved in *TbIP₃R*. Other features of trypanosomatid IP_3Rs have been described before (Prole and Taylor, 2011).

In addition to the studies on permeabilized trypanosomes (Huang *et al.*, 2013) functional analyses of *TbIP₃R* (Huang *et al.*, 2013) and *TcIP₃R* (Hashimoto *et al.*, 2013) were also done by stable transfection of the respective genes in a chicken B lymphocyte cell line (DT40, R23-11) in which the genes for all three vertebrate IP₃Rs have been stably ablated (DT40-3KO) (Miyakawa *et al.*, 1999). Both *TbIP₃R* and *TcIP₃R* localized to the ER of DT40-KO cells, and Ca²⁺ release by IP₃ was investigated using permeabilized cells (Huang *et al.*, 2013), microsomal vesicles, or intact cells stimulated by anti-B cell receptor monoclonal antibodies (Hashimoto *et al.*, 2013). Microsomal vesicles from DT40-KO cells expressing *TcIP₃R* also exhibited IP₃ binding activity (Hashimoto *et al.*, 2013). *TcIP₃R* was also expressed in HeLa cells, where it localized to the ER, and these permeabilized cells also exhibited Ca²⁺ release in the presence of IP₃ (Hashimoto *et al.*, 2013). *TbIP₃R* was found to be considerably less sensitive to IP₃ than the rat IP₃R1 (*RnIP₃R*1) transfected in DT40-3KO cells (Huang *et al.*, 2013). Ca²⁺ release by IP₃ was also investigated in live *T. brucei* procyclic trypomastigotes loaded with Fluo 4-AM with caged IP₃ (Huang *et al.*, 2013). In cells loaded with caged IP₃ there were rapid Ca²⁺ increases after UV flashes to release free IP₃ (Fig. 5), and these increases were considerably reduced when using trypanosomes in which the expression of *TbIP₃R* was downregulated by RNAi (Huang *et al.*, 2013).

To study the importance of *TbIP₃R* and *TcIP₃R* in the biology of trypanosomes, several strategies were used. Knockdown of *TbIP₃R* expression was done by induction of RNAi and resulted in growth defects in both bloodstream and procyclic trypomastigotes (Huang *et al.*, 2013). Knockdown of the expression of *TbIP₃R* in procyclic forms by RNAi reduced the ability of IP₃ to release Ca²⁺ from permeabilized cells and reduced the virulence of bloodstream forms *in vivo* (Huang *et al.*, 2013). Knockdown of *TcIP₃R* was done by single-knockout in epimastigotes of the Tulahuen strain. Attempts to obtain *null* mutants in this or in the Esmeraldo strain were unsuccessful, suggesting the essentiality of this gene (Hashimoto *et al.*, 2013). *TcIP₃R* knockdown resulted in deficient growth of epimastigotes, deficient metacyclogenesis (transformation of epimastigotes into metacyclic trypomastigotes), deficient host cell invasion by trypomastigotes associated with reduced Ca²⁺ release upon their attachment to the host cells, deficient replication of amastigotes, increased transformation of amastigotes into trypomastigotes, and defects in virulence *in vivo* (Hashimoto *et al.*, 2013). Overexpression of *TcIP₃R* also resulted in deficient growth of epimastigotes and amastigotes, and deficient metacyclogenesis, suggesting that an appropriate level of this receptor is necessary for these processes (Hashimoto *et al.*, 2013). In contrast, overexpression of *TcIP₃R* resulted in increased host cell invasion by trypomastigotes associated with increased Ca²⁺ release upon their attachment to host cells, and decreased transformation of amastigotes into trypomastigotes, with no changes in virulence *in vivo* except for an early appearance of parasitemia (Hashimoto *et al.*, 2013).

In conclusion, these works (Hashimoto *et al.*, 2013; Huang *et al.*, 2013) clearly established the presence of a functional IP₃ receptor in *T. brucei* and *T. cruzi*, and together with previous reports (Docampo and Pignataro, 1991; Moreno *et al.*, 1992a; Moreno *et al.*, 1992b; Furuya *et al.*, 2000; Okura *et al.*, 2005; de Paulo Martins *et al.*, 2010; Martins *et al.*, 2010), the function of a complete IP₃/diacylglycerol pathway in trypanosomes.

Although acidocalcisomes were initially described almost 20 years ago in *T. brucei* (Vercesi *et al.*, 1994), the mechanism for Ca^{2+} release from these organelles was unknown until now. The localization of the IP_3 receptor in acidocalcisomes (Huang *et al.*, 2013) provides the long sought mechanism for Ca^{2+} release from these organelles.

4.2 Intracellular Ca^{2+} channels in *Toxoplasma gondii*

Measurements of intracellular Ca^{2+} levels have been done in *T. gondii* extracellular tachyzoites using the Ca^{2+} dye Fura 2-AM (Fura 2/acetomethoxy) and values of 60–100 nM were obtained (Moreno and Zhong, 1996). The ER and acidocalcisomes were identified as the largest Ca^{2+} stores. A SERCA-type Ca^{2+} -ATPase, which is present in the ER (Nagamune *et al.*, 2007) and inhibited by thapsigargin (Moreno and Zhong, 1996), is the main Ca^{2+} uptake mechanism in this compartment. In *T. gondii*, the presence of intracellular Ca^{2+} stores responsive to IP_3 , ryanodine (Lovett *et al.*, 2002) and cADPR (Chini *et al.*, 2005; Nagamune *et al.*, 2008), have been described but there is no genetic evidence for the presence of IP_3 or ryanodine gated channels (Nagamune and Sibley, 2006; Plattner and Verkhratsky, 2013). This is despite the evidence for the presence of enzymes involved in the generation of some of these second messengers such as a phosphoinositide phospholipase C (Fang *et al.*, 2006) and cADPR cyclase and hydrolase activities (Chini *et al.*, 2005). Acidocalcisomes of *T. gondii* possess a PMCA-type Ca^{2+} -ATPase (TgA1) for Ca^{2+} uptake (Luo *et al.*, 2001; Rohloff *et al.*, 2011), but their mechanism of Ca^{2+} release is unknown. Evidence for a $\text{Ca}^{2+}/\text{H}^+$ exchanger in these acidocalcisomes has also been reported (Rohloff *et al.* 2011). The recently described plant-like vacuole (PLV) (Miranda *et al.*, 2010) was also found to be rich in Ca^{2+} and also possesses the PMCA-type Ca^{2+} -ATPase (TgA1) for Ca^{2+} uptake. A $\text{Ca}^{2+}/\text{H}^+$ exchanger is also present in the PLV (Miranda *et al.*, 2010), but the mechanism of Ca^{2+} release is also unknown. There is no genetic evidence for the presence of a mitochondrial calcium uniporter (MCU) in any Apicomplexan parasite (Bick *et al.*, 2012) and the role of the mitochondria in Ca^{2+} regulation is not clear. A $\text{Ca}^{2+}/\text{H}^+$ antiporter is apparently present in the mitochondria of *T. gondii* (Guttery *et al.*, 2013). Ca^{2+} entry into tachyzoites is regulated (Pace *et al.*, 2013). Other unexplored sources of Ca^{2+} in *T. gondii* are Golgi complex, apicoplast, inner membrane complex (IMC), and secretory organelles.

T. gondii is unique among parasitic protozoa in possessing an ortholog to two pore channels (TPCs), for which there are no orthologs in other Apicomplexans (Prole and Taylor, 2011). This TPC has substantial similarity to mammalian TPCs in the pore region responsible for ion conduction, suggesting that it may act as Ca^{2+} -permeable channel (Prole and Taylor, 2011). However, the localization and function of this channel have not been reported. Mammalian TPCs can be Ca^{2+} channels gated by NAADP, and localized in lysosome-like compartments (Brailoiu *et al.*, 2009; Calcraft *et al.*, 2009; Zong *et al.*, 2009). Their function as Ca^{2+} channels and their stimulation by NAADP have been disputed (Wang *et al.*, 2012; Cang *et al.*, 2013), although the reason for this discrepancy is apparently that tagging the N-terminal region of the channel abolishes its sensitivity to NAADP (Churamani *et al.*, 2013). Fig. 6 shows a schematic representation of Ca^{2+} distribution in tachyzoites.

4.3 Mitochondrial calcium uniporter

Mammalian mitochondria have been shown to have a central role in intracellular Ca^{2+} homeostasis, and it is well established that intramitochondrial Ca^{2+} concentration can rapidly reach tens or hundreds micromolar values upon cytosolic Ca^{2+} rises of a few micromolar (Rizzuto *et al.*, 1993; Montero *et al.*, 2000). This is because mitochondria are exposed to microdomains of high Ca^{2+} concentration in proximity to sites of Ca^{2+} release at the ER, or to Ca^{2+} channels at the plasma membrane (Rizzuto *et al.*, 1993; Rizzuto *et al.*, 1998; Csordas *et al.*, 1999; Montero *et al.*, 2000; Csordas *et al.*, 2010; Giacomello *et al.*, 2010). This Ca^{2+} uptake is important for shaping the amplitude and spatio-temporal patterns of cytosolic Ca^{2+} increases (Boitier *et al.*, 1999; Hajnoczky *et al.*, 1999; Tinel *et al.*, 1999) and for regulating the activity of three intramitochondrial dehydrogenases that result in increased ATP generation (Denton and McCormack, 1990; McCormack *et al.*, 1990; Hajnoczky *et al.*, 1995; Jouaville *et al.*, 1999; Voronina *et al.*, 2010), as well as in stimulating the activity of the mitochondrial ATP synthase (Balaban, 2009). Ca^{2+} also regulates mitochondrial transporters in the inner membrane (Lasorsa *et al.*, 2003; Satrustegui *et al.*, 2007). Excessive Ca^{2+} uptake, however, favors the formation of the “permeability transition pore” leading to the release of pro-apoptotic factors in the cytosol and cell death (Kroemer *et al.*, 2007).

The ability of mitochondria to take up Ca^{2+} was discovered more than 50 years ago (De Luca and Engstrom, 1961; Vasington and Murphy, 1962) and the channel biophysical properties were well characterized in a patch clamp study of mitoplasts (mitochondria devoid of the outer mitochondrial membrane) (Kirichok *et al.*, 2004). However, the molecular nature of the uniporter remained unknown for decades.

For many years after the discovery of the MCU in mammalian mitochondria, it was thought that less complex life forms such as plants, insects and other invertebrates, or unicellular organisms such as yeast, lacked a specific mitochondrial calcium uptake pathway (Carafoli and Lehninger, 1971). However, in 1989 it was reported (Docampo and Vercesi, 1989a, b) that epimastigotes of *T. cruzi* possesses a MCU with characteristics similar to those described in mammalian mitochondria: electrogenic transport, sensitivity to ruthenium red, and low affinity for the cation. The evidence of the presence of a MCU in trypanosomes but its absence in yeast was the key to the discovery of the molecular identity of MCU. An elegant study by Perocchi and coworkers (Perocchi *et al.*, 2010) first identified a gene called *mitochondrial calcium uptake 1* or *MICU1* as encoding a potential regulator of the uniporter. The study was based on the observation that the Ca^{2+} uniporter was detected in *T. cruzi* (Docampo and Vercesi, 1989a, b) and *Leishmania donovani* (Vercesi and Docampo, 1992) yet not measurable in the yeast *Saccharomyces cerevisiae* (Balcavage *et al.*, 1973). From a library of 1,000 mouse mitochondrial proteins, 18 candidate genes were identified that have homologues in vertebrates and trypanosomes but not in yeast (Perocchi *et al.*, 2010). Using short hairpin (sh)RNA silencing of 13 selected genes in a commercially available HeLa cell line that stably expresses a mitochondria-targeted aequorin (mt-AEQ) as a reporter of Ca^{2+} uptake, they identified MICU1 as an important component in Ca^{2+} uptake. Based on the finding of Perocchi *et al.* (Perocchi *et al.*, 2010), De Stefani *et al.* (De Stefani *et al.*, 2011) and Baughman *et al.* (Baughman *et al.*, 2011) used a similar strategy of comparing between

mitochondrial genomes of trypanosomes and yeast and performing RNAi experiments of the identified genes and found a gene encoding a protein with all the characteristics of the mitochondrial calcium uniporter (MCU).

The MCU was also found in other trypanosomatids including *T. brucei* (Docampo and Lukes, 2012). The finding of a MCU uniporter in the bloodstream stage of *T. brucei* (Vercesi *et al.*, 1992) was surprising because these stages lack a respiratory chain. However, mitochondrial Ca^{2+} uptake can also be energized by ATP in the absence of respiration, in which case it is inhibited by oligomycin, and not by inhibitors of the respiratory chain (Lehninger *et al.*, 1963). This phenomenon also occurs in bloodstream trypomastigotes: the mitochondrial membrane potential is dependent on hydrolysis of ATP by the ATP synthase which acts as an ATPase (Nolan and Voorheis, 1992; Vercesi *et al.*, 1992; Schnauffer *et al.*, 2005; Brown *et al.*, 2006) allowing for Ca^{2+} to still be electrophoretically transported by the MCU (Vercesi *et al.*, 1992). Ca^{2+} uptake by bloodstream forms of *T. brucei* has three characteristics: 1) it occurs until the ambient free Ca^{2+} concentration is lowered to 0.6–0.7 μM ; 2) it is inhibited by oligomycin; and 3) it is associated with the depolarization of the inner membrane energized by ATP. These results indicate that Ca^{2+} uptake is mediated by the ATPase-dependent energization of the inner mitochondrial membrane (Vercesi *et al.*, 1992).

Although present in trypanosomatids and other protozoa, such as *Tetrahymena thermophila*, and *Naegleria gruberi*, the MCU is absent in Apicomplexan parasites such as *T. gondii* or malaria parasites (Bick *et al.*, 2012).

The roles of mitochondrial Ca^{2+} in trypanosomes are apparently more limited than in mammalian cells. None of the dehydrogenases stimulated by Ca^{2+} in vertebrates have been studied in detail in trypanosomatids and there is no evidence of their stimulation by Ca^{2+} (Docampo and Lukes, 2012). Experiments using aequorin targeted to the mitochondria of *T. brucei* procyclic trypomastigotes revealed that intramitochondrial Ca^{2+} concentrations can reach values much higher than cytosolic Ca^{2+} rises when Ca^{2+} influx through the plasma membrane or Ca^{2+} release from acidocalcisomes are stimulated (Xiong *et al.*, 1997). These results suggest a very close proximity of these organelles and the presence of microdomains of high Ca^{2+} concentration in the vicinity of the plasma membrane and acidocalcisomes (Xiong *et al.*, 1997) and are in agreement with the presence of an IP_3R in acidocalcisomes (Huang *et al.*, 2013). Because the ER-type Ca^{2+} -ATPase (SERCA) of *T. brucei* is insensitive to thapsigargin, a microdomain of high Ca^{2+} concentration between the ER and the mitochondria could not be established in those studies (Xiong *et al.*, 1997). However, these results suggest that one of the main functions of the MCU in procyclic trypomastigotes would be to shape the amplitude and spatio-temporal patterns of cytosolic Ca^{2+} increases. No similar studies have been done with bloodstream trypomastigotes. Fig. 7 shows a scheme of the close contact that would exist between acidocalcisomes and the unique mitochondrion of trypanosomes.

Mitochondrial Ca^{2+} could also be a contributor to programmed cell death, or apoptosis-like death, in trypanosomatids. Trypanosomatids lack some of the key regulatory or effector molecules involved in apoptosis in mammalian cells, such as the tumor necrosis factor

(TNF)-related family of receptors, Bcl-2 family members, and caspases (Ridgley *et al.*, 1999; Smirlis *et al.*, 2010; Kaczanowski *et al.*, 2011). Mitochondrial Ca^{2+} overload with changes in mitochondrial membrane potential, reactive oxygen species (ROS) generation and release of cytochrome *c* have been observed upon different triggers of cell death in some trypanosomatids (Smirlis and Soteriadou, 2011). In *T. brucei* procyclic trypomastigotes, the production of ROS impairs mitochondrial Ca^{2+} transport, leading to its accumulation in the nucleus, and causing cell death (Ridgley *et al.*, 1999).

In summary, mitochondrial Ca^{2+} uptake in trypanosomes appears to have a role in shaping the amplitude of cytosolic Ca^{2+} increases after influx through the plasma membrane or release from acidocalcisomes, and in apoptosis-like death, but it is not known whether it has a role in the regulation of ATP production.

4.4 Pharmacology of CRC s in parasitic protozoa

TbIP₃R does not respond to 10 μM NAADP or 1 μM cADPR when expressed in DT40-3KO cells (Huang *et al.*, 2013). cADPR was also ineffective in producing Ca^{2+} release in permeabilized HeLa cells transfected with *TcIP₃R* (Hashimoto *et al.*, 2013). In contrast, the IP_3 agonist adenophostin A (0.5 μM) showed a Ca^{2+} -release that was comparable to that of 10 μM IP_3 in the *TbIP₃R*-expressing DT40-3KO cells or in permeabilized *T. brucei* procyclic forms (Huang *et al.*, 2013).

There is pharmacological evidence for the presence of channels responsive to IP_3 in *T. gondii* and malaria parasites (Passos and Garcia, 1998; Lovett and Sibley, 2003; Alves *et al.*, 2011) although there are no gene orthologs to the mammalian IP_3R or RyR in any of the Apicomplexan genomes (Nagamune and Sibley, 2006; Plattner *et al.*, 2012). There is an ER Ca^{2+} pool, sensitive to thapsigargin, and the Ca^{2+} -ATPase involved in pumping Ca^{2+} from the cytosol into the ER has been characterized (Nagamune *et al.*, 2007).

4.5 Calcium signaling and function in parasitic protozoa

Although Ca^{2+} signaling appears to be important for several functions in *T. cruzi* such as host cell invasion (Moreno *et al.*, 1994), multiplication and differentiation (Lammel *et al.*, 1996), osmoregulation (Rohloff *et al.*, 2003), and programmed cell death (Irigoin *et al.*, 2009), there is much less information on the role of Ca^{2+} in *T. brucei*. Based on the use of Ca^{2+} ionophores, roles for Ca^{2+} in the release of the bloodstream stage surface coat (Bowles and Voorheis, 1982), and in the maintenance of the cytoskeleton (Selzer *et al.*, 1991) have been proposed. Changes in cytosolic Ca^{2+} levels have also been reported during differentiation from bloodstream to procyclic stages of *T. brucei* (Stojdl and Clarke, 1996). Results obtained with *TbIP₃R* (Huang *et al.*, 2013) and *TcIP₃R* (Hashimoto *et al.*, 2013) knockdowns indicate that Ca^{2+} signaling through the trypanosome IP_3Rs has roles in growth *in vitro* and *in vivo*, as well as in cell differentiation. The localization of the *TbIP₃R* in acidocalcisomes (Huang *et al.*, 2013) also supports a role for acidocalcisomes in Ca^{2+} signaling.

Several studies have looked into the role of Ca^{2+} signaling during the lytic cycle of *T. gondii*. Using Ca^{2+} ionophores, Ca^{2+} -chelating agents and ethanol, a link between Ca^{2+} and

conoid extrusion was demonstrated (Mondragon *et al.*, 1994; Mondragon and Frixione, 1996; Del Carmen *et al.*, 2009), although no direct parallel Ca^{2+} measurements were reported. The effect of Ca^{2+} on gliding motility was studied on trypsin-permeabilized tachyzoites (Mondragon and Frixione, 1996) or by measuring Ca^{2+} oscillations in extracellular tachyzoites loaded with the Ca^{2+} dye Fluo 4-AM and studying their correlation with gliding motility (Lovett and Sibley, 2003; Wetzel *et al.*, 2004). The role of Ca^{2+} in microneme secretion was studied in extracellular tachyzoites following the effects of Ca^{2+} ionophores or chelators on protein secretion as evaluated by western blot analyses (Carruthers *et al.*, 1999). A role for Ca^{2+} signaling in invasion was postulated on the basis of the analysis of changes in Ca^{2+} occurring in parasites loaded with Fura 2-AM upon their attachment to host cells (Vieira and Moreno, 2000) or by detecting the cessation of Ca^{2+} oscillations in extracellular tachyzoites loaded with Fluo 4-AM (Lovett and Sibley, 2003). These and more recent studies have provided indirect evidence that Ca^{2+} signaling is part of the pathways that result in the stimulation of conoid extrusion, gliding motility, microneme secretion, and invasion. These traits, which are important steps of the lytic cycle of the parasite are enhanced by extracellular Ca^{2+} (Pace *et al.*, 2013). Ca^{2+} signaling was also proposed to be part of the pathway leading to parasite egress from the host cells based on the effect of Ca^{2+} ionophores, which stimulate egress, although it was never demonstrated that $[\text{Ca}^{2+}]_i$ increases in tachyzoites before egress (Endo *et al.*, 1982; Garrison *et al.*, 2012). In summary, most of the studies on Ca^{2+} signaling were done with extracellular tachyzoites (conoid extrusion, gliding motility, microneme secretion, invasion) measuring indirectly the involvement of Ca^{2+} (using Ca^{2+} chelators or ionophores) or detecting Ca^{2+} changes with fluorescent dyes independently of the phenomena examined.

5. Conclusions and Perspectives

On the one hand there are many similarities in Ca^{2+} signaling in the protozoa discussed here, but on the other hand there also occur considerable differences. Differences concern the drug sensitivity of the SERCA, the absence or the occurrence of a calmodulin-BD in the PMCA (typical for higher eukaryotes) and the occasional occurrence of this pump in Ca^{2+} -storage organelles. Interestingly, a SOCE mechanism has been verified with some protozoa analyzed so far, based in part on the activation of RyR-LPs (Ladenburger *et al.*, 2009). There is, however, no evidence of the presence of Orai or Stim orthologs in either *Paramecium* (Ladenburger and Plattner, 2011) or parasitic protozoa (EuPathDB.com). Acidocalcisomes are present in some (trypanosomatids, Apicomplexan) but not in other parasitic protozoa (like *Giardia*, *Trichomonas*, *Entamoeba*) (Docampo *et al.*, 2005) while it is not yet known if they are present in *Paramecium* (Plattner *et al.*, 2012). IP_3Rs orthologs are present in trypanosomatids (Huang *et al.*, 2013; Hashimoto *et al.*, 2013) and *Paramecium* (Plattner and Verkhatsky, 2013) but unknown in Apicomplexa.

The range of CRC molecules, type $\text{IP}_3\text{Rs/RyR}$ in *Paramecium* and of IP_3Rs in *Monosiga brevicollis*, and trypanosomatids indicates that protozoa already have evolved this Ca^{2+} -signaling pathway. Except in ciliates, only IP_3Rs have been unambiguously identified; none have been identified in Apicomplexa – in striking contrast to their response to IP_3 (which, thus, still awaits elucidation). In ciliates some IP_3Rs and RyR-LPs display mixed features of the two. Concomitantly, in *Paramecium* some of the CRCs can be clearly attributed to either

IP₃Rs and to RyR-LPs, all CRCs are differentially localized and – as far as analyzed – can be differentially activated. In *Paramecium* and other protozoa the selectivity filter in both types of CRCs, IP₃R- and RyR-like channels (or their putative homologues) is identical for most paralogues (Landenburger and Plattner, 2011; Plattner and Verkhatsky, 2013). By contrast the parasitic flagellates have the same amino acid sequence in the selectivity filter of their IP₃R (Huang *et al.*, 2013) as occurring in metazoa (Boehning, 2010). Evidently selective enrichment of Ca²⁺ in the stores can suffice to guarantee selective release of Ca²⁺ as a second messenger. A RyR-type CRC seems to be absent from Apicomplexa; none has been identified in choanoflagellates and in the parasitic flagellates discussed here, whereas their occurrence in other protists has to be left open for the time being (Landenburger and Plattner, 2011; Plattner and Verkhatsky, 2013). Also the number of transmembrane domains in RyR-LPs, five or six being proposed depending on the database analysis method applied, requires experimental verification.

The presence of the mitochondrial calcium uniporter in a number of protozoa indicates that mitochondrial regulation of Ca²⁺ signaling is also an early appearance in evolution. There is only circumstantial or sequence evidence for the occurrence of TPCs, TRPCs and of intracellular mechano-sensitive Ca²⁺-channels in *Paramecium* and in parasitic protozoa. This remains a vast field for further explorations.

Acknowledgments

Work by R.D. was supported by U.S. National Institutes of Health (NIH), grant AI077538. Work by S.N.J.M. was supported by U.S. NIH, grant AI096836. Work by H.P. has been supported by Deutsche Forschungsgemeinschaft, grant PL 78/21.

References

- Allan CY, Fisher PR. In vivo measurements of cytosolic calcium in *Dictyostelium discoideum*. *Methods Mol Biol.* 2009; 571:291–308. [PubMed: 19763975]
- Allen RD, Fok AK. Membrane trafficking and processing in *Paramecium*. *Int Rev Cytol.* 2000; 198:277–318. [PubMed: 10804465]
- Alves E, Bartlett PJ, Garcia CR, Thomas AP. Melatonin and IP₃-induced Ca²⁺ release from intracellular stores in the malaria parasite *Plasmodium falciparum* within infected red blood cells. *J Biol Chem.* 2011; 286:5905–5912. [PubMed: 21149448]
- Aury JM, Jaillon O, Duret L, Noel B, Jubin C, Porcel BM, Segurens B, Daubin V, Anthouard V, Aiach N, Arnaiz O, Billaut A, Beisson J, Blanc I, Bouhouche K, Camara F, Dharcourt S, Guigo R, Gogendeau D, Katinka M, Keller AM, Kissmehl R, Klotz C, Koll F, Le Mouel A, Lepere G, Malinsky S, Nowacki M, Nowak JK, Plattner H, Poulain J, Ruiz F, Serrano V, Zagulski M, Dessen P, Betermier M, Weissenbach J, Scarpelli C, Schachter V, Sperling L, Meyer E, Cohen J, Wincker P. Global trends of whole-genome duplications revealed by the ciliate *Paramecium tetraurelia*. *Nature.* 2006; 444:171–178. [PubMed: 17086204]
- Balaban RS. The role of Ca²⁺ signaling in the coordination of mitochondrial ATP production with cardiac work. *Biochim Biophys Acta.* 2009; 1787:1334–1341. [PubMed: 19481532]
- Balcavage WX, Lloyd JL, Mattoon JR, Ohnishi T, Scarpa A. Cation movements and respiratory response in yeast mitochondria treated with high Ca²⁺ concentrations. *Biochim Biophys Acta.* 1973; 305:41–51. [PubMed: 4578276]
- Bangs JD, Uyetake L, Brickman MJ, Balber AE, Boothroyd JC. Molecular cloning and cellular localization of a BiP homologue in *Trypanosoma brucei*. Divergent ER retention signals in a lower eukaryote. *J Cell Sci.* 1993; 105:1101–1113. [PubMed: 8227199]

- Baughman JM, Perocchi F, Girgis HS, Plovanich M, Belcher-Timme CA, Sancak Y, Bao XR, Strittmatter L, Goldberger O, Bogorad RL, Kotliansky V, Mootha VK. Integrative genomics identifies MCU as an essential component of the mitochondrial calcium uniporter. *Nature*. 2011; 476:341–345. [PubMed: 21685886]
- Berridge MJ, Bootman MD, Roderick HL. Calcium signalling: dynamics, homeostasis and remodelling. *Nat Rev Mol Cell Biol*. 2003; 4:517–529. [PubMed: 12838335]
- Berridge MJ, Lipp P, Bootman MD. The versatility and universality of calcium signalling. *Nat Rev Mol Cell Biol*. 2000; 1:11–21. [PubMed: 11413485]
- Besteiro S, Tonn D, Tetley L, Coombs GH, Mottram JC. The AP3 adaptor is involved in the transport of membrane proteins to acidocalcisomes of *Leishmania*. *J Cell Sci*. 2008; 121:561–570. [PubMed: 18252798]
- Bezprozvanny I. The inositol 1,4,5-trisphosphate receptors. *Cell Calcium*. 2005; 38:261–272. [PubMed: 16102823]
- Bick AG, Calvo SE, Mootha VK. Evolutionary diversity of the mitochondrial calcium uniporter. *Science*. 2012; 336:886. [PubMed: 22605770]
- Boehning D, Mak DO, Foskett JK, Joseph SK. Molecular determinants of ion permeation and selectivity in inositol 1,4,5-trisphosphate receptor Ca^{2+} channels. *J Biol Chem*. 2001; 276:13509–13512. [PubMed: 11278266]
- Boehning DF. Molecular architecture of the inositol 1,4,5-trisphosphate receptor pore. *Curr Top Membr*. 2010; 66C:191–207. [PubMed: 22353481]
- Boitier E, Rea R, Duchen MR. Mitochondria exert a negative feedback on the propagation of intracellular Ca^{2+} waves in rat cortical astrocytes. *J Cell Biol*. 1999; 145:795–808. [PubMed: 10330407]
- Bootman MD, Berridge MJ. The elemental principles of calcium signaling. *Cell*. 1995; 83:675–678. [PubMed: 8521483]
- Bosanc I, Alattia JR, Mal TK, Chan J, Talarico S, Tong FK, Tong KI, Yoshikawa F, Furuichi T, Iwai M, Michikawa T, Mikoshiba K, Ikura M. Structure of the inositol 1,4,5-trisphosphate receptor binding core in complex with its ligand. *Nature*. 2002; 420:696–700. [PubMed: 12442173]
- Bowles DJ, Voorheis HP. Release of the surface coat from the plasma membrane of intact bloodstream forms of *Trypanosoma brucei* requires Ca^{2+} FEBS Lett. 1982; 139:17–21. [PubMed: 6804262]
- Brailoiu E, Churamani D, Cai X, Schrlau MG, Brailoiu GC, Gao X, Hooper R, Boulware MJ, Dun NJ, Marchant JS, Patel S. Essential requirement for two-pore channel 1 in NAADP-mediated calcium signaling. *J Cell Biol*. 2009; 186:201–209. [PubMed: 19620632]
- Brown SV, Hosking P, Li J, Williams N. ATP synthase is responsible for maintaining mitochondrial membrane potential in bloodstream form *Trypanosoma brucei*. *Eukaryot Cell*. 2006; 5:45–53. [PubMed: 16400167]
- Cai X. Unicellular Ca^{2+} signaling ‘toolkit’ at the origin of metazoa. *Mol Biol Evol*. 2008; 25:1357–1361. [PubMed: 18385221]
- Calcraft PJ, Ruas M, Pan Z, Cheng X, Arredouani A, Hao X, Tang J, Rietdorf K, Teboul L, Chuang KT, Lin P, Xiao R, Wang C, Zhu Y, Lin Y, Wyatt CN, Parrington J, Ma J, Evans AM, Galione A, Zhu MX. NAADP mobilizes calcium from acidic organelles through two-pore channels. *Nature*. 2009; 459:596–600. [PubMed: 19387438]
- Cang C, Zhou Y, Navarro B, Seo YJ, Aranda K, Shi L, Battaglia-Hsu S, Nissim I, Clapham DE, Ren D. mTOR regulates lysosomal ATP-sensitive two-pore Na^{+} channels to adapt to metabolic state. *Cell*. 2013; 152:778–790. [PubMed: 23394946]
- Carafoli E, Lehninger AL. A survey of the interaction of calcium ions with mitochondria from different tissues and species. *Biochem J*. 1971; 122:681–690. [PubMed: 5129264]
- Cardenas C, Miller RA, Smith I, Bui T, Molgo J, Muller M, Vais H, Cheung KH, Yang J, Parker I, Thompson CB, Birnbaum MJ, Hallows KR, Foskett JK. Essential regulation of cell bioenergetics by constitutive InsP_3 receptor Ca^{2+} transfer to mitochondria. *Cell*. 2010; 142:270–283. [PubMed: 20655468]
- Carruthers VB, Moreno SN, Sibley LD. Ethanol and acetaldehyde elevate intracellular $[\text{Ca}^{2+}]$ and stimulate microneme discharge in *Toxoplasma gondii*. *Biochem J*. 1999; 342:379–386. [PubMed: 10455025]

- Case RM, Eisner D, Gurney A, Jones O, Muallem S, Verkhatsky A. Evolution of calcium homeostasis: from birth of the first cell to an omnipresent signalling system. *Cell Calcium*. 2007; 42:345–350. [PubMed: 17574670]
- Cheek TR, Barry VA. Stimulus-secretion coupling in excitable cells: a central role for calcium. *J Exp Biol*. 1993; 184:183–196. [PubMed: 8270855]
- Chini EN, Nagamune K, Wetzel DM, Sibley LD. Evidence that the cADPR signalling pathway controls calcium-mediated microneme secretion in *Toxoplasma gondii*. *Biochem J*. 2005; 389:269–277. [PubMed: 15773818]
- Christensen KA, Myers JT, Swanson JA. pH-dependent regulation of lysosomal calcium in macrophages. *J Cell Sci*. 2002; 115:599–607. [PubMed: 11861766]
- Churamani D, Hooper R, Rahman T, Brailoiu E, Patel S. The N-terminal region of two-pore channel 1 regulates trafficking and activation by NAADP. *Biochem J*. 2013; 453:147–151. [PubMed: 23634879]
- Clapham DE. Calcium signaling. *Cell*. 2007; 131:1047–1058. [PubMed: 18083096]
- Csordas G, Thomas AP, Hajnoczky G. Quasi-synaptic calcium signal transmission between endoplasmic reticulum and mitochondria. *EMBO J*. 1999; 18:96–108. [PubMed: 9878054]
- Csordas G, Varnai P, Golenar T, Roy S, Purkins G, Schneider TG, Balla T, Hajnoczky G. Imaging interorganelle contacts and local calcium dynamics at the ER-mitochondrial interface. *Mol Cell*. 2010; 39:121–132. [PubMed: 20603080]
- De Luca HF, Engstrom GW. Ca^{2+} uptake by rat kidney mitochondria. *Proc Natl Acad Sci USA*. 1961; 47:1744–1750. [PubMed: 13885269]
- de Paulo Martins V, Okura M, Maric D, Engman DM, Vieira M, Docampo R, Moreno SN. Acylation-dependent export of *Trypanosoma cruzi* phosphoinositide-specific phospholipase C to the outer surface of amastigotes. *J Biol Chem*. 2010; 285:30906–30917. [PubMed: 20647312]
- De Stefani D, Raffaello A, Teardo E, Szabo I, Rizzuto R. A forty-kilodalton protein of the inner membrane is the mitochondrial calcium uniporter. *Nature*. 2011; 476:336–340. [PubMed: 21685888]
- Del Carmen MG, Mondragon M, Gonzalez S, Mondragon R. Induction and regulation of conoid extrusion in *Toxoplasma gondii*. *Cell Microbiol*. 2009; 11:967–982. [PubMed: 19416276]
- Denton RM, McCormack JG. Ca^{2+} as a second messenger within mitochondria of the heart and other tissues. *Annu Rev Physiol*. 1990; 52:451–466. [PubMed: 2184763]
- Docampo R, de Souza W, Miranda K, Rohloff P, Moreno SN. Acidocalcisomes - conserved from bacteria to man. *Nat Rev Microbiol*. 2005; 3:251–261. [PubMed: 15738951]
- Docampo R, Lukes J. Trypanosomes and the solution to a 50-year mitochondrial calcium mystery. *Trends Parasitol*. 2012; 28:31–37. [PubMed: 22088944]
- Docampo R, Moreno SN, Vercesi AE. Effect of thapsigargin on calcium homeostasis in *Trypanosoma cruzi* trypomastigotes and epimastigotes. *Mol Biochem Parasitol*. 1993; 59:305–313. [PubMed: 8341327]
- Docampo R, Pignataro OP. The inositol phosphate/diacylglycerol signalling pathway in *Trypanosoma cruzi*. *Biochem J*. 1991; 275:407–411. [PubMed: 2025225]
- Docampo R, Scott DA, Vercesi AE, Moreno SN. Intracellular Ca^{2+} storage in acidocalcisomes of *Trypanosoma cruzi*. *Biochem J*. 1995; 310:1005–1012. [PubMed: 7575396]
- Docampo R, Vercesi AE. Ca^{2+} transport by coupled *Trypanosoma cruzi* mitochondria in situ. *J Biol Chem*. 1989a; 264:108–111. [PubMed: 2491844]
- Docampo R, Vercesi AE. Characteristics of Ca^{2+} transport by *Trypanosoma cruzi* mitochondria in situ. *Arch Biochem Biophys*. 1989b; 272:122–129. [PubMed: 2500059]
- Douzery EJ, Snell EA, Baptiste E, Delsuc F, Philippe H. The timing of eukaryotic evolution: does a relaxed molecular clock reconcile proteins and fossils? *Proc Natl Acad Sci U S A*. 2004; 101:15386–15391. [PubMed: 15494441]
- Endo T, Sethi KK, Piekarski G. *Toxoplasma gondii*: calcium ionophore A23187-mediated exit of trophozoites from infected murine macrophages. *Exp Parasitol*. 1982; 53:179–188. [PubMed: 6800836]

- Erxleben C, Klauke N, Flotenmeyer M, Blanchard MP, Braun C, Plattner H. Microdomain Ca^{2+} activation during exocytosis in *Paramecium* cells. Superposition of local subplasmalemmal calcium store activation by local Ca^{2+} influx. *J Cell Biol.* 1997; 136:597–607. [PubMed: 9024690]
- Erxleben C, Plattner H. Ca^{2+} release from subplasmalemmal stores as a primary event during exocytosis in *Paramecium* cells. *J Cell Biol.* 1994; 127:935–945. [PubMed: 7525605]
- Fang J, Marchesini N, Moreno SN. A *Toxoplasma gondii* phosphoinositide phospholipase C (TgPI-PLC) with high affinity for phosphatidylinositol. *Biochem J.* 2006; 394:417–425. [PubMed: 16288600]
- Fessenden JD, Feng W, Pessah IN, Allen PD. Amino acid residues Gln4020 and Lys4021 of the ryanodine receptor type 1 are required for activation by 4-chloro-m-cresol. *J Biol Chem.* 2006; 281:21022–21031. [PubMed: 16737973]
- Fok AK, Aihara MS, Ishida M, Nolte KV, Steck TL, Allen RD. The pegs on the decorated tubules of the contractile vacuole complex of *Paramecium* are proton pumps. *J Cell Sci.* 1995; 108:3163–3170. [PubMed: 7593277]
- Furuya T, Kashuba C, Docampo R, Moreno SN. A novel phosphatidylinositol-phospholipase C of *Trypanosoma cruzi* that is lipid modified and activated during trypomastigote to amastigote differentiation. *J Biol Chem.* 2000; 275:6428–6438. [PubMed: 10692446]
- Furuya T, Okura M, Ruiz FA, Scott DA, Docampo R. TcSCA complements yeast mutants defective in Ca^{2+} pumps and encodes a Ca^{2+} -ATPase that localizes to the endoplasmic reticulum of *Trypanosoma cruzi*. *J Biol Chem.* 2001; 276:32437–32445. [PubMed: 11382780]
- Galione A, Evans AM, Ma J, Parrington J, Arredouani A, Cheng X, Zhu MX. The acid test: the discovery of two-pore channels (TPCs) as NAADP-gated endolysosomal Ca^{2+} release channels. *Pflugers Arch.* 2009; 458:869–876. [PubMed: 19475418]
- Galione A, Morgan AJ, Arredouani A, Davis LC, Rietdorf K, Ruas M, Parrington J. NAADP as an intracellular messenger regulating lysosomal calcium-release channels. *Biochem Soc Trans.* 2010; 38:1424–1431. [PubMed: 21118101]
- Galvani A, Sperling L. RNA interference by feeding in *Paramecium*. *Trends Genet.* 2002; 18:11–12. [PubMed: 11750689]
- Gamberucci A, Fulceri R, Marcolongo P, Pralong WF, Benedetti A. Histones and basic polypeptides activate Ca^{2+} /cation influx in various cell types. *Biochem J.* 1998; 331:623–630. [PubMed: 9531506]
- Garrison E, Treeck M, Ehret E, Butz H, Garbuz T, Oswald BP, Settles M, Boothroyd J, Arrizabalaga G. A forward genetic screen reveals that calcium-dependent protein kinase 3 regulates egress in *Toxoplasma*. *PLoS Pathog.* 2012; 8:e1003049. [PubMed: 23209419]
- Giacomello M, Drago I, Bortolozzi M, Scorzeto M, Gianelle A, Pizzo P, Pozzan T. Ca^{2+} hot spots on the mitochondrial surface are generated by Ca^{2+} mobilization from stores, but not by activation of store-operated Ca^{2+} channels. *Mol Cell.* 2010; 38:280–290. [PubMed: 20417605]
- Grover JE, Rope AF, Kaneishiro ES. The occurrence of biogenic struvite ($\text{Mg,Ca}\text{NH}_4\text{PO}_4\cdot 6\text{H}_2\text{O}$) as intracellular crystals in *Paramecium*. *J Eukaryot Microbiol.* 1997; 44:366–373.
- Guttery DS, Pittman JK, Frenal K, Poulin B, McFarlane LR, Slavic K, Wheatley SP, Soldati-Favre D, Krishna S, Tewari R, Staines HM. The *Plasmodium berghei* $\text{Ca}^{2+}/\text{H}^+$ exchanger, PbCAX, is essential for tolerance to environmental Ca^{2+} during sexual development. *PLoS Pathog.* 2013; 9:e1003191. [PubMed: 23468629]
- Hajnoczky G, Hager R, Thomas AP. Mitochondria suppress local feedback activation of inositol 1,4,5-trisphosphate receptors by Ca^{2+} . *J Biol Chem.* 1999; 274:14157–14162. [PubMed: 10318833]
- Hajnoczky G, Robb-Gaspers LD, Seitz MB, Thomas AP. Decoding of cytosolic calcium oscillations in the mitochondria. *Cell.* 1995; 82:415–424. [PubMed: 7634331]
- Hamilton SL. Ryanodine receptors. *Cell Calcium.* 2005; 38:253–260. [PubMed: 16115682]
- Hardt M, Plattner H. Sub-second quenched-flow/X-ray microanalysis shows rapid Ca^{2+} mobilization from cortical stores paralleled by Ca^{2+} influx during synchronous exocytosis in *Paramecium* cells. *Eur J Cell Biol.* 2000; 79:642–652. [PubMed: 11043405]
- Harumoto T, Miyake A. Defensive function of trichocysts in *Paramecium*. *J Exp Zool.* 1991; 260:84–92.

- Hashimoto M, Enomoto M, Morales J, Kurebayashi N, Sakurai T, Hashimoto T, Nara T, Mikoshiba K. Inositol 1,4,5-trisphosphate receptor regulates replication, differentiation, infectivity and virulence of the parasitic protist *Trypanosoma cruzi*. *Mol Microbiol*. 2013; 87:1133–1150. [PubMed: 23320762]
- Hay JC. Calcium: a fundamental regulator of intracellular membrane fusion? *EMBO Rep*. 2007; 8:236–240. [PubMed: 17330068]
- Horn T, Ullrich ND, Egger M. 'Eventless' InsP₃-dependent SR-Ca²⁺ release affecting atrial Ca²⁺ sparks. *J Physiol*. 2013; 591:2103–2111. [PubMed: 23381902]
- Huang G, Bartlett PJ, Thomas AP, Moreno SN, Docampo R. Acidocalcisomes of *Trypanosoma brucei* have an inositol 1,4,5-trisphosphate receptor that is required for growth and infectivity. *Proc Natl Acad Sci U S A*. 2013; 110:1887–1892. [PubMed: 23319604]
- Irigoin F, Inada NM, Fernandes MP, Piacenza L, Gadelha FR, Vercesi AE, Radi R. Mitochondrial calcium overload triggers complement-dependent superoxide-mediated programmed cell death in *Trypanosoma cruzi*. *Biochem J*. 2009; 418:595–604. [PubMed: 19053945]
- Jouaville LS, Pinton P, Bastianutto C, Rutter GA, Rizzuto R. Regulation of mitochondrial ATP synthesis by calcium: evidence for a long-term metabolic priming. *Proc Natl Acad Sci U S A*. 1999; 96:13807–13812. [PubMed: 10570154]
- Kaczanowski S, Sajid M, Reece SE. Evolution of apoptosis-like programmed cell death in unicellular protozoan parasites. *Parasit Vectors*. 2011; 4:44. [PubMed: 21439063]
- King N, Westbrook MJ, Young SL, Kuo A, Abedin M, Chapman J, Fairclough S, Hellsten U, Isogai Y, Letunic I, Marr M, Pincus D, Putnam N, Rokas A, Wright KJ, Zuzow R, Dirks W, Good M, Goodstein D, Lemons D, Li W, Lyons JB, Morris A, Nichols S, Richter DJ, Salamov A, Sequencing JG, Bork P, Lim WA, Manning G, Miller WT, McGinnis W, Shapiro H, Tjian R, Grigoriev IV, Rokhsar D. The genome of the choanoflagellate *Monosiga brevicollis* and the origin of metazoans. *Nature*. 2008; 451:783–788. [PubMed: 18273011]
- Kirichok Y, Kravitsinsky G, Clapham DE. The mitochondrial calcium uniporter is a highly selective ion channel. *Nature*. 2004; 427:360–364. [PubMed: 14737170]
- Klauke N, Blanchard M, Plattner H. Polyamine triggering of exocytosis in *Paramecium* involves an extracellular Ca²⁺/(polyvalent cation)-sensing receptor, subplasmalemmal Ca-store mobilization and store-operated Ca²⁺-influx via unspecific cation channels. *J Membr Biol*. 2000; 174:141–156. [PubMed: 10742458]
- Klauke N, Plattner H. Imaging of Ca²⁺ transients induced in *Paramecium* cells by a polyamine secretagogue. *J Cell Sci*. 1997; 110:975–983. [PubMed: 9152023]
- Klauke N, Plattner H. Caffeine-induced Ca²⁺ transients and exocytosis in *Paramecium* cells. A correlated Ca²⁺ imaging and quenched-flow/freeze-fracture analysis. *J Membr Biol*. 1998; 161:65–81. [PubMed: 9430622]
- Klee CB, Crouch TH, Richman PG. Calmodulin. *Annu Rev Biochem*. 1980; 49:489–515. [PubMed: 6250447]
- Kroemer G, Galluzzi L, Brenner C. Mitochondrial membrane permeabilization in cell death. *Physiol Rev*. 2007; 87:99–163. [PubMed: 17237344]
- Ladenburger EM, Korn I, Kasielke N, Wassmer T, Plattner H. An Ins(1,4,5)P₃ receptor in *Paramecium* is associated with the osmoregulatory system. *J Cell Sci*. 2006; 119:3705–3717. [PubMed: 16912081]
- Ladenburger EM, Plattner H. Calcium-release channels in *Paramecium*. Genomic expansion, differential positioning and partial transcriptional elimination. *PLoS One*. 2011; 6:e27111. [PubMed: 22102876]
- Ladenburger EM, Sehring IM, Korn I, Plattner H. Novel types of Ca²⁺ release channels participate in the secretory cycle of *Paramecium* cells. *Mol Cell Biol*. 2009; 29:3605–3622. [PubMed: 19380481]
- Lammel EM, Barbieri MA, Wilkowsky SE, Bertini F, Isola EL. *Trypanosoma cruzi*: involvement of intracellular calcium in multiplication and differentiation. *Exp Parasitol*. 1996; 83:240–249. [PubMed: 8682192]

- Lange S, Klauke N, Plattner H. Subplasmalemmal Ca^{2+} stores of probable relevance for exocytosis in *Paramecium*. Alveolar sacs share some but not all characteristics with sarcoplasmic reticulum. *Cell Calcium*. 1995; 17:335–344. [PubMed: 7553786]
- Lapatsina L, Brand J, Poole K, Daumke O, Lewin GR. Stomatin-domain proteins. *Eur J Cell Biol*. 2012; 91:240–245. [PubMed: 21501885]
- Lasorsa FM, Pinton P, Palmieri L, Fiermonte G, Rizzuto R, Palmieri F. Recombinant expression of the Ca^{2+} -sensitive aspartate/glutamate carrier increases mitochondrial ATP production in agonist-stimulated Chinese hamster ovary cells. *J Biol Chem*. 2003; 278:38686–38692. [PubMed: 12851387]
- Lehninger AL, Rossi CS, Greenawalt JW. Respiration-dependent accumulation of inorganic phosphate and Ca ions by rat liver mitochondria. *Biochem Biophys Res Commun*. 1963; 10:444–448. [PubMed: 13929376]
- Leondaritis G, Sarri T, Dafnis I, Efstathiou A, Galanopoulou D. Biochemical and genetic evidence for the presence of multiple phosphatidylinositol- and phosphatidylinositol 4,5-bisphosphate-specific phospholipases C in *Tetrahymena*. *Eukaryot Cell*. 2011; 10:412–422. [PubMed: 21169416]
- Lovett JL, Marchesini N, Moreno SN, Sibley LD. *Toxoplasma gondii* microneme secretion involves intracellular Ca^{2+} release from inositol 1,4,5-triphosphate (IP_3)/ryanodine-sensitive stores. *J Biol Chem*. 2002; 277:25870–25876. [PubMed: 12011085]
- Lovett JL, Sibley LD. Intracellular calcium stores in *Toxoplasma gondii* govern invasion of host cells. *J Cell Sci*. 2003; 116:3009–3016. [PubMed: 12783987]
- Lu HG, Zhong L, de Souza W, Benchimol M, Moreno S, Docampo R. Ca^{2+} content and expression of an acidocalcisomal calcium pump are elevated in intracellular forms of *Trypanosoma cruzi*. *Mol Cell Biol*. 1998; 18:2309–2323. [PubMed: 9528801]
- Luo S, Rohloff P, Cox J, Uyemura SA, Docampo R. *Trypanosoma brucei* plasma membrane-type Ca^{2+} -ATPase 1 (*TbPMC1*) and 2 (*TbPMC2*) genes encode functional Ca^{2+} -ATPases localized to the acidocalcisomes and plasma membrane, and essential for Ca^{2+} homeostasis and growth. *J Biol Chem*. 2004; 279:14427–14439. [PubMed: 14724285]
- Luo S, Vieira M, Graves J, Zhong L, Moreno SN. A plasma membrane-type Ca^{2+} -ATPase co-localizes with a vacuolar H^+ -pyrophosphatase to acidocalcisomes of *Toxoplasma gondii*. *EMBO J*. 2001; 20:55–64. [PubMed: 11226155]
- Luzio JP, Gray SR, Bright NA. Endosome-lysosome fusion. *Biochem Soc Trans*. 2010; 38:1413–1416. [PubMed: 21118098]
- Mackrill JJ. Ryanodine receptor calcium release channels: an evolutionary perspective. *Adv Exp Med Biol*. 2012; 740:159–182. [PubMed: 22453942]
- Malsam J, Kreye S, Sollner TH. Membrane fusion: SNAREs and regulation. *Cell Mol Life Sci*. 2008; 65:2814–2832. [PubMed: 18726177]
- Martins V, de P, Galizzi M, Salto ML, Docampo R, Moreno SN. Developmental expression of a *Trypanosoma cruzi* phosphoinositide-specific phospholipase C in amastigotes and stimulation of host phosphoinositide hydrolysis. *Infect Immun*. 2010; 78:4206–4212. [PubMed: 20643853]
- McCormack JG, Halestrap AP, Denton RM. Role of calcium ions in regulation of mammalian intramitochondrial metabolism. *Physiol Rev*. 1990; 70:391–425. [PubMed: 2157230]
- Miranda K, Pace DA, Cintron R, Rodrigues JC, Fang J, Smith A, Rohloff P, Coelho E, de Haas F, de Souza W, Coppens I, Sibley LD, Moreno SN. Characterization of a novel organelle in *Toxoplasma gondii* with similar composition and function to the plant vacuole. *Mol Microbiol*. 2010; 76:1358–1375. [PubMed: 20398214]
- Miyakawa T, Maeda A, Yamazawa T, Hirose K, Kurosaki T, Iino M. Encoding of Ca^{2+} signals by differential expression of IP_3 receptor subtypes. *EMBO J*. 1999; 18:1303–1308. [PubMed: 10064596]
- Mondragon R, Frixione E. Ca^{2+} -dependence of conoid extrusion in *Toxoplasma gondii* tachyzoites. *J Eukaryot Microbiol*. 1996; 43:120–127. [PubMed: 8720941]
- Mondragon R, Meza I, Frixione E. Divalent cation and ATP dependent motility of *Toxoplasma gondii* tachyzoites after mild treatment with trypsin. *J Eukaryot Microbiol*. 1994; 41:330–337. [PubMed: 8087104]

- Montero M, Alonso MT, Carnicero E, Cuchillo-Ibanez I, Albillos A, Garcia AG, Garcia-Sancho J, Alvarez J. Chromaffin-cell stimulation triggers fast millimolar mitochondrial Ca^{2+} transients that modulate secretion. *Nat Cell Biol.* 2000; 2:57–61. [PubMed: 10655583]
- Moreno SN, Docampo R, Vercesi AE. Calcium homeostasis in procyclic and bloodstream forms of *Trypanosoma brucei*. Lack of inositol 1,4,5-trisphosphate-sensitive Ca^{2+} release. *J Biol Chem.* 1992a; 267:6020–6026. [PubMed: 1556113]
- Moreno SN, Silva J, Vercesi AE, Docampo R. Cytosolic-free calcium elevation in *Trypanosoma cruzi* is required for cell invasion. *J Exp Med.* 1994; 180:1535–1540. [PubMed: 7931085]
- Moreno SN, Vercesi AE, Pignataro OP, Docampo R. Calcium homeostasis in *Trypanosoma cruzi* amastigotes: presence of inositol phosphates and lack of an inositol 1,4,5-trisphosphate-sensitive calcium pool. *Mol Biochem Parasitol.* 1992b; 52:251–261. [PubMed: 1620163]
- Moreno SN, Zhong L. Acidocalcisomes in *Toxoplasma gondii* tachyzoites. *Biochem J.* 1996; 313:655–659. [PubMed: 8573106]
- Nagamune K, Beatty WL, Sibley LD. Artemisinin induces calcium-dependent protein secretion in the protozoan parasite *Toxoplasma gondii*. *Eukaryot Cell.* 2007; 6:2147–2156. [PubMed: 17766463]
- Nagamune K, Hicks LM, Fux B, Brossier F, Chini EN, Sibley LD. Abscissic acid controls calcium-dependent egress and development in *Toxoplasma gondii*. *Nature.* 2008; 451:207–210. [PubMed: 18185591]
- Nagamune K, Sibley LD. Comparative genomic and phylogenetic analyses of calcium ATPases and calcium-regulated proteins in the Apicomplexa. *Mol Biol Evol.* 2006; 23:1613–1627. [PubMed: 16751258]
- Neher E. The use of Fura-2 for estimating Ca buffers and Ca fluxes. *Neuropharmacology.* 1995; 34:1423–1442. [PubMed: 8606791]
- Neher E. Vesicle pools and Ca^{2+} microdomains: new tools for understanding their roles in neurotransmitter release. *Neuron.* 1998; 20:389–399. [PubMed: 9539117]
- Nolan DP, Voorheis HP. The mitochondrion in bloodstream forms of *Trypanosoma brucei* is energized by the electrogenic pumping of protons catalysed by the F_1F_0 -ATPase. *Eur J Biochem.* 1992; 209:207–216. [PubMed: 1327770]
- Oberholzer M, Langousis G, Nguyen HT, Saada EA, Shimogawa MM, Jonsson ZO, Nguyen SM, Wohlschlegel JA, Hill KL. Independent analysis of the flagellum surface and matrix proteomes provides insight into flagellum signaling in mammalian-infectious *Trypanosoma brucei*. *Mol Cell Proteomics.* 2011; 10:M111 010538. [PubMed: 21685506]
- Okura M, Fang J, Salto ML, Singer RS, Docampo R, Moreno SN. A lipid-modified phosphoinositide-specific phospholipase C (TcPI-PLC) is involved in differentiation of trypomastigotes to amastigotes of *Trypanosoma cruzi*. *J Biol Chem.* 2005; 280:16235–16243. [PubMed: 15710612]
- Pace DA, Liu J, McNight C, Jimenez V, Moreno SN. Calcium entry induced by depletion of intracellular stores enhances invasion-linked traits in *Toxoplasma gondii*. 2013 Submitted.
- Parekh AB, Putney JW Jr. Store-operated calcium channels. *Physiol Rev.* 2005; 85:757–810. [PubMed: 15788710]
- Passos AP, Garcia CR. Inositol 1,4,5-trisphosphate induced Ca^{2+} release from chloroquine-sensitive and -insensitive intracellular stores in the intraerythrocytic stage of the malaria parasite *P. chabaudi*. *Biochem Biophys Res Commun.* 1998; 245:155–160. [PubMed: 9535800]
- Patel S, Docampo R. In with the TRP channels: intracellular functions for TRPM1 and TRPM2. *Sci Signal.* 2009; 2:pe69. [PubMed: 19887679]
- Patel S, Docampo R. Acidic calcium stores open for business: expanding the potential for intracellular Ca^{2+} signaling. *Trends Cell Biol.* 2010; 20:277–286. [PubMed: 20303271]
- Perocchi F, Gohil VM, Girgis HS, Bao XR, McCombs JE, Palmer AE, Mootha VK. MICU1 encodes a mitochondrial EF hand protein required for Ca^{2+} uptake. *Nature.* 2010; 467:291–296. [PubMed: 20693986]
- Plattner H. How to design a highly organized cell: An unexpectedly high number of widely diversified SNARE proteins positioned at strategic sites in the ciliate, *Paramecium tetraurelia*. *Protist.* 2010; 161:497–516. [PubMed: 20646960]
- Plattner H. The contractile vacuole complex – its expanding protein inventory. *Int Rev Cell Mol Biol.* 2013 in press.

- Plattner H, Hentschel J. Sub-second cellular dynamics: time-resolved electron microscopy and functional correlation. *Int Rev Cytol.* 2006; 255:133–176. [PubMed: 17178466]
- Plattner H, Klauke N. Calcium in ciliated protozoa: sources, regulation, and calcium-regulated cell functions. *Int Rev Cytol.* 2001; 201:115–208. [PubMed: 11057832]
- Plattner H, Sehring IM, Mohamed IK, Miranda K, De Souza W, Billington R, Genazzani A, Ladenburger EM. Calcium signaling in closely related protozoan groups (Alveolata): non-parasitic ciliates (*Paramecium*, *Tetrahymena*) vs. parasitic Apicomplexa (*Plasmodium*, *Toxoplasma*). *Cell Calcium.* 2012; 51:351–382. [PubMed: 22387010]
- Plattner H, Sehring IM, Schilde C, Ladenburger EM. Pharmacology of ciliated protozoa--drug (in)sensitivity and experimental drug (ab)use. *Int Rev Cell Mol Biol.* 2009; 273:163–218. [PubMed: 19215905]
- Plattner H, Verkhatsky A. Calcium signaling early in evolution – all but primitive. *J Cell Sci.* 2013; 126:2141–2150. [PubMed: 23729741]
- Prole DL, Taylor CW. Identification of intracellular and plasma membrane calcium channel homologues in pathogenic parasites. *PLoS One.* 2011; 6:e26218. [PubMed: 22022573]
- Ramachandran S, Chakraborty A, Xu L, Mei Y, Samso M, Dokholyan NV, Meissner G. Structural determinants of skeletal muscle ryanodine receptor gating. *J Biol Chem.* 2013; 288:6154–6165. [PubMed: 23319589]
- Reuter AT, Stuermer CA, Plattner H. Identification, localization, and functional implications of the microdomain-forming stomatin family in the ciliated protozoan *Paramecium tetraurelia*. *Eukaryot Cell.* 2013; 12:529–544. [PubMed: 23376944]
- Ricklefs RE, Outlaw DC. A molecular clock for malaria parasites. *Science.* 2010; 329:226–229. [PubMed: 20616281]
- Ridgley EL, Xiong ZH, Ruben L. Reactive oxygen species activate a Ca^{2+} -dependent cell death pathway in the unicellular organism *Trypanosoma brucei brucei*. *Biochem J.* 1999; 340:33–40. [PubMed: 10229656]
- Rizo J, Chen X, Arac D. Unraveling the mechanisms of synaptotagmin and SNARE function in neurotransmitter release. *Trends Cell Biol.* 2006; 16:339–350. [PubMed: 16698267]
- Rizzuto R, Brini M, Murgia M, Pozzan T. Microdomains with high Ca^{2+} close to IP_3 -sensitive channels that are sensed by neighboring mitochondria. *Science.* 1993; 262:744–747. [PubMed: 8235595]
- Rizzuto R, Pinton P, Carrington W, Fay FS, Fogarty KE, Lifshitz LM, Tuft RA, Pozzan T. Close contacts with the endoplasmic reticulum as determinants of mitochondrial Ca^{2+} responses. *Science.* 1998; 280:1763–1766. [PubMed: 9624056]
- Rodrigues CO, Scott DA, Docampo R. Characterization of a vacuolar pyrophosphatase in *Trypanosoma brucei* and its localization to acidocalcisomes. *Mol Cell Biol.* 1999; 19:7712–7723. [PubMed: 10523660]
- Rohloff P, Miranda K, Rodrigues JC, Fang J, Galizzi M, Plattner H, Hentschel J, Moreno SN. Calcium uptake and proton transport by acidocalcisomes of *Toxoplasma gondii*. *PLoS One.* 2011; 6:e18390. [PubMed: 21541023]
- Rohloff P, Rodrigues CO, Docampo R. Regulatory volume decrease in *Trypanosoma cruzi* involves amino acid efflux and changes in intracellular calcium. *Mol Biochem Parasitol.* 2003; 126:219–230. [PubMed: 12615321]
- Satrústegui J, Pardo B, Del Arco A. Mitochondrial transporters as novel targets for intracellular calcium signaling. *Physiol Rev.* 2007; 87:29–67. [PubMed: 17237342]
- Schnauffer A, Clark-Walker GD, Steinberg AG, Stuart K. The F_1 -ATP synthase complex in bloodstream stage trypanosomes has an unusual and essential function. *EMBO J.* 2005; 24:4029–4040. [PubMed: 16270030]
- Schonemann B, Bledowski A, Sehring IM, Plattner H. A set of SNARE proteins in the contractile vacuole complex of *Paramecium* regulates cellular calcium tolerance and also contributes to organelle biogenesis. *Cell Calcium.* 2013; 53:204–216. [PubMed: 23280185]
- Scott DA, de Souza W, Benchimol M, Zhong L, Lu HG, Moreno SN, Docampo R. Presence of a plant-like proton-pumping pyrophosphatase in acidocalcisomes of *Trypanosoma cruzi*. *J Biol Chem.* 1998; 273:22151–22158. [PubMed: 9705361]

- Selzer PM, Webster P, Duszenko M. Influence of Ca^{2+} depletion on cytoskeleton and nucleolus morphology in *Trypanosoma brucei*. *Eur J Cell Biol.* 1991; 56:104–112. [PubMed: 1666039]
- Sherwood MW, Prior IA, Voronina SG, Barrow SL, Woodsmith JD, Gerasimenko OV, Petersen OH, Tepikin AV. Activation of trypsinogen in large endocytic vacuoles of pancreatic acinar cells. *Proc Natl Acad Sci U S A.* 2007; 104:5674–5679. [PubMed: 17363470]
- Smirlis D, Duszenko M, Ruiz AJ, Scoulica E, Bastien P, Fasel N, Soteriadou K. Targeting essential pathways in trypanosomatids gives insights into protozoan mechanisms of cell death. *Parasit Vectors.* 2010; 3:107. [PubMed: 21083891]
- Smirlis D, Soteriadou K. Trypanosomatid apoptosis: ‘Apoptosis’ without the canonical regulators. *Virulence.* 2011; 2:253–256. [PubMed: 21566464]
- Solovyova N, Verkhatsky A. Neuronal endoplasmic reticulum acts as a single functional Ca^{2+} store shared by ryanodine and inositol-1,4,5-trisphosphate receptors as revealed by intra-ER [Ca^{2+}] recordings in single rat sensory neurones. *Pflugers Arch.* 2003; 446:447–454. [PubMed: 12764616]
- Stelly N, Mauger JP, Claret M, Adoutte A. Cortical alveoli of *Paramecium*: a vast submembranous calcium storage compartment. *J Cell Biol.* 1991; 113:103–112. [PubMed: 1848863]
- Stock C, Gronlien HK, Allen RD. The ionic composition of the contractile vacuole fluid of *Paramecium* mirrors ion transport across the plasma membrane. *Eur J Cell Biol.* 2002; 81:505–515. [PubMed: 12416727]
- Stojdl DF, Clarke MW. *Trypanosoma brucei*: analysis of cytoplasmic Ca^{2+} during differentiation of bloodstream stages in vitro. *Exp Parasitol.* 1996; 83:134–146. [PubMed: 8654542]
- Taylor CW, da Fonseca PC, Morris EP. IP_3 receptors: the search for structure. *Trends Biochem Sci.* 2004; 29:210–219. [PubMed: 15082315]
- Tinel H, Cancela JM, Mogami H, Gerasimenko JV, Gerasimenko OV, Tepikin AV, Petersen OH. Active mitochondria surrounding the pancreatic acinar granule region prevent spreading of inositol trisphosphate-evoked local cytosolic Ca^{2+} signals. *EMBO J.* 1999; 18:4999–5008. [PubMed: 10487752]
- Tjondrokoesoemo A, Li N, Lin PH, Pan Z, Ferrante CJ, Shirokova N, Brotto M, Weisleder N, Ma J. Type 1 inositol 1,4,5-trisphosphate receptor activates ryanodine receptor 1 to mediate calcium spark signaling in adult mammalian skeletal muscle. *J Biol Chem.* 2013; 288:2103–2109. [PubMed: 23223241]
- Ulrich PN, Jimenez V, Park M, Martins VP, Atwood J 3rd, Moles K, Collins D, Rohloff P, Tarleton R, Moreno SN, Orlando R, Docampo R. Identification of contractile vacuole proteins in *Trypanosoma cruzi*. *PLoS One.* 2011; 6:e18013. [PubMed: 21437209]
- Vasington FD, Murphy JV. Ca ion uptake by rat kidney mitochondria and its dependence on respiration and phosphorylation. *J Biol Chem.* 1962; 237:2670–2677. [PubMed: 13925019]
- Vercesi AE, Docampo R. Ca^{2+} transport by digitonin-permeabilized *Leishmania donovani*. Effects of Ca^{2+} , pentamidine and WR-6026 on mitochondrial membrane potential in situ. *Biochem J.* 1992; 284:463–467. [PubMed: 1376113]
- Vercesi AE, Docampo R, Moreno SN. Energization-dependent Ca^{2+} accumulation in *Trypanosoma brucei* bloodstream and procyclic trypomastigotes mitochondria. *Mol Biochem Parasitol.* 1992; 56:251–257. [PubMed: 1484549]
- Vercesi AE, Moreno SN, Bernardes CF, Meinicke AR, Fernandes EC, Docampo R. Thapsigargin causes Ca^{2+} release and collapse of the membrane potential of *Trypanosoma brucei* mitochondria in situ and of isolated rat liver mitochondria. *J Biol Chem.* 1993; 268:8564–8568. [PubMed: 8473301]
- Vercesi AE, Moreno SN, Docampo R. $\text{Ca}^{2+}/\text{H}^{+}$ exchange in acidic vacuoles of *Trypanosoma brucei*. *Biochem J.* 1994; 304:227–233. [PubMed: 7998937]
- Vieira MC, Moreno SN. Mobilization of intracellular calcium upon attachment of *Toxoplasma gondii* tachyzoites to human fibroblasts is required for invasion. *Mol Biochem Parasitol.* 2000; 106:157–162. [PubMed: 10743619]
- Voronina SG, Barrow SL, Simpson AW, Gerasimenko OV, da Silva Xavier G, Rutter GA, Petersen OH, Tepikin AV. Dynamic changes in cytosolic and mitochondrial ATP levels in pancreatic acinar cells. *Gastroenterology.* 2010; 138:1976–1987. [PubMed: 20102715]

- Wang X, Zhang X, Dong XP, Samie M, Li X, Cheng X, Goschka A, Shen D, Zhou Y, Harlow J, Zhu MX, Clapham DE, Ren D, Xu H. TPC proteins are phosphoinositide- activated sodium-selective ion channels in endosomes and lysosomes. *Cell*. 2012; 151:372–383. [PubMed: 23063126]
- Wassmer T, Sehring IM, Kissmehl R, Plattner H. The V-ATPase in *Paramecium*: functional specialization by multiple gene isoforms. *Pflugers Arch*. 2009; 457:599–607. [PubMed: 18228038]
- Westerblad H, Andrade FH, Islam MS. Effects of ryanodine receptor agonist 4-chloro-m-cresol on myoplasmic free Ca^{2+} concentration and force of contraction in mouse skeletal muscle. *Cell Calcium*. 1998; 24:105–115. [PubMed: 9803311]
- Wetzel DM, Chen LA, Ruiz FA, Moreno SN, Sibley LD. Calcium-mediated protein secretion potentiates motility in *Toxoplasma gondii*. *J Cell Sci*. 2004; 117:5739–5748. [PubMed: 15507483]
- Williams K. Modulation and block of ion channels: a new biology of polyamines. *Cell Signal*. 1997; 9:1–13. [PubMed: 9067625]
- Xiong ZH, Ridgley EL, Enis D, Olness F, Ruben L. Selective transfer of calcium from an acidic compartment to the mitochondrion of *Trypanosoma brucei*. Measurements with targeted aequorins. *J Biol Chem*. 1997; 272:31022–31028. [PubMed: 9388251]
- Yoshikawa F, Morita M, Monkawa T, Michikawa T, Furuichi T, Mikoshiba K. Mutational analysis of the ligand binding site of the inositol 1,4,5-trisphosphate receptor. *J Biol Chem*. 1996; 271:18277–18284. [PubMed: 8663526]
- Zalk R, Lehnart SE, Marks AR. Modulation of the ryanodine receptor and intracellular calcium. *Annu Rev Biochem*. 2007; 76:367–385. [PubMed: 17506640]
- Zampese E, Pizzo P. Intracellular organelles in the saga of Ca^{2+} homeostasis: different molecules for different purposes? *Cell Mol Life Sci*. 2012; 69:1077–1104. [PubMed: 21968921]
- Zerial M, McBride H. Rab proteins as membrane organizers. *Nat Rev Mol Cell Biol*. 2001; 2:107–117. [PubMed: 11252952]
- Zong X, Schieder M, Cuny H, Fenske S, Gruner C, Rotzer K, Griesbeck O, Harz H, Biel M, Wahl-Schott C. The two-pore channel TPCN2 mediates NAADP-dependent Ca^{2+} -release from lysosomal stores. *Pflugers Arch*. 2009; 458:891–899. [PubMed: 19557428]

Fig. 1.

Distribution of CRCs in the *P. tetraurelia* cell (*PtCRC*), subfamilies I to VI (and subtypes analyzed in some detail given in arabic numbers), as localized by immuno-fluorescence. Additional sites of *PtCRC* localization are not included in the scheme. These are *PtCRC*-V-4, occurring along the oral cavity (together with *PtCRC*-VI-3) and along the cleavage furrow; remarkably *PtCRC*-V-4 also occurs in the parasomal sacs (clathrin coated pits at the ciliary basis) membranes. Beyond structures identified below the scheme, the following abbreviations are used: in the contractile vacuole (*cv*) system, *a* marks ampullae (with extending radial/collecting canals), *ds* = decorated spongiome and *ss* = smooth spongiome. Other structures are: *ac* = acidosomes (vesicles of endosomal origin contributing to phagosome/food vacuole formation), *cf* = cytopharyngeal fibers; *ci* = cilia; *cp* = cytoproct, *dv* = discoidal vesicles, *rv* = recycling vesicles, *mac* = macronucleus, *oc* = oral cavity, *pof* = postoral fibers, *ps* = parasomal sacs (clathrin-coated pits), *tr* = trichocysts and their “ghosts” (*gh*) occurring after contents release. Compiled according to (Ladenburger and Plattner, 2011).

Fig. 2.

Examples of immuno-localization of different *PtCRCs*. **(A)** *PtCRC-II* (IP₃R) is localized to the star-shaped contractile vacuole complex. **(B)** *PtCRC-III-4* is localized to the phagocytic pathway, including recycling vesicles. Here, phagosomes (food vacuoles) show up in dark due to the addition of Indian ink in the medium. For better identification of structures labeled with antibodies against *PtCRC-III-4* (green) red antibodies have been applied to stain microtubules. Here, *PtCRC-III-4* traffic with recycling vesicles along microtubules that are associated with the postoral fibers (*pof*) which originate from the oral apparatus (*oa*). By contrast, the contractile vacuole complex (*cvc*) is stained only for tubulin, but not for

PtCRC-III-4. Also indicated is the number of z-stacks used for a pile-up image in **(B)**. See Fig. 1 for further details. **(C)** Immunolocalization of *PtCRC-IV-1* (RyR-LP) by *green* fluorescence. *Red* staining comes from labeling of protein disulfide isomerase (PDI), an ER-specific marker. Note the localization of *PtCRC-IV-1* preferably to the alveolar sacs (Ca^{2+} -stores, represented by the green patches in the cell cortex) and less to the ER. Abbreviations are: *oc* = oral cavity, *cp* = cytoproct. **(D)** *PtCRC-V-4* localizes to micronuclei (*mic*), i.e. their envelopes, as well as to cell surface components (see Fig. 1 for details). From (Ladenburger *et al.*, 2009). Scale bars = 10 μm .

Fig. 3.

(A) Molecular modeling of the inositol 1,4,5-trisphosphate-BD of *PtCRC-II* (designated *PtIP₃R_N*) in comparison to type 1 IP₃R of the mouse (*MmIP₃R1*). The IP₃-BD of the *Paramecium* molecule has been modeled by comparison with that of *MmIP₃R1*. Numbers in parenthesis indicate the amino acid sequence included in the modeling procedure. Note considerable coincidence of the motifs in both IP₃Rs, with some additional loops (of yet unexplained significance) in the IP₃-BD of *Paramecium*. From (Ladenburger *et al.*, 2006).

(B) Exocytosis performance (percent of cells performing exocytosis [ordinate] to a different extent indicated in the abscissa) and the effect of *PtCRC-IV-1* (RyR-LP) silencing. Two types of control cells have been evaluated: (i) Cells mock-silenced with the vector (pPD) containing only a GFP sequence served as a negative control; (ii) cells silenced in the exocytosis-relevant *ND7* gene served as a positive control. The relevant experiment was carried out with a vector containing a sequence appropriate to silence *PtCRC-IV*. Note considerable depression of AED-stimulated exocytosis after ND7 and *PtCRC-IV* silencing, respectively. From (Ladenburger *et al.*, 2009). (C) Effect of silencing of *PtCRC-IV-1* (RyR-LP) on Ca^{2+} -signaling. Cells were exceptionally contained in a medium with $[Ca^{2+}]_o$ reduced to a calculated value of ~ 30 nM, i.e. slightly below internal resting level ($[Ca^{2+}]_i \sim 50$ nM, thus excluding signals from Ca^{2+} -influx. The Ca^{2+} signal generated during

exocytosis stimulation was evaluated by ratio imaging (f/f_0). *Black lines*: non-silenced cells, *red lines*: silenced cells. Exocytosis was stimulated by aminoethyl dextrane (AED), 4-chloro-m-cresol (4CmC) and caffeine, respectively. For further details, see text. From (Ladenburger *et al.*, 2009).

Fig. 4.

Localization of *TbIP*₃*R* in procyclic trypomastigotes. **(A)** *TbIP*₃*R* partially co-localizes with *TbVP*₁ in acidocalcisomes (Pearson's correlation coefficient of 0.874), as shown by immunofluorescence microscopy analysis. The merge images show the co-localization in yellow. **(B)** Lack of co-localization of *TbIP*₃*R* with *TbBiP* in the endoplasmic reticulum (Pearson's correlation coefficient of 0.156). Scale bars, 10 μ m. DIC, differential interference contrast. From (Huang *et al.*, 2013).

Fig. 5.

Caged IP_3 -dependent Ca^{2+} release in *T. brucei*. (**A–B**) Representative traces of Ca^{2+} responses to UV flash in control cells in the absence (**A**) and presence (**B**) of caged IP_3 , respectively (first flash 3 pulses, second flash 6 pulses). Nigericin (Nig; 5 μM) and Ionomycin (IO; 5 μM) were added where indicated. From (Huang *et al.*, 2013).

Fig. 6.

Schematic representation of the distribution of Ca^{2+} in a *T. gondii* tachyzoite. Ca^{2+} entry is probably through Ca^{2+} channels (a). Once inside the cells, Ca^{2+} can be translocated back to the extracellular environment, primarily by the action of the PMCA (b). In addition, Ca^{2+} will interact with binding proteins (CBPs) or become sequestered by the ER by the action of the SERCA- Ca^{2+} -ATPase (c), passively sequestered by the mitochondrion (d), or sequestered by the acidocalcisome (e) or the PLV (f) by the action of a Ca^{2+} -ATPase (TgA1). Ca^{2+} appears to diffuse freely into the nucleus. Calcium could also be released into

the cytoplasm from the internal stores, such as the ER, through an uncharacterized channel, which appears to respond to ryanodine and caffeine (*g*). It may also be released from the PLV and the acidocalcisome (AC) through a $\text{Ca}^{2+}/\text{H}^{+}$ exchanger (*h*). Acidic compartments such as the acidocalcisome and the PLV contain enzymes involved in their acidification e.g. the H^{+} -ATPase (*i*) and the vacuolar- H^{+} -pyrophosphatase (*j*). Mitochondrial Ca^{2+} release is through a $\text{Ca}^{2+}/\text{H}^{+}$ exchanger (*k*).

Fig. 7.

Scheme of the potential contact between acidocalcisomes and mitochondrion in trypanosomes. The scheme depicts the molecules mediating Ca^{2+} influx (MCU, MICU1) and efflux ($\text{Ca}^{2+}/\text{H}^+$ exchanger, CAX) across the inner mitochondrial membrane (IMM) at an area of acidocalcisome (ACCSOME)-mitochondrial association. The shades of gray represent the $[\text{Ca}^{2+}]$: dark gray: $> 500 \mu\text{M}$; white, 100 nM ; PMCA, plasma membrane-type Ca^{2+} -ATPase; OMM, outer mitochondrial membrane; VDAC, voltage-dependent anion-selective channel; IP_3R , inositol 1,4,5-trisphosphate receptor.