Published in final edited form as:

Eur J Neurosci. 2010 January ; 31(1): 79–89.

Impact of cocaine on adult hippocampal neurogenesis in an animal model of differential propensity to drug abuse

M. Julia García-Fuster, Javier A. Perez, Sarah M. Clinton, Stanley J Watson, and Huda Akil

Molecular and Behavioral Neuroscience Institute, University of Michigan, Ann Arbor, MI, USA

Abstract

Hippocampal plasticity (e.g., neurogenesis) likely plays an important role in maintaining addictive behavior and/or relapse. This study assessed whether rats with differential propensity to drug-seeking behavior, bred Low-Responders (bLR) and bred High-Responders (bHR) to novelty, show differential neurogenesis regulation after cocaine exposure. Using specific immunological markers, we labeled distinct populations of adult stem cells in the dentate gyrus at different time-points of the cocaine sensitization process; Ki-67 for newly born cells, NeuroD for cells born partway, and BrdU for older cells born prior to sensitization. Results show that: (1) bHRs exhibited greater psychomotor response to cocaine than bLRs. (2) Acute cocaine did not alter cell proliferation in bLR/bHR rats. (3) Chronic cocaine decreased cell proliferation in bLRs only, which became amplified through the course of abstinence. (4) Neither chronic cocaine nor cocaine abstinence affected the survival of immature neurons in either phenotype. (5) Cocaine abstinence decreased survival of mature neurons in bHRs only, an effect that paralleled the greater psychomotor response to cocaine. (6) Cocaine treatment did not affect the ratio of neurons to glia in bLR/bHR rats as most cells differentiated into neurons in both lines. Thus, cocaine exerts distinct effects on neurogenesis in bLR versus bHR rats, with a decrease in the birth of new progenitor cells in bLRs and a suppression of the survival of new neurons in bHRs which likely leads to an earlier decrease in formation of new connections. This latter effect in bHRs could contribute to their enhanced degree of cocaine-induced psychomotor behavioral sensitization.

Keywords

psychomotor sensitization; cell genesis; dentate gyrus; hippocampus; bLR/bHR rats

Introduction

Cocaine is a highly abused drug that poses an enormous health, social and economic toll on society. A major challenge of addiction research is to identify factors that predispose individuals' addiction in order to develop novel and better-tailored treatment options. Our laboratory has selectively-bred Sprague-Dawley rats for high (High-Responder, bHR) and low (Low-Responder, bLR) novelty-seeking behavior, a trait that predicts a variety of differences in emotional reactivity (Perez et al., 2009), including propensity to acquire

cocaine self-administration (Piazza et al., 1989; Piazza and Le Moal, 1996; Davis et al., 2008) and behavioral sensitization (Hooks et al., 1991; 1992).

Drugs of abuse produce neuroadaptations in particular brain areas implicated in reward and learning that likely underlie their long-lasting effects (Robinson and Kolb, 2004; Kalivas, 2007). It is well known that drugs of abuse such as cocaine are rewarding by virtue of their ability to increase dopamine concentrations in ventral tegmental area projections, such as the NAc (Gold et al., 1989). However, beyond modulating specific transmitter systems, initial propensity to drug abuse (i.e., bLR vs. bHR rats) may have more to do with environmental reactivity and stress responsiveness at other anatomical levels. In particular, the hippocampus (HC), which is highly interconnected with limbic structures that regulate mood and cognition, can influence the learning component of addiction and is associated with many psychostimulant effects, including contextual conditioning (Shen et al., 2006) and behavioral sensitization (Lodge and Grace, 2008). Indeed, neonatal lesions of the ventral HC produced enhanced cocaine sensitization in adulthood (Chamber and Taylor, 2004), providing a direct association between sensitization and altered hippocampal (HCal) circuits. Moreover, some of the known bLR/bHR neuroadaptations that may contribute to their drug-taking differences occur in the HC (Maccari et al., 1991; Kabbaj et al., 2000; Turner et al., 2008).

Adult neurogenesis represents one form of HCal plasticity that may contribute to the brain's ability to process, adapt, and respond to stimuli (e.g., brain dysfunction in drug addiction). Adult HCal neurogenesis is a highly dynamic process where mature functionally-integrated granular neurons are generated from progenitor cells in the dentate gyrus (DG) (Ming and Song, 2005). Each stage of neurogenesis is demarcated by distinct molecular markers (Von Bohlen and Halbach, 2007) and is regulated by a variety of factors, including adverse environmental conditions (Schmidt and Duman, 2007) and exposure to addictive substances (Eisch and Harburg, 2006). Particularly, experimenter-administered cocaine impaired cell proliferation in adult rat HC (Yamaguchi et al., 2004) without altering the survival and growth of immature cells (Dominguez-Escriba et al., 2006), suggesting that cocaine differentially modulates cells at different stages of neurogenesis. Thus, altered neurogenesis represents one drug-regulated form of neuroplasticity that may contribute to the maintenance of addictive behavior and/or relapse.

Against this background, the present study attempts to understand how experimenter-administered cocaine may alter the cellular composition of the HC in an animal model of differential propensity to drug abuse (bLR/bHR rats), and to ascertain which stages of adult HCal neurogenesis (e.g., cell proliferation, survival, differentiation) may be regulated by cocaine within each phenotype.

Materials and methods

Experimental design, animals, and drug treatment

The first part of the study (Study I) characterized the bLR/bHR animal model of differential propensity to drug abuse to examine their behavioral responses to novelty (i.e., what defines the phenotype) and cocaine (i.e., different psychomotor response to cocaine). Once the

animal model was well defined, the second part of the study (Study II) assessed the impact of cocaine on adult HCal neurogenesis in bLR/bHR rats. The experimental design, animals used, and drug treatments are described in detail below.

Study I: Characterization of bLR and bHR rats

1. bLR/bHR differences in locomotor response to novelty—For the last several years our laboratory has selectively-bred rats (bLR and bHR lines) on the basis of locomotor reactivity to a novel environment, demonstrating reliable differences across multiple behavioral and neurobiological dimensions. A description of the breeding strategy and initial behavioral characterization of the bHR and bLR lines was previously published (Stead et al, 2006) and has been extended in the present study (see Results and Figure 1).

Each generation of male rats was screened for locomotor activity around 60 days of age (see Stead et al., 2006). Note that Figure 1A represents bLR and bHR rats' inborn basal locomotor differences from generations F1-F20, although the present studies only used rats from generations F15 and F20. Horizontal and rearing activity was monitored by computer in 5 min intervals over 60 min by placing animals into clear acrylic 43×21.5×25.5 cm (high) cages equipped with infrared photocell emitters mounted 2.3 and 6.5 cm above the grid floor in a different room from where the animals had been housed. The locomotion-testing rig and motion-recording software were created in-house at the University of Michigan. All testing was performed between 8:00 and 11:30 am. Final locomotion scores for each rat were calculated by adding the total number of horizontal and rearing movements. bHR animals whose scores fell one standard deviation below the bHR group average and bLR animals whose scores fell one standard deviation above the group average were not used for the subsequent cocaine-treatment study. The present series of studies used two sets of 25 bLR/bHR pairs from two different generations (F15 and F20). Animals were maintained in accordance with the University of Michigan Committee on the Use and Care of Animals (UCUCA).

2. bLR/bHR differences in psychomotor sensitization to cocaine—It is known that in commercially-available outbred LR/HR rats, the animals' reactivity to novelty predicts the degree of cocaine sensitization, with HR rats showing a greater psychomotor response to cocaine than LR rats (Hooks et al., 1992). Thus, the idea behind this experiment was to extend these earlier findings showing that our selectively-bred bLR/bHR rats also show differences in psychomotor sensitization to cocaine.

For this purpose, following bLR/bHR phenotype characterization (described above), we used 7 pairs of adult bLR and bHR males from the 20th generation weighting 225-250 g (n=7 per group). On the first day of testing, rats were transferred from the colony room and placed into the activity monitoring chambers described above. Due to the known bLR/bHR differences in motor activity, animals were allowed to habituate for 1 h before receiving a cocaine injection (15 mg/kg, i.p., see Figure 1B) followed by 1 h during which locomotion (horizontal and rearing activity) was scored. After the first day of testing, rats were returned to their home cages. Once a day for the next 6 days, rats were injected in an adjacent novel room with 15 mg/kg of cocaine (around 10:00 hours). On day 7, rats were returned to the

activity chambers where they were again allowed to habituate before receiving a single cocaine injection followed by 1 hour behavioral scoring. After 14 days of abstinence, rats were transferred to the activity chambers; following 1 h habituation, animals were challenged with 15 mg/kg dose of cocaine to assess its impact on locomotion for an additional 1 h.

For each rat we calculated locomotion scores (total, lateral, and rearing activity) as the value following cocaine injection (stimulated) minus the background (defined as activity occurring during the last 30 minutes of habituation). This method allowed us to correct for the basal inborn differences in bLR/bHR rats and to compare both groups following cocaine treatments.

Study II: bLR/bHR differences in neurogenesis following experimenter-administered cocaine

Once we characterized animals according to their bLR/bHR phenotype, the aim of the next study was to investigate potential differences in adult HCal neurogenesis (recent cell proliferation, early and late cell survival, and cell differentiation) following cocaine treatments. This second study consisted of two sets of experiments detailed below. The initial pilot study assessed the effect of cocaine on cell proliferation in bLR/bHR rats, and later led to a second more complete study which included neurogenesis regulation at different stages.

1. bLR/bHR differences in cell proliferation following experimenter-administered cocaine

—We used bLR and bHR rats (F15, n=5 per phenotype per experimental group) to determine the effects of acute, chronic and early cocaine abstinence on adult HCal cell proliferation as measured by Ki-67 immunohistochemistry (IHC) in the DG (Figure 2A). The chronic cocaine treatment is known to induce behavioral sensitization (Gosnell, 2005), and produces bLR/bHR differences in psychomotor response to the drug (see Figure 1B). Thus, one group of bLR/bHR rats received daily injections of 15 mg/kg cocaine (i.p.), whereas the control group received daily injections of saline for 7 consecutive days. The acute cocaine group received daily injections of saline for 6 days and an acute cocaine injection (15 mg/kg, i.p.) on day 7. All animals were transferred to an adjacent room to receive their daily injections (around 10:00 hours) and were then returned to their colony room. On day 7 of cocaine administration, animals were either sacrificed 45 min or 3 days (early abstinence; García-Fuster et al., 2009) following the last cocaine injection.

2. bLR/bHR differences in neurogenesis following experimenter-administered cocaine

—We used bLR/bHR rats (F20, n=6 per phenotype per experimental group) to determine the effects of experimenter-administered cocaine on the different stages of adult HCal neurogenesis (see Figure 3A). The chosen cellular markers enabled us to look back in time from the day of the animal's death and study cells born at different stages along the sensitization process —i.e., new cells that proliferated immediately prior to death, young immature neurons born partway through the sensitization process, or mature neurons that were born prior to the first cocaine or saline administration and survived and differentiated during the course of the treatment. As depicted in Figure 3, rats were pretreated for 3 days

with BrdU (50 mg/kg, i.p.; Calbiochem) delivered at 12 h intervals 4 days before the treatment started (Thomas et al., 2007). This protocol avoids marker dilution with cell division (Dayer et al., 2003) and labels newly generated cells that will differentiate and after 28 days of survival (Brown et al., 2003) be stably incorporated within the HC (Kempermann, 2002; Kempermann et al., 2004). This enabled us to measure cocaine effects on late cell survival (BrdU) and cell differentiation (BrdU/NeuN/GFAP) separately from measures of recent cell proliferation (Ki-67) and early cell survival (NeuroD) that can be achieved with postmortem markers. Following the same paradigm as described earlier, some bLR/bHR rats received injections of 15 mg/kg cocaine (7 days, i.p.) whereas the control group received saline for 7 consecutive days (Figure 3B). All animals were transferred to an adjacent room to receive their daily injections (around 10:00 hours) and were then returned to their colony room. On day 7 of cocaine administration, animals were either sacrificed 45 min or 14 days (intermediate abstinence) following the last cocaine injection. In an attempt to minimize the number of animals used for the study, the control abstinence group was not run due to the fact that we had observed no differences in neurogenesis between the control group and the control abstinence group (see Figure 2, Results of previous experiment).

Tissue collection and immunohistochemistry studies

At the times indicated above, animals were killed by decapitation and the brains removed. The left half-brain was fast frozen and stored at -80°C until use for IHC experiments. For each animal, a series of every 8th section was cryostat cut at 30 μm and slide-mounted throughout the entire extent of the HC.

As mentioned earlier, we utilized molecular tools to examine at a given point in time the effects of experimenter-administered cocaine on cells at different stages of adult HCal neurogenesis within bLR vs. bHR rats. Distinct cellular markers signified date of birth, thus serving as an index of cocaine's effects on recent cell generation, early cell survival and late cell survival at the time of killing the animals.

To determine the rate of recent cell proliferation at the time of animals' killing, we performed IHC labeling of Ki-67 (Figures 2 and 3), an intrinsic marker of ongoing cell proliferation present during the active phases of the cell cycle (G_1 -M) (Scholzen and Gerdes, 2000; Kee et al., 2002). As described in detail by Perez et al., 2009, sections were postfixed in 4% paraformaldehyde (PFA) for 1 h, followed by 1 h incubation in 10% sodium citrate pH 6.0 at 90°C for epitope retrieval. Sections were then rinsed with PBS, washed in 0.3% peroxide, blocked with BSA containing 1% goat serum and 0.05% Triton X-100, and incubated overnight with polyclonal rabbit anti-Ki-67 (1:40,000; University of Michigan). After PBS washes, sections were incubated in biotinylated anti-rabbit secondary antibody 1:1,000 (Vector Laboratories, Burlingame, CA) followed by Avidin/Biotin complex (Vectastain Elite ABC kit; Vectors Laboratories) and diaminobenzidine (DAB) as chromogen.

The survival of developing neurons (cells born during the course of the sensitization process) was visualized with the marker NeuroD (Figure 3), a transcription factor with a transient expression restricted to cells as they become committed to a neuronal lineage during differentiation within the DG in the HC (Lee et al., 1997; Miyata et al., 1999; Amrein

et al., 2004; Hevner et al., 2006). Sections were postfixed in 4% PFA, incubated in endogenous peroxidase blocking reagent (0.1% peroxide) and blocked with BSA containing 1% donkey serum and 0.05% Triton X-100. Sections were then incubated overnight with goat polyclonal anti-NeuroD (1:2,500; Santa Cruz Biotechnology). The next day, following PBS washes, sections were incubated in biotinylated anti-goat secondary antibody 1:1,000 (Vector Labs) followed by Avidin/Biotin complex and DAB reaction with Nickel Chloride for visualization of signal.

The survival of cells born just prior to the sensitization process (rate of late cell survival) was determined by labeling of BrdU (Figure 3), a synthetic thymidine analog that was injected before cocaine administration. BrdU is incorporated during the S-phase of the cell cycle into newly synthesized DNA strands of actively proliferating cells (Wojtowicz and Kee, 2006), and the number of these new cells that survive throughout the treatment can then be measured at the end of the study. Tissue was postfixed in 4% PFA for 1 h, rinsed in PBS and washed in 0.3% peroxide. Sections were then incubated in 50% formamide—2× SSC at 65°C for 2 h followed by rinses in 2X SSC. Slides were placed for 30 min in 2N HCl at 37°C for DNA denaturation and 10 min in 0.1 M boric acid at room temperature followed by rinsing in PBS and blocking with BSA. Sections were incubated overnight with polyclonal anti-BrdU (1:20,000; University of Michigan) followed by anti-rabbit secondary antibody 1:1,000 (Vector labs), Avidin/Biotin complex amplification and DAB reaction for visualization of signal.

For all antibodies, controls were performed by blocking the specific signal with the respective blocking peptide (data not shown). Cresyl violet staining was performed for Ki-67 and BrdU protocols. All sections were dehydrated through graded alcohols followed by immersion in xylene and then coverslipped with Permount® mounting medium.

For determining the effects of cocaine treatments on cell differentiation (phenotype of surviving cells) in the DG of the HC, we used NeuN antibody to identify mature neurons (Mullen et al., 1992) and GFAP antibody to label mature glial cells. Briefly, for fluorescent triple labeling, following the same protocol as for BrdU labeling we first incubated the sections overnight with a combination of polyclonal anti-BrdU (1:20,000; University of Michigan) with monoclonal anti-GFAP (1:1,000; Millipore Bioscience Research Reagents). The next day the sections were incubated with fluorescent secondary antibodies: Alexa 594 goat anti-rat and Alexa-647 goat anti-mouse (Invitrogen). For NeuN labeling, we used a primary monoclonal mouse anti-NeuN antibody tagged with an Alexa 488 fluor (Millipore Bioscience Research Reagents).

Cell counting

To ensure that a blind observer performed all cell counts for quantifying Ki-67, NeuroD and BrdU-positive cells, slides were initially coded and the code was not broken until counts were analyzed. Thus, immunostained positive cells were counted with a Leica DMR light microscope within the DG focusing through the thickness of the section using a 63× oil objective. We used a modified unbiased stereological procedure (Malberg and Duman, 2003) in which every 8th section was counted throughout the extent of the HC, and the sum was multiplied by the sampling factor 8 to provide an estimate of the total number of

positive cells per animal. In triple labeling experiments, in order to determine the percentage of BrdU positive cells that co-labeled with NeuN or GFAP, we based our quantification methodology on previous studies which followed the same procedures and obtained reliable measures (Warner-Smith and Duman, 2007; Perez et al., 2009) and examined at least 30 BrdU-positive cells per subject using a laser scanning confocal microscope (Olympus BX62). Laser scans of 0.5 μ m serial Z-section planes were visualized using a 63 $\times$ oil objective. To obtain an estimate of the total number of new neurons and new glial cells in the DG of each animal, the percentage of BrdU-positive cells co-labeled with NeuN, GFAP or neither was calculated and multiplied by the total number of BrdU-positive cells for each animal.

Data and statistical analysis

All data were analyzed with GraphPad Prism, Version 5. Results are expressed as mean values $\pm$ standard error of the mean (SEM). One-way or two-way ANOVA followed by a multiple comparison test (Bonferroni) and Student's two-tailed *t*-test were used for the statistical evaluations. The level of significance was chosen as $p < 0.05$.

Drugs and chemicals

Cocaine-HCl was obtained from Mallinckrodt Inc (St. Louis, MO) and BrdU (5-Bromo-2'-deoxyuridine) was purchased from Calbiochem. Other materials were purchased from Santa Cruz Biotechnology, Sigma-Aldrich and Vector Laboratories.

Results

Study I: Characterization of bLR and bHR rats

1. bLR/bHR differences in locomotor response to novelty—As shown in Figure 1A, selective breeding across several generations (F1-F20) produced a line of bHR rats, relative to bLR rats, that show high levels of locomotion activity (lateral movement, rearing, and total activity which is the sum of lateral and rearing movement) when placed in a novel environment. By the 10th generation of breeding, we could predict with over 95% certainty the bLR or bHR classification of our animals based on an animals' parentage (i.e. 95% of bLR offspring behaved reliably as bLRs, and 95% of bHR offspring showed typical bHR behavior). Baseline differences in novelty-induced locomotor activity dramatically diverged with increasing rounds of selection (Figure 1A). Note that for the present set of experiments, we used bLR/bHR rats from generation 15 and 20 (F15, F20) which clearly show marked basal differences in novelty-induced motor activity, both lateral movement and rearing.

2. bLR/bHR differences in psychomotor sensitization to cocaine—As mentioned earlier, following bLR/bHR phenotype characterization (Figure 1A), we used 7 pairs of adult bLR and bHR males from the 20th generation to ascertain bLR/bHR differences in psychomotor sensitization to cocaine. Figure 1B shows the effects of repeated cocaine administration and 14 days of cocaine abstinence on locomotor activity on test days 1, 7 and 21 in bLR and bHR rats. To analyze bLR and bHR differences in psychomotor response to cocaine we first calculated the background levels of total locomotion (last 30 minutes of habituation) for bLR (day 1: 29 $\pm$ 9; day 7: 14 $\pm$ 9; day 21: 56 $\pm$ 5) and bHR (day 1: 494 $\pm$ 53;

day 7: 390 ± 64 ; day 21: 530 ± 79) rats. We then used these background locomotion values to correct the values post cocaine stimulation for each phenotype providing the graphs shown in Figure 1B. The same was done for lateral (bLR, day 1: 11 ± 4 ; day 7: 4 ± 2 ; day 21: 20 ± 3 , and bHR, day 1: 150 ± 23 ; day 7: 120 ± 25 ; day 21: 145 ± 26) and rearing locomotion (bLR, day 1: 18 ± 6 ; day 7: 10 ± 7 ; day 21: 36 ± 4 , and bHR, day 1: 345 ± 38 ; day 7: 270 ± 47 ; day 21: 385 ± 63). When analyzing total locomotion, two-way ANOVA followed by Bonferroni's *post-hoc* test revealed a significant effect of day ($F_{2,30}=12.05$, $p=0.0001$) and group ($F_{1,30}=11.36$, $p=0.0021$) with no interaction day $\times$ group ($F_{2,30}=0.54$, $p=0.589$). *Post-hoc* comparisons revealed a significant difference between groups after the challenge dose at day 21 ($p<0.05$), with bHR rats showing increased total locomotion compared to bLR rats. When lateral and rearing locomotion were analyzed separately, two-way ANOVA revealed the following differences: lateral locomotion (effect of day, $F_{2,30}=9.09$, $p=0.0008$; no group effect, $F_{1,30}=0.66$, $p=0.424$; no interaction, $F_{2,30}=0.01$, $p=0.989$), and rearing locomotion (no day effect, $F_{2,30}=2.23$, $p=0.124$; effect of group, $F_{1,30}=11.10$, $p=0.0023$; no interaction, $F_{2,30}=0.63$, $p=0.538$). *Post-hoc* comparisons revealed a significant difference between groups only for rearing locomotion after the challenge dose at day 21 ($p<0.05$), with bHR rats showing increased rearing locomotion compared to bLR rats. These results indicate that although both groups of animals exhibited a certain degree of psychomotor sensitization to cocaine (e.g. day 1 versus day 21 for each phenotype both when analyzing total and lateral locomotion), the enhanced drug response seen in bHRs is mainly driven by differences in rearing locomotion, a hallmark of sensitization development (Szumlinski et al., 2003).

Study II: bLR/bHR differences in neurogenesis following experimenter-administered cocaine

1. bLR/bHR differences in cell proliferation following experimenter-administered cocaine

The effects of experimenter-administered cocaine on recent cell proliferation (within a cell cycle time of 25 h; Cameron and McKay, 2001) as measured by Ki-67-positive cells in the DG was ascertained in bLR/bHR rats (F15, $n=5$ per phenotype per experimental group, see Figure 2A). Note that two groups —Vehicle and Vehicle Abstinence— were pooled as a single control group due to the lack of significant differences in Ki-67 positive cell number between them (*t*-test: bLR, $F_{1,8}=10.46$, $p=0.019$, n.s.; bHR, $F_{1,8}=1.76$, $p=0.26$, n.s.). As shown in Figure 2B-C, one-way ANOVA detected a significant effect of cocaine treatment only in bLR rats (ANOVA followed by Bonferroni's test: LR rats: $F_{3,21}=3.247$, $p=0.04$; HR rats: $F_{3,21}=0.855$, $p=0.480$, n.s.). Acute cocaine treatment did not alter cell proliferation (bLR rats: $16 \pm 12\%$ increase, n.s.; bHR rats: $1 \pm 14\%$ change, n.s.). However, after chronic cocaine treatment, cell proliferation was decreased for both phenotypes, although it reached significance only in the bLR animals (bLR rats: $32 \pm 11\%$ decrease, $p<0.05$ when compared with acute cocaine; bHR rats: $16 \pm 8\%$ decrease, n.s.). This effect, although it lost significance, persisted after 3 days of abstinence (bLR rats: $18 \pm 14\%$ decrease, n.s.; bHR rats: $20 \pm 12\%$ decrease, n.s.). The results suggest that bLR and bHR animals exhibit different patterns of cell proliferation as measured by Ki-67 cells in the DG, with bLR rats being more reactive to the effects of cocaine on recent cell proliferation.

2. bLR/bHR differences in neurogenesis following experimenter-administered cocaine

bLR/bHR rats from the second cohort of animals (F20, $n=6$ per phenotype per

experimental group) were used to examine the effects of chronic cocaine exposure and 14 days cocaine abstinence on adult HCal neurogenesis. As shown in Figure 3 and described above, the three measures quantified were: a) cells that were recently born at the time of killing the animals (in the last 25 h; recent cell proliferation); b) cells born midway through the cocaine sensitization process (early cell survival); and c) cells born at the beginning of the experiment (~14-28 days old; late cell survival).

Before analyzing the effects of cocaine on different stages of neurogenesis, the basal rates of cells marked for each phenotype were compared between vehicle-treated bLR and bHR groups. It should be noted that as previously described in Perez et al. (2009), bLR/bHR rats showed similar basal rates of cell proliferation as measured by Ki-67 IHC ($6\pm 9\%$ difference; bLR: $8,088\pm 547$ versus bHR: $8,544\pm 748$; $p=0.637$), and a non-significant tendency to exhibit different basal rates of early ($19\pm 7\%$ difference; bLR: $26,648\pm 1901$ versus bHR: $31,653\pm 1766$; $p=0.08$) and late cell survival ($16\pm 7\%$ difference; bLR: $10,547\pm 675$ versus bHR: $12,229\pm 705$; $p=0.116$). The combination of data from the Ki-67, NeuroD and BrdU markers (see Figure 3C-H) suggests that bLR and bHR lines tend to have some basal differences in the turnover of adult-born cells in the HC, with bLRs showing similar birth rates but lower rates of survival (Perez et al., 2009).

When studying the effects of cocaine treatments on recent cell proliferation in bLR rats, Ki-67 immunostaining revealed that chronic cocaine reduced the number of Ki-67-positive cells in the DG by $26\pm 5\%$ relative to the bLR vehicle treatment group, and this effect was even greater ($44\pm 6\%$ decrease) after 14 days of abstinence (ANOVA: $F_{2,15}=13.52$, $p=0.0004$; Figure 3C). For bHR rats, one-way ANOVA did not reveal significant changes ($F_{2,15}=3.072$, $p=0.08$, Figure 3D) in Ki-67 cell number in the DG following chronic cocaine ($1\pm 12\%$ change) when compared to the bHR vehicle treatment group. However, although it did not reach statistical significance, following two weeks of cocaine abstinence, recent cell proliferation was decreased by $26\pm 5\%$ in the DG of bHR rats. These results strengthen the previous findings (F15 generation, Figure 2B) and suggest that following chronic cocaine treatments, bLR and bHR animals exhibit different patterns of recent cell proliferation as measured by Ki-67 cell number in the DG, with bLR rats showing a greater effect of cocaine on the rate of cell birth (decreased after chronic cocaine and abstinence). Taken together, the results show that chronic cocaine exposure has a negative impact on recent cell proliferation in the DG of bLR rats.

To study the effects of experimenter-administered cocaine on the survival of cells born midway through the sensitization process (see Figure 3A), we examined NeuroD, a neuronal marker transiently expressed by immature neurons (approximately 4 to 10 days of age). As shown in Figure 3E-F, quantitative analysis of NeuroD-positive cells revealed no significant change in the number of young immature neurons in the DG following cocaine treatments in bLR/bHR rats (ANOVA followed by Bonferroni's test: bLR rats: $F_{2,15}=1.6$, $p=0.234$, n.s.; bHR rats: $F_{2,15}=0.350$, $p=0.71$, n.s.). Therefore, these results suggest that experimenter administered cocaine did not alter the survival of cells that were already born (young immature neurons) at the time of killing the animals. This lack of effect suggests that less mature cells may not respond to cocaine treatment because they are a resistant cell population.

In order to study the effects of cocaine treatments on the survival of cells born at the beginning of the experiment (late phase of cell survival), BrdU (50 mg/kg, i.p., twice a day for 3 days) was administered under basal conditions before the animals were exposed to vehicle or cocaine (see Figure 3B; based on Thomas et al., 2007). The animals were killed either 14 or 28 days after BrdU pretreatment. Thus, determination of BrdU-positive cells in the DG yields a measure of the impact of chronic exposure to cocaine, with or without subsequent abstinence, on the survival rate of the cells born at the time of BrdU labeling.

For bLR rats, Figure 3G shows that BrdU incorporation in the DG did not differ significantly between treatment groups ($F_{2,15}=0.858$, $p=0.444$), suggesting that in bLRs, cocaine did not compromise the viability of the cells that were generated in the DG prior to treatment and then exposed to cocaine. Interestingly, for bHR rats, we found that the number of surviving BrdU positive cells was reduced by $30\pm 8\%$ after 2 weeks of abstinence (28 days after BrdU labeling) following chronic cocaine treatments ($F_{2,15}=7.256$, $p<0.01$ vs. vehicle, $p<0.05$ vs. chronic cocaine). Thus, bLR and bHR animals exhibited different patterns of cell survival as measured by BrdU cell in the DG, with bHR rats showing a main effect on the late phase of survival of cells that were in the final stages of maturation during the course of cocaine abstinence.

Given the differential response in late cell survival following cocaine abstinence for bLR and bHR rats, we evaluated the pattern of cell differentiation of the surviving BrdU-labeled cells and obtained an estimate of the total number of neurons and astrocytes generated in the DG. As detailed in Figure 4A, injections of BrdU were administered 28 days before animals were killed, providing us with a measure of the survival rate of the cells born at the time of BrdU labeling and showing us possible effects of cocaine abstinence on basal neurogenesis. Such a protocol allows the follow-up of newborn cells with the analysis of their maturation dynamics and the process of phenotype acquisition. Figure 4B illustrates that the large majority of newborn BrdU positive cells in the DG expressed the neuronal marker NeuN (bLR, $91.7\pm 2.6\%$; bHR, $92.9\pm 1.4\%$), whereas a minority was either double-labeled with GFAP, an astroglial marker (bLR, $3.8\pm 1.7\%$; bHR, $4.6\pm 1.4\%$), or not identified (bLR, $4.6\pm 1.5\%$; bHR, $2.5\pm 0.9\%$). These ratios were not significantly different between groups (NeuN, $F_{1,10}=1.088$, $p=0.766$; GFAP, $F_{1,10}=1.440$, $p=0.585$; Neither, $F_{1,10}=2.531$, $p=0.215$). These results show that the difference in late cell survival shown by bLR and bHR rats following cocaine abstinence did not correlate with a change in their maturation dynamics, as most of the surviving cells expressed the neuronal phenotype. It is therefore tempting to speculate that the decrease in late cell survival (mature cells, 28 days old) observed for bHR rats might translate in a decrease in mature neurons in the DG.

Discussion

The present findings provide a detailed characterization of cocaine's effect on adult hippocampal neurogenesis in rats selectively-bred for low (bLR) vs. high (bHR) novelty-seeking behavior. In general, cocaine decreased HCal neurogenesis in both groups but at different control points. bLR rats primarily showed an effect of chronic cocaine and cocaine abstinence on cell proliferation, with no apparent impact on the survival or differentiation of preexisting adult stem cells. Moreover, this effect appears to become amplified over time,

exhibiting a steady decrease in proliferation from acute to chronic cocaine exposure, and from chronic exposure to abstinence. By contrast, bHR rats exhibited abstinence-specific changes in the late survival of cells that were born under basal conditions before the cocaine treatment was initiated.

Specifically, the results for Study I regarding the characterization of the bLR/bHR animal model showed that: (1) selective breeding for the novelty-seeking trait across generations produced a line of bHR rats that show high levels of locomotion compared to their bLR counterparts. (2) The enhanced psychomotor response to cocaine seen in bHR rats is mainly driven by differences in rearing locomotion, a hallmark of sensitization development. For Study II, regarding the differential effects of cocaine on adult HCal neurogenesis in bLR/bHR rats, we showed that: (1) acute cocaine administration did not alter recent cell proliferation as measured by Ki-67-positive cells in the DG of either bLR or bHR rats. (2) Repeated cocaine administration induced a decrease in cell proliferation, an effect only seen in bLR rats which persisted and became amplified through the course of abstinence (14 days). This effect was replicated in two separate generations of bLR/bHR rats (F15 and F20) in separate sets of experiments. (3) Neither chronic cocaine administration nor cocaine abstinence affected the survival of cells born midway through the sensitization process and expressing a marker of immature neurons (NeuroD) both in bLR and bHR rats. (4) Cocaine abstinence was associated with decreased survival of adult mature neurons specifically in bHR rats. These cells were born under basal conditions before the cocaine treatment started and were 28 days old when the animals were killed. (5) The course of treatment did not affect the ratio of neurons to glia in either bLR or bHR, with most of the surviving cells acquiring the neuronal phenotype.

The LR-HR rat model provides a useful tool to examine potential molecular mechanisms that underlie their differential behavioral response to cocaine, although, notably, the LR/HR trait does not appear to predict the transition to compulsive drug use (see Belin et al., 2008). Among other differences in drug-seeking behavior, commercially available (non-selectively-bred) HR rats will self-administer psychostimulants more readily than their LR counterparts (Piazza et al., 1989; Piazza and Le Moal, 1996) and will display exaggerated behavioral sensitization to these drugs (Hooks et al., 1991; 1992). Consistent with these studies in commercially-available LR/HR rats, selective-breeding has amplified many of the differences seen in commercially-available LR/HR animals that appear directly relevant to the enhanced differences in initial propensity to drug abuse. In this line, bHRs acquired cocaine self-administration more rapidly than bLR rats (Davis et al., 2008). Moreover, a recent study from our lab (Waselus et al., 2009) examined multiple measures indicative of behavioral sensitization (i.e., locomotion, darting, immobility, left-right head movements and rearing) to identify differences between cocaine-treated bHR and bLR rats. Similar to our results, the study demonstrated that while both phenotypes showed evidence of sensitization, bHRs exhibited an enhanced drug response. Along these lines, our differences in motor response were mainly driven by changes in rearing locomotion, which has been described as a landmark of cocaine sensitization (Szumlinski et al., 2003).

After demonstrating the bLR/bHR differences in cocaine-induced psychomotor sensitization we examined adult HCal neurogenesis as a potential molecular mechanism differentially

altered by cocaine within each phenotype. Note that individual correlations between the behavioral (i.e., psychomotor response to cocaine) and the molecular data (i.e., neurogenesis) were not assessed because these experiments were performed in separate groups of rats. However, the experiments were run in parallel to ensure that brains assessed for possible neurogenesis changes were exposed to an identical cocaine treatment regimen known to evoke psychomotor differences in bLR/bHR rats.

When studying neurogenesis, it is critical to establish adequate protocols to appropriately distinguish dividing cells such that cell proliferation and cell survival can be clearly differentiated. The present study evaluated the impact of cocaine on cells born at different time points, i.e., cells at different stages of maturation along the neurogenesis process when the animals were killed. Ki-67 was used to mark recently divided cells (Scholzen and Gerdes, 2000; Kee et al., 2002); NeuroD indicated immature young neurons (von Bohlen, 2007); and BrdU demarcated mature cells (Dayer et al., 2003; Thomas et al., 2007) that differentiated into glial, neuronal or an undefined cell type. Remarkably, it has been recently described that Ki-67 and NeuroD do not colocalize in the same population of cells (Larsen et al., 2007), and that Ki-67 and BrdU only colocalize from 2 h until 4 days post BrdU injection (Dayer et al., 2003). This indicates that new BrdU-labeled daughter cells are added throughout this period of approximately four cell cycles (see Figure 3 for experimental design), and that cell proliferation and survival can be clearly differentiated using these two molecular markers. Moreover, as no co-localization of NeuroD was observed following 14 and 28 days of BrdU injections (Kawai et al., 2004), these molecular markers allowed us to study the effects of early and late cell survival separately.

Our results showed that for bLR rats, repetitive but not single administration of cocaine decreased cell proliferation as measured by Ki-67-positive cells in adult rat HC without influencing cell survival and cell differentiation. Thus, experimenter-administered cocaine in bLR rats affected different cell populations depending on their stage of maturity along the neurogenesis process; specifically, cocaine appears to decrease the rate of cell birth and proliferation without altering the maturation of cells that were already born at the time of killing the animals (i.e. NeuroD-positive immature neurons and BrdU-positive mature neurons). Similar results were previously described for commercially-available Sprague-Dawley rats (Yamaguchi et al., 2004; Dominguez-Escriba et al., 2006). Indeed, newborn cells could be undergoing plasticity changes before they reach full maturity, and their functions may well be different from those of mature granule neurons (Canales, 2007). It is believed that the adult DG only has a limited supply of dividing progenitor cells (von Bohlen and Halbach, 2007); thus, decreased cell proliferation might reflect a direct action of cocaine on the progenitor cell population (Eisch and Harburg, 2006), leading to a cessation, reduction, or otherwise alteration of cell cycle division, or perhaps enhanced cell death. Furthermore, decreased cell proliferation might also lead to a delayed reduction of cell survival, i.e., less cell birth giving rise to a lesser number of mature neurons. Or, as described by Noonan et al., 2008, the effects of cocaine administration on cell proliferation may become evident in a later change in their maturity. Future experiments will address these hypotheses by administering BrdU immediately before the end of the abstinence period and assessing cocaine effects on cell maturation and cell survival (14-28 days later).

On the other end, for bHR rats, experimenter administration of cocaine (acute and chronic) did not affect the rate of recent cell proliferation (cell birth during the last cell cycle, 25 h), and did not influence the survival of cells born halfway (immature neurons) or at the beginning of the sensitization process (mature neurons). However, during the course of abstinence, there were specific changes in bHR rats, i.e., a 26% non-significant decrease in the rate of cell birth but most importantly, a decrease in the number of mature surviving neurons when compared to vehicle or chronic cocaine groups. Notably, the survival of immature cells labeled with NeuroD was not compromised following cocaine abstinence. It has previously been shown that distinct populations of cells in the adult DG will respond to certain stimulus depending on the cell's age at the time of exposure to the stimulus (Cameron and Gould, 1996). An important thing to remember when studying the regulation of neurogenesis is that the generation of new neurons is the end product of a series of steps consisting of cell proliferation, survival, migration, differentiation, and establishment of functional connections with other neurons (Lledo et al., 2006). Having this in mind, the results showed that during abstinence to chronic cocaine, bHR rats appeared to have a critical window of neurogenesis regulation from immature neurons to mature integrated neurons in which the number of surviving mature neurons is decreased, possibly due to more cell death. The critical time point in postnatal cell genesis when the decision toward neuronal differentiation is made and the crucial restriction "point of no return" is reached remains unclear (von Bohlen and Halbach, 2007). Our results suggest that in the context of cocaine sensitization, in bHR rats, the critical window of neurogenesis regulation occurs during abstinence at the level of decreasing the number of functionally mature neurons. Interestingly, this window of decreased neurogenesis regulation parallels the time when bLR and bHR rats show distinct psychomotor responses to cocaine, with bHR rats exhibiting greater drug-induced rearing locomotion. Thus, it is reasonable to hypothesize that the regulation of adult HCal neurogenesis may be involved in the emergence of certain symptoms manifested during psychomotor sensitization. More importantly, our results support the idea previously described by Chamber and Taylor (2004), which associated an impairment of HCal function with a higher behavioral sensitization to cocaine. In the present study, the impairment of HCal function was manifested as a decrease in the number of mature neurons (lesser connections in the circuitry) in conjunction with a greater psychomotor response to cocaine in bHR rats. Other studies also showed that adult neurogenesis plays an important role in HCal function (Cameron and McKay, 2001), and specifically in drug-taking and drug-seeking behaviors via its projections to the reward-related limbic regions (Taepavarapruk et al., 2000; Canales, 2007).

To conclude, the present results suggest that bLR and bHR rat brains react differently to drugs of abuse, with each selectively-bred line exhibiting a distinct psychomotor response to cocaine and showing different patterns and time-points of neurogenesis regulation following experimenter-administered cocaine. Indeed, cocaine targets different populations of cells in bLR/bHR rats: the progenitor neural stem cells causing decreased birth of new cells in bLR rats, and the number of mature neurons in bHR rats. Several factors could account for this differential regulation. For example, the absence of bLR/bHR behavioral differences after chronic cocaine and their obvious difference in cell proliferation could be due to a disconnection between these factors and a connection between other factors known to differ

in bLR/bHR (e.g., Perez et al., 2009) and the manifestation of changes in cell proliferation. Another important factor to consider involves a matter of timing; less cell birth in bLR rats will eventually lead to decreased number of mature neurons, suggesting a delayed effect over neurogenesis regulation and reaffirming the delayed or blunted response to drugs of abuse apparent for bLR vs. bHR animals. The exact role of new neurons in adult HCal function remains unknown, but it is tempting to speculate that the earlier decrease in new functional connections in bHR rats could contribute at least in part, to their enhanced psychomotor response following chronic cocaine administration. These results support the growing literature regarding altered neurogenesis as one drug-regulated form of plasticity. Moreover, the use of selective breeding highlights the importance of identifying the neuroadaptations that might confer enhanced drug responses, because they might help develop novel and better-tailored treatment options. Thus, in future studies it will be of great interest to ascertain how neurogenesis is differentially regulated in bLR and bHR rats when exposed to a paradigm of long-term addiction (i.e., extended access cocaine self-administration).

Acknowledgments

This study was supported by NIDA: 5P01DA021633-02; NIMH Conte Center Grant #L99MH60398; Office of Naval Research (ONR) N00014-02-1-0879 to HA & SJW; and The Pritzker Neuropsychiatric Research Foundation. The authors would like to thank Sharon Burke, Jennifer Fitzpatrick and James Beals for excellent technical assistance.

Abbreviations

bLR	bred Low-Responder
bHR	bred High-Responder
BrdU	5-bromo-2'-deoxyuridine
BSA	Bovine Serum Albumina
DG	Dentate Gyrus
HC	Hippocampus
HCal	Hippocampal
IHC	Immunohistochemistry
PFA	paraformaldehyde

References

- Amrein I, Slomianka L, Poletaeva II, Bologova NV, Lipp HP. Marked species and age-dependent differences in cell proliferation and neurogenesis in the hippocampus of wild-living rodents. *Hippocampus*. 2004; 14(8):1000–1010. [PubMed: 15390172]
- Belin D, Mar AC, Dalley JW, Robbins TW, Everitt BJ. High impulsivity predicts the switch to compulsive cocaine-taking. *Science*. 2008; 320(5881):1352–1355. [PubMed: 18535246]
- Brown JP, Couillard-Despres S, Cooper-Kuhn CM, Winkler J, Aigner L, Kuhn HG. Transient expression of doublecortin during adult neurogenesis. *J Comp Neurol*. 2003; 467(1):1–10. [PubMed: 14574675]

- Cameron HA, Gould E. Distinct populations of cells in the adult dentate gyrus undergo mitosis or apoptosis in response to adrenalectomy. *J Comp Neurol.* 1996; 369(1):56–63. [PubMed: 8723702]
- Cameron HA, McKay RD. Adult neurogenesis produces a large pool of new granule cells in the dentate gyrus. *J Comp Neurol.* 2001; 435(4):406–417. [PubMed: 11406822]
- Canales JJ. Adult neurogenesis and the memories of drug addiction. *Eur Arch Psychiatry Clin Neurosci.* 2007; 257(5):261–270. [PubMed: 17401727]
- Chambers RA, Taylor JR. Animal modeling dual diagnosis schizophrenia: Sensitization to cocaine in rats with neonatal ventral hippocampal lesions. *Biol Psychiatry.* 2004; 56(5):308–316. [PubMed: 15336512]
- Davis BA, Clinton SM, Akil H, Becker JB. The effects of novelty-seeking phenotypes and sex differences on acquisition of cocaine self-administration in selectively-bred High-Responder and Low-Responder rats. *Pharmacol Biochem Behav.* 2008; 90(3):331–338. [PubMed: 18445506]
- Dayer AG, Ford AA, Cleaver KM, Yassaee M, Cameron HA. Short-term and long-term survival of new neurons in the rat dentate gyrus. *J Comp Neurol.* 2003; 460(4):563–572. [PubMed: 12717714]
- Dominguez-Escriba L, Hernandez-Rabaza V, Soriano-Navarro M, Barcia JA, Romero FJ, Garcia-Verdugo JM, et al. Chronic cocaine exposure impairs progenitor proliferation but spares survival and maturation of neural precursors in adult rat dentate gyrus. *Eur J Neurosci.* 2006; 24(2):586–594. [PubMed: 16903860]
- Eisch AJ, Harburg GC. Opiates, psychostimulants, and adult hippocampal neurogenesis: Insights for addiction and stem cell biology. *Hippocampus.* 2006; 16(3):271–286. [PubMed: 16411230]
- García-Fuster MJ, Clinton SM, Watson SJ, Akil H. Effect of cocaine on Fas-associated protein with death domain in the rat brain: individual differences in a model of differential vulnerability to drug abuse. *Neuropsychopharmacology.* 2009; 34(5):1123–1134. [PubMed: 18580876]
- Gold LH, Geyer MA, Koob GF. Neurochemical mechanisms involved in behavioral effects of amphetamines and related designer drugs. *NIDA Res Monogr.* 1989; 94:101–126. [PubMed: 2514360]
- Gosnell BA. Sucrose intake enhances behavioral sensitization produced by cocaine. *Brain Res.* 2005; 1031(2):194–201. [PubMed: 15649444]
- Hevner RF, Hodge RD, Daza RA, Englund C. Transcription factors in glutamatergic neurogenesis: conserved programs in neocortex, cerebellum, and adult hippocampus. *Neurosci Res.* 2006; 55(3):223–233. [PubMed: 16621079]
- Hodge RD, Kowalczyk TD, Wolf SA, Encinas JM, Rippey C, Enikolopov G, et al. Intermediate progenitors in adult hippocampal neurogenesis: Tbr2 expression and coordinate regulation of neuronal output. *J Neurosci.* 2008; 28(14):3707–3717. [PubMed: 18385329]
- Hooks MS, Jones GH, Neill DB, Justice JB Jr. Individual differences in locomotor activity and sensitization. *Pharmacol Biochem Behav.* 1992; 38(2):467–470. [PubMed: 2057515]
- Hooks MS, Jones GH, Smith AD, Neill DB, Justice JB Jr. Individual differences in amphetamine sensitization: dose-dependent effects. *Pharmacol Biochem Behav.* 1991; 41(1):203–210. [PubMed: 1539070]
- Kabbaj M. Individual differences in vulnerability to drug abuse: the high responders/low responders model. *CNS Neurol Disord Drug Targets.* 2006; 5(5):513–520. [PubMed: 17073654]
- Kabbaj M, Devine DP, Savage VR, Akil H. Neurobiological correlates of individual differences in novelty-seeking behavior in the rat: differential expression of stress-related molecules. *J Neurosci.* 2000; 20(18):6983–6988. [PubMed: 10995843]
- Kalivas PW. Neurobiology of cocaine addiction: implications for new pharmacotherapy. *Am J Addict.* 2007; 16(2):71–78. [PubMed: 17453607]
- Kawai T, Takagi N, Miyake-Takagi K, Okuyama N, Mochizuki N, Takeo S. Characterization of BrdU-positive neurons induced by transient global ischemia in adult hippocampus. *J Cereb Blood Flow Metab.* 2004; 24(5):548–555. [PubMed: 15129187]
- Kee N, Sivalingam S, Boonstra R, Wojtowicz JM. The utility of Ki-67 and BrdU as proliferative markers of adult neurogenesis. *J Neurosci Methods.* 2002; 115(1):97–105. [PubMed: 11897369]
- Kempermann G. Why new neurons? Possible functions for adult hippocampal neurogenesis. *J Neurosci.* 2002; 22(3):635–638. [PubMed: 11826092]

- Kempermann G, Wiskott L, Gage FH. Functional significance of adult neurogenesis. *Curr Opin Neurobiol.* 2004; 14(2):186–191. [PubMed: 15082323]
- Larsen MH, Rosenbrock H, Sams-Dodd F, Mikkelsen JD. Expression of brain derived neurotrophic factor, activity-regulated cytoskeleton protein mRNA, and enhancement of adult hippocampal neurogenesis in rats after sub-chronic and chronic treatment with the triple monoamine re-uptake inhibitor tesofensine. *Eur J Pharmacol.* 2007; 555(2-3):115–121. [PubMed: 17112503]
- Lee JE. Basic helix-loop-helix genes in neural development. *Curr Opin Neurobiol.* 1997; 7(1):13–20. [PubMed: 9039799]
- Lledo PM, Alonso M, Grubb MS. Adult neurogenesis and functional plasticity in neuronal circuits. *Nat Rev Neurosci.* 2006; 7(3):179–193. [PubMed: 16495940]
- Lodge DJ, Grace AA. Amphetamine activation of hippocampal drive of mesolimbic dopamine neurons: a mechanism of behavioral sensitization. *J Neurosci.* 2008; 28(31):7876–7882. [PubMed: 18667619]
- Maccari S, Piazza PV, Deminiere JM, Angelucci L, Simon H, Le Moal M. Hippocampal type I and type II corticosteroid receptor affinities are reduced in rats predisposed to develop amphetamine self-administration. *Brain Res.* 1991; 548(1-2):305–309. [PubMed: 1868340]
- Malberg JE, Duman RS. Cell proliferation in adult hippocampus is decreased by inescapable stress: reversal by fluoxetine treatment. *Neuropsychopharmacology.* 2003; 28(9):1562–1571. [PubMed: 12838272]
- Ming GL, Song H. Adult neurogenesis in the mammalian central nervous system. *Annu Rev Neurosci.* 2005; 28:223–250. [PubMed: 16022595]
- Miyata T, Maeda T, Lee JE. NeuroD is required for differentiation of the granule cells in the cerebellum and hippocampus. *Genes Dev.* 1999; 13(13):1647–1652. [PubMed: 10398678]
- Mullen RJ, Buck CR, Smith AM. NeuN, a neuronal specific nuclear protein in vertebrates. *Development.* 1992; 116(1):201–211. [PubMed: 1483388]
- Noonan MA, Choi KH, Self DW, Eisch AJ. Withdrawal from cocaine self-administration normalizes deficits in proliferation and enhances maturity of adult-generated hippocampal neurons. *J Neurosci.* 2008; 28(10):2516–2526. [PubMed: 18322096]
- Perez JA, Clinton SM, Turner CA, Watson SJ, Akil H. A new role for FGF2 as an endogenous inhibitor of anxiety. *J Neurosci.* 2009; 29(19):6379–6387. [PubMed: 19439615]
- Piazza PV, Deminiere JM, Le Moal M, Simon H. Factors that predict individual vulnerability to amphetamine self-administration. *Science.* 1989; 245(4925):1511–1513. [PubMed: 2781295]
- Piazza PV, Le Moal ML. Pathophysiological basis of vulnerability to drug abuse: role of an interaction between stress, glucocorticoids, and dopaminergic neurons. *Annu Rev Pharmacol Toxicol.* 1996; 36:359–378. [PubMed: 8725394]
- Robinson TE, Kolb B. Structural plasticity associated with exposure to drugs of abuse. *Neuropharmacology.* 2004; 47(Suppl 1):33–46. [PubMed: 15464124]
- Schmidt HD, Duman RS. The role of neurotrophic factors in adult hippocampal neurogenesis, antidepressant treatments and animal models of depressive-like behavior. *PNAS.* 2007; 104(11):4647–4652. [PubMed: 17360578]
- Scholzen T, Gerdes J. The Ki-67 protein: from the known and the unknown. *J Cell Physiol.* 2000; 182(3):311–322. [PubMed: 10653597]
- Shen F, Meredith GE, Napier TC. Amphetamine-induced place preference and conditioned motor sensitization requires activation of tyrosine kinase receptors in the hippocampus. *J Neurosci.* 2006; 26(43):11041–11051. [PubMed: 17065446]
- Stead JD, Clinton S, Neal C, Schneider J, Jama A, Miller S, et al. Selective breeding for divergence in novelty-seeking traits: heritability and enrichment in spontaneous anxiety-related behaviors. *Behav Genet.* 2006; 36(5):697–712. [PubMed: 16502134]
- Szumliński KK, Frys KA, Kalivas PW. Pretreatment with serotonin 5-HT₃ receptor antagonists produces no observable blockade of long-term motor sensitization to cocaine in rats. *Psychopharmacology (Berl).* 2003; 165:329–336. [PubMed: 12451437]
- Taepavarapruk P, Floresco SB, Phillips AG. Hyperlocomotion and increased dopamine efflux in the rat nucleus accumbens evoked by electrical stimulation of the ventral subiculum: role of ionotropic

- glutamate and dopamine D1 receptors. *Psychopharmacology (Berl)*. 2000; 151(2-3):242–251. [PubMed: 10972471]
- Thomas RM, Hotsenpiller G, Peterson DA. Acute psychosocial stress reduces cell survival in adult hippocampal neurogenesis without altering proliferation. *J Neurosci*. 2007; 27(11):2734–2743. [PubMed: 17360895]
- Turner CA, Flagel SB, Clinton SM, Akil H, Watson SJ. Cocaine interacts with the novelty-seeking trait to modulate FGFR1 gene expression in the rat. *Neurosci Lett*. 2008; 446(2-3):105–107. [PubMed: 18824081]
- von Bohlen, Halbach O. Immunohistological markers for staging neurogenesis in adult hippocampus. *Cell Tissue Res*. 2007; 329(3):409–420. [PubMed: 17541643]
- Warner-Schmidt JL, Duman RS. VEGF is an essential mediator of the neurogenic and behavioral actions of antidepressants. *Nat Protoc*. 2007; 1(3):1399–1405.
- Waselus, M.; Flagel, SB.; Jedynak, J.; Akil, H.; Robinson, TE.; Watson, SJ. Individual differences in cocaine sensitization: Effects on structural plasticity in the nucleus accumbens; Neuroscience Meeting Planner; Chicago, IL: Society for Neuroscience. 2009; Program No. 649.2Online
- Wojtowicz JM, Kee N. BrdU assay for neurogenesis in rodents. *Nat Protoc*. 2006; 1(3):1399–1405. [PubMed: 17406427]
- Yamaguchi M, Suzuki T, Seki T, Namba T, Juan R, Arai H, et al. Repetitive cocaine administration decreases neurogenesis in adult rat hippocampus. *Ann N Y Acad Sci*. 2004; 1025:351–362. [PubMed: 15542736]

FIG. 1.

Characterizing the bLR/bHR rat phenotype: Selective breeding for novelty seeking trait. **(A)** bLR/bHR differences in locomotor response to novelty across generations (F1-F20).

Differences in mean locomotion (total activity, which is the sum of lateral movement and rearing, lateral movement, rearing) between bLR and bHR animals are shown for twenty generations of selective breeding (F1-F20). Mean locomotion scores diverge dramatically between the two lines with increasing rounds of selection. All data are mean $\pm$ standard deviation. **(B)** bLR/bHR differences in psychomotor sensitization to cocaine (F20).

Experimental design: Effects of chronic cocaine administration (15 mg/kg, i.p., 7 days) on locomotor activity on day 1, 7 and 21 of testing in bLR and bHR rats (7 bLR/bHR pairs). On testing days rats were habituated for 1 h to the locomotor cage before receiving cocaine. Locomotion (total, lateral, rearing) was calculated for each rat as the locomotion scores after cocaine injection (stimulated) minus the background (last 30 minutes of habituation) (mean $\pm$ SEM). * $p < 0.05$

FIG. 2.

bLR/bHR differences in cell proliferation following experimenter-administered cocaine. **(A)** bLR and bHR rats (F15, 25 pairs, $n=5$ per group) were used to determine the effects of acute, chronic and early cocaine abstinence on adult HCal cell proliferation as measured by Ki-67 IHC in the DG. Groups of treatment: vehicle (0.9% NaCl, i.p., 7 days), acute cocaine (acute, 6 days of saline and on day 7, 15 mg/kg of cocaine), chronic cocaine (15 mg/kg for 7 days), chronic cocaine followed by abstinence (3 days), and vehicle abstinence control group. The pool of Ki-67-positive cells being labeled is indicated by a rectangle coded for each treatment group and represents the cells recently born in the last 25 h at the time of killing the animals (within a cell cycle; Cameron and McKay, 2001). **(B-C)** Chronic cocaine decreased cell proliferation only in bLR rats. Columns are means $\pm$ SEM per group and expressed as Ki-67 cell number in the DG. The groups vehicle and vehicle abstinence were pooled as a single control group due to no significant Ki-67 cell number differences between them. $*p<0.05$ acute vs. chronic cocaine. Note that there is a $32\pm11\%$ decrease in Ki-67 cell number when comparing vehicle with chronic cocaine even though this effect did not reach statistical significance.

FIG. 3.

bLR/bHR differences in adult HCal neurogenesis following experimenter-administered cocaine. **(A)** Stages of adult HCal neurogenesis showing the molecular markers used to label each pool of cells: recent cell proliferation (Ki-67, cells born in the last 25 h); early cell survival (NeuroD, ~4-10 days old immature neurons); late cell survival and cell differentiation (BrdU, ~14-28 days old mature neurons). Figure adapted from Schmidt and Duman (2007). **(B)** Experimental design: bLR/bHR rats (F20, 18 pairs, $n=6$ per group) were pretreated for 3 days with BrdU delivery (50 mg/kg, i.p.) at 12 h intervals 4 days before the treatment began. Groups of treatment: vehicle (0.9% NaCl, i.p., 7 days), chronic cocaine (15 mg/kg for 7 days), chronic cocaine followed by 14 days of abstinence. **(C)** Quantitative analysis of Ki-67 positive cells revealed a decrease in cell proliferation following chronic cocaine in bLR rats. This effect persisted during the course of abstinence. **(D)** bHR rats showed no significant change in cell proliferation following chronic cocaine. Note that two weeks of cocaine abstinence decreased cell proliferation by $26 \pm 5\%$ in the DG although it did not reach statistical significance. **(E-F)** Quantitative analysis of NeuroD-positive cells revealed no significant change in the number of short-term surviving cells in the DG following cocaine treatments. **(G)** Quantitative analysis of BrdU-positive cells revealed no significant change in the number of long-term surviving cells in the DG following cocaine

treatments in bLR rats. **(H)** bHRs rats showed cocaine abstinence-specific changes as measured by a decrease in the number of BrdU-positive cells (28 days old) in the DG. Data is presented as the mean $\pm$ SEM. ** p <0.01 vs. vehicle; Ψ p <0.05 vs. chronic cocaine.

FIG. 4.

Effect of cocaine abstinence on cell differentiation of adult-born cells in the dentate gyrus of bLR/bHR rats. **(A)** Experimental design. **(B)** Data is presented as the mean $\pm$ SEM. The large majority of the newborn BrdU-positive cells in the DG expressed the neuronal marker NeuN (bLR, 91.7 $\pm$ 2.6%; bHR, 92.9 $\pm$ 1.4%), whereas a minority was either double labeled with GFAP, an astroglial marker (bLR, 3.8 $\pm$ 1.7%; bHR, 4.6 $\pm$ 1.4%), or not identified (bLR, 4.6 $\pm$ 1.5%; bHR, 2.5 $\pm$ 0.9%). These ratios were not significantly different between groups. **(C)** Representative confocal images of BrdU (red labeling), NeuN (green labeling) and GFAP (blue labeling). The merged image, BrdU-labeled cell co-labeled with NeuN (merge) was confirmed in the x, y/x, z axis (scale bar, 10 μ m) and included 8 optical sections. Voxel size (x = 0.265 μ m/pixel; y = 0.265 μ m/pixel; z = 0.44 μ m/pixel).