CORRESPONDENCE

Headache in School Children: Prevalence and Risk Factors

by Prof. Dr. med. Andreas Straube, Prof. Dr. med. Florian Heinen, PD Dr. med. Friedrich Ebinger, Prof. Dr. med. von Rüdiger Kries in volume 48/2013

School Headache

Among school children, the anteflexion or school headache described by Gutmann in 1967 is a rather common finding if this type of headache is included in the differential diagnosis. It represents a special type of cervicocranial pain, occurring with prolonged anteflexion of the head while working at a school desk. This condition is caused by hypermobility in the craniovertebral joints with overstretching of the transverse ligament of the atlas, as demonstrated by targeted functional radiography. The affected school children are experiencing headache with increasing intensity as their school day progresses which leads to inattentiveness. Typically the pain can be triggered by forced anteflexion of the head. Physical findings include tenderness of the arch and transverse process of the atlas as well as the dens of the axis. On a reflex level, the manual therapist finds blockages at the atlanto-occipital joint. Treatment aims at eliminating these functional problems and strengthening the postural muscles with physiotherapy. Most importantly, however, prolonged anteflexion of the head should be prevented by using an oblique desk-up device when writing, reading, drawing at school and at home, and to avoid forward rolls during physical education. An oblique desk-up device can be either purchased or easily built at home. In my experience, headache syndromes can be challenging to diagnose. Therefore, functional therapy should only be started after an extensive interdisciplinary morphological workup.

However, should postural school headache be eventually diagnosed, the treatment outlined above is very effective.

DOI: 10.3238/arztebl.2014.0328a

REFERENCES

1. Lewit K, Sachse J, Janda V: Manuelle Medizin. Leipzig: Johann Ambrosius Barth 1983: 417–9.
2. Laser T: Muskelverspannung und Rückenschmerz. Stuttgart: Georg Thieme 1996.
3. Taubert K: Der Schmerzpatient in der Praxis. Balingen. Spitta Verlag GmbH 1997; 1.
4. Straube A, Heinen F, Ebinger F, von Kries R: Headache in school children: prevalence and risk factors. Dtsch Arztebl Int 2013; 110: 811–8.

Dr. med. Jürgen Fege
Weißborn/OT Berthelsdorf, Germany
juergen.fege@gmx.de

Functional Block

As an ENT specialist and manual medicine practitioner (retired), I have frequently seen children and adolescents with headache who were referred to me to exclude an underlying ENT condition. In most cases, a cervical syndrome was present, i.e. a functional block of the craniovertebral joints (phenomenon of the

jammed drawer) (1–4). In most cases, the treatment success was surprising for the parents. Unfortunately, not many health professionals are aware of the fact that children can have cervical syndromes too. Cervical syndromes are rarely diagnosed because

- children seldom complain about occipital headache
- no laboratory findings can verify this diagnosis
- radiography shows no abnormalities in the absence of osteoarthritic changes of the cervical spine
- the condition can only be diagnosed based on the manual palpation findings and the treatment success (“ex juvantibus”).

However, in their paper the authors provide very important medical history information indicative of cervical syndrome (5):

- back pain
- neck / shoulder pain
- dizziness (cervical vertigo)
- little physical exercise

In almost all patients with cervical syndrome, one side is more affected than the other because facet joints are paired joints. Consequently, one-sided complaints are strongly indicative of an underlying cervical problem. The one-sided block of paired facet joints is the only sensible explanation for one-sided and same-sided headache. The medical literature offers no other plausible explanation for one-sided chronic, recurrent headache.

Of course, there are other physical and psychological causes overlapping and modifying a cervical syndrome and therefore all the different diagnostic and therapeutic approaches have their value. But when no attention is paid to the cervicocranial joints in the diagnosis and treatment of headache in children, including school children, therapy will certainly go the wrong way.

DOI: 10.3238/arztebl.2014.0328b

REFERENCES

1. Falkenau HA: Das Zervikalsyndrom des Kindes in der Hals-Nasen-Ohrenheilkunde. HNO 1978; 26: 384–5.
2. Graf M: Manuelle Diagnostik und Therapie im Kopfgebirgsbereich. Manuelle Medizin 2007; 45: 232–41.
3. Frese A, Schilgen M, et al.: Pathophysiologie und Klinik zervikogener Kopfschmerzen. Der Schmerz 2003; 17: 125–30.
4. Lohse-Busch H: Manuelle Medizin bei kindlichen muskuloskelettalen Schmerzen. Manuelle Medizin 2002; 40: 32–40.
5. Straube A, Heinen F, Ebinger F, von Kries R: Headache in school children: prevalence and risk factors. Dtsch Arztebl Int 2013; 110: 811–8.

Dr. med. Hans Alex Falkenau
Hoya, Germany
drfalkenau@t-online.de

Considering the Eyes

In my ophthalmology practice with focus on strabology and binocular vision disorders, headache is one of the most commonly reported complaints, besides specific reading disorder/dyslexia and AD(H)D, but often it takes a targeted question to learn about it. It is so common for school children to experience headaches that they do not consider it necessary to mention them. Headaches always go along with uncorrected impairments of binocular vision in the sense of associated heterophoria.

Vision, and binocular vision in particular, is an extremely complex process, comparable with two linked autofocus cameras which have to send every second the enormous amount of 10 MB of data—in conjunction, simultaneously and for both sides identically—generated by the complex retinas to the brain, including the motor centers in the diencephalon, the nuclei that govern eye movements, the vestibular system, the cerebellum, and the autonomous nervous system. In addition, they have to react to the responses of these structures. All functions involve muscle activity: pupillomotor activity to control the amount of light and for contrast vision, the ciliary muscle for accommodation, the extraocular muscles for convergence and eye movements. Only if all of these elements are perfectly tuned up, fatigue-free fusion with stereopsis can be achieved without spending extra energy. If this is not the case, the motor deficits will shift to deeper regions and result in unnatural compensatory positions of the head with associated headache.

This is the case in many adolescent headache patients. Here, correction with glasses under cycloplegia and prismatic correction of the associated heterophoria, preferably measured using Haase's measurement and correction method along with intact fusion, can help and will lastingly stop the headaches from occurring.

With headache, it is always best to first think of the eyes, before initiating demanding neurological or even psychiatric measures.

DOI: 10.3238/arztebl.2014.0328c

REFERENCES

1. Heymann W, Köneke A, Gorzny F: Kranio-mandibuläre Dysfunktion, assoziierte Heterophorie und auditive Verarbeitungs- und Wahrnehmungsstörungen. *Manuelle Medizin* 2–20190.
2. Augustin A: *Augenheilkunde*. 3rd edition. Berlin Heidelberg: Springer Verlag 2007; 574 ff.
3. Rüssmann W, Kommerell G: Heterophorie und Asthenopie. In: Kaufmann H (ed.): *Strabismus*. 4th edition. Stuttgart: Thieme Verlag 2012; 199 ff.
4. Straube A, Heinen F, Ebinger F, von Kries R: Headache in school children: prevalence and risk factors. *Dtsch Arztebl Int* 2013; 110: 811–8.

Dr. med. Fritz Gorzny

Augenarzt, Koblenz, Germany, Vizepräsident der Internationalen Vereinigung für Binokulares Sehen (IVBS)
dr.fritz_gorzny@web.de

In Reply

In school children, primary headache is by far the most common cause of recurrent headache. However, symptomatic headaches of cervical, dental or eye-related origin have repeatedly been discussed as well.

The colleagues Dr. Fege and Dr. Falkenau draw our attention to the potential role of cervical structures in the etiology of these headaches. Besides hypermobility of the craniovertebral joints, also caused by prolonged anteflexion of the head during school hours, a functional block of these joints is regarded as an etiological factor. So far, however, there is a lack of meaningful studies evaluating this approach, one reason being that no definition of cervicogenic headache syndrome is available. Cervicogenic headache meeting to the criteria described by Sjaastad (1) has been evaluated most frequently. However, even using these criteria it is not possible to clearly define this condition, as evidenced by the significant variability in the published prevalence rates (0.17 – 13.8 %) (2, 3). It comes as no surprise that the substantially less clear criteria applied by manual therapy have not

led to a definition of a distinct syndrome which could be used in studies.

A fundamental aspect to be considered here is that the primary cause of neck pain is not necessarily related to an abnormality in this region, since the anatomical convergence of cervical and trigeminal afferences in the caudal trigeminal nucleus will result in a projection of the pain to the neck region, even if the primary activation is purely trigeminal. It is crucial to prepare and verify operationalized criteria which are independent of the examiner for potentially primary cervicogenic headache. Here, the “Who proposes it must prove it” motto has to be strictly applied. It is necessary to caution not to overinterpret pain with a maximum intensity in the neck in a one-side, monodimensional fashion as “cervicogenic”; this also applies to dizziness. It has to be taken into consideration that any type of stress will result in an unconscious increase in pericranial muscle tension (mimic as well as masticatory and neck muscles) which may trigger musculo-skeletal pain.

Dr. Gorzny highlights the potential relationship between binocular vision disorders and headache. Any impairment of binocular coordination may result in increased fusion work in the sense of latent, still compensated heterophoria which, on the one hand, may give rise to early fatigue and, on the other hand, lead to intermittent double vision; this also applies to very different refractive anomalies. Here, the aim should be to achieve the best possible correction. Reliable epidemiological studies about the occurrence of heterophoria and headache are not available. In addition, the published prevalence rates for heterophoria in the general population vary widely (1–6%, up to 70% max.) (4). Furthermore, chronic daily headache is not as common among school children as in adults and the majority of those affected suffer from paroxysmal headache, a fact that cannot easily be explained by chronic fusion abnormalities. In cases of chronic daily headache, strabological investigations should be considered.

The take-home message is – as long as attention is paid to the red flags, i.e. headaches with acute onset or acute changes in character and accompanying neurological symptoms or signs of inflammation – not to overdiagnose or overinterpret alleged “organic” causes as this will impede the path to successful treatment based on the available guidelines.

DOI: 10.3238/arztebl.2014.0329

REFERENCES

1. Sjaastad O, Fredriksen TA, Pfaffenrath V: Cervicogenic headache: diagnostic criteria. The Cervicogenic Headache International Study Group. *Headache* 1998; 38: 442–5.
2. Knackstedt H, Bansevicius D, Aaseth K, Grande RB, Lundqvist C, Russell MB: Cervicogenic headache in the general population: the Akershus study of chronic headache. *Cephalalgia* 2010; 30: 1468–76.
3. Pfaffenrath V, Kaube H: Diagnostics of cervicogenic headache. *Funct Neurol* 1990; 5: 159–64.
4. Williams C, Northstone K, Howard M, Harvey I, Harrad RA, Sparrow JM: Prevalence and risk factors for common vision problems in children: data from the ALSPAC study. *Br J Ophthalmol* 2008; 92: 959–64.
5. Straube A, Heinen F, Ebinger F, von Kries R: Headache in school children: prevalence and risk factors. *Dtsch Arztebl Int* 2013; 110: 811–8.

Prof. Dr. med. Andreas Straube

Neurologie, Campus Großhadern
Universität München, Germany
astraube@nefo.med.uni-muenchen.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.