Published in final edited form as:

J Clin Psychiatry. 2011 February ; 72(2): 175–182. doi:10.4088/JCP.09m05407blu.

Lifetime prevalence of mood and anxiety disorders in fragile X premutation carriers

James A. Bourgeois, OD MD¹, Andreea L. Seritan, MD¹, E. Melina Casillas, BA^{2,4}, David Hessel, PhD^{1,2}, Andrea Schneider, PhD^{1,2}, Ying Yang, MS³, Inderjeet Kaur, MBBS², Jennifer B. Cogswell, BA^{2,4}, Danh V. Nguyen, PhD³, and Randi J. Hagerman, MD^{2,4}

¹Department of Psychiatry and Behavioral Sciences, University of California, Davis Medical Center, Sacramento, CA

²M.I.N.D. Institute, University of California, Davis Medical Center

³Department of Public Health Sciences, University of California, Davis, Davis CA

⁴Department of Pediatrics, University of California, Davis Medical Center

Abstract

Objective—The authors studied the lifetime prevalence of DSM-IV-TR psychiatric disorders in a population of adults with the fragile X premutation.

Methods—Structured Clinical Interview for DSM-IV (SCID) were conducted on 85 individuals with the fragile X premutation, 47 with the fragile X-associated tremor/ataxia syndrome (FXTAS; 33 male, 14 female, mean age 66) and 38 without FXTAS (16 male, 22 female, mean age 52). Lifetime prevalence for mood and anxiety disorders among carriers with and without FXTAS was compared to available age-specific population estimates from the National Comorbidity Survey Replication (NCS-R).

Results—Among subjects with FXTAS, 30 cases (65%) met lifetime DSM-IV-TR criteria for mood disorder; 24 cases (52%) met lifetime DSM-IV-TR criteria for anxiety disorder. Among the non-FXTAS subjects, there were 15 cases (42%) of lifetime mood disorder and 18 cases (47%) of lifetime anxiety disorder. When compared to age-specific NCS-R data, the lifetime prevalence of any mood disorder, major depressive disorder, any anxiety disorder, panic disorder, specific phobia, and PTSD were significantly higher in subjects with FXTAS. The lifetime rates of social phobia in individuals with the premutation without FXTAS were significantly higher than NCS-R data.

Conclusions—This sample of carriers of the fragile X premutation had a notably high lifetime risk of mood and anxiety disorders. Mood and anxiety disorders may be part of the clinical phenotype of the fragile X premutation conditions, especially in carriers with FXTAS. Clinicians encountering these patients are advised to consider FXTAS as a neuropsychiatric syndrome as well as a neurological disorder.

Keywords

Fragile X syndrome; fragile X-associated tremor/ataxia syndrome; *FMR1* gene; (FXTAS); mood disorders; anxiety disorders

Introduction

Fragile X syndrome (FXS) is a trinucleotide repeat disorder caused by an expansion of greater than 200 CGG repeats in the 5' untranslated region of the fragile X mental retardation 1 (*FMR1*) gene on the X chromosome.¹ Depending on the number of CGG repeats, patients are classified as having normal alleles (5–44), premutation alleles (55–200), or full mutation alleles (>200). Alleles between 45–54 CGG repeats are considered the “gray zone” because the risk for expansion or clinical involvement is unclear or “gray”.²

The *FMR1* gene codes for *FMR1* protein (FMRP), which is believed to be involved in dendritic transport of mRNAs, regulation of translation, and synaptic plasticity.³ It has been demonstrated that *FMR1* mRNA levels in peripheral blood leukocytes are two- to eight-fold higher than normal in premutation subjects, despite a normal or slightly low level of FMRP.⁴ The excess *FMR1* mRNA in premutation carriers may result in a “toxic gain of function” effect leading to dysregulation of lamin A/C and several heat shock proteins, with subsequent accumulation, along with the *FMR1* mRNA, in intranuclear inclusions found in post-mortem studies of FXTAS.^{5, 6}

Fragile X Premutation: Clinical Aspects

Premutation expansions are common; as common as 1 per 113–259 females and 1 per 260–810 males.^{7–11} Patients with the premutation can present with a characteristic late-onset neurodegenerative disorder, fragile X-associated tremor/ataxia syndrome (FXTAS), described below. FXTAS is estimated to occur in 40% of premutation males and 8% of premutation females over 50 years of age who are identified through families of children with fragile X syndrome.¹² Primary ovarian insufficiency (POI) with early menopause affects up to 20% of premutation females.¹³ Co-morbid psychiatric illness is also seen in a subgroup of premutation carriers.^{14–17}

Psychopathology in Premutation Carriers without FXTAS

Initial studies of psychiatric illness in premutation carriers focused on women because they were more available in the evaluation of their FXS-affected children. Females may be more protected from RNA toxicity, as they have a normal X chromosome in addition to the X chromosome with the premutation.¹⁵ Initial reports did not find significant psychopathology in female carriers compared to controls.¹⁸ Although the stress of raising a child with FXS could certainly add to the risk for psychiatric illness, many female carriers report the lifetime experience of anxiety and depression *prior* to giving birth to children who have FXS.¹⁹

Franke et al.¹⁶ reported a lifetime diagnosis of depressive disorders in 21.3%, major depression in 19.7%, and bipolar disorder in 11.5% of women with the premutation who were mothers of children with FXS. Thompson et al.²⁰ studied 14 women with the

premutation (and 5 with the full mutation) and found 78% to have had a lifetime history of major depression; 33% had had recurrent mood disorder. Sobesky et al.²¹ found lifetime prevalence in the following psychiatric disorders in females with the premutation: major depression in 42%, dysthymic disorder in 38%, social phobia in 38%, and generalized anxiety disorder in 11%. A more recent study revealed that the abnormal elevation of mRNA correlated with severity of psychiatric symptoms in males with the premutation (with and without FXTAS); this association was not observed in females with the premutation.¹⁵

Roberts et al.¹⁹ studied 93 females with the premutation versus data from 2,159 age-adjusted female controls from the National Comorbidity Survey Replication (NCSR). They completed the Structured Clinical Interview for the DSM-IV (SCID-I) for lifetime and current history of mood and anxiety disorders. Compared to NCS-R control prevalences, the authors found an elevated risk of lifetime major depressive disorder (43.0% vs. 31.9%), lifetime panic disorder without agoraphobia (8.6% vs. 2.3%), and current agoraphobia without panic disorder (3.2% vs. 0.7%) in the premutation sample. The risk of mood disorder in women with the premutation was associated with *decreased* CGG repeat length and was not associated with variables pertinent to number of children with FXS or child behavioral problems, suggesting that the social stress of raising a FXS child may not account for the incidence of mood disorders. In distinction from the finding on mood disorders, the prevalence of anxiety disorders in the premutation group was correlated with number of children with FXS and problematic child behaviors.

Various clinical findings consistent with anxiety disorders, such as anxiety, social avoidance, interpersonal sensitivity, shyness, and avoidance of eye contact have been reported in carriers of the premutation, although prevalences and psychiatric diagnostic specificity are not clear at this time.^{14, 22, 23} Franke et al.¹⁶ reported a lifetime prevalence of anxiety disorder of 41.0% in mothers of children with FXS with the premutation; this included a high lifetime risk of social phobia (18.0%) and panic disorder (11.5%). This study also demonstrated a high prevalence of social phobia that was also seen in sisters of these women with the premutation, who did not have children with FXS.

FXTAS

FXTAS is a late onset (generally > 50 years) neuropsychiatric and neurologic degenerative illness, occurring predominantly in males with the premutation; only 8% of females with the premutation are affected.^{24–29} As such, FXTAS may be one of the more common late-onset, progressive neurological diseases resulting from a single gene mutation.³⁰ The major motor findings in FXTAS are kinetic, intention or postural tremor, cerebellar gait and limb ataxia, and parkinsonism.^{31, 32} Age of onset of motor symptoms and degree of brain atrophy correlate with CGG repeat length.^{33–35} The clinical features of FXTAS are shown in the diagnostic criteria in Table 1. FXTAS clinical staging is found in Table 2.

Psychopathology in FXTAS

As is common in neurodegenerative conditions, either concurrent or sequential clinical expression of psychiatric illnesses may occur in FXTAS. Indeed, the conceptualization for

CNS-derived symptoms and signs in such illnesses is that of “neuropsychiatric” illness. Patients may experience symptoms on several psychiatric dimensions adding further conceptual and diagnostic imprecision. In FXTAS, this dispersion of psychiatric symptoms was illustrated by Hessel et al.¹⁵ in a study of subjects with the premutation both with and without FXTAS. On the Symptom Checklist-90-Revised (SCL-90-R), relative to published norms, elevated scores on somatization, obsessive compulsive, interpersonal sensitivity, psychoticism, and the global severity index were found for carriers of the premutation (both genders); increased scores on depression and phobic anxiety were found in males with the premutation relative to published norms. These symptoms were present in carriers who did not have FXTAS suggesting that the RNA toxicity to the limbic system may begin in early adult life or perhaps has a neurodevelopmental component as suggested by Farzin et al.¹⁷

With the onset of FXTAS the symptoms of anxiety and depression may worsen and dementia may develop.^{36–38} Dementia may present with frontal lobe features (disinhibition, inappropriate social behavior, poor executive functioning, perseveration, irritability, mood disturbances) and subcortical features (psychomotor slowing, bradyphrenia, attention and concentration difficulties).^{31, 36, 37} Onset of the dementia usually follows the onset of movement disorder, and is rare before age 50.³¹ Upon formal assessment, patients with FXTAS are found to suffer a larger decrement in performance than verbal IQ.³¹ A recent paper of 15 subjects with FXTAS revealed dementia in 7; 7 were also diagnosed with mood or anxiety disorders. Twelve had cognitive impairment on formal cognitive testing.³⁸

The course of progression of cognitive impairment in FXTAS dementia remains obscure, although it appears likely that FXTAS dementia is insidiously progressive. Some cognitive deficits of FXTAS dementia are of severity comparable to those in Alzheimer’s disease, albeit in different domains.³⁹ The behavioral disturbances in FXTAS dementia have been demonstrated on the Neuropsychiatric Inventory (NPI) to include apathy, depression, and agitation as more common than in age-matched controls.³⁷

Methods

Patients were recruited from 3 studies and each involved informed consent approved by the IRB at University of California at Davis Medical Center. The first study recruited individuals with the premutation who had neurological symptoms including tremor and ataxia. They were assessed for the presence of FXTAS and were subsequently recruited into a study of neuroprotective agents for FXTAS. The second study evaluated young men with the premutation without any neurological symptoms and male siblings without the premutation. The third study assessed all family members with the premutation of a proband with the full mutation.

All subjects underwent a neurological examination to clarify the presence or absence of the motor signs of FXTAS utilizing diagnostic criteria as reported by Jacquemont et al.³¹ The subjects also completed a psychiatric evaluation, which included the Structured Clinical Interview for DSM-IV (SCID). The SCID is a structured clinical interview for DSM-IV psychiatric disorders. The format for the SCID begins with a general medical and psychiatric history, and then proceeds with diagnostic criteria for mood, psychotic,

substance use, anxiety, somatoform, and adjustment disorders. The interviewer asks a prescribed series of questions that correspond to the DSM-IV diagnostic criteria for the various psychiatric disorders. The format of the SCID includes breakdown for current and past psychiatric disorders, duration of each episode, age of onset of first episode, severity of functional impairment, and other details. To render a formal diagnosis of a psychiatric illness based on SCID-derived data requires the validation of DSM-IV criteria for each illness. Current and lifetime illness experience may be tabulated separately. Interview data is then gleaned from the interview document for analysis. The SCID interviews were completed by three of the authors (JAB, ALS and AS), all of whom are experienced at conducting this interview for research programs. As this is a detailed structured clinical interview, test-retest reliability is high and this instrument is considered a “gold standard” for psychiatric epidemiology.^{40–41} One of its major benefits, as the interviewer is required to progress throughout major categories of psychiatric disorders in the conduct of the interview, is its utility in helping the interviewer to document psychiatric co-morbidity. Of relevance in our cohort are psychiatric co-morbidity, both “across” psychiatric disorder boundaries (e.g., a patient with independent history of mood and anxiety disorders) and “within” psychiatric disorder boundaries (e.g., a patient with two or more distinct anxiety disorders).

In interpretation of SCID interview data, the examiner seeks to describe patients’ illness in terms of “caseness” (e.g., a patient with a history of major depression and panic disorder is considered a “case” of each illness), so that multiple diagnoses (“cases”) in a single patient are allowed. Weaknesses of the SCID include the failure to include criteria for cognitive, personality, and childhood onset disorders. An additional weakness of the SCID paradigm, particularly in the analysis of older patients with risk of cognitive impairment, is the requirement of intact memory to recall prior illness experiences. Our primary analyses focused on mood and anxiety disorders. All other analyses are considered secondary and exploratory. We compared the lifetime prevalence of mood and anxiety disorders to available age-specific population estimates from the National Comorbidity Survey Replication (NCS-R).⁴² This is a nationally-representative sample that used both the Composite International Diagnostic Interview (CIDI) and the SCID. For NCS-R, 9,282 adult interviews were completed with statistical weighting models applied to demographic variables. Lifetime prevalence of psychiatric illness was reported stratified by age grouping and gender.

The CIDI is mostly used by the World Health Organization (WHO) in World Mental Health surveys, and is based on the International Statistical Classification of Diseases and Related Health Problems 10th Revision (ICD-10). This classification system is comparable to the DSM-IV-TR and focuses on the coding of diseases, social circumstances and external causes of injury or illnesses. Several studies have been published so far about the comparison between the diagnostic specificity of the CIDI and SCID.^{43–45} Especially for the diagnoses of mood and anxiety disorders, Kessler et al.^{46, 47} described an excellent concordance of both instruments.

Comparisons of the current study’s prevalence estimates to NCS-R’s age-specific prevalence figures were based on the exact binomial method (two-sided test at level 0.05).

P-values were adjusted for multiple testing the false discovery rate (FDR) criterion and a significant p-value after FDR adjustment is indicated by an asterisk in Table 4 and 5. There was one missing data point for mood disorders and anxiety in the FXTAS group and 2 missing data points in the group without FXTAS. This is a one-sample comparison of the proportion to published prevalence figures for age groups similar to the age of our study premutation cohort with and without FXTAS. Patient characteristics (Table 3), including molecular measures (CGG repeat number, *FMRI* mRNA), were based on t-tests continuous variables and Fisher's exact test for categorical variables.

Results

Patient characteristics, including demographics, CGG, *FMRI* mRNA, and cognitive testing results are summarized in Table 3 by FXTAS status. Mean age of carriers with FXTAS ($n = 47$) was 66 years old (SD 7.9; range 54–87) and for carriers without FXTAS (nonFXTAS, $n = 38$), the mean age was 51.6 (SD 11.7, range 35–76). See Table 3 for details. Subjects with the premutation with no neurologic signs and Stage 1 FXTAS ($N = 11$) were combined as one group since stage 1 FXTAS has no definite tremor or ataxia and it does not include a definitive diagnosis of FXTAS (Table 2), while Stages 2–5 FXTAS subjects were combined as the FXTAS group. The relationship between FXTAS stage and SCID anxiety and mood disorders were not significant. In addition, we did not find a significant association between mRNA levels and number of CGG repeats to the presence of a SCID-diagnosed disorder.

To compare age-specific lifetime prevalences of mood and anxiety disorders, we selected NCS-R figures for the age categories of 45–59 and 60+ to compare our samples of nonFXTAS and FXTAS, respectively. These NCS-R age-specific categories contain the mean age of our study groups. When we examined lifetime prevalence of any mood disorder, both carriers with and without FXTAS had higher prevalences compared to NCS-R general population estimates. However, this prevalence in carriers without FXTAS was not significant after p-value adjustment. Observed lifetime prevalence of major depressive disorder in premutation carriers without FXTAS was similar to the NCS-R general population figure. However, among patients with FXTAS, lifetime prevalence of major depressive disorder was significantly higher. There was no difference with NCS-R figures with respect to the lifetime prevalence of dysthymic disorder and bipolar disorders. The higher prevalence estimates for major depressive disorder and any mood disorder in FXTAS remained unchanged for males with FXTAS ($n = 32$, after excluding females with FXTAS).

Details are provided in Table 4.

With respect to anxiety disorders, the subjects with FXTAS had higher lifetime prevalence of any anxiety disorder, panic disorder, specific phobia, social phobia, and PTSD. However, the increased prevalence of social phobia was not significantly higher after p-value adjustment. Premutation carriers without FXTAS also had higher lifetime prevalence of social phobia, although lifetime prevalence of other anxiety disorders and the aggregate of any anxiety disorder were not different from NCS-R general population estimates. The higher prevalence estimates for any anxiety disorder and specific phobia in FXTAS remained significant for males with FXTAS ($n = 32$, after excluding females with FXTAS).

However, panic disorder and PTSD prevalences were not significant among males with FXTAS after excluding females. Details are provided in Table 5.

Discussion

This project examined the lifetime prevalences of DSM-IV psychiatric disorders in a cohort of 85 carriers of the fragile X premutation. To examine psychiatric disorders in the premutation cohort with and without FXTAS, the analysis was accordingly stratified by the groups. These lifetime prevalences were compared to corresponding age-specific general population estimates (NCS-R). Because of our low numbers of female subjects, analysis controlling for gender was not done. When compared to age-specific NCS-R data, the lifetime prevalence of any mood disorder, major depressive disorder, any anxiety disorder, panic disorder, specific phobia, and PTSD were significantly higher in the subjects with FXTAS.

The analysis of these lifetime prevalences did not include assessment of associated and antecedent events, which may be of importance in particular in the increased prevalence of PTSD. It may be postulated that patients with FXTAS, by virtue of their increased experience of anxiety symptoms in general, may be somewhat more disposed to develop PTSD when exposed to traumatic experiences. This could be a productive area of future study. The lifetime prevalence of social phobia in premutation subjects without FXTAS was significantly higher than NCS-R data. The lifetime prevalences of mood and anxiety disorder diagnoses for both the premutation group without FXTAS and the group with FXTAS are both high compared to general population studies using similar structured interviews.

In addition to the NCS-R study, several other studies using DSM-referenced diagnostic interviews to assess the lifetime prevalence of mood and anxiety disorders in adult population-based cohorts have been conducted.^{44, 48–58} While employing different diagnostic instruments, these comparable studies reported lifetime prevalences of mood (8% - 21%) and anxiety (15% - 29%) disorders at rates similar to those in the NCS-R study.

Previous studies of psychopathology in premutation carriers had included only female subjects and found significant lifetime prevalences of mood and/or anxiety disorders.^{16, 19–21} Our study is the first large study of psychopathology in premutation carriers to include a large percentage of male premutation patients (70% of the FXTAS group, 43% of the nonFXTAS group). As males in the general population tend to have a lower prevalence of psychiatric illness (particularly major depression), our findings are all the more striking.

It is important to emphasize that the current study examined the lifetime (not necessarily current) prevalence of mood and anxiety disorders in individuals with the fragile X premutation. As we did not, in this study, assess the age of onset of psychiatric disorders, the temporal sequencing of “psychiatric illness first, motor symptoms second” is suggested because we saw a higher prevalence of certain psychiatric problems in premutation carriers

in those without FXTAS compared to the general population. Further studies of the age of onset of psychiatric syndromes are underway to address this issue specifically.

It is hypothesized that vulnerability to lifetime mood and anxiety disorders may be a condition of the fragile X premutation itself, rather than purely as a consequence of social stressors, such as the experience of a degenerative neuropsychiatric condition or the responsibility of caring for a child with FXS. While responsibility for the care of a neurodevelopmentally impaired child with FXS can clearly represent a profound chronic social stressor, many of our subjects were diagnosed with premutation carrier status as the result of pedigree analysis of an affected grandchild, and had not been primarily responsible for care for an impaired child. Further studies examining age of onset of the first episode of psychiatric disorders and social stressors (such as raising a FXS child) in premutation carriers are now underway to address these potential contributing factors.

We note here that recent brain imaging studies document that men with the premutation without FXTAS and who are not raising affected children have abnormal reduction in amygdala and hippocampal activation related to both abnormal elevation of *FMRI* mRNA and psychiatric symptoms.^{59, 60} Additional reports of progressive decline in executive function, such as working memory and attention in males with the premutation beginning in the third decade of life may point toward a neurophysiological basis of psychiatric disorder in this population.⁶¹ Evidence of psychopathology and neurophysiological deficits before the onset of FXTAS is not surprising because evidence of RNA toxicity including inclusions develop in the premutation mouse throughout the limbic system including the amygdala and hippocampus earlier than expected and before the onset of FXTAS in the mouse.⁶²

There is an apt comparison available between the psychopathology in fragile X premutation conditions and Parkinson's disease. Indeed, it has been the experience of our research group that many patients with FXTAS, by virtue of their progressive motor symptoms, are often initially diagnosed with variants of Parkinson's disease. While the research methodologies differ, the literature to date assessing lifetime mood and anxiety disorders prevalence in Parkinson's disease are similar to our findings, especially in patients with FXTAS. Lifetime prevalence of mood disorder as high as 60% and anxiety disorders (including simple phobia and panic disorder) as high as 40% in Parkinson's disease.^{63–68} As such, the possibility exists that Parkinson's disease with mood and anxiety disorder co-morbidity may share a common neuroanatomic pathway (at least in functional terms) with fragile X premutation conditions. Further research studies enrolling fragile X premutation patients and Parkinson's disease patients in the same research protocol would be desirable to further examine this comparison.

A potential confounder in the analysis of lifetime history of mood and anxiety disorders in patients with the premutation may be related to the co-morbidity of dementing illness in patients with FXTAS. Mood and anxiety disorders are common co-morbidities in dementing illnesses, thus primary FXTAS dementia may eventually lead to dementia-associated mood and anxiety disorders, rather than a temporal primacy of mood and/or anxiety disorders in patients who later develop dementing illness. Secondly, patients with clinical dementia may, by dint of retrograde amnesia, have difficulty in accurately describing previously

experienced psychiatric illness in sufficient detail to render an accurate DSM-IV diagnosis in the SCID interview. In our cohort, however, the mean MMSE score was 28.6 for the FXTAS group and 29.8 for nonFXTAS, scores consistent with intact cognitive function. Our subjects' intact cognitive status is reinforced by IQ results, with mean full scale IQ of 107 in the FXTAS group and 112 in nonFXTAS. Despite high MMSE and IQ scores on average, two of the subjects met criteria for dementia and one had mild cognitive impairment. When performing SCID interviews with these subjects, we obtained collateral information from their spouses in completing the SCIDs. Therefore, it is unlikely that cognitive function introduced bias in the vast majority of our subjects' recall of their psychiatric histories. Similarly, due to the small number of patients with cognitive impairment, the possible number of dementia-associated mood and anxiety disorders in our cohort is quite low.

Although the majority of patients with FXTAS were clinic-referred for their movement disorder, none were referred due to psychiatric concerns. In addition, younger individuals with the premutation without FXTAS seen in the study were not clinically referred but were recruited from pedigrees of families affected by FXS, also unrelated to psychiatric concerns. Despite these efforts to minimize selection bias, we recognize that potential selection bias may still exist and, therefore, may affect the prevalence estimates of psychiatric disorders observed in our sample. Future studies examining psychiatric conditions in carriers of the premutation should attempt to control for various sources of bias and confounding factors, and employ well-matched control groups. In addition, further study of the sequencing of the experience of psychiatric disorders in these population by examination of the age of onset of the first episode of the various SCID-diagnosed psychiatric illness compared to the age of onset of motor symptoms are planned to elucidate the issue of primary versus secondary illness episodes.

Despite the imprecision of attribution of the genesis of mood and anxiety disorders in this population, clinicians encountering patients with a family history of FXS (or any autism-spectrum disorder) and/or dementing illness or movement disorder should consider *FMR1* DNA testing as part of the clinical evaluation. In many cases, positive identification of the premutation in an older adult with the aforementioned motor or cognitive signs has led to first diagnosis of FXS in grandchildren or other family members, and as such genetic counseling is strongly recommended in these cases. Prompt clinical intervention with medication and/or psychotherapy for mood and anxiety disorders should be offered for these patients. Vigilance for cognitive and motor status is also highly recommended.

Conclusions

Lifetime prevalences of mood and anxiety disorders were found to be high in individuals with the fragile X premutation with FXTAS. Subjects with the premutation without FXTAS had a high prevalence of lifetime mood disorders overall (although the lifetime prevalences of major depression, dysthymic disorder, and bipolar disorder were not individually significantly more common) and social phobia. This study is consistent with an emerging neuropsychiatric phenotype in fragile X carriers where genetically-determined mood and anxiety disorders appear to be substantial clinical components of this disorder. This illness may well represent a relatively common neuropsychiatric disorder that is attributable to a

single gene mutation. Clinicians encountering mood and anxiety disorders in patients with a family history of intellectual disability apparent in childhood and/or dementia, particularly dementia with a movement disorder, may wish to obtain confirmatory DNA testing for the fragile X premutation in these patients.

Acknowledgments

This work was supported by the National Fragile X Foundation and the National Institute of Health (NIH) grants HD036071, HD056031, NS044299, HD02274, MH77554, MH078041 and also by a NIH Roadmap Interdisciplinary Research Consortium Grant (NIA RL1 AG032115, NINDS RL1 NS062412, NIA RL1 AG032119, and NIDCR DE019583) and the M.I.N.D. Institute, University of California, Davis. The work was also supported in part by the Intramural Research Program, NIH.

I notice that the my grant added to the acknowledgement on May 15 was not included in the final version. When you get the proofs, if you could please add my grant support acknowledgement:

UL1 RR024146 (National Center for Research Resources)

Thanks very much.

References

1. Hagerman RJ, Rivera SM, Hagerman PJ. The fragile X family of disorders: A model for autism and targeted treatments. *Current Pediatric Reviews*. 2008; 4:40–52.
2. Nolin SL, Brown WT, Glicksman A, et al. Expansion of the fragile X CGG repeat in females with premutation or intermediate alleles. *Am J Hum Genet*. 2003; 72:454–464. [PubMed: 12529854]
3. Bassell GJ, Warren ST. Fragile X syndrome: loss of local mRNA regulation alters synaptic development and function. *Neuron*. 2008; 60:201–214. [PubMed: 18957214]
4. Tassone F, Hagerman RJ, Taylor AK, et al. Elevated levels of *FMR1* mRNA in carrier males: a new mechanism of involvement in fragile X syndrome. *Am J Hum Genet*. 2000; 66:6–15. [PubMed: 10631132]
5. Greco CM, Hagerman RJ, Tassone F, et al. Neuronal intranuclear inclusions in a new cerebellar tremor/ataxia syndrome among fragile X carriers. *Brain*. 2002; 125:1760–1771. [PubMed: 12135967]
6. Greco CM, Berman RF, Martin RM, et al. Neuropathology of fragile X-associated tremor/ataxia syndrome (FXTAS). *Brain*. 2006; 129(Pt 1):243–255. [PubMed: 16332642]
7. Dawson AJ, Chodirker BN, Chudley AE. Frequency of *FMR1* premutations in a consecutive newborn population by PCR screening of Guthrie blood spots. *Biochem Mol Med*. 1995; 56:63–69. [PubMed: 8593539]
8. Dombrowski C, Levesque ML, Morel ML, et al. Premutation and intermediatesize *FMR1* alleles in 10572 males from the general population: loss of an AGG interruption is a late event in the generation of fragile X syndrome alleles. *Hum Mol Genet*. 2002; 11:371–378. [PubMed: 11854169]
9. Rousseau F, Rouillard P, Morel ML, et al. Prevalence of carriers of premutationsize alleles of the *FMRI* gene--and implications for the population genetics of the fragile X syndrome. *Am J Hum Genet*. 1995; 57:1006–1018. [PubMed: 7485149]
10. Toledano-Alhadeif H, Basel-Vanagaite L, Magal N, et al. Fragile-X carrier screening and the prevalence of premutation and full-mutation carriers in Israel. *Am J Hum Genet*. 2001; 69:351–360. [PubMed: 11443541]
11. Hagerman PJ. The Fragile X prevalence paradox. *J Med Gen*. 2008 April 15 [epub ahead of print] (PMID: 18413371).
12. Jacquemont S, Hagerman RJ, Leehey MA, et al. Penetrance of the fragile X-associated tremor/ataxia syndrome in a premutation carrier population. *JAMA*. 2004; 291:460–469. [PubMed: 14747503]
13. Sullivan AK, Marcus M, Epstein MP, et al. Association of *FMR1* repeat size with ovarian dysfunction. *Hum Reprod*. 2005; 20:402–412. [PubMed: 15608041]

14. Hagerman RJ. Lessons from fragile X regarding neurobiology, autism, and neurodegeneration. *J Dev Behav Pediatr.* 2006; 27:63–74. [PubMed: 16511373]
15. Hessel D, Tassone F, Loesch DZ, et al. Abnormal elevation of *FMR1* mRNA is associated with psychological symptoms in individuals with the fragile X premutation. *Am J Med Genet B Neuropsychiatr Genet.* 2005; 139:115–121. [PubMed: 16184602]
16. Franke P, Leboyer M, Gansicke M, et al. Genotype-phenotype relationship in female carriers of the premutation and full mutation of *FMR-1*. *Psychiatry Res.* 1998; 80:113–127. [PubMed: 9754690]
17. Farzin F, Perry H, Hessel D, et al. Autism spectrum disorders and attention deficit/hyperactivity disorder in boys with the fragile X premutation. *J Dev Behav Pediatr.* 2006; 27(2 Suppl):S137–S144. [PubMed: 16685180]
18. Reiss AL, Freund L, Abrams MT, et al. Neurobehavioral effects of the fragile X premutation in adult women: a controlled study. *Am J Hum Genet.* 1993; 52:884–894. [PubMed: 8488838]
19. Roberts JE, Bailey DB Jr, Mankowski J, et al. Mood and anxiety disorders in females with the *FMR1* premutation. *Am J Med Genet B Neuropsychiatr Genet.* 2009; 150B:130–139. [PubMed: 18553360]
20. Thompson NM, Rogeness GA, McClure E, et al. Influence of depression on cognitive functioning in fragile X females. *Psychiatry Res.* 1996; 64:97–104. [PubMed: 8912951]
21. Sobesky WE, Pennington BF, Porter D, et al. Emotional and neurocognitive deficits in fragile X. *Am J Med Gen.* 1994; 51:378–385.
22. Hagerman RJ, Hagerman PJ. The fragile X premutation: into the phenotypic fold. *Curr Opin Genet Dev.* 2002; 12:278–283. [PubMed: 12076670]
23. Hagerman PJ, Hagerman RJ. The fragile-X premutation: a maturing perspective. *Am J Hum Genet.* 2004; 74:805–816. [PubMed: 15052536]
24. Hagerman RJ, Leavitt BR, Farzin F, et al. Fragile-X-associated tremor/ataxia syndrome (FXTAS) in females with the *FMR1* premutation. *Am J Hum Genet.* 2004; 74:1051–1056. [PubMed: 15065016]
25. Jacquemont S, Hagerman RJ, Hagerman PJ, Leehey MA. Fragile-X syndrome and fragile X-associated tremor/ataxia syndrome: two faces of *FMR1*. *Lancet Neurol.* 2007; 6:45–55. [PubMed: 17166801]
26. Berry-Kravis E, Sumis A, Hervey C, et al. Open-label treatment trial of lithium to target the underlying defect in fragile X syndrome. *J Dev Behav Pediatr.* in press.
27. Leehey MA, Berry-Kravis E, Min SJ, et al. Progression of tremor and ataxia in male carriers of the *FMR1* premutation. *Mov Disord.* 2007; 22:203–206. [PubMed: 17133502]
28. Willemsen R, Mientjes E, Oostra BA. FXTAS: a progressive neurologic syndrome associated with fragile X premutation. *Curr Neurol Neurosci Rep.* 2005; 5:405–410. [PubMed: 16131424]
29. Hagerman PJ, Hagerman RJ. Fragile X-associated tremor/ataxia syndrome—an older face of the fragile X gene. *Nat Clin Pract Neurol.* 2007; 3:107–112. [PubMed: 17279084]
30. Jacquemont S, Leehey MA, Hagerman RJ, et al. Size bias of fragile X premutation alleles in late-onset movement disorders. *J Med Genet.* 2006; 43:804–809. [PubMed: 16723388]
31. Jacquemont S, Hagerman RJ, Leehey M, et al. Fragile X premutation tremor/ataxia syndrome: molecular, clinical, and neuroimaging correlates. *Am J Hum Genet.* 2003; 72:869–878. [PubMed: 12638084]
32. Jacquemont S, Farzin F, Hall D, et al. Aging in individuals with the *FMR1* mutation. *Am J Ment Retard.* 2004; 109:154–164. [PubMed: 15000674]
33. Tassone F, Adams J, Berry-Kravis EM, et al. CGG repeat length correlates with age of onset of motor signs of the fragile X-associated tremor/ataxia syndrome (FXTAS). *Am J Med Genet B Neuropsychiatr Genet.* 2007; 144:566–569. [PubMed: 17427188]
34. Cohen S, Masyn K, Adams J, et al. Molecular and imaging correlates of the fragile X-associated tremor/ataxia syndrome. *Neurology.* 2006; 67:1426–1431. [PubMed: 17060569]
35. Adams JS, Adams PE, Nguyen D, et al. Volumetric brain changes in females with fragile X-associated tremor/ataxia syndrome (FXTAS). *Neurology.* 2007; 69:851–859. [PubMed: 17724287]
36. Bourgeois JA, Farzin F, Brunberg JA, et al. Dementia with mood symptoms in a fragile X premutation carrier with the fragile X-associated tremor/ataxia syndrome: clinical intervention

- with donepezil and venlafaxine. *J Neuropsychiatry Clin Neurosci*. 2006; 18:171–177. [PubMed: 16720793]
37. Bacalman S, Farzin F, Bourgeois JA, et al. Psychiatric phenotype of the fragile X-associated tremor/ataxia syndrome (FXTAS) in males: newly described frontosubcortical dementia. *J Clin Psychiatry*. 2006; 67:87–94. [PubMed: 16426093]
 38. Bourgeois JA, Cogswell JB, Hessel D, et al. Cognitive, anxiety and mood disorders in the fragile X-associated tremor/ataxia syndrome. *Gen Hosp Psychiatry*. 2007; 29:349–356. [PubMed: 17591512]
 39. Seritan AL, Nguyen D, Farias S, et al. Dementia in fragile X-associated tremor/ataxia syndrome (FXTAS): comparison with Alzheimer's disease. *Am J Med Genet B: Neuropsychiatr Genet*. 2008; 147:1138–1144. [PubMed: 18384046]
 40. Spitzer RL, Williams JB, Gibbon M, First MB. The Structured Clinical Interview for *DSM-III-R* (SCID). I. History, rationale, and description. *Arch Gen Psychiatry*. 1992; 49:624–629. [PubMed: 1637252]
 41. Williams JB, Gibbon M, First MB, et al. The Structured Clinical Interview for *DSM-III-R* (SCID). II. Multisite test-retest reliability. *Arch Gen Psychiatry*. 1992; 49:630–636. [PubMed: 1637253]
 42. National Comorbidity Survey Replication (NCS-R). <http://www.hcp.med.harvard.edu/ncs/>.
 43. Kessler RC, Abelson J, Demler O, et al. Clinical calibration of *DSM-IV* diagnoses in the World Mental Health (WMH) version of the World Health Organization (WHO) Composite International Diagnostic Interview (WMH-CIDI). *Int J Methods Psychiatr Res*. 2004; 13:122–139. [PubMed: 15297907]
 44. Kessler RC, Berglund P, Demler O, et al. Lifetime prevalence and age-of-onset distributions of *DSM-IV* disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2005; 62:593–602. [PubMed: 15939837]
 45. Haro JM, Arbabzadeh-Bouchez S, Brugha TS, et al. Concordance of the Composite International Diagnostic Interview Version 3.0 (CIDI 3.0) with standardized clinical assessments in the WHO World Mental Health surveys. *Int J Methods Psychiatr Res*. 2006; 15:167–180. [PubMed: 17266013]
 46. Kessler RC, Akiskal HS, Angst J, et al. Validity of the assessment of bipolar spectrum disorders in the WHO CIDI 3.0. *J Affect Disord*. 2006; 96:259–269. [PubMed: 16997383]
 47. Kessler RC, Chiu WT, Jin R, et al. The epidemiology of panic attacks, panic disorder, and agoraphobia in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2003; 63:415–424. [PubMed: 16585471]
 48. Bijl RV, Ravelli A, van Zessen G. Prevalence of psychiatric disorder in the general population: results of the Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Soc Psychiatry Psychiatr Epidemiol*. 1998; 33:587–595. [PubMed: 9857791]
 49. Kessler RC, Berglund P, Demler O, et al. The epidemiology of major depressive disorder: Results from the National Comorbidity Survey Replication (NCS-R). *JAMA*. 2003; 289:3095–3105. [PubMed: 12813115]
 50. Weissman MM, Leaf PJ, Tischler GL, et al. Affective disorders in five United States communities. *Psychol Med*. 1988; 18:141–153. [PubMed: 3363034]
 51. Kessler RC, McGonagle KA, Zhao S, et al. Lifetime and 12-month prevalence of *DSM-III-R* psychiatric disorders in the United States: results from the National Comorbidity Survey. *Arch Gen Psychiatry*. 1994; 51:8–19. [PubMed: 8279933]
 52. Kessler RC, Rubinow DR, Holmes C, et al. The epidemiology of *DSM-III-R* bipolar I disorder in a general population survey. *Psychol Med*. 1997; 27:1079–1089. [PubMed: 9300513]
 53. Regier DA, Rae DS, Narrow WE, et al. Prevalence of anxiety disorders and their comorbidity with mood and addictive disorders. *Br J Psychiatry*. 1998; 173(Suppl 34):24–28.
 54. Weissman MM, Myers JK. Affective disorders in a US urban community: the use of Research Diagnostic Criteria in an epidemiological survey. *Arch Gen Psychiatry*. 1978; 35:1304–1311. [PubMed: 708194]
 55. Magee WJ, Eaton WW, Wittchen H-U, et al. Agoraphobia, simple phobia, and social phobia in the National Comorbidity Survey. *Arch Gen Psychiatry*. 1996; 53:159–168. [PubMed: 8629891]

56. Grant BF, Hasin DS, Stinson FS, et al. The epidemiology of DSM-IV-panic disorder and agoraphobia in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *J Clin Psychiatry*. 2006; 67:363–374. [PubMed: 16649821]
57. Grant BF, Hasin DS, Stinson FS, et al. Prevalence, correlates, co-morbidity, and comparative disability of DSM-IV generalized anxiety disorder in the USA: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Psychol Med*. 2005; 35:1747–1759. [PubMed: 16202187]
58. Grant BF, Hasin DS, Blanco C, et al. The epidemiology of social anxiety disorder in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *J Clin Psychiatry*. 2005; 66:1351–1361. [PubMed: 16420070]
59. Hessl D, Rivera S, Koldewyn K, et al. Amygdala dysfunction in men with the fragile X premutation. *Brain*. 2007; 130:404–416. [PubMed: 17166860]
60. Koldewyn K, Hessl D, Adams J, et al. Reduced hippocampal activation during recall is associated with elevated *FMRI* mRNA and psychiatric symptoms in men with the fragile X premutation. *Brain Imaging Behav*. 2008; 2:105–116. [PubMed: 19430586]
61. Cornish KM, Kogan CS, Li L, et al. Lifespan changes in working memory in fragile X premutation males. *Brain Cogn*. 2009; 69:551–558. [PubMed: 19114290]
62. Kogan CS, Turk J, Hagerman RJ, et al. Impact of the Fragile X mental retardation 1 (*FMRI*) gene premutation on neuropsychiatric functioning in adult males without fragile X-associated Tremor/Ataxia syndrome: a controlled study. *Am J Med Genet B Neuropsychiatr Genet*. 2008; 147B:859–872. [PubMed: 18165971]
63. Voon V, Saint-Cyr J, Lozano AM, et al. Psychiatric symptoms in patients with Parkinson disease presenting for deep brain stimulation. *J Neurosurg*. 2005; 103:246–251. [PubMed: 16175853]
64. Nuyen J, Schellevis FG, Santariano WA, et al. Comorbidity was associated with neurologic and psychiatric diseases: a general practice-based controlled study. *J Clin Epidemiol*. 2006; 59:1274–1284. [PubMed: 17098570]
65. Lauterbach EC, Freeman A, Vogel RL. Differential DSM-III psychiatric disorder prevalence profiles in dystonia and Parkinson's disease. *J Neuropsychiatry Clin Neurosci*. 2004; 16:29–36. [PubMed: 14990756]
66. Cole SA, Woodard JL, Juncos JL, et al. Depression and disability in Parkinson's disease. *J Neuropsychiatry Clin Neurosci*. 1996; 8:20–25. [PubMed: 8845697]
67. Leentjens AFG, Van der Akker M, Metsemakers JFM, et al. Higher incidence of depression preceding the onset of Parkinson's disease: a register study. *Mov Disord*. 2003; 18:414–417. [PubMed: 12671948]
68. Lauterbach EC, Freeman A, Vogel RL. Correlates of generalized anxiety and panic attacks in dystonia and Parkinson disease. *Cog Behav Neurol*. 2003; 16:225–233.

TABLE 1

Diagnostic criteria for FXTAS*

MOLECULAR		CGG repeat 55 – 200
CLINICAL		
Major		Intention tremor
		Cerebellar gait ataxia
Minor		Parkinsonism
		Moderate to severe short term memory deficit
		Executive function deficit
RADIOLOGICAL		
Major		MRI white matter lesions involving middle cerebellar peduncles
		MRI lesions involving cerebral white matter
Minor		Moderate to severe generalized brain atrophy
Diagnostic Categories		
DEFINITE	PROBABLE	POSSIBLE
One major clinical, and	Two major clinical, or	One major clinical, and
One major radiological, or	One minor clinical, and	One minor radiological
Presence of FXTAS inclusions	One major radiological	

* Adapted from Jacquemont et al.(32).

TABLE 2

FXTAS Stage Description

STAGE	DESCRIPTION
0	Normal function
1	Subtle or questionable signs such as subtle tremor or mild balance problems, with no interference in ADLs
2	Minor, but clear, tremor and/or balance problems with minor interference with ADLs
3	Moderate tremor and/or balance problems and occasional falls with significant interference with ADLs
4	Severe tremor and/or balance problems; uses cane or walker
5	Uses wheelchair on a daily basis
6	Bedridden

Table 3

Characteristics of patients with premutation with and without FXTAS

Variable	FXTAS (N=47)			nonFXTAS (N=38)			P-value
	N	Mean	SD	N	Mean	SD	
Age	47	66.40	7.88	38	51.61	11.67	<0.0001
Male	33	70%		16	43%		0.0154
Education	46	14.98	2.89	31	15.42	1.82	0.4528
CGG	43	90.26	20.45	25	101.28	63.90	0.2987
FMR/mRNA	37	3.00	0.99	23	2.91	1.55	0.7848
Verbal IQ	40	108.00	21.42	37	114.30	13.91	0.1336
Performance IQ	39	103.59	16.91	37	112.38	16.21	0.0236
Full scale IQ	38	107.29	14.00	37	111.89	22.58	0.2908
MMSE	46	28.57	2.46	19	29.79	0.42	0.0353
BDS2	42	15.60	4.83	35	21.23	3.54	<0.0001
Stroop	41	43.98	9.27	14	45.93	4.12	0.4508

Table 4
Lifetime Prevalence of Mood Disorders, nonFXTAS and FXTAS, Compared to Age Adjusted NCS Data

Mood Disorders	Any mood disorder			Major depressive disorder			Dysthymic disorder			Bipolar disorders		
	%	n/N	P-value	%	n/N	P-value	%	n/N	P-value	%	n/N	P-value
nonFXTAS	41.7	15/36	0.031	22.2	8/36	0.88	11.1	4/36	0.094	8.3	3/36	0.295
NCS (age category 45-59)	24.2			20.1			3.8			3.7		
FXTAS males	62.5	20/32	<.0001*	31.3	10/32	0.013*	0	0/32	--	6.3	2/32	0.805
FXTAS	65.2	30/46	<.0001*	43.5	20/46	<.0001*	2.2	1/46	0.905	4.4	2/46	0.241
NCS (age category 60+)	12.2			10.7			1.3			1.3		

Table 5

Lifetime Prevalence of Anxiety Disorders, nonFXTAS and FXTAS vs. Age-Adjusted NCS Data

Anxiety Disorders	Any anxiety disorder			Panic disorder			Specific phobia			Social phobia			Generalized anxiety disorder			PTSD			Obsessive-compulsive disorder		
	%	n/N	P-value	%	n/N	P-value	%	n/N	P-value	%	n/N	P-value		n/N	P-value		n/N	P-value	n/N	P-value	
nonFXTAS	47.4	18/38	0.128	13.2	5/38	0.142	23.7	9/38	0.173	34.2	13/38	0.001*	2.6	1/38	0.409	2.6	1/38	0.248	0	0/38	--
NCS (age category 45-59)	34.2			5.9			14.4			12.6			7.6			9.2			2.4		
FXTAS males	50	16/32	0.005*	3.1	1/32	1		7/32	<.0001*	15.6	5/32	0.558	12.5	4/32	0.088	9.4	3/32	0.214	0	0/32	NA
FXTAS	52.2	24/46	<.0001*	10.9	5/46	0.006*	26.1	12/46	0.0003*	17.4	8/46	0.023	10.9	5/46	0.072	13	6/46	0.004*	0	0/46	--
NCS (age category 60+)	17.8			2.1			7.7			6.8			4			2.8			0.6		