Published in final edited form as:

Curr Opin Biotechnol. 2012 October ; 23(5): 672–678. doi:10.1016/j.copbio.2012.02.002.

Cell-free biology: exploiting the interface between synthetic biology and synthetic chemistry

D. Calvin Harris^{1,2} and Michael C. Jewett^{1,2,3,†}

¹Department of Chemical and Biological Engineering, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208, USA

²Chemistry of Life Processes Institute, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208, USA

³Member, Robert H. Lurie Comprehensive Cancer Center, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208, USA

Abstract

Just as synthetic organic chemistry once revolutionized the ability of chemists to build molecules (including those that did not exist in nature) following a basic set of design rules, cell-free synthetic biology is beginning to provide an improved toolbox and faster process for not only harnessing but also expanding the chemistry of life. At the interface between chemistry and biology, research in cell-free synthetic systems is proceeding in two different directions: using synthetic biology for synthetic chemistry and using synthetic chemistry to reprogram or mimic biology. In the coming years, the impact of advances inspired by these approaches will make possible the synthesis of non-biological polymers having new backbone compositions, new chemical properties, new structures, and new functions.

1. Introduction

Biology is unparalleled in its ability to produce complex molecules and polymers from simple building blocks. By harnessing biosynthetic pathways and macromolecular machines, biological systems convert monomer precursor molecules into a variety of products with atomic-scale resolution over composition, architecture, and functionality. These products are not only precise (defined here as being stereo- and regio-specific), but are also produced with remarkable fidelity, efficiency, and yield. Both the extraordinary synthetic capability of biological systems, as well as their incredible versatility, provides tremendous opportunities for serving society. For instance, biological ensembles have already been used as factories to produce polyketide antibiotics, fatty acid fuels, and protein therapeutics [1–3].

© 2012 Elsevier Ltd. All rights reserved.

[†]Correspondence to m-jewett@northwestern.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The products of biological systems are governed by the chemistry of life. This chemistry is limited to natural building blocks (*e.g.*, nucleic acids, ~20 naturally occurring amino acids, etc.), from which the structures of living organisms are constructed. Strikingly, while biological polymers are structurally narrow, they are functionally diverse. In contrast, chemical polymers have much more structural diversity, but limited functionality (relative to the size of the monomer repertoire) [4]. The potent functionalities and higher order architectures that arise from the relatively simple monomers found in biology inspires us to expand the chemistry of life to include diverse, synthetic monomers that are not limited to those found in nature [5,6].

Indeed, efforts to incorporate non-natural monomers in biological systems are expanding at an accelerated pace. Dominant amongst frontier applications is the manufacturing of proteins with non-canonical amino acids [5,7] and efforts to expand the genetic alphabet [8]. While the lion's share of this work has been conducted *in vivo*, we focus this review on the role of cell-free systems in these synthetic biology efforts. We begin by defining cell-free biology and its advantages. We then describe recent applications that exploit the cell's protein synthesis machinery for synthetic chemistry. Finally, we consider how synthetic chemistry is used for cell-free synthetic biology.

2. Cell-free biology introduction

Cell-free biology is the activation of complex biological processes without using intact living cells [9–11]. Bypassing cell walls, one can access and manipulate biology directly. In contrast to *in vivo* engineering efforts, this direct relationship with biocatalytic enzymes provides the ability to focus metabolism on the production of a single compound, removes physical barriers (allowing easy substrate addition, product removal, and rapid sampling), and eliminates the requirement for cellular viability. Furthermore, removal of native genomic regulation means that microbial growth and engineering design objectives do not conflict. Figure 1 depicts the basic concept of cell-free biology and lists key advantages. The most prominent example of cell-free biology is cell-free protein synthesis (CFPS). Originally used to decipher the genetic code [12], CFPS has emerged as a powerful platform for microscale to manufacturing scale synthesis of complex proteins, opening the way to new applications, such as expanding the chemistry of life [10].

3. Using cell-free synthetic biology for synthetic chemistry

The synthesis of proteins with an expanded repertoire of non-genetically encoded amino acids (ngeAAs) requires (re-)assignment of codons to ngeAAs, ngeAA-transfer RNA (tRNA) substrates (typically created through an orthogonal tRNA–aminoacyl-tRNA synthetase pair), and ribosome selection of these non-natural substrates into the catalytic center [13]. Currently, ~70 ngeAAs have been site-specifically incorporated into proteins [5]. In one exemplary example, incorporating a uniquely reactive ngeAA for site-specific conjugation of polyethylene glycol resulted in a modified human growth hormone with increased potency and reduced injection frequency [14]. This work and others, like a recent breakthrough showing the synthesis of phosphoproteins [15], illustrates the utility of ngeAA incorporation.

Efforts to use cell-free protein synthesis (CFPS) for site-specific incorporation of ngeAAs are beginning to grow [10,16]. The driving force behind this recent growth is two-fold. First, a technical renaissance in CFPS has inspired new applications [10]. Underpinning these developments have been game-changing increases in cost-effective, high-level protein synthesis in both crude extracts and in the Protein synthesis Using Recombinant Elements (PURE) system [10,17]. Jewett *et al.*, for example, demonstrated the ability to activate central metabolism and oxidative phosphorylation to synthesize up to 1.2 mg mL⁻¹ protein in 2 hours [18]. More recently, Dong-Myung Kim's lab used a polymeric carbohydrate to enable the synthesis of 1.7 g L⁻¹ protein in an *E. coli* CFPS system over 10 hours, the highest known reported batch yield to our knowledge [19]. Second, CFPS has now been demonstrated at the manufacturing scale [20]. Zawada *et al.* synthesized 700 mg L⁻¹ of human granulocyte-macrophage colony-stimulating factor (rhGM-CSF) in 10 hours at the 100 liter scale [20]. This pioneering advance has transformed CFPS systems from a foundational tool used solely for bench-top research into an enabling technology at the industrial scale.

Technological improvements in CFPS have opened the way to exploiting advantages over *in vivo* methods for incorporating ngeAAs into proteins (*e.g.*, avoiding transport limitations for getting ngeAAs into the cell). The first design decision is choosing between a crude extract or a purified system [11,17]. Typically, this decision considers trade-offs between cost and productivity versus freedom of design. For example, the cost of the PURE system is prohibitive for most commercial applications, leaving crude extract systems the clear current choice [11]. In addition, cell lysate systems produce more protein per ribosome [21]. On the other hand, the PURE system opens a whole new world of opportunities for co-opting nearly any codon for the incorporation of ngeAAs into a single peptide or protein without recoding organisms because of the ability to omit components (*e.g.*, tRNAs) [7]. Equally important, the PURE system allows for mutating and engineering the components of the translation machinery to enhance ngeAA incorporation without background native components present. Efforts such as a recent report from Wang *et al.* may help to close the cost gap between crude extract and purified systems [22]. Here, the authors report the use of an advanced genome engineering technology [23] to simultaneously tag multiple components from the translation apparatus in a single strain for co-purification and reconstitution *in vitro*.

The Swartz lab has recently demonstrated the power of using CFPS crude extract systems for site-specific incorporation of ngeAAs. In one case, efficient incorporation of p-azido-L-phenylalanine (pAz) was accomplished with high-level production of active chloramphenicol acetyltransferase and dihydrofolate reductase, ~400–600 mg L⁻¹ [16]. Showcasing the freedom of design in adjusting cell-free system components by direct addition to the reaction, both the orthogonal synthetase and tRNA were synthesized in the extract. More recently, Bundy and Swartz leveraged cell-free systems to avoid solubility and transport limitations typically encountered *in vivo* to incorporate the tyrosine analog *p*-propargyloxyphenylalanine (pPa) as well as pAz for demonstrating a one-step, site-specific direct protein-protein conjugation using copper(I)-catalyzed azide-alkyne [3 + 2] cycloaddition (CuAAC) [24]. In another example from the same group, cell-free systems were used to show global replacement of a natural amino acid with an ngeAA analog

without requiring an auxotrophic strain [25]. This approach enabled the synthesis of decorated virus-like particles that could function as potential vaccines and imaging agents. CFPS systems have also been used to site-specifically incorporate ngeAAs at multiple sites in a protein without having to modify the source strain [26] or selectively removing a tagged release factor-1 (RF-1) to remove competition for the amber codon [27]. Moving forward, recent advances to improve ngeAA incorporation efficiency *in vivo*, such as deletion of RF-1 [27–29] or reassigning natural codons to expand the number of co-opted codons [30], should advance *in vitro* efforts that use crude extracts.

Transitioning to the PURE system avoids some key limitations observed in crude extracts [31]. Indeed, genetic code reprogramming is revolutionizing our ability to incorporate multiple and diverse types of ngeAAs into proteins [7]. In pioneering works, Suga and coworkers have demonstrated the ability to synthesize a variety of peptides featuring ngeAAs that would be difficult to prepare by traditional translation or synthetic techniques [32–36]. In a landmark report, they demonstrated the ability to synthesize backbone-cyclized peptides (Figure 2) [34]. This was the first report of ribosomal synthesis of such peptides and highlights the utility of being able to incorporate multiple ngeAAs in a single peptide.

In another exemplary example from the same group, exotic amino acids were expressed by reprogramming of the translation initiation system [33]. Specifically, Flexizyme (a synthetic ribozyme that enables a ngeAA to be charged onto a desired tRNA [33]) was used to charge initiator tRNAs with short peptides, termed Xpeps, comprised of D-amino acids, β -amino acids, and N-methyl amino acids (Figure 2), rather than individual amino acids. Then, the Xpep-tRNA complex was used to initiate ribosomal protein synthesis. Eleven different Xpep sequences (with 2–5 different amino acids) were used, with varying efficiencies of incorporation (12% to 133%) relative to the natural starting residue for all proteins, formyl-methionine. This report opens the way to the synthesis of a wide variety of unusual peptides. Further developments of CFPS coupled with *in vitro* display technologies promise the ability to screen vastly diverse libraries using Darwinian evolution to accelerate the discovery of peptidomimetic drugs. However, low efficiencies of incorporation of non-natural monomers by the ribosome must be addressed [37]. Ribosome engineering methods could be one solution [38].

4. Using synthetic chemistry for cell-free synthetic biology

Complementary to efforts to synthesize protein products containing synthetic monomers, are efforts to re-program interactions of biology using synthetic chemistry. In pushing the limits of chemical design for natural biology, chemists are yielding a deep understanding of natural biology and providing opportunities for expanding the capabilities of canonical systems. Central to this endeavor is the expansion of the genetic alphabet by increasing the capacity of nucleic acids to store, recall, and propagate information. This is a rich and diverse field [39]. Here, we focus our discussion on modulating biological function by exploring base composition. While many researchers, such as Kool and colleagues, have tried to develop a genetic alphabet on a nucleic acid-like system that lacks hydrogen bonding altogether [39], we focus on work that diverges minimally from natural biology. Indeed, such efforts have provided a powerful approach to overcome the limits of natural translational biochemistry

for making new codons for ngeAAs. For example, the creation of post-transcriptionally modified bases [40] or other non-standard bases [41–43] have been used to expand the genetic code (Figure 3).

Now thirty years old, groundbreaking work by Benner and coworkers showed that novel Watson-Crick base pairings could be realized by re-orientating the hydrogen-bond donors and acceptors of cytosine and guanine [44]. This led to the creation of a “65th” codon:anticodon pair that efficiently supported the incorporation of a ngeAA in CFPS from the resulting nucleobases, iso-Cytosine (iso-C) and iso-Guanine (iso-G) (Figure 3) [43]. However, it was observed that during polymerase chain reactions featuring the iso-C and iso-G base pairs, tautomerization of iso-G caused occasional mispairings with thymidine [45]. Furthermore, it was noted that when iso-G and iso-C were assembled into a DNA duplex, they lacked electron rich substituents in the minor-groove, which is important for polymerase function [46].

Recently, the structures of Benner’s synthetic nucleobases have been optimized with respect to their chemical stability and ability to be accepted by various enzymes routinely used in molecular biology for a variety of applications [47]. Specifically, these issues were addressed by the creation of an “Artificially Expanded Genetic Information System (AEGIS)” that deviates less from existing biological design [8]. AEGIS increases the number of independent Watson-Crick bases from four to twelve by rearranging hydrogen bond donor and acceptor groups on the nucleobases [48]. A key recent focus has been on the development of the Z and P nucleobase pair [49]. Z features an electron rich, exocyclic oxygen attached to C2 and P features a nitrogen at the position analogous to N3 of purines. Additionally, Z and P are not susceptible to oxidation and epimerization. Just last year, an expansion to a six base genetic alphabet (G:C, A:T, and Z:P) has been demonstrated with polymerases capable of PCR amplification of a virtually unlimited number of sequences including sequences with multiple, consecutive non-natural Z and P’s [50]. It is tantalizing to consider an unnatural genetic system for cell-free synthetic biology (or even an artificial form of life) with six different nucleotides (GACTZP), or 216 codons.

Beyond new genetic languages, another approach that exploits synthetic chemistry for synthetic biology focuses on mimicking and adapting templated polymerizations for making biomolecules [51]. Here, conventional biological scaffolds are modified with strategic functionality. This results in molecules that leverage natural biological interactions to participate in synthetic biology operations.

Notably, DNA-templated synthesis combines the self-assembly properties of nucleic acids with traditional organic synthesis [52–57]. DNA oligonucleotides are functionalized with reactive synthetic precursors. Thereby, hybridization of complementary oligonucleotides places the reactants in close proximity, boosting their effective molarity (Figure 4). In this manner, multiple reactions can be controlled in a “single pot.” An added benefit of this technique is that large libraries of compounds can be generated by successive addition of assorted complements. Examining the template sequence can then identify the identity of each compound. Looking forward, scalability issues must be addressed for larger adoption of DNA templating technologies in the commercial arena.

Templating synthetic molecules with biology has received much attention in the field of polymer chemistry, where chemists are on a relentless pursuit to exert control over polydispersity and sequence. Once again, nucleobase recognition is being investigated as a mechanism to bridle polymerization reactions [58,59]. In one embodiment, a polymer template featuring thymine nucleobases is synthesized (Figure 4). Monomer featuring adenine is then introduced and subsequently self-assembles with the template. A polymerization reaction then yields a polymer with a narrow molecular weight distribution and chain length equivalent to the template [58]. Future efforts in synthetic polymer chemistry will likely continue to focus on producing more precise templates, with the aim of producing polymers of predefined, exact molecular weights with controllable sequences.

5. Conclusion and outlook

Synthetic chemistry and synthetic biology have been major driving forces in advancing biotechnology products beyond their natural limitations. For this to continue, the interface between synthetic chemistry and synthetic biology must continue to be explored and tinkered with. Immediate challenges include methods for improving the efficiency of incorporation of ngeAAs by the ribosome and the development of robust strategies for expanding the number of available codons for engineering in extract-based systems. Two key areas for growth are likely to include efforts to coordinately tune and evolve the translation apparatus for ngeAA incorporation and strategies to enable the chemically templated synthesis of arbitrary polymers of defined sequence. Addressing these challenges will enable new technologies with applications in therapeutics, practical diagnostics, and advanced materials.

Acknowledgments

The authors would like to thank Rachel K. O'Reilly for her advice and discussions. We also gratefully acknowledge funding from the National Institutes of Health (Grant Number R00GM081450), the National Academies Keck Futures Initiative (Grant Number NAFKI-SB5), the National Science Foundation (Grant Number MCB-0943393), the Office of Naval Research (Grant Number N00014-11-1-0363), the DARPA YFA Program (Grant Number N66001-11-1-4137), and the ARMY Research Office (Grant Number W911NF-11-1-044).

References

- * of special interest
- ** of outstanding interest
- 1. Khosla C. Structures and Mechanisms of Polyketide Synthases. *The Journal of Organic Chemistry*. 2009; 74:6416–6420. [PubMed: 19711990]
- 2. Liu T, Vora H, Khosla C. Quantitative analysis and engineering of fatty acid biosynthesis in *E. coli*. *Metabolic Engineering*. 2010; 12:378–386. [PubMed: 20184964]
- 3. Leader B, Baca QJ, Golan DE. Protein therapeutics: a summary and pharmacological classification. *Nat Rev Drug Discov*. 2008; 7:21–39. [PubMed: 18097458]
- 4. O'Reilly RK. Using controlled radical polymerisation techniques for the synthesis of functional polymers containing amino acid moieties. *Polymer International*. 2010; 59:568–573.
- 5. Liu CC, Schultz PG. Adding New Chemistries to the Genetic Code. *Annual Review of Biochemistry*. 2010; 79:413–444.
- 6. Fritz BR, Timmerman LE, Daringer NM, Leonard JN, Jewett MC. Biology by Design: From Top to Bottom and Back. *Journal of Biomedicine and Biotechnology*. 2010; 2010:11.

7. Ohta A, Yamagishi Y, Suga H. Synthesis of biopolymers using genetic code reprogramming. *Current Opinion in Chemical Biology*. 2008; 12:159–167. [PubMed: 18249198]
8. Benner SA. Understanding Nucleic Acids Using Synthetic Chemistry. *Accounts of Chemical Research*. 2004; 37:784–797. [PubMed: 15491125]
9. Swartz JR. Transforming biochemical engineering with cell-free biology. *AIChE Journal*. 2012; 58:5–13.
10. Carlson ED, Gan R, Hodgman CE, Jewett MC. Cell-free protein synthesis: Applications come of age. *Biotechnology Advances*.
11. Hodgman CE, Jewett MC. Cell-free synthetic biology: Thinking outside the cell. *Metabolic Engineering*.
12. Nirenberg M. Historical review: Deciphering the genetic code - a personal account. *Trends in Biochemical Sciences*. 2004; 29:46–54. [PubMed: 14729332]
13. Noren C, Anthony-Cahill S, Griffith M, Schultz P. A general method for site-specific incorporation of unnatural amino acids into proteins. *Science*. 1989; 244:182–188. [PubMed: 2649980]
14. Cho H, Daniel T, Buechler YJ, Litzinger DC, Maio Z, Putnam A-MH, Kraynov VS, Sim B-C, Bussell S, Javahishvili T, et al. Optimized clinical performance of growth hormone with an expanded genetic code. *Proceedings of the National Academy of Sciences*. 2011; 108:9060–9065.
15. Park H-S, Hohn MJ, Umehara T, Guo L-T, Osborne EM, Benner J, Noren CJ, Rinehart J, Söll D. Expanding the Genetic Code of *Escherichia coli* with Phosphoserine. *Science*. 2011; 333:1151–1154. This new technology shows the critical importance of tuning the entire translation apparatus for incorporation of ngeAAs. Further, it demonstrates directed protein phosphorylation. [PubMed: 21868676]
16. Goerke AR, Swartz JR. High-level cell-free synthesis yields of proteins containing site-specific non-natural amino acids. *Biotechnology and Bioengineering*. 2009; 102:400–416. CFPS used to site-specifically incorporate non-natural amino acid for participation in copper-assisted click chemistry. [PubMed: 18781689]
17. Ohashi H, Kanamori T, Shimizu Y, Ueda T. A Highly Controllable Reconstituted Cell-Free System -a Breakthrough in Protein Synthesis Research. *Current Pharmaceutical Biotechnology*. 2010; 11:267–271. [PubMed: 20210745]
18. Jewett MC, Calhoun KA, Voloshin A, Wu JJ, Swartz JR. An integrated cell-free metabolic platform for protein production and synthetic biology. *Molecular Systems Biology*. 2008;4.
19. Kim H-C, Kim T-W, Kim D-M. Prolonged production of proteins in a cell-free protein synthesis system using polymeric carbohydrates as an energy source. *Process Biochemistry*. 2011; 46:1366–1369.
20. Zawada JF, Yin G, Steiner AR, Yang J, Naresh A, Roy SM, Gold DS, Heinsohn HG, Murray CJ. Microscale to manufacturing scale-up of cell-free cytokine production – a new approach for shortening protein production development timelines. *Biotechnology and Bioengineering*. 2011; 7:1570–1578. CFPS demonstrated at the manufacturing scale, 700 mg L⁻¹ of rhGM-CSF synthesized in 10 hours at the 100 liter scale. [PubMed: 21337337]
21. Jewett MC, Forster AC. Update on designing and building minimal cells. *Current Opinion in Biotechnology*. 2010; 21:697–703. [PubMed: 20638265]
22. Po-Yi, Wang HH.; Xu, George; Haas, Wilhelm; Marblestone, Adam; Li, Jun; Gygi, Steven; Forster, Anthony; Jewett, Michael; Church, George. Multiplexed in vivo His-tagging of enzyme pathways for in vitro single-pot multi-enzyme catalysis. *ACS Synthetic Biology*. 2012 Just accepted.
23. Wang HH, Isaacs FJ, Carr PA, Sun ZZ, Xu G, Forest CR, Church GM. Programming cells by multiplex genome engineering and accelerated evolution. *Nature*. 2009; 460:894–898. [PubMed: 19633652]
24. Bundy BC, Swartz JR. Site-Specific Incorporation of p-Propargyloxyphenylalanine in a Cell-Free Environment for Direct Protein–Protein Click Conjugation. *Bioconjugate Chemistry*. 2010; 21:255–263. [PubMed: 20099875]
25. Patel KG, Swartz JR. Surface Functionalization of Virus-Like Particles by Direct Conjugation Using Azide–Alkyne Click Chemistry. *Bioconjugate Chemistry*. 2011; 22:376–387. Synthesis of

VLPs by the incorporation of a copper-assisted, click chemistry non-natural amino acid. [PubMed: 21355575]

26. Ozawa K, Loscha KV, Kuppan KV, Loh CT, Dixon NE, Otting G. High-yield cell-free protein synthesis for site-specific incorporation of unnatural amino acids at multiple sites. *Biochemical and Biophysical Research Communications*.
27. Loscha KV, Herlt AJ, Qi R, Huber T, Ozawa K, Otting G. Multiple-Site Labeling of Proteins with Unnatural Amino Acids. *Angewandte Chemie International Edition*. 2012;n/a–n/a.
- 28•. Johnson DBF, Xu J, Shen Z, Takimoto JK, Schultz MD, Schmitz RJ, Xiang Z, Ecker JR, Briggs SP, Wang L. RF1 knockout allows ribosomal incorporation of unnatural amino acids at multiple sites. *Nat Chem Biol*. 2011; 7:779–786. Deletion of RF1, and a more promiscuous RF2 generates a strain of bacteria capable of efficiently producing proteins containing ngeAAs at multiple places. [PubMed: 21926996]
29. Mukai T, Hayashi A, Iraha F, Sato A, Ohtake K, Yokoyama S, Sakamoto K. Codon reassignment in the *Escherichia coli* genetic code. *Nucleic Acids Research*. 2010; 38:8188–8195. [PubMed: 20702426]
- 30•. Isaacs FJ, Carr PA, Wang HH, Lajoie MJ, Sterling B, Kraal L, Tolonen AC, Gianoulis TA, Goodman DB, Reppas NB, et al. Precise Manipulation of Chromosomes in Vivo Enables Genome-Wide Codon Replacement. *Science*. 2011; 333:348–353. Genome wide codon editing changes across complete genome. Awaiting completed host strain integrating TAG edits that will likely allow the incorporation of ngeAAs at much higher efficiencies than currently possible. [PubMed: 21764749]
31. Forster AC, Tan Z, Nalam MNL, Lin H, Qu H, Cornish VW, Blacklow SC. Programming peptidomimetic syntheses by translating genetic codes designed de novo. *Proceedings of the National Academy of Sciences*. 2003; 100:6353–6357.
32. Kawakami T, Murakami H, Suga H. Messenger RNA-Programmed Incorporation of Multiple N-Methyl-Amino Acids into Linear and Cyclic Peptides. *Chemistry & Biology*. 2008; 15:32–42. [PubMed: 18215771]
33. Goto Y, Suga H. Translation Initiation with Initiator tRNA Charged with Exotic Peptides. *Journal of the American Chemical Society*. 2009; 131:5040–5041. [PubMed: 19301866]
- 34•. Kawakami T, Ohta A, Ohuchi M, Ashigai H, Murakami H, Suga H. Diverse backbone-cyclized peptides via codon reprogramming. *Nat Chem Biol*. 2009; 5:888–890. Genetic code reprogramming used synthesize backbone-cyclized peptides through the incorporation of a unique, reactive peptide sequence featuring a glycolidic ngeAA. [PubMed: 19915537]
35. Goto Y, Katoh T, Suga H. Flexizymes for genetic code reprogramming. *Nat Protocols*. 2011; 6:779–790.
36. Ohshiro Y, Nakajima E, Goto Y, Fuse S, Takahashi T, Doi T, Suga H. Ribosomal Synthesis of Backbone-Macrocyclic Peptides Containing γ -Amino Acids. *ChemBioChem*. 2011; 12:1183–1187. [PubMed: 21506233]
37. Pavlov MY, Watts RE, Tan Z, Cornish VW, Ehrenberg M, Forster AC. Slow peptide bond formation by proline and other N-alkylamino acids in translation. *Proceedings of the National Academy of Sciences*. 2009; 106:50–54.
38. Neumann H, Wang K, Davis L, Garcia-Alai M, Chin JW. Encoding multiple unnatural amino acids via evolution of a quadruplet-decoding ribosome. *Nature*. 2010; 464:441–444. [PubMed: 20154731]
39. Krueger AT, Kool ET. Redesigning the Architecture of the Base Pair: Toward Biochemical and Biological Function of New Genetic Sets. *Chemistry & Biology*. 2009; 16:242–248. [PubMed: 19318205]
40. Karijolich J, Yu Y-T. Converting nonsense codons into sense codons by targeted pseudouridylation. *Nature*. 2011; 474:395–398. [PubMed: 21677757]
41. Hirao I, Ohtsuki T, Fujiwara T, Mitsui T, Yokogawa T, Okuni T, Nakayama H, Takio K, Yabuki T, Kigawa T, et al. An unnatural base pair for incorporating amino acid analogs into proteins. *Nat Biotech*. 2002; 20:177–182.

42. Hirao I, Harada Y, Kimoto M, Mitsui T, Fujiwara T, Yokoyama S. A Two-Unnatural-Base-Pair System toward the Expansion of the Genetic Code. *Journal of the American Chemical Society*. 2004; 126:13298–13305. [PubMed: 15479084]
43. Bain JD, Switzer C, Chamberlin R, Bennert SA. Ribosome-mediated incorporation of a non-standard amino acid into a peptide through expansion of the genetic code. *Nature*. 1992; 356:537–539. [PubMed: 1560827]
44. Piccirilli JA, Benner SA, Krauch T, Moroney SE. Enzymatic incorporation of a new base pair into DNA and RNA extends the genetic alphabet. *Nature*. 1990; 343:33–37. [PubMed: 1688644]
45. Sismour AM, Benner SA. The use of thymidine analogs to improve the replication of an extra DNA base pair: a synthetic biological system. *Nucleic Acids Research*. 2005; 33:5640–5646. [PubMed: 16192575]
46. Joyce CM, Steitz TA. Function and Structure Relationships in DNA Polymerases. *Annual Review of Biochemistry*. 1994; 63:777–822.
47. Yang Z, Chen F, Chamberlin SG, Benner SA. Expanded Genetic Alphabets in the Polymerase Chain Reaction. *Angewandte Chemie International Edition*. 2010; 49:177–180.
- 48•. Yang Z, Hutter D, Sheng P, Sismour AM, Benner SA. Artificially expanded genetic information system: a new base pair with an alternative hydrogen bonding pattern. *Nucleic Acids Research*. 2006; 34:6095–6101. Expansion of codon alphabet to a six letter system utilizing AEGIS shown to be highly effective when used in PCR. [PubMed: 17074747]
49. Hutter D, Benner SA. Expanding the Genetic Alphabet: Non-Epimerizing Nucleoside with the pyDDA Hydrogen-Bonding Pattern. *The Journal of Organic Chemistry*. 2003; 68:9839–9842. [PubMed: 14656120]
50. Yang Z, Chen F, Alvarado JB, Benner SA. Amplification, Mutation, and Sequencing of a Six-Letter Synthetic Genetic System. *Journal of the American Chemical Society*. 2011; 133:15105–15112. [PubMed: 21842904]
- 51•. McHale R, Patterson JP, Zetterlund PB, O'Reilly RK. Biomimetic Radical Polymerization via Cooperative Assembly of Segregating Templates. *Nature Chemistry*. 2012 (in press). Biological mimicking polymer bearing nucleobases is capable of self-assembling into higher-ordered structures.
52. Gartner ZJ, Tse BN, Grubina R, Doyon JB, Snyder TM, Liu DR. DNA-Templated Organic Synthesis and Selection of a Library of Macrocycles. *Science*. 2004; 305:1601–1605. [PubMed: 15319493]
53. Gartner ZJ, Kanan MW, Liu DR. Multistep Small-Molecule Synthesis Programmed by DNA Templates. *Journal of the American Chemical Society*. 2002; 124:10304–10306. [PubMed: 12197733]
- 54•. He Y, Liu DR. A Sequential Strand-Displacement Strategy Enables Efficient Six-Step DNA-Templated Synthesis. *Journal of the American Chemical Society*. 2011; 133:9972–9975. The multistep synthesis of an oligamide without the addition of chemical activators or the cleavage of linkers and protective groups by combining DNA-templated synthesis with a DNA walker. [PubMed: 21657248]
55. He Y, Liu DR. Autonomous multistep organic synthesis in a single isothermal solution mediated by a DNA walker. *Nat Nano*. 2010; 5:778–782.
- 56••. Brudno Y, Birnbaum ME, Kleiner RE, Liu DR. An in vitro translation, selection and amplification system for peptide nucleic acids. *Nat Chem Biol*. 2010; 6:148–155. The authors developed a process in which peptide nucleic acids may be used for in vitro evolution. [PubMed: 20081830]
57. McKee ML, Milnes PJ, Bath J, Stulz E, Turberfield AJ, O'Reilly RK. Multistep DNA-Templated Reactions for the Synthesis of Functional Sequence Controlled Oligomers. *Angewandte Chemie International Edition*. 2010; 49:7948–7951.
58. Lo PK, Sleiman HF. Nucleobase-Templated Polymerization: Copying the Chain Length and Polydispersity of Living Polymers into Conjugated Polymers. *Journal of the American Chemical Society*. 2009; 131:4182–4183. [PubMed: 19275231]
59. Spijker HJ, Dirks AJ, van Hest JCM. Unusual rate enhancement in the thymine assisted ATRP process of adenine monomers. *Polymer*. 2005; 46:8528–8535.

Highlights

- We focus this review on the role of cell-free systems in these synthetic biology efforts.
- Cell-free biology is examined along with its advantages.
- We then describe recent applications of exploiting the cell's protein synthesis machinery for synthetic chemistry.
- Finally, we consider how synthetic chemistry is used for cell-free synthetic biology.

Figure 1.

Cell-free biology is a rapidly developing technology for making advanced materials, sustainable biomolecules, and life-saving medicines. This approach provides numerous advantages for exploiting the interface between synthetic biology and synthetic chemistry.

Figure 2.

Exploiting cell-free protein synthesis for producing non-natural peptides. (A) Flexizyme (blue) charges an initiator tRNA (red) with an Xpep (multi-colored), which is expressed by the PURE system. (B) Flexizyme (blue) charges a tRNA (red) with glycolic acid (black). Glycolic acid is then expressed adjacent to a proline (red pentagon) and cysteine (blue circle) residue. (C) An intramolecular rearrangement occurs, resulting in the formation of a dkp-thioester and loss of the glycolic acid portion of the peptide. The resulting dkp-thioester can then react with the N-terminal amine to form a backbone-cyclic peptide.

Figure 3.

An expanded genetic alphabet for synthetic biology. (A) Iso-C and iso-G with shuffled hydrogen-bond donors and acceptors in red [43]. (B) Z and P nucleobases, note electron rich substituents on minor groove face [49]. (C) The thymidine isostere dF, with weak hydrogen-bond acceptor fluorine and hydrogen-bond donating carbon-hydrogen bond in red [39]. (D) Alternate base-pairing thiophene-purine derivative S and imidazolinone derivative I [42].

Figure 4.

(A) DNA-templated synthesis of small-molecules. An oligonucleotide with terminus functionalized with a reactive precursor (black triangle) is subjected to a partially complementary oligonucleotide bearing a second reactant (red circle). (B) After the first reaction is complete, a fully complementary oligonucleotide bearing a third reactant (blue hexagon) is introduced. In this manner, small-molecules can be synthesized with precision. (C) A polymer template featuring thymine nucleobase repeat units (red polygons) is subjected to adenine nucleobase monomers (black). (D) Upon self-assembly of the template-monomer complexes, a polymerization reaction yields a polymer with similar structure.