CASE REPORT

Syringocystadenoma papilliferum of the vulva: a rarity in gynaecology

Oleksandr Steshenko, Nirmala Chandrasekaran, Frank Lawton

Department of Obstetrics and Gynaecology, King's College Hospital, London, UK

Correspondence to
Dr Oleksandr Steshenko,
asteshenko@nhs.net

Accepted 9 May 2014

SUMMARY

A case of syringocystadenoma papilliferum of the vulva in a 36-year-old woman is reported. The patient presented with a single cystic lesion on the left labia minora of 8 years duration with a recent increase in size and redness around the lesion. Examination revealed a polypoid cystic lesion with no regional lymphadenopathy. An excision biopsy was performed under general anaesthesia and the specimen was subjected to histological examination. Histopathology revealed closely excised syringocystadenoma papilliferum. On follow-up, the patient was asymptomatic and the wound had healed well.

CASE PRESENTATION

A 36-year-old woman presented with a single cystic lesion on the left labia minora of 8 years duration. The patient reported increase in redness around the cyst and increase in size which she noticed over the last few days.

On examination of the vulva, it was noticed that on the normal surface of the skin there was a 6 × 3 mm polypoid tan lesion with a 5 mm cyst filled content. The patient was consented for excision of the lesion which was performed under general anaesthesia in our hospital. The lesion was closely excised and the specimen was sent for histological examination. The wound was closed with Vicryl sutures.

OUTCOME AND FOLLOW-UP

The patient had uneventful recovery and discharged home the same day. She was asymptomatic and her wound healed well when we followed her up 3 months after the operation.

DISCUSSION

Syringocystadenoma papilliferum (SP) is a rare neoplasm which described in the literature as a benign hamartomatous adnexal tumour with unclear histopathogenesis.¹ The most common sites for presentation of the SP are the head and neck (up to 75%), the trunk (20.0–53.5%), on the extremities (5.0–33.8%) and (12.5%) on the genitalia.^{2–7} It is less likely to develop on the abdomen as just two cases have been reported in the literature.^{8,9} This neoplasm can be of various sizes from 5 to 160 mm¹ in a form of single or multiple polypoid, cystic, nodulated lesion which could be brown or erythematous in colour^{1,7} presenting at birth or develop during puberty.^{6,7,10}

The origin of the SP remains unclear. There is a possibility that SP can resemble lesions of the

breasts deriving from the supernumerary mammary tissues from milk ridges.^{11,12} In some individuals those ridges may not regress completely during embryological development forming breast-like tissue.¹³ This can be explained by evidence of previously reported cases of patients with the presence of lactating mammary tissues in the vulva.^{13–17} Other possibilities are that SP may develop from 'sweat' glands of the skin and mucosa.¹⁸ Evidence to that could be its close association with tumours from hair follicles, sebaceous and apocrine glands.¹⁹

Although benign, SP can be associated with malignant tumours such as basal cell carcinoma, metaplastic adenocarcinoma and ductal carcinoma.^{20,21} There has been report in the literature of pure intraductal carcinoma of the breast and invasive carcinoma of ectopic breast tissue of the vulva detected simultaneously in the same patient.²² SP may also be linked to its malignant counterpart syringocystadenoma carcinoma papilliferum.²⁰ It is difficult to determine what the estimated risk of malignant transformation of the SP is due to its rarity.

Diagnosis of SP based on clinical examination can be a challenging task in view of its variety in presentation and similar appearance with molluscum contagiosum, syringoma, basal cell carcinoma.

Histopathological examination of SP shows as in our case the cystic component, ducts with intraluminal papillary proliferation of the lining epithelium in two layers with apocrine differentiation and sclerotic fibroconnective tissue (figures 1–4).

Figure 1 Histological examination showing sections of vulval skin with reactive hyperplasia of the epidermis which contained a fairly circumscribed tumour composed of cystic component and tubules arranged in lobular pattern.

To cite: Steshenko O, Chandrasekaran N, Lawton F. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-203902

Figure 2 The cystic component involved ducts with intraluminal papillary proliferation of the epithelial lining in two layers with apocrine differentiation.

Using immunohistochemical staining, which is positive to gross cystic disease fluid protein 15 (GCDPF-15), carcinoembryonic antigen and epithelial membrane antigen can support the diagnosis of SP.^{1 20}

SP in the vulvar region is an extremely rare condition in gynaecological practice with a few cases identified in the literature.^{2–5} These cases were reported in women between 27 and 64 years old.

Figure 3 Adjacent to the cystic component irregular tubules lined by two layers of cells were seen with intervening sclerotic fibroconnective tissue and lymphoplasmic cell infiltration.

Figure 4 Although the tumour involved epidermis in addition to the dermis, there was no evidence of malignancy.

In view of possible malignant transformation of SP, its complete excisional biopsy with a 3–5 mm margin of healthy skin is recommended.⁸ It is unclear from cases reported in the literature regarding the extent and depth of excision.

We recommend including SP in the differential diagnosis of vulvar lesions. The excisional biopsy seems to be an effective treatment of this condition.

The guidelines for the treatment of SP remain unavailable because of the rarity of this condition. We believe that this type of tumour should be excised because of the possibility of its transformation into adenocarcinoma.

Learning points

- Syringocystadenoma papilliferum in the vulvar region is a rare condition presenting as papule lesion in the age group of 27–64 years.
- Gynaecological assessment has to be performed to exclude potential for malignancy.
- Excisional biopsy is a definitive method of treatment.

Acknowledgements The authors acknowledge the Department of Obstetrics and Gynaecology, King's College Hospital, London. The authors also thank Dr Hizbullah Shaikh, Consultant Hystopathologist, for providing the histology slides.

Contributors OS was involved in the literature search, preparation of the manuscript, concept and design of the case report. NC was involved in the definition of intellectual content, conducting literature search, preparation and editing of the manuscript. FL was involved in editing of the manuscript.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Yap FBB, Lee BR, Baba R. Syringocystadenoma papilliferum in an unusual location beyond the head and neck region. A case report and review of literature *Dermatol Online J* 2010;16:4.
- 2 Tseung J, Russell P. 'Breast-like' lesions in the vulva: histology, origin and significance. *Pathology* 2008;40:321–6.
- 3 Al-Brahim N, Daya D, Alowami S. A 64-year-old woman with vulvar papule. *Arch Pathol Lab Med* 2005;129:e126–7.
- 4 Nishie W, Sawamura D, Mayuzumi M, et al. Hidradenoma papilliferum with mixed histopathologic features of syringocystadenoma papilliferum and anogenital mammary-like glands. *J Cutan Pathol* 2004;31:561–4.
- 5 Stewart CJR. Syringocystadenoma papilliferum-like lesion of the vulva. *Pathology* 2008;40:638–9.
- 6 Karg E, Korom I, Varga E, et al. Congenital syringocystadenoma papilliferum. *Pediatr Dermatol* 2008;25:132–3.
- 7 Mammino J, Vidmar D. Syringocystadenoma papilliferum. *Int J Dermatol* 1991;30:763–6.
- 8 Xu D, Bi T, Lan H, et al. Syringocystadenoma papilliferum in the right lower abdomen: a case report and review of literature. *Oncol Targets Ther* 2013;6:233–6.
- 9 Suzuki T, Ikeda H, Hamasaki Y, et al. Syringocystadenoma papilliferum associated with apocrine poroma. *J Dermatol* 2006;33:249–51.
- 10 Böni R, Xin H, Hohl D, et al. Syringocystadenoma papilliferum: a study of potential tumor suppressor genes. *Am J Dermatopathol* 2001;23:87–9.
- 11 Barua R. *Tumors of the lower genital tract*. Berlin: Springer Verlag, 1990.
- 12 Van der Putte SCJ. Mammary-like glands of the vulva and their disorders. *Int J Gynecol Pathol* 1994;13:150.
- 13 Deaver JB, McFarland J. *The breast, its anomalies, its diseases and their treatment*. London: W Heinemann, 1918.
- 14 Levin N, Diener RL. Bilateral ectopic breast of the vulva. *Obstet Gynaecol* 1968;32:274–6.
- 15 Bell JW. Supernumerary breast near labium. *Am J Obstet Gynecol* 1926;11:507–9.
- 16 Irvin WP, Cathro HP, Grosh WW, et al. Primary breast carcinoma of the vulva: a case report and literature review. *Gynecol Oncol* 1999;73:155–9.

- 17 Pfeifer JD, Barr RJ, Wick MR. Ectopic breast tissue and breast-like sweat gland metaplasias: an overlapping spectrum of lesions. *J Cutan Pathol* 1999;26:190–6.
- 18 Van der Putte SCJ. Anogenital 'sweat' glands. Histology and pathology of a gland that may mimic mammary glands. *Am J Dermatopathol* 1991;13:557–67.
- 19 Mazoujian G, Margolis R. Immunohistochemical gross cystic disease fluid protein (GCDPF-15) in 65 benign sweat gland tumors of the skin. *Am J Dermatopathol* 1988;10:28–35.
- 20 Arai Y, Kusakabe H, Kiyokane K. A case of syringocystadenocarcinoma papilliferum in situ occurring partially in syringocystadenoma papilliferum. *J Dermatol* 2003;30:146–50.
- 21 Diaz RC, Babu SC. Ductal carcinoma arising from syringocystadenoma papilliferum in the external auditory canal. *Otol Neurotol* 2007;28:873–4.
- 22 Intra M, Maggioni A, Sonzogni A, et al. A rare association of synchronous intraductal carcinoma of the breast and invasive carcinoma of ectopic breast tissue of the vulva: case report and literature review. *Int J Gynecol Cancer* 2006;16 (Suppl 1):428–33.

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow