Published in final edited form as:

Psychopharmacology (Berl). 2014 June ; 231(12): 2417–2433. doi:10.1007/s00213-013-3407-z.

History of Childhood Adversity is Positively Associated with Ventral Striatal Dopamine Responses to Amphetamine

Lynn M. Oswald^{a,b}, Gary S. Wand^{b,c}, Hiroto Kuwabara^d, Dean F. Wong^{c,d,e,f}, Shijun Zhu^a, and James R. Brasic^d

^aDepartment of Family and Community Health, University of Maryland School of Nursing, Baltimore, MD 21201, USA

^bDepartment of Medicine, The Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

^cDepartment of Psychiatry and Behavioral Sciences, The Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

^dDepartment of Radiology, The Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

^eDepartment of Neuroscience, The Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

^fDepartment of Environmental Health Sciences, The Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

Abstract

Childhood exposure to severe or chronic trauma is an important risk factor for the later development of adult mental health problems, such as substance abuse. Even in nonclinical samples of healthy adults, persons with a history of significant childhood adversity seem to experience greater psychological distress than those without this history. Evidence from rodent studies suggests that early life stress may impair dopamine function in ways that increase risks for drug abuse. However, the degree to which these findings translate to other species remains unclear. This study was conducted to examine associations between childhood adversity and dopamine and subjective responses to amphetamine in humans. Following intake assessment, 28 healthy male and female adults, ages 18–29 years, underwent two consecutive 90-minute positron emission tomography (PET) studies with high specific activity [¹¹C]raclopride. The first scan was preceded by intravenous saline; the second by amphetamine (AMPH 0.3 mg/kg). Consistent with prior literature, findings showed positive associations between childhood trauma and current levels of perceived stress. Moreover, greater number of traumatic events and higher levels of perceived stress were each associated with higher ventral striatal dopamine responses to AMPH. Findings of mediation analyses further showed that a portion of the relationship between childhood trauma and dopamine release may be mediated by perceived stress. Overall, results are

consistent with preclinical findings suggesting that early trauma may lead to enhanced sensitivity to psychostimulants and that this mechanism may underlie increased vulnerability for drug abuse.

Keywords

Dopamine; Positron Emission Tomography (PET); Human; Amphetamine; Childhood Adversity; Stress

1. Introduction

Childhood adversity can be defined as negative events occurring during childhood or adolescence which are 1) outside of the control of the child, 2) have the potential to alter normal development, and 3) may impair the child's physical and/or psychological well-being (Burgermeister. 2007). In recent years, there has been increasing awareness that adverse childhood experiences (ACEs) may contribute to the development of a wide range of mental health problems in adulthood (Benjet et al. 2010; Cuijpers et al. 2011; Gordon. 2002; Huang et al. 2011; Husted et al. 2012; Matheson et al. 2012; Nanni et al. 2012; Nelson et al. 2002; Weber et al. 2008). Greater severity, frequency, and duration of trauma are generally associated with greater likelihood of such problems and with worse prognosis (D'Andrea et al. 2012; Edwards et al. 2003; Felitti et al. 1998; Finkelhor et al. 2007; Ford et al. 2010; Kessler et al. 2010). Moreover, it has been shown that even children who do not go on to develop adult psychopathology may suffer long-term consequences of childhood maltreatment. In non-clinical samples of adults, individuals with a history of ACEs often report greater psychological distress, emotional reactivity, and sensitivity to stress than those without this history (Chu et al. 2013; Edwards et al. 2003; Glaser et al. 2006; Hyman et al. 2007; Marx and Sloan. 2003).

One potentially modifiable consequence of childhood adversity is elevated risks for substance abuse. Numerous studies have documented associations between childhood adversity and the development of substance use disorders in adults (Carswell et al. 2008; Kendler et al. 2000; Pilowsky et al. 2009; Rothman et al. 2008). Although no single pathway can be expected to entirely explain this complex relationship, findings showing that substance abusers with a history of ACEs report greater psychological distress than those without this history suggest that alcohol and/or drugs may be used as a form of self-medication by these groups (Medrano et al. 2002; Wang et al. 2010). There is also a large body of research showing that perceived stress and current daily hassles are also positively associated with substance use and related problems (Back et al. 2008; Cole et al. 1990; Corbin et al. 2013; Kaplan. 2013; Nakajima and al' Absi. 2012; Waldrop et al. 2007).

A large body of preclinical literature suggests that both acute and chronic stress may influence vulnerability for drug abuse by altering mesocorticolimbic dopamine (DA) neurotransmission (Belujon and Grace. 2011; Kosten et al. 2003; Lu et al. 2003; Mangiavacchi et al. 2001; Marinelli and Piazza. 2002; Miczek et al. 2011; Saal et al. 2003; Shimamoto et al. 2011; Yavich and Tiisonen. 2000). Interestingly, there is also preclinical evidence showing that early life stress may lead to long-term derangements in DA neurotransmission that persist into adulthood. Rodents exposed to early life stress exhibit

altered ventral striatal (VS) and nucleus accumbens (NAcc) DA responses to stress in later life (Brake et al. 2004; Chocyk et al. 2011; Fulford and Marsden. 1998; Jahng et al. 2010; Meaney et al. 2002; Hall et al. 1999), as well as enhanced DA responses to psychostimulants (Hall et al. 1999; Kosten et al. 2005; Moffett et al. 2006) and increased alcohol/drug consumption (Kosten et al. 2005; Moffett et al. 2007; Zhang et al. 2005). These preclinical findings suggest that early life trauma has sustained effects on brain DA systems, which may lead to altered drug sensitivity and greater vulnerability for drug abuse in later life (Rodrigues et al. 2011).

Clinical studies examining the impact of childhood trauma on brain DA function and associated risks for drug abuse are sparse. In one functional magnetic resonance imaging (fMRI) study, Dillon et al. (2009) showed dysfunction in left basal ganglia reward pathways in adults who were abused as children. Mehta and colleagues (2010) similarly reported that adolescents who experienced severe global deprivation early in life exhibited hyporesponsivity of the VS and, to a lesser extent, the caudate nucleus during anticipation of monetary reward. Although it is likely that this hyporesponsiveness was related to altered DA activity, interpretation of the precise nature of the underlying dysfunction was limited by the lack of clear understanding about relationships between fMRI blood-oxygen-level dependent (BOLD) response and DA neural function. Using positron emission tomography (PET) imaging, Pruessner and colleagues (2004) found that persons who reported low maternal care had greater stress-induced changes in VS [^{11}C]raclopride binding potential (BP_{ND}) than individuals who reported high maternal care. In the present study, we selected intravenous (IV) amphetamine (AMPH) to interrogate mesolimbic DA function, conducting the first examination of relationships among childhood adversity, perceived stress, and DA and subjective drug responses in humans. Participants were healthy young adults who did not meet diagnostic criteria for a major DSM-IV psychiatric Axis I disorder (American Psychiatric Association. 2000). The advantage of conducting the study in healthy adults without psychopathology was that it allowed us to rule out alternative explanations for hypothesized relationships between childhood trauma and DA functioning. Because of prior evidence that the NAcc plays a central role in the reinforcing effects of psychostimulants and in reward-based behaviors (Di Chiara et al. 2004; McGinty et al. 2013; Robison et al. 2013; Wise. 1996), the VS was our primary region of interest in this study. Our primary hypotheses were that childhood trauma would be positively associated with 1) VS AMPH-induced DA release, 2) pleasant drug effects, and 3) current levels of perceived stress.

2. Materials and methods

2.1 Participants

Twenty-eight healthy male ($n = 19$) and female ($n = 9$) participants, aged 18–29 years, were recruited from the Baltimore metropolitan area by newspaper advertisements, fliers, and internet postings. Respondents who appeared to qualify for the study on the basis of a brief telephone screen were invited to our research offices at the University of Maryland to complete an intake session. All volunteers provided written informed consent after being given a complete description of the study. The research was approved by the University of Maryland and the Johns Hopkins Medicine Institutional Review Boards.

Psychiatric diagnostic screening was conducted using the Structured Clinical Interview for DSM Disorders (SCID-I/NP) (First et al. 2002). Alcohol consumption and drug use were further evaluated by 90-day time line follow back (Sobell and Sobell. 1992). Additional screening measures included the Beck Depression Inventory (Beck et al. 1996), Shipley Institute of Living Scale-2 (Shipley et al. 2009), Fagerstrom Test for Nicotine Dependence (Heatherton et al. 1991), alcohol breathalyzer test, and urine toxicology. Participants also completed a medical evaluation that consisted of a history and physical examination, complete blood count, comprehensive metabolic panel (including renal and hepatic function tests), coagulation tests, pregnancy screen/progesterone levels (women), electrocardiogram, and urinalysis. The medical evaluation was conducted at the University of Maryland School of Medicine General Clinical Research Center (GCRC).

Exclusion criteria included: (a) younger than 18 or older than 30 years; (b) lifetime history of a major DSM-IV psychiatric Axis I disorder, including attention deficit hyperactivity disorder (ADHD) or conduct disorder; (c) BDI score > 18; (d) currently in need of psychiatric treatment; (e) illicit drug use during the past 30 days or lifetime stimulant use; (f) positive urine drug screen; (g) nicotine dependence or smokes > 10 cigarettes weekly; (h) heavy alcohol consumption; (i) oral contraceptive use, pregnant or lactating; (j) active medical conditions; (k) treatment (past 12 months) with antidepressants, neuroleptics, sedative/hypnotics, appetite suppressants, opiates, antihypertensives, dopamine or serotonergic agents, glucocorticoids, or propecia; (l) claustrophobia or fear of needles; (m) BMI < 20 or > 31; and (n) less than 5th grade reading level.

Qualified participants were scheduled to complete MRI and PET scans at the Johns Hopkins Hospital Department of Radiology. For women, PET scans were scheduled during the follicular phase of their menstrual cycle, confirmed by progesterone levels drawn on the day of the session. Women with progesterone levels < 2 ng/ml were classified as being in the follicular phase of the menstrual cycle.

2.2 Baseline Assessment Measures

Self-report measures administered during the intake assessment to evaluate primary and secondary hypotheses of the study included the Perceived Stress Scale (Cohen. 1994) and the Early Trauma Inventory Short Form (Bremner et al. 2007).

Perceived Stress Scale—The PSS is a 10-item questionnaire that measures the degree to which situations are appraised as stressful. In contrast to life event measures, which assess objective life events, the PSS is a global measure designed to tap how unpredictable, uncontrollable, and overloaded respondents find their lives. Items are answered on a 5-point scale from “never” to “very often”, e.g., “In the last month, how often have you found that you could not cope with all the things that you had to do?” The PSS has shown adequate internal consistency and test-retest reliability (Cohen et al. 1983), as well as criterion-related and construct validity (Cohen. 1994).

Early Trauma Inventory Short Form—The 27-item ETISR-SF is a self-report measure that assesses traumatic events that occurred before the age of 18 years in each of four domains: general trauma (11 items), physical punishment (5 items), emotional abuse (5

items), and sexual abuse (6 items). The ETI “total score” is derived by summing the number of items positively endorsed in each domain, which gives an index of the total extent of abuse. The ETI-SF was adapted from the earlier 62-item self-report version of the measure (ETI-SR) through factor analysis and other correlational methods and has been shown to have similarly acceptable psychometric properties of validity and internal consistency in healthy and clinical populations (Bremner et al. 2007; Plaza et al. 2011; Tonmyr et al. 2011; Wang et al. 2010).

2.3 MRI Acquisition

A spoiled gradient (SPGR) sequence using the 3 T Siemens Trio MRI was obtained on each subject for anatomical identification of the structures of interest using the following parameters: Repetition time, 35 ms; echo time, 6 ms; flip angle, 45°; slice thickness, 1.5 mm with no gap; field of view, 24 × 18 cm²; image acquisition matrix, 256 × 192, reformatted to 256 × 256.

2.4 PET Experiments

2.4.1 General procedures—Participants were asked to arrive at our research office at Johns Hopkins at 08:30 on the day of the scans fasting, except for water. A brief medical evaluation, urine toxicology screen, alcohol breathalyzer test, hematocrit level, and serum pregnancy test (women) were conducted. Serum progesterone levels were also obtained for women. A caffeine-free, calorie controlled breakfast and lunch were provided. PET data acquisition commenced at 13:00. Both scans were conducted on the same day as described below.

2.4.2 PET data acquisition—Upon arrival at the PET center, a venous catheter was placed in the antecubital vein for the radioligand injection and saline/AMPH administration. Subjects were positioned in the scanner with their heads restrained by a custom-made thermoplastic mask to reduce head motion. A 6-min transmission scan was acquired using a rotating Cs-137 source for attenuation correction. Each subject had two scans performed on the High Resolution Research Tomograph scanner (HRRT, CPS Innovations, Inc., Knoxville, TN) with a high specific activity IV bolus injection of [¹¹C]raclopride, an antagonist radioligand for dopamine D₂/D₃ receptors, following bolus IV injection of equal volumes of either saline or AMPH (0.3 mg/Kg). Dynamic PET acquisition was performed in a three-dimensional list mode for 90 minutes following each injection of [¹¹C]raclopride. The [¹¹C]raclopride was prepared with minor changes in purification and formulation according to published procedure (Ehrin et al. 1985). Participants had approximately 45 minutes between the two scans to get up to stretch and void if needed. All participants were under continuous cardiovascular monitoring during the scans. Because of potential carry-over effects of AMPH, the order of drug administration is routinely fixed when both scans are completed on the same day; saline was administered during the first scan and AMPH during the second in all cases. One participant did not complete both scans on the same day due to equipment problems; however, the saline scan was done on one day and the AMPH scan on the following day. All participants were blind to the order of drug administration.

2.4.3 Reconstruction of PET data—Emission PET scans were reconstructed using the iterative ordered-subset expectation-maximization algorithm correcting for attenuation, scatter, and dead-time (Rahmim et al. 2005). The radioactivity was corrected for physical decay to the injection time and re-binned to 30 dynamic PET frames of 256 (left-to-right) by 256 (nasion-to-inion) by 207 (neck-to-cranium) voxels of 1.2188 mm cubic dimensions. The frame schedules were four 15-s, four 30-s, three 1-min, two 2-min, five 4-min, and twelve 5-min frames. The final spatial resolution is expected to be less than 2 mm full-width at half-maximum in three directions (Rahmim et al. 2005).

2.4.4 Subjective assessments—Visual analog scales of subjective drug effects were administered at scheduled intervals following saline and AMPH administration as previously described (Oswald et al. 2005). The measure included five scales asking about pleasant drug effects (i.e., good effects, high, rush, liking, and desire for drug) and six asking about negative effects (i.e., bad effects, fidgety, anxious, dizziness, dry mouth, and distrust). Each effect was rated on a scale ranging from 0 “not at all” to 10 “extremely”.

2.4.5 Cortisol assays—Measurements of baseline plasma cortisol levels were obtained prior to each scan (–25 and –5 min). Plasma concentrations were measured as previously described (Oswald et al. 2005).

2.5 PET Data Analysis

2.5.1 Volumes of interest (VOIs)—VOIs for putamen, caudate nucleus, and cerebellum were defined on MRI using the 3-D interactive-segmentation mode of a locally developed VOI defining tool (VOILand), as previously reported (Oswald et al. 2005). Striatal VOIs were subdivided into motor (posterior putamen), associative striatum (anterior putamen and anterior and posterior caudate nucleus) and limbic ventral striatum subdivisions using a semi-automated method that incorporated anatomical guidance based on postmortem human materials (Baumann et al. 1999; Oswald et al. 2005). VOIs were transferred from MRI to PET space according to MRI-to-PET coregistration parameters obtained with the coregistration module SPM5 (The Statistical Parametric Mapping 5; The Wellcome Trust Centre for Neuroimaging; available at www.fil.ion.ac.uk/spm) (Ashburner and Friston. 2003), and applied to PET frames to obtain regional time (radio-)activity curves (TACs). PET frames were aligned to the frame of the highest counts (decay-uncorrected) using the SPM coregistration module to correct head motion during the scan (Kumar et al. 2007; Montgomery. 2006). VOIs of two sides were unified after confirming no side-to-side differences in PET outcome variables described below.

2.5.2 Derivation of PET outcome variables—Binding potential (BP_{ND}) (Innis et al. 2007) of [^{11}C]raclopride was obtained by the reference tissue graphical analysis (RTGA) (Logan et al. 1996) for striatum subdivisions. Then, intrasynaptic DA release, which represents the displacement of [^{11}C]raclopride by endogenous DA (DA_{rel} in %) (Innis et al. 1992), was obtained using the following formula: $(BP_{ND}[S] - BP_{ND}[A]) / BP_{ND}[S] \times 100$, where [S] and [A] stands for BP_{ND} of saline and AMPH scans.

2.6 Statistical Analyses

Associations between participant characteristics and primary outcome variables were examined using Pearson Product Moment correlations for continuous variables and independent samples *t*-tests for gender. Paired *t*-tests were used to examine differences in [¹¹C]raclopride BP_{ND} and in subjective drug effects between the saline and AMPH sessions. Drug effect ratings were summarized as area under the time response curve (AUC), calculated by trapezoidal approximation from 3 to 85 minutes following drug administration. The AUC values for the five measures of pleasant effects were highly intercorrelated within each session, internal reliability was moderately high, and findings of factor analysis confirmed that one saline and one active session variable could be extracted (eigenvalues > 1). Therefore, for purposes of data reduction, we derived composite AMPH and saline session ‘pleasant effects’ scores for each subject by averaging his/her AUC values on these five items. Findings of factor analysis did not support similar derivation of composite scores for the six scales measuring unpleasant drug effects. Because we were interested primarily in individual differences in pleasant subjective responses to AMPH and to avoid running multiple tests, no further analyses were conducted on unpleasant effects.

Evaluation of progesterone levels suggested that two of the female participants were not in the follicular phase of their menstrual cycle on the day of the scans (level > 2 ng/ml). However, *t*-test findings indicated that VS DA release values for these two women did not differ from those of the other women who completed the scans. Therefore, data from all nine women were included in subsequent analyses.

Linear regression models were used to evaluate each of the primary hypotheses. The independent variable of interest in each regression was the ETI total score. Outcome variables for the three respective hypotheses were as follows: 1) VS DA release defined as the % change in (saline-AMPH PET)/saline PET BP_{ND}; 2) AMPH scan composite pleasant drug effect ratings (saline scan ratings were used as a covariate for baseline adjustment), and 3) PSS scores. Regression models were also used to evaluate whether PSS scores predicted DA release or subjective drug responses. Measures of VS DA release or pleasant drug effects were treated as outcome variables; the primary independent variable was PSS scores. Although brain DA neurotransmission has previously been shown to vary as a function of age (Cervenka et al. 2008; Del Arco et al. 2001; Inoue et al. 2001; Tupala et al. 2003), we did not covary for age in the regression analyses since this variable was already controlled by the study design and findings of preliminary correlational analyses showed no relationships between age and any of the outcome variables. However, because of the lack of control for gender in the design and prior evidence of sex differences in DA and subjective responses to AMPH (Abi-Dargham et al. 2003; Evans. 2007; Munro et al. 2006; White et al. 2002), perceived stress (Allen et al. 2011; Andreou et al. 2011), and responses to ACEs (Heffner et al. 2011; Kendler et al. 2002; Samplin et al. 2013), we did evaluate the relevance of gender by controlling for it and testing interactions with each of the primary independent variables of interest. When interactions were significant, we conducted *post hoc* analyses to examine the models separately in each gender. For parsimony, the interaction term was dropped if not significant. Saline scan [¹¹C]raclopride BP_{ND} was used as the outcome variable in parallel analyses that examined associations between VS D2 receptor

availability and ETI or PSS scores. Although our primary region of interest was the VS, we conducted exploratory regression analyses of four additional striatal regions to evaluate the specificity of associations between VS DA release or BP_{ND} and ETI or PSS scores. The additional regions included anterior and posterior subdivisions of the caudate nucleus and the putamen as described in the section on PET Data Analysis. Mean baseline cortisol levels were obtained by averaging the values of specimens obtained prior to the placebo and amphetamine scans. Relationships between mean baseline cortisol levels and DA release, ETI scores, and PSS scores were evaluated using gender-adjusted regression analyses.

Based on associations observed in the regression analyses, we also tested whether perceived stress may mediate the relationship between childhood trauma and DA release. Mediation analysis consisted of four steps, as discussed in the classical approach (Baron and Kenny, 1986). First, we assessed associations between (1) ETI and PSS scores (a path); (2) PSS scores and VS DA release (b path); (3) ETI scores and VS DA release (c path); and (4) ETI scores and VS DA release (c' path) controlling for PSS score. The mediation effect of perceived stress was examined with the following two regressions using the path model: (a) PSS on ETI scores and (b) VS DA release on PSS scores after controlling for the ETI scores. The mediation effect (indirect effect) was estimated by the product of estimates of parameters (a*b). Since Sobel test is underpowered in this data set, bias-corrected Bootstrapping method, which is especially effective for relatively small sample sizes and relaxes the normality assumption of the outcome variables (MacKinnon, 2008; MacKinnon et al. 2002; MacKinnon et al. 2004; Shrout and Bolger, 2002), was used to estimate the significance of the indirect effect. The significance of the mediation effect was tested using the 95% confidence interval. Since findings of secondary analyses showed that ETI scores were not significantly associated with DA release in other striatal regions, no mediation analyses were performed for other regions.

3. Results

3.1 Sample Characteristics

Participants were between the ages of 18 and 29 years, predominantly male (67.9%), and light to moderate drinkers, with the exception of two who reported no alcohol use. Twenty-six participants were non-smokers; two were non-dependent occasional smokers, one reported < 1 and the other < 5 cigarettes/weekly. Demographic characteristics and mean scores on self-report measures are displayed in Table 1. Mean scores on the PSS were slightly lower than norms previously reported for this age group (Cohen et al. 1983). Mean ETI total and subscale scores were generally consistent with those reported by Bremner and colleagues in healthy adults without a history of psychiatric disorder (Bremner et al. 2007). Dopamine release, PSS, and ETI total scores were normally distributed in the sample. Although pleasant drug effect ratings were not, skewness was within an acceptable range and the boxplot showed no outliers. No associations were found between scores on outcome measures and demographic characteristics of age, BMI, drinking history, or gender. None of the participants were clinically depressed and no relationships were observed between BDI scores and DA release, BP_{ND}, or pleasant drug effects. However, BDI scores were positively

associated with both perceived stress ($r = .540$; $p = .003$) and trauma ratings ($r = .578$; $p = .001$).

3.2 Associations with Childhood Trauma

3.2.1 DA release—A significant decrease in VS [^{11}C]raclopride BP_{ND} ($p < 0.001$) was noted from the post saline ($M=3.06$, $SD=0.31$) to the post AMPH ($M=2.64$, $SD=0.30$) scan, indicating significant AMPH-induced DA release. A positive relationship was found between ETI scores and VS DA release in the sample as a whole (Table 2), suggesting that individuals who experience a greater number of adverse events in childhood have greater VS DA responses to AMPH. No significant interaction was observed between ETI scores and gender. The unadjusted relationship is shown in Figure 1. Images of BP_{ND} (saline minus AMPH scans) are displayed in Figure 2 to further illustrate the relationship using subjects in the highest and lowest quartiles of ETI scores.

No significant relationships were observed between ETI scores and DA release in the other striatal regions (ACN, PCN, APU, and PPU) examined in exploratory analyses (Table 3).

3.2.2 D2 receptor availability—Findings indicated that associations between ETI scores and VS D2 receptor availability (BP_{ND}) differed in males and females. Gender-stratified *post hoc* analyses showed a positive relationship between ETI scores and VS BP_{ND} among males, but a marginally negative relationship among females (Table 2).

Interactions between ETI scores and gender were also significant in exploratory analyses of APU and ACN BP_{ND} , showing that males with higher ETI scores had higher VS D2 receptor availability (BP_{ND}). Although not significant, the direction of the relationships was negative in females, consistent with what was observed between ETI scores and BP_{ND} in the VS (Table 3).

3.2.3 Pleasant drug effects—Ratings on all five scales of pleasant drug effects were significantly higher during the AMPH than during the saline scan ($p < 0.001$ in all cases). Mean scores on four of these measures are shown in Figure 3. A significant interaction was observed between ETI scores and gender in the full model (Table 2). Although findings of subsequent gender-stratified *post hoc* analyses were not significant in either gender, the direction of the relationship between childhood trauma and pleasant drug effects was noted to be positive among males and negative among females.

3.2.4 Perceived stress—ETI scores were positively related to scores on the PSS in the sample as a whole (Table 2), suggesting that individuals who experience a greater number of ACEs also experience higher levels of perceived stress as adults. No significant interaction was found between ETI scores and gender. The unadjusted relationship between ETI and PSS is shown in Figure 4.

3.3 Associations with Perceived Stress

3.3.1 DA release—A positive association was observed between PSS scores and VS DA release in the sample as a whole; higher perceived stress was associated with greater VS DA

responses to AMPH (Table 4). The relationship between perceived stress and VS DA release did not differ in males and females.

A positive relationship was observed between PSS scores and AMPH-induced DA release in the ACN in exploratory analyses (Table 5). Effects were not modified by gender. Scores on the PSS were not associated with DA release in any of the other striatal subdivisions.

3.3.2 D2 receptor availability—No relationships were observed between PSS scores and VS D2 receptor availability (BP_{ND}) in the primary analyses (Table 4).

A trend was observed in exploratory analyses, which suggested that levels of perceived stress are positively associated with ACN BP_{ND} in males and negatively associated with ACN BP_{ND} in females (Table 5).

3.3.3 Pleasant drug effects—Examination of the relationship between PSS scores and pleasant subjective effects of AMPH revealed a significant interaction between PSS scores and gender (Table 4). Findings of gender-stratified analyses showed that PSS scores were positively related to pleasant drug effects among males; whereas, no association was observed among females.

3.4 Associations with Baseline Cortisol

No associations were observed between baseline cortisol levels and DA release, total ETI scores, or PSS scores.

3.5 Mediation Analysis

The first regression indicated that ETI scores were associated with PSS scores (path a in Figure 5), which is consistent with the previous findings when controlling for gender. In the second regression, ETI scores were positively associated with VS DA release (path c). The third regression revealed that the effect of ETI scores on DA release was not significant after controlling for PSS scores, suggesting that the relationship between ETI and DA was mediated by PSS (path c'). The alternative bias-corrected bootstrap method confirmed the mediating role of PSS on the relationship between ETI and DA. The production of $a*b$ ($0.67*0.43$) is 0.285. The 95% bias-corrected bootstrapped confidence interval of this mediation effect was within the range of 0.008–0.856, indicating the presence of mediation.

4. Discussion

A major gap that still remains in our knowledge about relationships between childhood adversity and the etiology of substance use disorders is lack of understanding about biological mechanisms that may underlie the relationship. To our knowledge, this study represents the first examination of relationships between childhood adversity and mesolimbic DA responses to AMPH in humans. Results indicated that the number of ACEs that participants experienced was positively associated with their DA responses to AMPH, as well as with levels of perceived stress that they experienced as adults. Current levels of perceived stress were also positively associated with DA responses to AMPH in the sample as a whole and partially mediated the relationship between childhood trauma and AMPH-

induced DA release, suggesting that ACEs may have both direct and indirect effects on DA neurotransmission. Although associations between childhood trauma and DA release did not differ in males and females, relationships between childhood trauma and both D2 receptor availability and pleasant drug effects were significantly modified by gender. Childhood trauma was positively associated with D2 receptor availability and in males; whereas, a trend for a negative relationship was observed in females. Although not significant, similar gender differences in directionality were observed in relationships between ACEs and pleasant drug effects (i.e., positive in males and negative in females). Perceived stress was also positively associated with pleasant drug effects in males; whereas, no relationship was observed in females. No associations were found between perceived stress and D2 receptor availability. Overall, our results are consistent with notions that early life experiences may have an enduring influence on neural systems that mediate responses to stress and drugs of abuse in humans.

Findings from observational studies have provided compelling evidence that a history of childhood adversity may increase risks for substance abuse in both adolescents and adults (Carswell et al. 2008; Fishbein et al. 2011; Hamburger et al. 2008; Kerr et al. 2009; Rothman et al. 2008; Sansone et al. 2009; Shin et al. 2013; Simpson and Miller. 2002; Strine et al. 2012). It is logical to speculate that one biological mechanism that may underlie this relationship is mesolimbic DA neurotransmission. First, a voluminous body of preclinical (Bonci et al. 2003; Doyon et al. 2003; Koob. 2003; Wise. 1998) and human neuroimaging (Drevets et al. 2001; Leyton et al. 2002; Martinez et al. 2003; Oswald et al. 2005; Urban et al. 2010; Volkow et al. 1999; Volkow et al. 2002; Yoder et al. 2007) literature has established the importance of mesocorticolimbic DA pathways in reinforcement, reward, and addiction. Second, there is preclinical evidence that early life stress may lead to profound and long-lasting alterations in DA pathways in animals. During critical developmental periods when plasticity is heightened, the brain may be particularly sensitive to the destructive effects of adverse experiences on neural function (Rodrigues et al. 2011). In rodents, early life stress is associated with increased tissue levels and reduced rate of DA clearance in the striatum (Matthews et al. 2001; Womersley et al. 2011), as well as altered levels of D2 receptor expression, which vary as a function of D2 polymorphisms (Lovic et al. 2013). Several investigators have reported enhanced striatal DA responses to later stress in rodents exposed to early life adversity (Brake et al. 2004; Fulford and Marsden. 1998; Meaney et al. 2002; Miura et al. 2002), although blunting of DA responses has also been found ((Brake et al. 2004; Fulford and Marsden. 1998; Jahng et al. 2010; Meaney et al. 2002; Miura et al. 2002). Enhanced DA responses to psychostimulants (Hall et al. 1999; Kosten et al. 2003) and increased propensity for drug self-administration have also been observed (Brake et al. 2004; Kosten et al. 2005; Kosten et al. 2006; Meaney et al. 2002; Moffett et al. 2007; Ploj et al. 2003; Zhang et al. 2005).

Exposure to glucocorticoids during early life also has enduring effects on the organization and epigenetic control of midbrain DA systems (McArthur et al. 2007; Niwa et al. 2013), suggesting that the neurobiological mechanisms by which early life stress alters DA function may involve the hypothalamic-pituitary-adrenal (HPA) axis. Elevated levels of glucocorticoids induced by environmental stress during adolescence were recently shown to be associated with augmented NAcc DA levels following methamphetamine challenge in a

transgenic mouse model with a relevant genetic risk factor (Niwa et al. 2013). The enhancement in DA levels in that study was shown to be a consequence of cortisol-induced epigenetic methylation of the tyrosine hydroxylase gene whose gene product is an important enzyme in the biosynthesis of DA. Although no associations were observed between DA release and baseline cortisol levels in our sample, the lack of associations does not preclude the possibility that prolonged elevations in glucocorticoid levels *during the period of trauma exposure* may lead to alterations in DA function that persist into adulthood. This assumption fits a body of literature showing crucial time windows in development where neuroplasticity can be altered (Rosa et al. 2013).

The first evidence that some of these findings may translate to humans was provided by Pruessner and colleagues (2004) in an [^{11}C]raclopride PET study. Acute psychosocial stress was associated with a significant increase in stress-induced DA release in subjects who reported low maternal care, but not in those who reported high maternal care. Low maternal care humans also had higher anxiety and lower self-esteem scores. In the current study, we extended this line of research by showing that ACEs are also associated with enhanced VS DA sensitivity to AMPH in humans. Our findings parallel those of preclinical studies showing that rodents exposed to early life stress have augmented VS DA responses to psychostimulants (Hall et al. 1999; Kosten et al. 2003). Findings of the exploratory analyses suggested that relationships between ETI scores and D2 receptor availability in the anterior putamen (APU) and anterior caudate (ACN) may be analogous to those observed with D2 receptor availability in the VS. Although it is well-established that the NAcc region of the VS plays a key role in appetitive behaviors and in the reinforcing effects of drugs of abuse, there has been growing evidence in recent years that the dorsal striatum is fundamental in the transition from voluntary to habitual or compulsive drug-seeking (Everitt and Robbins. 2013). Thus, concurrent subversion of DA function in both ventral and dorsal regions of the striatum by early life stress could be expected to have a significant impact on the broad-based integration of learning and motivational behaviors across the various phases of addiction.

Somewhat unexpectedly, relationships between ACEs and both baseline D2 receptor availability and pleasant drug effects were found to be significantly modified by gender. The underlying processes and clinical relevance of these sexual dimorphisms are not currently clear, but the findings are consistent with prior evidence that males may have greater cortisol reactivity to stress (Strine et al. 2012; Uhart et al. 2006) and greater DA and subjective responses to AMPH than females (Gabbay. 2005; Munro et al. 2006). Gender-specific differences in the function of brain monoamine systems early trauma have previously been reported in rodents exposed to early life stress (Chocyk et al. 2011; Festa et al. 2004; Matthews et al. 2001; Nazarian et al. 2009; Walker et al. 2006). Interestingly, maternal separation has been shown to be associated with ethanol consumption and preference in male, but not female rats (Gustafsson et al. 2005; Jaworski et al. 2005; Moffett et al. 2007; Roman et al. 2004). It is possible that our gender-related findings may have been different if the women were in the luteal phase of the menstrual cycle. Differences in D2 receptor availability have been reported in female cynomolgus monkeys as a function of menstrual cycle; levels were significantly higher in the luteal as compared to the follicular phase in

both the caudate nucleus and putamen (Czoty et al. 2009). Nevertheless, the current findings related to gender should be interpreted cautiously since the sample included only 9 females.

A positive correlation was found between childhood adversity and perceived stress in this study. Because each of these variables was measured at the same time, the findings might reflect a recall bias, i.e., people who feel less stressed may not report as many negative events as those who feel more stressed. However, the observed relationship between childhood adversity and perceived stress was analogous to findings of prior research showing associations between ACEs and psychological distress and/or reactivity to acute laboratory stressors in both healthy adults (Cagampang et al. 2011; Edwards et al. 2003; Glaser et al. 2006; Kendler et al. 2004; McFarlane et al. 2005; Pierrehumbert et al. 2009; Sesar et al. 2010; Tyrka et al. 2008) and substance abusers (Back et al. 2008; Medrano et al. 2002; Wang et al. 2010). A vast body of findings from preclinical, human laboratory, and clinical observational studies has shown that stress may contribute to the pathogenesis and progression of substance use disorders (Back et al. 2010; Breese et al. 2011; Erb et al. 1996; Karlsgodt et al. 2003; Kreek et al. 2005; Laudet et al. 2004; Lu et al. 2003; Sinha. 2001; Soderpalm and de Wit. 2002; Tidey and Miczek. 1997). Recent preclinical findings indicate that both acute and chronic stress impact DA neurotransmission and that the DA system may be an important vulnerability substrate in the relationship between stress and drug abuse (Cuadra et al. 2001; Kosten and Ambrosio. 2002; Piazza and Le Moal. 1996; Rouge-Pont et al. 1993; Tidey and Miczek. 1996; Tsukada et al. 2011; Yavich and Tiihonen. 2000). Although results of the few human studies that have examined effects of acute psychological stress on brain DA levels have been mixed (Lataster et al. 2011; Montgomery et al. 2006; Pruessner et al. 2004), our group and others have previously shown that cortisol responses to either psychosocial stress (TSST; Trier Social Stress Test)(Kirschbaum et al. 1993) or AMPH are positively associated with AMPH-induced DA release in humans (Pruessner et al. 2004; Oswald et al. 2005; Wand et al. 2007). Our current findings further revealed that levels of perceived stress (which may reflect stress reactivity) were also positively associated with DA release. Findings of the preliminary correlational analyses, which gave no evidence of associations between BDI scores and DA release or BP_{ND}, also support inferences that observed effects on DA neurotransmission were linked to stress as opposed to depression.

Results indicated that the relationship between childhood trauma and DA sensitivity to AMPH was partially mediated by perceived stress. Thus, exposure to childhood trauma may not directly influence DA function in some individuals unless it leads to elevated levels of psychological distress in adulthood. Choy et al. reported that a combination of maternal deprivation and later stress had added effects on disruption of prepulse inhibition by AMPH in young adult rats (Choy et al. 2009). Psychological distress, avoidant coping, and mood/anxiety disorders have also been shown to mediate the relationship between ACEs and adult alcohol or drug problems (DeWit et al. 1999; Douglas et al. 2010; Fishbein et al. 2011; Min et al. 2007; Simpson and Miller. 2002; Strine et al. 2012; White and Widom. 2008). Results from a study conducted by Douglas et al. (2010) indicated that effects of childhood trauma on risks for substance dependence are cumulative, based on the number of events experienced.

Our design has both strengths and limitations. None of the subjects met diagnostic criteria for a DSM-IV Axis I disorder and the number of ACEs reported were generally low. An important strength of this design is that the absence of comorbid depression or anxiety strengthens our supposition that associations with DA release were related to childhood trauma rather than to depression or anxiety. A limitation of the design is that effect sizes may be underestimated given the restricted range of adverse events that subjects reported. We suggest that our finding in this sample of healthy adults represents an under-estimation of the true relationship, which may have emerged even stronger in a more clinically diverse group. However, this hypothesis remains to be examined. Prior evidence that DA signaling modulates brain function in an inverted U-shaped manner (Gjedde et al. 2010; Tian et al. 2013) leaves open the possibility that our findings may not generalize to persons with a psychiatric diagnosis. Another limitation of the study is related to the interpretation of findings. Although our findings indicate that childhood adversity is associated with enhanced DA responses to AMPH, the cross-sectional nature of the study and the fact that none of the subjects met diagnostic criteria for alcohol or drug abuse/dependence preclude definitive causal conclusions about the predictive relevance of the findings. The extent to which findings related to AMPH will generalize to alcohol and other drugs of abuse is also undetermined. Nevertheless, the vast body of preclinical literature showing close interactions between stress, mesocorticolimbic DA function, and drug reinforcement supports such extrapolation. One final caveat to consider is that not all children who have been exposed to ACEs go on to develop a substance use disorder. Individual differences in susceptibility to the effects of childhood trauma on DA-dependent processes and related psychopathology could be the result of variations in the duration or developmental stage of stress exposure, interactions of early stress with the genome, epigenetic programming effects of early life stress, or later environmental experiences, which were beyond the scope of this study to examine (Das et al. 2011; Enoch. 2011; Kim-Cohen and Turkewitz. 2012; Heim and Binder. 2012; Tyrka et al. 2012; Lovic et al. 2013).

Acknowledgments

This work was supported by NIH grants R01 DA022433 (LMO), M01 RR016500 (UMSOM GCRC), and R01MH078175 (DFW). Portions of the work were presented at the American College on Neuropsychopharmacology 2011 Annual Meeting. The authors thank Mark Streer for his technical assistance with preparation of the manuscript.

References

- Abi-Dargham A, Kegeles LS, Martinez D, Innis RB, Laruelle M. Dopamine mediation of positive reinforcing effects of amphetamine in stimulant naive healthy volunteers: results from a large cohort. *Eur Neuropsychopharmacol.* 2003; 13:459–468. [PubMed: 14636962]
- Allen MT, Bocek CM, Burch AE. Gender differences and the relationships of perceived background stress and psychological distress with cardiovascular responses to laboratory stressors. *Int J Psychophysiol.* 2011; 81:209–217.10.1016/j.ijpsycho.2011.06.013 [PubMed: 21729723]
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. Washington DC: 2000.
- Andreou E, Alexopoulos EC, Lionis C, Varvogli L, Gnardellis C, Chrousos GP, Darviri C. Perceived Stress Scale: reliability and validity study in Greece. *Int J Environ Res Public Health.* 2011; 8:3287–3298.10.3390/ijerph8083287 [PubMed: 21909307]

- Ashburner, J.; Friston, KJ. Rigid body registration. In: Frackowiak, RSJ.; Friston, KJ.; Frith, C.; Dolan, R.; Price, CJ.; Zeki, S.; Ashburner, J.; Penny, WD., editors. *Human Brain Function*. Academic Press; 2003. p. 635-654.
- Back SE, Hartwell K, DeSantis SM, Saladin M, McRae-Clark AL, Price KL, Moran-Santa Maria MM, Baker NL, Spratt E, Kreek MJ, Brady KT. Reactivity to laboratory stress provocation predicts relapse to cocaine. *Drug Alcohol Depend*. 2010; 106:21–27.10.1016/j.drugalcdep.2009.07.016 [PubMed: 19726138]
- Back SE, Brady KT, Waldrop AE, Yeatts SD, McRae AL, Spratt E. Early life trauma and sensitivity to current life stressors in individuals with and without cocaine dependence. *Am J Drug Alcohol Abuse*. 2008; 34:389–396.10.1080/00952990802122614 [PubMed: 18584568]
- Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol*. 1986; 51:1173–1182. [PubMed: 3806354]
- Baumann B, Danos P, Krell D, Diekmann S, Leschinger A, Stauch R, Wurthmann C, Bernstein HG, Bogerts B. Reduced volume of limbic system-affiliated basal ganglia in mood disorders: preliminary data from a postmortem study. *J Neuropsychiatry Clin Neurosci*. 1999; 11:71–78. [PubMed: 9990559]
- Beck, AT.; Steer, RA.; Brown, GK. BDI-II: Beck depression inventory. Psychological Corp; San Antonio [Tex.]: 1996.
- Belujon P, Grace AA. Hippocampus, amygdala, and stress: interacting systems that affect susceptibility to addiction. *Ann N Y Acad Sci*. 2011; 1216:114–121.10.1111/j.1749-6632.2010.05896.x [PubMed: 21272015]
- Benjet C, Borges G, Medina-Mora ME. Chronic childhood adversity and onset of psychopathology during three life stages: childhood, adolescence and adulthood. *J Psychiatr Res*. 2010; 44:732–740.10.1016/j.jpsychires.2010.01.004 [PubMed: 20144464]
- Bonci A, Bernardi G, Grillner P, Mercuri NB. The dopamine-containing neuron: maestro or simple musician in the orchestra of addiction? *Trends Pharmacol Sci*. 2003; 24:172–177.10.1016/S0165-6147(03)00068-3 [PubMed: 12707003]
- Brake WG, Zhang TY, Diorio J, Meaney MJ, Gratton A. Influence of early postnatal rearing conditions on mesocorticolimbic dopamine and behavioural responses to psychostimulants and stressors in adult rats. *Eur J Neurosci*. 2004; 19:1863–1874.10.1111/j.1460-9568.2004.03286.x [PubMed: 15078560]
- Breese GR, Sinha R, Heilig M. Chronic alcohol neuroadaptation and stress contribute to susceptibility for alcohol craving and relapse. *Pharmacol Ther*. 2011; 129:149–171.10.1016/j.pharmthera.2010.09.007 [PubMed: 20951730]
- Bremner JD, Bolus R, Mayer EA. Psychometric properties of the Early Trauma Inventory-Self Report. *J Nerv Ment Dis*. 2007; 195:211–218.10.1097/01.nmd.0000243824.84651.6c [PubMed: 17468680]
- Burgermeister D. Childhood adversity: a review of measurement instruments. *J Nurs Meas*. 2007; 15:163–176. [PubMed: 18232616]
- Cagampang FR, Poore KR, Hanson MA. Developmental origins of the metabolic syndrome: body clocks and stress responses. *Brain Behav Immun*. 2011; 25:214–220.10.1016/j.bbi.2010.09.005 [PubMed: 20851177]
- Carswell K, Chen Y, Nair RC, Shaw AK, Speechley KN, Barrera M, Maunsell E. Smoking and binge drinking among Canadian survivors of childhood and adolescent cancers: a comparative, population-based study. *Pediatr Blood Cancer*. 2008; 51:280–287.10.1002/pbc.21568 [PubMed: 18386781]
- Cervenka S, Halldin C, Farde L. Age-related diurnal effect on D2 receptor binding: a preliminary PET study. *Int J Neuropsychopharmacol*. 2008; 11:671–678.10.1017/S1461145707008358 [PubMed: 18184446]
- Chocyk A, Przyborowska A, Dudys D, Majcher I, Mackowiak M, Wedzony K. The impact of maternal separation on the number of tyrosine hydroxylase-expressing midbrain neurons during different stages of ontogenesis. *Neuroscience*. 2011; 182:43–61.10.1016/j.neuroscience.2011.03.008 [PubMed: 21396433]

- Choy KH, de Visser YP, van den Buuse M. The effect of 'two hit' neonatal and young-adult stress on dopaminergic modulation of prepulse inhibition and dopamine receptor density. *Br J Pharmacol*. 2009; 156:388–396.10.1111/j.1476-5381.2008.00008.x [PubMed: 19154431]
- Chu DA, Williams LM, Harris AW, Bryant RA, Gatt JM. Early life trauma predicts self-reported levels of depressive and anxiety symptoms in nonclinical community adults: Relative contributions of early life stressor types and adult trauma exposure. *J Psychiatr Res*. 2013; 47:23–32.10.1016/j.jpsychires.2012.08.006 [PubMed: 23020924]
- Cohen, S. Perceived stress scale. Mind Garden; Palo Alto, CA: 1994.
- Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav*. 1983; 24:385–396. [PubMed: 6668417]
- Cole G, Tucker L, Friedman GM. Relationships among measures of alcohol drinking behavior, life-events and perceived stress. *Psychol Rep*. 1990; 67:587–591. [PubMed: 2263712]
- Corbin WR, Farmer NM, Nolen-Hoekesma S. Relations among stress, coping strategies, coping motives, alcohol consumption and related problems: A mediated moderation model. *Addict Behav*. 2013; 38:1912–1919. <http://dx.doi.org/10.1016/j.addbeh.2012.12.005>. [PubMed: 23380486]
- Cuadra G, Zurita A, Gioino G, Molina V. Influence of different antidepressant drugs on the effect of chronic variable stress on restraint-induced dopamine release in frontal cortex. *Neuropsychopharmacology*. 2001; 25:384–394.10.1016/S0893-133X(01)00234-2 [PubMed: 11522466]
- Cuijpers P, Smit F, Unger F, Stikkelbroek Y, Ten Have M, de Graaf R. The disease burden of childhood adversities in adults: a population-based study. *Child Abuse Negl*. 2011; 35:937–945.10.1016/j.chiabu.2011.06.005 [PubMed: 22099144]
- Czoty PW, Riddick NV, Gage HD, Sandridge M, Nader SH, Garg S, Bounds M, Garg PK, Nader MA. Effect of menstrual cycle phase on dopamine D2 receptor availability in female cynomolgus monkeys. *Neuropsychopharmacology*. 2009; 34:548–554.10.1038/npp.2008.3 [PubMed: 18256593]
- D'Andrea W, Ford J, Stolbach B, Spinazzola J, van der Kolk BA. Understanding interpersonal trauma in children: why we need a developmentally appropriate trauma diagnosis. *Am J Orthopsychiatry*. 2012; 82:187–200.10.1111/j.1939-0025.2012.01154.x [PubMed: 22506521]
- Das D, Cherbuin N, Tan X, Anstey KJ, Eastale S. DRD4-exonIII-VNTR moderates the effect of childhood adversities on emotional resilience in young-adults. *PLoS One*. 2011; 6:e20177.10.1371/journal.pone.0020177 [PubMed: 21637770]
- Del Arco A, Segovia G, Mora F. Dopamine release during stress in the prefrontal cortex of the rat decreases with age. *Neuroreport*. 2001; 12:4019–4022. [PubMed: 11742231]
- DeWit DJ, MacDonald K, Offord DR. Childhood stress and symptoms of drug dependence in adolescence and early adulthood: social phobia as a mediator. *Am J Orthopsychiatry*. 1999; 69:61–72. [PubMed: 9990437]
- Di Chiara G, Bassareo V, Fenu S, De Luca MA, Spina L, Cadoni C, Acquas E, Carboni E, Valentini V, Lecca D. Dopamine and drug addiction: the nucleus accumbens shell connection. *Neuropharmacology*. 2004; 47(Suppl 1):227–241.10.1016/j.neuropharm.2004.06.032 [PubMed: 15464140]
- Dillon DG, Holmes AJ, Birk JL, Brooks N, Lyons-Ruth K, Pizzagalli DA. Childhood adversity is associated with left basal ganglia dysfunction during reward anticipation in adulthood. *Biol Psychiatry*. 2009; 66:206–213.10.1016/j.biopsych.2009.02.019 [PubMed: 19358974]
- Douglas KR, Chan G, Gelernter J, Arias AJ, Anton RF, Weiss RD, Brady K, Poling J, Farrer L, Kranzler HR. Adverse childhood events as risk factors for substance dependence: partial mediation by mood and anxiety disorders. *Addict Behav*. 2010; 35:7–13.10.1016/j.addbeh.2009.07.004 [PubMed: 19720467]
- Doyon WM, York JL, Diaz LM, Samson HH, Czachowski CL, Gonzales RA. Dopamine activity in the nucleus accumbens during consummatory phases of oral ethanol self-administration. *Alcohol Clin Exp Res*. 2003; 27:1573–1582.10.1097/01.ALC.0000089959.66222.B8 [PubMed: 14574227]
- Drevets WC, Gautier C, Price JC, Kupfer DJ, Kinahan PE, Grace AA, Price JL, Mathis CA. Amphetamine-induced dopamine release in human ventral striatum correlates with euphoria. *Biol Psychiatry*. 2001; 49:81–96. [PubMed: 11164755]

- Edwards VJ, Holden GW, Felitti VJ, Anda RF. Relationship between multiple forms of childhood maltreatment and adult mental health in community respondents: results from the adverse childhood experiences study. *Am J Psychiatry*. 2003; 160:1453–1460. [PubMed: 12900308]
- Ehrin E, Farde L, de Paulis T, Eriksson L, Greitz T, Johnstrom P, Litton JE, Nilsson JL, Sedvall G, Stone-Elander S. Preparation of ¹¹C-labelled Raclopride, a new potent dopamine receptor antagonist: preliminary PET studies of cerebral dopamine receptors in the monkey. *Int J Appl Radiat Isot*. 1985; 36:269–273. [PubMed: 3874833]
- Enoch MA. The role of early life stress as a predictor for alcohol and drug dependence. *Psychopharmacology (Berl)*. 2011; 214:17–31.10.1007/s00213-010-1916-6 [PubMed: 20596857]
- Erb S, Shaham Y, Stewart J. Stress reinstates cocaine-seeking behavior after prolonged extinction and a drug-free period. *Psychopharmacology (Berl)*. 1996; 128:408–412. [PubMed: 8986011]
- Evans SM. The role of estradiol and progesterone in modulating the subjective effects of stimulants in humans. *Exp Clin Psychopharmacol*. 2007; 15:418–426.10.1037/1064-1297.15.5.418 [PubMed: 17924776]
- Everitt BJ, Robbins TW. From the ventral to the dorsal striatum: Devolving views of their roles in drug addiction. *Neurosci Biobehav Rev*. 201310.1016/j.neubiorev.2013.02.010
- Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V, Koss MP, Marks JS. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *Am J Prev Med*. 1998; 14:245–258. [PubMed: 9635069]
- Festa ED, Russo SJ, Gazi FM, Niyomchai T, Kemen LM, Lin SN, Foltz R, Jenab S, Quinones-Jenab V. Sex differences in cocaine-induced behavioral responses, pharmacokinetics, and monoamine levels. *Neuropharmacology*. 2004; 46:672–687.10.1016/j.neuropharm.2003.11.017 [PubMed: 14996545]
- Finkelhor D, Ormrod RK, Turner HA. Poly-victimization: a neglected component in child victimization. *Child Abuse Negl*. 2007; 31:7–26.10.1016/j.chiabu.2006.06.008 [PubMed: 17224181]
- First, MB.; Spitzer, RL.; Gibbon, M.; Williams, JBW. Biometrics Research. New York State Psychiatric Institute; New York: 2002. Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Non-patient Edition.
- Fishbein D, Novak SP, Krebs C, Warner T, Hammond J. The mediating effect of depressive symptoms on the relationship between traumatic childhood experiences and drug use initiation. *Addict Behav*. 2011; 36:527–531.10.1016/j.addbeh.2010.12.030 [PubMed: 21296505]
- Ford JD, Elhai JD, Connor DF, Frueh BC. Poly-victimization and risk of posttraumatic, depressive, and substance use disorders and involvement in delinquency in a national sample of adolescents. *J Adolesc Health*. 2010; 46:545–552.10.1016/j.jadohealth.2009.11.212 [PubMed: 20472211]
- Fulford AJ, Marsden CA. Effect of isolation-rearing on conditioned dopamine release in vivo in the nucleus accumbens of the rat. *J Neurochem*. 1998; 70:384–390. [PubMed: 9422385]
- Gabbay FH. Family history of alcoholism and response to amphetamine: sex differences in the effect of risk. *Alcohol Clin Exp Res*. 2005; 29:773–780. [PubMed: 15897722]
- Gjedde A, Kumakura Y, Cumming P, Linnet J, Moller A. Inverted-U-shaped correlation between dopamine receptor availability in striatum and sensation seeking. *Proc Natl Acad Sci U S A*. 2010; 107:3870–3875.10.1073/pnas.0912319107 [PubMed: 20133675]
- Glaser JP, van Os J, Portegijs PJ, Myin-Germeys I. Childhood trauma and emotional reactivity to daily life stress in adult frequent attenders of general practitioners. *J Psychosom Res*. 2006; 61:229–236.10.1016/j.jpsychores.2006.04.014 [PubMed: 16880026]
- Gordon HW. Early environmental stress and biological vulnerability to drug abuse. *Psychoneuroendocrinology*. 2002; 27:115–126. [PubMed: 11750773]
- Gustafsson L, Ploj K, Nylander I. Effects of maternal separation on voluntary ethanol intake and brain peptide systems in female Wistar rats. *Pharmacol Biochem Behav*. 2005; 81:506–516.10.1016/j.pbb.2005.03.016 [PubMed: 15927243]
- Hall FS, Wilkinson LS, Humby T, Robbins TW. Maternal deprivation of neonatal rats produces enduring changes in dopamine function. *Synapse*. 1999; 32:37–43. 2–4. [PubMed: 10188636]

- Hamburger ME, Leeb RT, Swahn MH. Childhood maltreatment and early alcohol use among high-risk adolescents. *J Stud Alcohol Drugs*. 2008; 69:291–295. [PubMed: 18299771]
- Heatherton TF, Kozlowski LT, Frecker RC, Fagerstrom KO. The Fagerstrom Test for Nicotine Dependence: a revision of the Fagerstrom Tolerance Questionnaire. *Br J Addict*. 1991; 86:1119–1127. [PubMed: 1932883]
- Heffner JL, Blom TJ, Anthenelli RM. Gender differences in trauma history and symptoms as predictors of relapse to alcohol and drug use. *Am J Addict*. 2011; 20:307–311.10.1111/j.1521-0391.2011.00141.x [PubMed: 21679261]
- Heim C, Binder EB. Current research trends in early life stress and depression: review of human studies on sensitive periods, gene-environment interactions, and epigenetics. *Exp Neurol*. 2012; 233:102–111.10.1016/j.expneurol.2011.10.032 [PubMed: 22101006]
- Huang S, Trapido E, Fleming L, Arheart K, Crandall L, French M, Malcolm S, Prado G. The long-term effects of childhood maltreatment experiences on subsequent illicit drug use and drug-related problems in young adulthood. *Addict Behav*. 2011; 36:95–102.10.1016/j.addbeh.2010.09.001 [PubMed: 20947260]
- Husted JA, Ahmed R, Chow EW, Brzustowicz LM, Bassett AS. Early environmental exposures influence schizophrenia expression even in the presence of strong genetic predisposition. *Schizophr Res*. 2012; 137:166–168.10.1016/j.schres.2012.02.009 [PubMed: 22381191]
- Hyman SM, Paliwal P, Sinha R. Childhood maltreatment, perceived stress, and stress-related coping in recently abstinent cocaine dependent adults. *Psychol Addict Behav*. 2007; 21:233–238.10.1037/0893-164X.21.2.233 [PubMed: 17563143]
- Innis RB, Cunningham VJ, Delforge J, Fujita M, Gjedde A, Gunn RN, Holden J, Houle S, Huang SC, Ichise M, Iida H, Ito H, Kimura Y, Koeppe RA, Knudsen GM, Knuuti J, Lammertsma AA, Laruelle M, Logan J, Maguire RP, Mintun MA, Morris ED, Parsey R, Price JC, Slifstein M, Sossi V, Suhara T, Votaw JR, Wong DF, Carson RE. Consensus nomenclature for in vivo imaging of reversibly binding radioligands. *J Cereb Blood Flow Metab*. 2007; 27:1533–1539.10.1038/sj.jcbfm.9600493 [PubMed: 17519979]
- Innis RB, Malison RT, al-Tikriti M, Hoffer PB, Sybirska EH, Seibyl JP, Zoghbi SS, Baldwin RM, Laruelle M, Smith EO. Amphetamine-stimulated dopamine release competes in vivo for [123I]IBZM binding to the D2 receptor in nonhuman primates. *Synapse*. 1992; 10:177–184.10.1002/syn.890100302 [PubMed: 1532675]
- Inoue M, Suhara T, Sudo Y, Okubo Y, Yasuno F, Kishimoto T, Yoshikawa K, Tanada S. Age-related reduction of extrastriatal dopamine D2 receptor measured by PET. *Life Sci*. 2001; 69:1079–1084. [PubMed: 11508650]
- Jahng JW, Ryu V, Yoo SB, Noh SJ, Kim JY, Lee JH. Mesolimbic dopaminergic activity responding to acute stress is blunted in adolescent rats that experienced neonatal maternal separation. *Neuroscience*. 2010; 171:144–152.10.1016/j.neuroscience.2010.08.063 [PubMed: 20828601]
- Jaworski JN, Francis DD, Brommer CL, Morgan ET, Kuhar MJ. Effects of early maternal separation on ethanol intake, GABA receptors and metabolizing enzymes in adult rats. *Psychopharmacology (Berl)*. 2005; 181:8–15.10.1007/s00213-005-2232-4 [PubMed: 15830234]
- Kaplan MS. Acute alcohol consumption as a contributing factor to suicidal behavior. *Am J Public Health*. 2013; 103:e2–3.10.2105/AJPH.2013.301422 [PubMed: 23865652]
- Karlsgodt KH, Lukas SE, Elman I. Psychosocial stress and the duration of cocaine use in non-treatment seeking individuals with cocaine dependence. *Am J Drug Alcohol Abuse*. 2003; 29:539–551. [PubMed: 14510039]
- Kendler KS, Kuhn JW, Prescott CA. Childhood sexual abuse, stressful life events and risk for major depression in women. *Psychol Med*. 2004; 34:1475–1482. [PubMed: 15724878]
- Kendler KS, Sheth K, Gardner CO, Prescott CA. Childhood parental loss and risk for first-onset of major depression and alcohol dependence: the time-decay of risk and sex differences. *Psychol Med*. 2002; 32:1187–1194. [PubMed: 12420888]
- Kendler KS, Bulik CM, Silberg J, Hettema JM, Myers J, Prescott CA. Childhood sexual abuse and adult psychiatric and substance use disorders in women: an epidemiological and cotwin control analysis. *Arch Gen Psychiatry*. 2000; 57:953–959. [PubMed: 11015813]

- Kerr T, Stoltz JA, Marshall BD, Lai C, Strathdee SA, Wood E. Childhood trauma and injection drug use among high-risk youth. *J Adolesc Health*. 2009; 45:300–302.10.1016/j.jadohealth.2009.03.007 [PubMed: 19699428]
- Kessler RC, McLaughlin KA, Green JG, Gruber MJ, Sampson NA, Zaslavsky AM, Aguilar-Gaxiola S, Alhamzawi AO, Alonso J, Angermeyer M, Benjet C, Bromet E, Chatterji S, de Girolamo G, Demyttenaere K, Fayyad J, Florescu S, Gal G, Gureje O, Haro JM, Hu CY, Karam EG, Kawakami N, Lee S, Lepine JP, Ormel J, Posada-Villa J, Sagar R, Tsang A, Ustun TB, Vassilev S, Viana MC, Williams DR. Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. *Br J Psychiatry*. 2010; 197:378–385.10.1192/bjp.bp.110.080499 [PubMed: 21037215]
- Kim-Cohen J, Turkewitz R. Resilience and measured gene-environment interactions. *Dev Psychopathol*. 2012; 24:1297–1306.10.1017/S0954579412000715 [PubMed: 23062298]
- Kirschbaum C, Pirke KM, Hellhammer DH. The ‘Trier Social Stress Test’--a tool for investigating psychobiological stress responses in a laboratory setting. *Neuropsychobiology*. 1993; 28:76–81. [PubMed: 8255414]
- Koob GF. Alcoholism: allostasis and beyond. *Alcohol Clin Exp Res*. 2003; 27:232–243.10.1097/01.ALC.0000057122.36127.C2 [PubMed: 12605072]
- Kosten TA, Zhang XY, Kehoe P. Heightened cocaine and food self-administration in female rats with neonatal isolation experience. *Neuropsychopharmacology*. 2006; 31:70–76.10.1038/sj.npp.1300779 [PubMed: 15956993]
- Kosten TA, Zhang XY, Kehoe P. Neurochemical and behavioral responses to cocaine in adult male rats with neonatal isolation experience. *J Pharmacol Exp Ther*. 2005; 314:661–667.10.1124/jpet.104.081216 [PubMed: 15845857]
- Kosten TA, Zhang XY, Kehoe P. Chronic neonatal isolation stress enhances cocaine-induced increases in ventral striatal dopamine levels in rat pups. *Brain Res Dev Brain Res*. 2003; 141:109–116.
- Kosten TA, Ambrosio E. HPA axis function and drug addictive behaviors: insights from studies with Lewis and Fischer 344 inbred rats. *Psychoneuroendocrinology*. 2002; 27:35–69. [PubMed: 11750769]
- Kreek MJ, Nielsen DA, Butelman ER, LaForge KS. Genetic influences on impulsivity, risk taking, stress responsivity and vulnerability to drug abuse and addiction. *Nat Neurosci*. 2005; 8:1450–1457.10.1038/nn1583 [PubMed: 16251987]
- Kumar A, Kuwabara H, Rousset O, Alexander M, Brasic J, Ye W, Wong D. Effects of head-motion correction and modeling approaches on SPM analyses. *J Nucl Med*. 2007; 48(41)
- Lataster J, Collip D, Ceccarini J, Haas D, Booij L, van Os J, Pruessner J, Van Laere K, Myin-Germeys I. Psychosocial stress is associated with in vivo dopamine release in human ventromedial prefrontal cortex: a positron emission tomography study using [(1)(8)F]fallypride. *Neuroimage*. 2011; 58:1081–1089.10.1016/j.neuroimage.2011.07.030 [PubMed: 21801840]
- Laudet AB, Magura S, Vogel HS, Knight EL. Perceived reasons for substance misuse among persons with a psychiatric disorder. *Am J Orthopsychiatry*. 2004; 74:365–375.10.1037/0002-9432.74.3.365 [PubMed: 15291712]
- Leyton M, Boileau I, Benkelfat C, Diksic M, Baker G, Dagher A. Amphetamine-induced increases in extracellular dopamine, drug wanting, and novelty seeking: a PET/[11C]raclopride study in healthy men. *Neuropsychopharmacology*. 2002; 27:1027–1035.10.1016/S0893-133X(02)00366-4 [PubMed: 12464459]
- Logan J, Fowler JS, Volkow ND, Wang GJ, Ding YS, Alexoff DL. Distribution volume ratios without blood sampling from graphical analysis of PET data. *J Cereb Blood Flow Metab*. 1996; 16:834–840.10.1097/00004647-199609000-00008 [PubMed: 8784228]
- Lovic V, Belay H, Walker CD, Burton CL, Meaney MJ, Sokolowski M, Fleming AS. Early postnatal experience and DRD2 genotype affect dopamine receptor expression in the rat ventral striatum. *Behav Brain Res*. 2013; 237:278–282.10.1016/j.bbr.2012.09.046 [PubMed: 23036844]
- Lu L, Shepard JD, Hall FS, Shaham Y. Effect of environmental stressors on opiate and psychostimulant reinforcement, reinstatement and discrimination in rats: a review. *Neurosci Biobehav Rev*. 2003; 27:457–491. [PubMed: 14505687]

- MacKinnon, DP. Introduction to Statistical Mediation Analysis. Routledge Academic; New York: 2008.
- Mackinnon DP, Lockwood CM, Williams J. Confidence Limits for the Indirect Effect: Distribution of the Product and Resampling Methods. *Multivariate Behav Res.* 2004; 39:99.10.1207/s15327906mbr3901_4 [PubMed: 20157642]
- MacKinnon DP, Lockwood CM, Hoffman JM, West SG, Sheets V. A comparison of methods to test mediation and other intervening variable effects. *Psychol Methods.* 2002; 7:83–104. [PubMed: 11928892]
- Mangiavacchi S, Masi F, Scheggi S, Leggio B, De Montis MG, Gambarana C. Long-term behavioral and neurochemical effects of chronic stress exposure in rats. *J Neurochem.* 2001; 79:1113–1121. [PubMed: 11752052]
- Marinelli M, Piazza PV. Interaction between glucocorticoid hormones, stress and psychostimulant drugs. *Eur J Neurosci.* 2002; 16:387–394. [PubMed: 12193179]
- Martinez D, Slifstein M, Broft A, Mawlawi O, Hwang DR, Huang Y, Cooper T, Kegeles L, Zarahn E, Abi-Dargham A, Haber SN, Laruelle M. Imaging human mesolimbic dopamine transmission with positron emission tomography. Part II: amphetamine-induced dopamine release in the functional subdivisions of the striatum. *J Cereb Blood Flow Metab.* 2003; 23:285–300. [PubMed: 12621304]
- Marx BP, Sloan DM. The effects of trauma history, gender, and race on alcohol use and posttraumatic stress symptoms in a college student sample. *Addict Behav.* 2003; 28:1631–1647. [PubMed: 14656550]
- Matheson SL, Shepherd AM, Pinchbeck RM, Laurens KR, Carr VJ. Childhood adversity in schizophrenia: a systematic meta-analysis. *Psychol Med.* 2012;1–13.10.1017/S0033291712000785
- Matthews KA, Gump BB, Owens JF. Chronic stress influences cardiovascular and neuroendocrine responses during acute stress and recovery, especially in men. *Health Psychol.* 2001; 20:403–410. [PubMed: 11714181]
- McArthur S, McHale E, Gillies GE. The size and distribution of midbrain dopaminergic populations are permanently altered by perinatal glucocorticoid exposure in a sex- region- and time-specific manner. *Neuropsychopharmacology.* 2007; 32:1462–1476.10.1038/sj.npp.1301277 [PubMed: 17164817]
- McFarlane A, Clark CR, Bryant RA, Williams LM, Niaura R, Paul RH, Hitsman BL, Stroud L, Alexander DM, Gordon E. The impact of early life stress on psychophysiological, personality and behavioral measures in 740 non-clinical subjects. *J Integr Neurosci.* 2005; 4:27–40. [PubMed: 16035139]
- McGinty VB, Lardeux S, Taha SA, Kim JJ, Nicola SM. Invigoration of reward seeking by cue and proximity encoding in the nucleus accumbens. *Neuron.* 2013; 78:910–922.10.1016/j.neuron.2013.04.010 [PubMed: 23764290]
- Meaney MJ, Brake W, Gratton A. Environmental regulation of the development of mesolimbic dopamine systems: a neurobiological mechanism for vulnerability to drug abuse? *Psychoneuroendocrinology.* 2002; 27:127–138. [PubMed: 11750774]
- Medrano MA, Hatch JP, Zule WA, Desmond DP. Psychological distress in childhood trauma survivors who abuse drugs. *Am J Drug Alcohol Abuse.* 2002; 28:1–13. [PubMed: 11853127]
- Mehta MA. Hyporesponsive reward anticipation in the basal ganglia following severe institutional deprivation early in life. *J Cogn Neurosci.* 2010; 22:2316.10.1162/jocn.2009.21394 [PubMed: 19929329]
- Miczek KA, Nikulina EM, Shimamoto A, Covington HE 3rd . Escalated or suppressed cocaine reward, tegmental BDNF, and accumbal dopamine caused by episodic versus continuous social stress in rats. *J Neurosci.* 2011; 31:9848–9857.10.1523/JNEUROSCI.0637-11.2011 [PubMed: 21734276]
- Min M, Farkas K, Minnes S, Singer LT. Impact of childhood abuse and neglect on substance abuse and psychological distress in adulthood. *J Trauma Stress.* 2007; 20:833–844.10.1002/jts.20250 [PubMed: 17955535]
- Miura H, Qiao H, Ohta T. Attenuating effects of the isolated rearing condition on increased brain serotonin and dopamine turnover elicited by novelty stress. *Brain Res.* 2002; 926:10–17. [PubMed: 11814401]

- Moffett MC, Vicentic A, Kozel M, Plotsky P, Francis DD, Kuhar MJ. Maternal separation alters drug intake patterns in adulthood in rats. *Biochem Pharmacol.* 2007; 73:321–330.10.1016/j.bcp.2006.08.003 [PubMed: 16962564]
- Moffett MC, Harley J, Francis D, Sanghani SP, Davis WI, Kuhar MJ. Maternal separation and handling affects cocaine self-administration in both the treated pups as adults and the dams. *J Pharmacol Exp Ther.* 2006; 317:1210–1218.10.1124/jpet.106.101139 [PubMed: 16517692]
- Montgomery AJ. Correction of head movement on PET studies: comparison of methods. *The Journal of Nuclear Medicine.* 2006; 47:1936.
- Montgomery AJ, Mehta MA, Grasby PM. Is psychological stress in man associated with increased striatal dopamine levels?: A [¹¹C]raclopride PET study. *Synapse.* 2006; 60:124–131.10.1002/syn.20282 [PubMed: 16715491]
- Munro CA, McCaul ME, Wong DF, Oswald LM, Zhou Y, Brasic J, Kuwabara H, Kumar A, Alexander M, Ye W, Wand GS. Sex differences in striatal dopamine release in healthy adults. *Biol Psychiatry.* 2006; 59:966–974.10.1016/j.biopsych.2006.01.008 [PubMed: 16616726]
- Nakajima M, al'Absi M. Predictors of risk for smoking relapse in men and women: a prospective examination. *Psychol Addict Behav.* 2012; 26:633–637.10.1037/a0027280 [PubMed: 22352701]
- Nanni V, Uher R, Danese A. Childhood maltreatment predicts unfavorable course of illness and treatment outcome in depression: a meta-analysis. *Am J Psychiatry.* 2012; 169:141–151. [PubMed: 22420036]
- Nazarian A, Sun WL, Zhou L, Kemen LM, Jenab S, Quinones-Jenab V. Sex differences in basal and cocaine-induced alterations in PKA and CREB proteins in the nucleus accumbens. *Psychopharmacology (Berl).* 2009; 203:641–650.10.1007/s00213-008-1411-5 [PubMed: 19052730]
- Nelson EC, Heath AC, Madden PA, Cooper ML, Dinwiddie SH, Bucholz KK, Glowinski A, McLaughlin T, Dunne MP, Statham DJ, Martin NG. Association between self-reported childhood sexual abuse and adverse psychosocial outcomes: results from a twin study. *Arch Gen Psychiatry.* 2002; 59:139–145. [PubMed: 11825135]
- Niwa M, Jaaro-Peled H, Tankou S, Seshadri S, Hikida T, Matsumoto Y, Cascella NG, Kano S, Ozaki N, Nabeshima T, Sawa A. Adolescent stress-induced epigenetic control of dopaminergic neurons via glucocorticoids. *Science.* 2013; 339:335–339.10.1126/science.1226931 [PubMed: 23329051]
- Oswald LM, Wong DF, McCaul M, Zhou Y, Kuwabara H, Choi L, Brasic J, Wand GS. Relationships among ventral striatal dopamine release, cortisol secretion, and subjective responses to amphetamine. *Neuropsychopharmacology.* 2005; 30:821–832.10.1038/sj.npp.1300667 [PubMed: 15702139]
- Piazza PV, Le Moal ML. Pathophysiological basis of vulnerability to drug abuse: role of an interaction between stress, glucocorticoids, and dopaminergic neurons. *Annu Rev Pharmacol Toxicol.* 1996; 36:359–378.10.1146/annurev.pa.36.040196.002043 [PubMed: 8725394]
- Pierrehumbert B, Torrisi R, Glatz N, Dimitrova N, Heinrichs M, Halfon O. The influence of attachment on perceived stress and cortisol response to acute stress in women sexually abused in childhood or adolescence. *Psychoneuroendocrinology.* 2009; 34:924–938.10.1016/j.psyneuen.2009.01.006 [PubMed: 19217212]
- Pilowsky DJ, Keyes KM, Hasin DS. Adverse childhood events and lifetime alcohol dependence. *Am J Public Health.* 2009; 99:258–263.10.2105/AJPH.2008.139006 [PubMed: 19059847]
- Plaza A, Torres A, Martin-Santos R, Gelabert E, Imaz ML, Navarro P, Bremner JD, Valdes M, Garcia-Esteve L. Validation and test-retest reliability of Early Trauma Inventory in Spanish postpartum women. *J Nerv Ment Dis.* 2011; 199:280–285.10.1097/NMD.0b013e31821245b9 [PubMed: 21451355]
- Ploj K, Roman E, Nylander I. Long-term effects of maternal separation on ethanol intake and brain opioid and dopamine receptors in male Wistar rats. *Neuroscience.* 2003; 121:787–799. [PubMed: 14568037]
- Pruessner JC, Champagne F, Meaney MJ, Dagher A. Dopamine release in response to a psychological stress in humans and its relationship to early life maternal care: a positron emission tomography study using [¹¹C]raclopride. *J Neurosci.* 2004; 24:2825–2831.10.1523/JNEUROSCI.3422-03.2004 [PubMed: 15028776]

- Rahmim A, Cheng JC, Blinder S, Camborde ML, Sossi V. Statistical dynamic image reconstruction in state-of-the-art high-resolution PET. *Phys Med Biol.* 2005; 50:4887–4912.10.1088/0031-9155/50/20/010 [PubMed: 16204879]
- Robison AJ, Vialou V, Mazei-Robison M, Feng J, Kourrich S, Collins M, Wee S, Koob G, Turecki G, Neve R, Thomas M, Nestler EJ. Behavioral and structural responses to chronic cocaine require a feedforward loop involving DeltaFosB and calcium/calmodulin-dependent protein kinase II in the nucleus accumbens shell. *J Neurosci.* 2013; 33:4295–4307.10.1523/JNEUROSCI.5192-12.2013 [PubMed: 23467346]
- Rodrigues AJ, Leao P, Carvalho M, Almeida OF, Sousa N. Potential programming of dopaminergic circuits by early life stress. *Psychopharmacology (Berl).* 2011; 214:107–120.10.1007/s00213-010-2085-3 [PubMed: 21088961]
- Roman E, Ploj K, Nylander I. Maternal separation has no effect on voluntary ethanol intake in female Wistar rats. *Alcohol.* 2004; 33:31–39.10.1016/j.alcohol.2004.04.002 [PubMed: 15353171]
- Rosa AM, Silva MF, Ferreira S, Murta J, Castelo-Branco M. Plasticity in the Human Visual Cortex: An Ophthalmology-Based Perspective. *Biomed Res Int.* 2013; 2013:568354.10.1155/2013/568354 [PubMed: 24205505]
- Rothman EF, Edwards EM, Heeren T, Hingson RW. Adverse childhood experiences predict earlier age of drinking onset: results from a representative US sample of current or former drinkers. *Pediatrics.* 2008; 122:e298–304.10.1542/peds.2007-3412 [PubMed: 18676515]
- Rouge-Pont F, Piazza PV, Kharouby M, Le Moal M, Simon H. Higher and longer stress-induced increase in dopamine concentrations in the nucleus accumbens of animals predisposed to amphetamine self-administration. A microdialysis study. *Brain Res.* 1993; 602:169–174. [PubMed: 8448654]
- Saal D, Dong Y, Bonci A, Malenka RC. Drugs of abuse and stress trigger a common synaptic adaptation in dopamine neurons. *Neuron.* 2003; 37:577–582. [PubMed: 12597856]
- Samplin E, Ikuta T, Malhotra AK, Szeszko PR, Derosse P. Sex differences in resilience to childhood maltreatment: Effects of trauma history on hippocampal volume, general cognition and subclinical psychosis in healthy adults. *J Psychiatr Res.* 2013;10.1016/j.jpsychires.2013.05.008
- Sansone RA, Whitecar P, Wiederman MW. The prevalence of childhood trauma among those seeking buprenorphine treatment. *J Addict Dis.* 2009; 28:64–67.10.1080/10550880802545101 [PubMed: 19197597]
- Sesar K, Simic N, Barisic M. Multi-type childhood abuse, strategies of coping, and psychological adaptations in young adults. *Croat Med J.* 2010; 51:406–416. [PubMed: 20960590]
- Shimamoto A, Debold JF, Holly EN, Miczek KA. Blunted accumbal dopamine response to cocaine following chronic social stress in female rats: exploring a link between depression and drug abuse. *Psychopharmacology (Berl).* 2011; 218:271–279.10.1007/s00213-011-2364-7 [PubMed: 21638221]
- Shin SH, Miller DP, Teicher MH. Exposure to childhood neglect and physical abuse and developmental trajectories of heavy episodic drinking from early adolescence into young adulthood. *Drug Alcohol Depend.* 2013; 127:31–38.10.1016/j.drugalcdep.2012.06.005 [PubMed: 22749563]
- Shipley, W.; Gruber, C.; Martin, T.; Klein, A. Shipley-2. Western Psychological Services; Los Angeles: 2009.
- Shrout PE, Bolger N. Mediation in experimental and nonexperimental studies: new procedures and recommendations. *Psychol Methods.* 2002; 7:422–445. [PubMed: 12530702]
- Simpson TL, Miller WR. Concomitance between childhood sexual and physical abuse and substance use problems. A review. *Clin Psychol Rev.* 2002; 22:27–77. [PubMed: 11793578]
- Sinha R. How does stress increase risk of drug abuse and relapse? *Psychopharmacology (Berl).* 2001; 158:343–359.10.1007/s002130100917 [PubMed: 11797055]
- Sobell, LC.; Sobell, MB. Timeline followback: A technique for assessing self-reported alcohol consumption. In: Litten, RZ.; Allen, J., editors. *Measuring alcohol consumption: Psychosocial and biological methods.* Humana Press; New Jersey: 1992. p. 41-72.
- Soderpalm AH, de Wit H. Effects of stress and alcohol on subjective state in humans. *Alcohol Clin Exp Res.* 2002; 26:818–826. [PubMed: 12068250]

- Strine TW, Dube SR, Edwards VJ, Prehn AW, Rasmussen S, Wagenfeld M, Dhingra S, Croft JB. Associations between adverse childhood experiences, psychological distress, and adult alcohol problems. *Am J Health Behav.* 2012; 36:408–423.10.5993/AJHB.36.3.11 [PubMed: 22370441]
- Tian T, Qin W, Liu B, Jiang T, Yu C. Functional Connectivity in Healthy Subjects Is Nonlinearly Modulated by the COMT and DRD2 Polymorphisms in a Functional System-Dependent Manner. *J Neurosci.* 2013; 33:17519–17526.10.1523/JNEUROSCI.2163-13.2013 [PubMed: 24174684]
- Tidey JW, Miczek KA. Acquisition of cocaine self-administration after social stress: role of accumbens dopamine. *Psychopharmacology (Berl).* 1997; 130:203–212. [PubMed: 9151353]
- Tidey JW, Miczek KA. Social defeat stress selectively alters mesocorticolimbic dopamine release: an in vivo microdialysis study. *Brain Res.* 1996; 721:140–149. [PubMed: 8793094]
- Tonmyr L, Draca J, Crain J, Macmillan HL. Measurement of emotional/psychological child maltreatment: a review. *Child Abuse Negl.* 2011; 35:767–782.10.1016/j.chiabu.2011.04.011 [PubMed: 22018520]
- Tsukada H, Ohba H, Nishiyama S, Kakiuchi T. Differential effects of stress on [(1)(1)C]raclopride and [(1)(1)C]MNPA binding to striatal D(2)/D(3) dopamine receptors: a PET study in conscious monkeys. *Synapse.* 2011; 65:84–89.10.1002/syn.20845 [PubMed: 20687105]
- Tupala E, Hall H, Bergstrom K, Mantere T, Rasanen P, Sarkioja T, Hiltunen J, Tiihonen J. Different effect of age on dopamine transporters in the dorsal and ventral striatum of controls and alcoholics. *Synapse.* 2003; 48:205–211.10.1002/syn.10206 [PubMed: 12687640]
- Tyrka AR, Price LH, Marsit C, Walters OC, Carpenter LL. Childhood adversity and epigenetic modulation of the leukocyte glucocorticoid receptor: preliminary findings in healthy adults. *PLoS One.* 2012; 7:e30148.10.1371/journal.pone.0030148 [PubMed: 22295073]
- Tyrka AR, Wier L, Price LH, Ross NS, Carpenter LL. Childhood parental loss and adult psychopathology: effects of loss characteristics and contextual factors. *Int J Psychiatry Med.* 2008; 38:329–344. [PubMed: 19069576]
- Uhart M, Chong RY, Oswald L, Lin PI, Wand GS. Gender differences in hypothalamic-pituitary-adrenal (HPA) axis reactivity. *Psychoneuroendocrinology.* 2006; 31:642–652.10.1016/j.psyneuen.2006.02.003 [PubMed: 16616815]
- Urban NB, Kegeles LS, Slifstein M, Xu X, Martinez D, Sakr E, Castillo F, Moadel T, O'Malley SS, Krystal JH, Abi-Dargham A. Sex differences in striatal dopamine release in young adults after oral alcohol challenge: a positron emission tomography imaging study with [(1)(1)C]raclopride. *Biol Psychiatry.* 2010; 68:689–696.10.1016/j.biopsych.2010.06.005 [PubMed: 20678752]
- Volkow ND, Wang GJ, Fowler JS, Thanos PP, Logan J, Gatley SJ, Gifford A, Ding YS, Wong C, Pappas N. Brain DA D2 receptors predict reinforcing effects of stimulants in humans: replication study. *Synapse.* 2002; 46:79–82.10.1002/syn.10137 [PubMed: 12211085]
- Volkow ND, Wang GJ, Fowler JS, Logan J, Gatley SJ, Gifford A, Hitzemann R, Ding YS, Pappas N. Prediction of reinforcing responses to psychostimulants in humans by brain dopamine D2 receptor levels. *Am J Psychiatry.* 1999; 156:1440–1443. [PubMed: 10484959]
- Waldrop AE, Back SE, Brady KT, Upadhyaya HP, McRae AL, Saladin ME. Daily stressor sensitivity, abuse effects, and cocaine use in cocaine dependence. *Addict Behav.* 2007; 32:3015–3025.10.1016/j.addbeh.2007.07.006 [PubMed: 17706887]
- Walker QD, Ray R, Kuhn CM. Sex differences in neurochemical effects of dopaminergic drugs in rat striatum. *Neuropsychopharmacology.* 2006; 31:1193–1202.10.1038/sj.npp.1300915 [PubMed: 16237396]
- Wand GS, Oswald LM, McCaul ME, Wong DF, Johnson E, Zhou Y, Kuwabara H, Kumar A. Association of amphetamine-induced striatal dopamine release and cortisol responses to psychological stress. *Neuropsychopharmacology.* 2007; 32:2310–2320.10.1038/sj.npp.1301373 [PubMed: 17342167]
- Wang Z, Du J, Sun H, Wu H, Xiao Z, Zhao M. Patterns of childhood trauma and psychological distress among injecting heroin users in China. *PLoS One.* 2010; 5:e15882.10.1371/journal.pone.0015882 [PubMed: 21209959]
- Weber K, Rockstroh B, Borgelt J, Awiszus B, Popov T, Hoffmann K, Schonauer K, Watzl H, Propster K. Stress load during childhood affects psychopathology in psychiatric patients. *BMC Psychiatry.* 2008; 8:63–244X-8-63.10.1186/1471-244X-8-63 [PubMed: 18651952]

- White HR, Widom CS. Three potential mediators of the effects of child abuse and neglect on adulthood substance use among women. *J Stud Alcohol Drugs*. 2008; 69:337–347. [PubMed: 18432375]
- White TL, Justice AJ, de Wit H. Differential subjective effects of D-amphetamine by gender, hormone levels and menstrual cycle phase. *Pharmacol Biochem Behav*. 2002; 73:729–741. [PubMed: 12213517]
- Wise RA. Drug-activation of brain reward pathways. *Drug Alcohol Depend*. 1998; 51:13–22. [PubMed: 9716927]
- Wise RA. Neurobiology of addiction. *Curr Opin Neurobiol*. 1996; 6:243–251. [PubMed: 8725967]
- Womersley JS, Hsieh JH, Kellaway LA, Gerhardt GA, Russell VA. Maternal separation affects dopamine transporter function in the spontaneously hypertensive rat: an in vivo electrochemical study. *Behav Brain Funct*. 2011; 7:49–9081-7-49.10.1186/1744-9081-7-49 [PubMed: 22133315]
- Yavich L, Tiihonen J. Ethanol modulates evoked dopamine release in mouse nucleus accumbens: dependence on social stress and dose. *Eur J Pharmacol*. 2000; 401:365–373. [PubMed: 10936495]
- Yoder KK, Constantinescu CC, Kareken DA, Normandin MD, Cheng TE, O'Connor SJ, Morris ED. Heterogeneous effects of alcohol on dopamine release in the striatum: a PET study. *Alcohol Clin Exp Res*. 2007; 31:965–973.10.1111/j.1530-0277.2007.00390.x [PubMed: 17428296]
- Zhang XY, Sanchez H, Kehoe P, Kosten TA. Neonatal isolation enhances maintenance but not reinstatement of cocaine self-administration in adult male rats. *Psychopharmacology (Berl)*. 2005; 177:391–399.10.1007/s00213-004-1963-y [PubMed: 15258719]

Fig. 1. Unadjusted relationship between amphetamine-induced ventral striatal (VS) dopamine release and total scores on the Early Trauma Inventory coded by gender. Dopamine release = percent change in [¹¹C]raclopride BP_{ND} from the placebo to the amphetamine scan.

Fig. 2.

Coronal images (10 mm anterior to the anterior commissure point) of BP_{ND} images (saline – amphetamine), spatially normalized to a standard MRI (middle panel), smoothed, and averaged across subjects who showed high (>5 ; Left) or low total scores (<2 ; Right) on the Early Trauma Inventory. Regions are putamen (Pu), caudate nucleus (CN), and ventral striatum (VS). A point in VS was indicated by a plus sign (+) to mark equivalent positions across images. Note that subjects with high total scores showed slightly greater BP_{ND} in the VS than subjects with low total scores.

Fig. 3. Unadjusted means of subjective drug effects (i.e., High, Like, Desire for Drug, and Rush) during the saline (circles) and amphetamine (squares) scans.

Fig. 4.
Unadjusted relationship between total scores on the Early Trauma Inventory (ETI) and Perceived Stress Scale (PSS) coded by gender.

Fig 5.

Model depicting the direct and indirect effects of childhood trauma on amphetamine-induced ventral striatal (VS) dopamine release. Note: the indirect effect from childhood trauma to dopamine release (mediated by perceived stress) was significant ($a*b = 0.285$, 95% CI: 0.008–0.856).

Table 1Subject Characteristics ^a

	Total	Male n=19	Female n=9	p-value
Race, no. (%)				0.272 ^b
Asian	3(10.7)	3(100.0)	0(0.0)	
Black	10(35.7)	6(60.0)	4(40.0)	
White	14(50.0)	10(71.4)	4(28.6)	
Other	1(3.6)	0(0.0)	1(100.0)	
Age, years	22.5(3.1)	22.8(3.1)	21.8(3.2)	0.430
BMI	24.1(2.8)	24.1(2.2)	24.2(2.2)	0.869
Drinks per week	1.7(3.0)	2.0(3.6)	1.3(1.7)	0.893 ^c
Beck Depression Inventory (BDI)	2.3(2.7)	2.4(2.8)	2.1(2.6)	0.707 ^c
Perceived Stress Scale (PSS)	10.4(4.8)	10.6(4.7)	9.8(5.3)	0.671
Early Trauma Inventory (ETI)-Total	3.9(3.1)	4.6(3.1)	2.6(3.0)	0.114
General Traumas	1.9(1.8)	2.1(1.6)	1.4(2.1)	0.235 ^c
Physical Punishment	1.4(1.5)	1.6(1.6)	1.0(1.1)	0.291
Emotional Abuse	0.64(0.91)	0.7(0.9)	0.4(1.0)	0.203 ^c
Sexual Events	0.07(0.26)	0.1(0.3)	0(0.0)	0.321 ^c

^aValues represent means (SD) except where otherwise indicated. Gender differences were compared by t-test except where otherwise indicated;

^bFisher's exact test;

^cWilcoxon-Mann-Whitney test.

Table 2
Effects of Early Trauma and Gender on Ventral Striatal Dopamine Function, Pleasant Drug Effects, and Perceived Stress

Outcomes	Models [†]			Females only		Males only	
	ETI total	Gender	ETI-by-gender interaction	ETI total	ETI total	ETI total	ETI total
	b1 (p-value)	b2(p-value)	b3(p-value)	b1 (p-value)	b1 (p-value)	b1 (p-value)	b1 (p-value)
DA release							
VS	0.79 (0.047)	0.59(0.816)	--	--	--	--	--
Pleasant drug effects [‡]							
Composite score	12.12(0.254)	-31.73(0.591)	-38.42(0.048)	-27.24(0.152)	12.18(0.257)		
Perceived stress							
PSS	0.70(0.024)	0.56(0.775)	--	--	--	--	--
Saline scan BP _{ND}							
VS	0.06(0.015)	-0.06(0.643)	-0.10(0.017)	-0.05(0.084)	0.06(0.031)		

ETI=Early Trauma Inventory, DA=dopamine, VS=ventral striatum, PSS=Perceived Stress Scale, BP_{ND} = [11C]raclopride binding potential.

[†] Full models included terms for ETI total scores, gender and their interaction. If ETI-by-gender interaction was not significant, the reduced models included ETI total scores and gender only.

[‡] Models of pleasant drug effects also controlled for drug effects during saline scan.

Table 3

Effects of Early Trauma and Gender on Regional Striatal Dopamine Function

Outcomes	Models [†]			Females only		Males only	
	ETI total	Gender	ETI-by-gender interaction	ETI total	b1 (p-value)	ETI total	b1 (p-value)
	b1 (p-value)	b2(p-value)	b3(p-value)				
DA release							
APU	0.28(0.471)	-2.17(0.392)	--	--	--	--	--
PPU	-0.38(0.276)	-5.1(0.032)	--	--	--	--	--
ACN	0.67(0.123)	-0.84(0.766)	--	--	--	--	--
PCN	0.35(0.509)	-0.98(0.778)	--	--	--	--	--
Saline scan BP _{ND}							
APU	0.064(0.023)	-0.22(0.153)	-0.13(0.016)	-0.06(0.118)	0.064(0.034)		
PPU	0.019(0.503)	-0.13(0.472)	--	--	--	--	--
ACN	0.077(0.001)	-0.16(0.217)	-0.13(0.002)	-0.057(0.161)	0.077(0.001)		
PCN	0.044(0.148)	-0.045(0.822)	--	--	--	--	--

ETI=Early Trauma Inventory, DA=dopamine, APU = anterior putamen, PPU = posterior putamen, ACN = anterior caudate nucleus, PCN = posterior caudate nucleus, BPND = [¹¹C]raclopride binding potential.

[†] Full models included terms for ETI total scores, gender and their interaction. If ETI-by-gender interaction was not significant, the reduced models included ETI total scores and gender only.

Table 4
Effects of Perceived Stress and Gender on Ventral Striatal Dopamine Function and Pleasant Drug Effects

Outcomes	Models [†]			Females only		Males only	
	PSS	Gender	PSS-by-gender interaction	PSS	b1 (p-value)	PSS	b1 (p-value)
DA release	b1 (p-value)	b2(p-value)	b3(p-value)				
VS	0.56(0.023)	-0.54(0.821)	--	--	--	--	--
Pleasant drug effects							
Composite score [‡]	17.89(0.012)	-12.55(0.809)	-24.63(0.027)	-6.84(0.549)		17.91(0.007)	
Saline scan BP _{ND}							
VS	0.004(0.75)	-0.03(0.827)	--	--	--	--	--

ETI=Early Trauma Inventory, DA=dopamine, VS=ventral striatum, PSS=Perceived Stress Scale, BP_{ND} = [11C]raclopride binding potential.

[†] Full models included terms for ETI total scores, gender and their interaction. If ETI-by-gender interaction was not significant, the reduced models included ETI total scores and gender only.

[‡] Models of pleasant drug effects also controlled for drug effects during saline scan.

Table 5
Effects of Perceived Stress and Gender on Regional Striatal Dopamine Function and Pleasant Drug Effects

Outcomes	Models [†]			Females only		Males only	
	PSS	Gender	PSS-by-gender interaction	PSS	b1 (p-value)	PSS	b1 (p-value)
	b1 (p-value)	b2(p-value)	b3(p-value)				
DA release							
APU	0.40(0.089)	-2.4(0.302)	--	--	--	--	--
PPU	0.029(0.894)	-4.3(0.061)	--	--	--	--	--
ACN	0.63(0.016)	-1.7(0.510)	--	--	--	--	--
PCN	0.30(0.356)	-1.4(0.665)	--	--	--	--	--
Saline scan BP _{ND}							
APU	-0.005(0.755)	-0.19(0.252)	--	--	--	--	--
PPU	-0.0003(0.988)	-0.17(0.339)	--	--	--	--	--
ACN	0.03(0.073)	-0.14(0.285)	-0.069(0.015)	-0.039(0.077)	0.03(0.089)		
PCN	0.0046(0.811)	-0.13(0.512)	--	--	--	--	--

PSS = Perceived Stress Scale, DA=dopamine, APU = anterior putamen, PPU = posterior putamen, ACN = anterior caudate nucleus, PCN = posterior caudate nucleus, BPND = [11C]raclopride binding potential.

[†] Full models included terms for ETI total scores, gender and their interaction. If ETI-by-gender interaction was not significant, the reduced models included ETI total scores and gender only.