of interstitial lung diseases. They found that patients walked longer distances in 6 min with the fast walk compared with the standard walk. We would like to emphasize that an important component of the applicability of the 6MWT is its ability to reflect activities of daily living over other walk tests.² In day-to-day life, patients are used to walking at their own pace and not fast. As a result, the reduction in distance of fast walk may not truly reflect an equivalent reduction in patients' activities of daily living. In addition, for measuring functional capacity, we believe that repeatability of the test, which the authors did not study, is more important.

Another important limitation of the study is the lack of a comparative arm of healthy subjects. The 6-min walk distance is compared with the predicted values obtained from the healthy population for making an interpretation. It is logical that healthy subjects will also walk more distance with the fast walk compared with the standard walk. Hence, it would have been interesting to see whether the increase in walk distance with the fast walk is also seen in terms of percent predicted. We hypothesize that percent predicted values might in fact be lower in patient populations with the fast walk compared with the standard walk.

Guidelines by the American Thoracic Society³ arbitrarily recommend giving a rest of 1 h between successive 6MWTs. Subjects were given a rest of about 15 min between tests in this study, which may potentially have an effect on subsequent tests. We believe that this effect will be larger on tests done after the fast walk compared with the standard walk. If that is true, this effect cannot be nullified by randomization of test instructions. Hence, we believe that it is important to statistically analyze and adjust the effect of order or sequence on the primary outcome.

We appreciate the efforts of the authors in bringing attention to the importance of standardized instructions for 6MWT and the potential effect different instructions can have on the distance walked. However, we believe that the increase in distance walked with the instruction "walk as fast as you can" does not truly reflect the activities of daily living of the patients. We also believe that the effects of previous tests on subsequent tests were not adequately adjusted in analysis to draw reliable conclusions from the study.

Nitin V. Vanjare, MSc
Rahul R. Kodgule, MD
Bill B. Brashier, MD
Pune, India

Affiliations: From the Department of Pulmonary Research, Chest Research Foundation.

Financial/nonfinancial disclosures: The authors have reported to *CHEST* that no potential conflicts of interest exist with any companies/organizations whose products or services may be discussed in this article.

Correspondence to: Nitin V. Vanjare, MSc, Department of Pulmonary Research, Chest Research Foundation, Marigold Complex Kalyaninagar, Pune, Maharashtra 411014, India; e-mail: vanjarenitin@crfindia.com

© 2014 American College of Chest Physicians. Reproduction of this article is prohibited without written permission from the American College of Chest Physicians. See online for more details. DOI: 10.1378/chest.14-0357

REFERENCES

1. Weir NA, Brown AW, Shlobin OA, et al. The influence of alternative instruction on 6-min walk test distance. *Chest*. 2013; 144(6):1900-1905.
2. Solway S, Brooks D, Lacasse Y, Thomas S. A qualitative systematic overview of the measurement properties of functional walk tests used in the cardiorespiratory domain. *Chest*. 2001; 119(1):256-270.

3. ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories. ATS statement: guidelines for the six-minute walk test. *Am J Respir Crit Care Med*. 2002;166(1): 111-117.

Response

To the Editor:

In our recent article in *CHEST*,¹ we showed that replacing "far" with "fast" in the instruction provided to patients resulted in an average improvement of about 50 m in the distance attained. It is intriguing that a one-letter substitution ("st" for "r") can stimulate such an improvement, as well as an ongoing debate. Dr Vanjare and colleagues question the value of optimizing the 6-min walk test (6MWT) distance as a reflection of a patient's ability to do daily activities. However, the 6MWT is a submaximal exercise test designed to determine the patient's capacity to perform activities of daily living.² The American Thoracic Society guidelines on the 6MWT state that "the self-paced 6MWT assesses the submaximal level of functional capacity."² Although some patients achieve their submaximal functional capacity in activities of daily living, many do not. It is the capacity that needs to be assessed and not the true activity level at home. Several studies that are investigating sedentary lifestyles and means to improve upon them address this very topic.^{3,4}

Our findings are pertinent to the very salient issue of the use of the 6MWT in clinical trials. Although the baseline 6MWT predicts outcomes, the change in the 6MWT distance does not.⁵ Therefore, the 6MWT has fallen into disfavor as a study end point. We hypothesized in our article that improving the 6MWT precision and reproducibility may be facilitated by maximizing the distance attained. Indeed, this concept is the foundation for our follow-up study (termed "the fast and the furious"), which is currently in the recruitment phase. Patients perform three 6MWTs on 2 separate days: (1) a standard 6MWT with the "far" instruction, (2) a 6MWT with the "fast" instruction, and (3) a 6MWT test with the fast instruction but a stronger ("furious") menu of scripted instructions each minute to encourage more effort. The hypothesis being tested is that the greater the effort and distance, the less variability in subsequent 6MWTs. We can report that there is a trend supporting this hypothesis among the 12 patients studied thus far, as depicted in Table 1.

Dr Vanjare and colleagues also raise the issue of whether sufficient time (15 min) was allowed between walks. To our knowledge, the American Thoracic Society recommendation for 1 h of rest between walks has never been tested or validated, and it is likely based on intuition and expert opinion. Unfortunately, adhering to this recommendation would increase the duration of four random walk tests in a day to 5 h or longer, thereby reducing the feasibility of our study. If patients were increasingly tired from

Table 1—Variation and Reproducibility in Walk Distance With Differing Instructions (n = 12)

6MWT	Distance, m		
	Visit 1	Visit 2	Difference
Standard	491 (91)	501 (88)	16.5 (11.3)
Fast	495 (71)	502 (83)	13 (13.5)
Furious	511 (55)	515 (91)	8 (8.5)

Data are presented as median (interquartile range). 6MWT = 6-min walk test.

Table 2—Difference in Distance by Sequence of Walk Type

Order of Tests	Median Distance, m		
	Fast Walk	Far Walk	Difference
Fast test first (n = 8)	576 (103)	521 (73)	40 (53)
Far test first (n = 6)	524 (57)	426 (87)	83 (35)

Data are presented as median (interquartile range).

prior walks, then one would expect less of a difference between the fast and far walks when the far test was performed first. To further test this hypothesis, we revisited the order sequence of the far and the fast walks and identified those who performed the fast test first (n = 8) or the far test first (n = 6) and then evaluated the difference between the fast and the far walks for the individual subjects (Table 2).

Although the numbers are small, outperformance with the fast instruction appears to be consistent regardless of the order of the tests. Therefore, based on this subgroup analysis, we feel that 15 min of rest is sufficient.

We thank Dr Vanjare and colleagues for their interest in our article and for fostering this discussion, which has enabled us to provide further insight into the performance of this conceptually simple, yet remarkably controversial, test.

Steven D. Nathan, MD, FCCP

A. Whitney Brown, MD

Nargues A. Weir, MD, FCCP

Falls Church, VA

Affiliations: From the Advanced Lung Disease and Transplant Program, Department of Medicine, Inova Fairfax Hospital.

Financial/nonfinancial disclosures: The authors have reported to *CHEST* the following conflicts of interest: Dr Nathan has served as a consultant for Intermune; Gilead Sciences, Inc; Actelion Pharmaceuticals Ltd; GeNO LLC; Boehringer-Ingelheim GmbH; Bayer; Genentech; and United Therapeutics Corporation. He has received grant funding from Intermune; Gilead Sciences, Inc; Actelion Pharmaceuticals Ltd; Boehringer-Ingelheim GmbH; and FibroGen, Inc. He has also served on speakers' bureaus for Gilead Sciences, Inc; Actelion Pharmaceuticals Ltd; and United Therapeutics Corporation. Dr Brown has served on an advisory board for Gilead Pharmaceuticals. Dr Weir is the recipient of grant funding from Gilead Pharmaceuticals.

Correspondence to: Steven D. Nathan, MD, FCCP, Advanced Lung Disease and Transplant Program, Department of Medicine, Inova Fairfax Hospital, 3300 Gallows Rd, Falls Church, VA 22042; e-mail: steven.nathan@inova.org

© 2014 American College of Chest Physicians. Reproduction of this article is prohibited without written permission from the American College of Chest Physicians. See online for more details.

DOI: 10.1378/chest.14-0720

REFERENCES

1. Weir NA, Brown AW, Shlobin OA, et al. The influence of alternative instruction on 6-min walk test distance. *Chest*. 2013;144(6):1900-1905.
2. ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories. ATS statement: guidelines for the six-minute walk test. *Am J Respir Crit Care Med*. 2002;166(1):111-117.
3. Moy ML, Weston NA, Wilson EJ, Hess ML, Richardson CR. A pilot study of an Internet walking program and pedometer in COPD. *Respir Med*. 2012;106(9):1342-1350.
4. Quon BS, Patrick DL, Edwards TC, et al. Feasibility of using pedometers to measure daily step counts in cystic fibrosis and

an assessment of its responsiveness to changes in health state.

J Cyst Fibros. 2012;11(3):216-222.

5. Gabler NB, French B, Strom BL, et al. Validation of 6-minute walk distance as a surrogate end point in pulmonary arterial hypertension trials. *Circulation*. 2012;126(3):349-356.

Anticytokeratin (CAM5.2) Reagent Identifies Cytokeratins 7 and 8, Not Cytokeratin 18

To the Editors:

We read with great interest the contribution by Mani and Zander published in an issue of *CHEST* (November 2012).¹ The authors presented a comprehensive and good quality review. However, one aspect of this article needs clarification, as it contains an incorrect message that we believe may misguide readers. The authors seem to have inadvertently stated that anticytokeratin (CAM5.2) identifies cytokeratins (CKs) 8 and 18.¹

The source of CAM5.2 is Becton, Dickinson and Co. This antibody is derived from mouse IgG2a, clone CAM5.2. On the basis of the company's datasheet, this clone reacts with CK7 and CK8.^{2,3} On the other hand, many manufacturers supplied the CK8/18 monoclonal antibody, which could be derived from the clone 5D3 (eg, Leica Microsystems GmbH, Thermo Fisher Scientific Inc, Imgenex Corp, Abcam plc, Novus Biologicals, GeneTex Inc); clones B22.1 and B23.1 (Novus Biologicals, Cell Marque Corp, GenWay Biotech Inc, Santa Cruz Biotechnology Inc, MyBiosciences.com); clone SP141 (eg, Santa Cruz Biotechnology Inc, Abcam plc, and GeneTex Inc); and clone Zym5.2 (UCD/PR-10.11; Invitrogen Corp). They reacted against CK8 and CK18.⁴

Sometimes CAM5.2 has been mistakenly regarded in both the clinical and anatomic pathologic communities as useful for identifying CKs 8 and 18, since previous results by Makin et al⁵ were attributed to breakdown products of CK8, revealing smaller molecular-weight fractions on immunoblot analysis. Becton, Dickinson and Co has revised the datasheet for CAM5.2 to indicate it has a primary reactivity with CK8 and a weaker but distinct reactivity with CK7. However, there is no reactivity with CK18 or other CKs.^{2,3} As a result, we clarify that CAM5.2 is not identical to antibodies CK8/18.

Kuang-Leei Chang, MD

Wan-Ru Chao, MD

Chih-Ping Han, MD, PhD

Taichung, Taiwan

Affiliations: From the Department of Emergency Medicine (Dr Chang), Department of Pathology (Drs Chao and Han), Clinical Trial Center (Dr Han), and Department of Obstetrics and Gynecology (Dr Han), Chung-Shan Medical University Hospital; and Institute of Medicine (Dr Chao) and School of Medicine (Drs Chao and Han), Chung-Shan Medical University.

Financial/nonfinancial disclosures: The authors have reported to *CHEST* that no potential conflicts of interest exist with any companies/organizations whose products or services may be discussed in this article.

Correspondence to: Chih-Ping Han, MD, PhD, Department of Obstetrics and Gynecology, Chung-Shan Medical University Hospital and School of Medicine, Chung-Shan Medical University, No. 110, Sec. 1, Jiangsu N Rd, Taichung, Taiwan 40201; e-mail: hanhaly@gmail.com.

© 2014 American College of Chest Physicians. Reproduction of this article is prohibited without written permission from the American College of Chest Physicians. See online for more details.

DOI: 10.1378/chest.14-0168