Published in final edited form as:

J Hosp Med. 2014 June ; 9(6): 379–383. doi:10.1002/jhm.2184.

Predictors of Rehospitalization after Admission for Pneumonia in the VA Health Care System

Victoria L. Tang, MD, Ethan A. Halm, MD, MPH, Michael J. Fine, MD, MSc, Christopher S. Johnson, MPH, Antonio Anzueto, MD, and Eric M. Mortensen, MD, MSc

VA North Texas Health Care System (VLT, EMM, CSJ). University of Texas Southwestern Medical Center (VLT, EMM, EAH, CSJ). South Texas Veterans Health Care System and University of Texas Health Science Center at San Antonio, San Antonio Texas (AA). VA Center for Health Equity Research and Promotion and VA Pittsburgh Healthcare System, and Division of General Internal Medicine, Department of Medicine, University of Pittsburgh (MJF)

Abstract

Introduction—Although some factors associated with rehospitalization after community-acquired pneumonia have been identified, other factors such as medical care utilization and medication usage have not been previously studied. We investigated novel predictors of rehospitalization in patients admitted with pneumonia.

Methods—Using Department of Veteran Affairs (VA) administrative data from October 2001 to September 2007, we examined a cohort of patients 65 years old and older who were hospitalized with pneumonia in 150 VA acute care hospitals. The cohort was randomly split into derivation and validation samples, and then used logistic regression models to identify and validate predictors of all-cause rehospitalization within 30 days.

Results—Of the 45,134 subjects, 13% were rehospitalized within 30-days. No significant differences were noted between the derivation and validation cohorts. Factors associated with readmission included age, being married, chronic renal disease, prior malignancy, nursing home residence, congestive heart failure, use of oral corticosteroids, number of emergency department visits in the prior year, number of outpatient clinic visits in the year prior, prior admission within 90-days, and length of hospital stay. The c-statistics for the derivation and validation models were 0.615 and 0.613, respectively.

Conclusions—Factors associated with readmission were largely unrelated to the underlying pneumonia but to demographics, comorbidities, healthcare utilization and length of stay on index admission.

Keywords

Readmission; rehospitalization; community-acquired pneumonia; pneumonia

Corresponding author: Eric Mortensen, MD, MSc, FACP, Dallas VA Medical Center, General Internal Medicine (111E), 4500 South Lancaster, Dallas, Texas 75216, Cell: 210-602-7316, Fax: 214-857-1575, Eric.Mortensen@UTSouthwestern.edu.

The views expressed in this article are those of the authors and do not necessarily represent the views of the Department of Veterans Affairs.

INTRODUCTION

Pneumonia remains a significant problem in the United States, both in terms of mortality as well as cost to the healthcare system. Each year, over a million patients are hospitalized for pneumonia with costs conservatively estimated at greater than \$7 billion in 2010 [1]. One contributing factor for these costs is that pneumonia is the second most frequent reason for rehospitalization [2]. Reprehospitalization after admission for pneumonia is now used as a marker for quality of care.

Given the cost and adverse outcomes of pneumonia, it is important to examine early rehospitalization to better understand what factors play a role. Studies have examined factors associated with readmission following an initial pneumonia-related hospitalization. Adamuz et al. showed that additional health care visits and rehospitalizations within 30 days of discharge from the hospital were common and associated with worsening of signs or symptoms of pneumonia and/or comorbidities [3]. Other studies demonstrated that “instability” on hospital discharge, as well as treatment failure, were associated with increased risk of rehospitalization [4, 5]. Jasti et al. found rehospitalizations following pneumonia were usually comorbidity related, primarily cardiopulmonary and/or neurologic disease, rather than due to the index pneumonia [6]. Many of these studies and others looking at age, activity of daily living score, socioeconomic status, and comorbidity characteristics were performed in relatively small cohorts. Predictors of rehospitalization have not been studied in a large cohort of patients in an integrated health care system.

Our study looks at factors not addressed in prior studies that have used administrative claims data to identify factors associated with early readmission. We also evaluated these admission risk factors in a Veteran population, whereas prior studies have primarily focused on those who receive Medicare. The purpose of this study was to examine predictors of early (30 days) readmission in the VA for patients age 65 years and older hospitalized for pneumonia. Our a priori hypothesis was that comorbid illnesses, such as congestive heart failure and chronic obstructive pulmonary disease, and patients with high medical complexity, such as high number of medications and/or prior hospitalizations and nursing home residence, are the primary factors associated with increased risk of rehospitalization.

METHODS

For this national cohort study, we used data from the VA Health Care System administrative and clinical databases that serve as repositories of clinical data from more than 150 VA hospitals and 850 outpatient clinics throughout the U.S. The Institutional Review Boards of the University of Texas Health Science Center at San Antonio and VA North Texas Health Care System approved this study. Details regarding the study design and methods were previously published [7].

Inclusion Criteria

Patients included in this study were hospitalized between October 2001 to September 2007, had a primary diagnosis of pneumonia/influenza (ICD-9 codes 480.0–483.99 or 485–487) or a secondary discharge diagnosis of pneumonia with a primary diagnosis of respiratory

failure (ICD-9 code 518.81) or sepsis (ICD-9 code 038.xx) [8], and were age 65 or older on the date of admission. To ensure patients obtained their care primarily at the VA, at least one VA outpatient clinic visit in the year preceding the index admission and at least one active and filled outpatient medication from a VA pharmacy within 90-days of admission were required for inclusion.

If a patient was admitted more than once during the study period, only the first hospitalization was included as the index admission.

Exclusion Criteria

Patients who died during the initial hospitalization were not included in this study. Patients admitted to hospitals with fewer than 25 reported hospitalizations during October 2001 to September 2007 were excluded as these hospitals are neither representative nor generalizable. This resulted in the removal of 59 patients from 5 different hospitals.

Data Sources and Definitions

We used inpatient and outpatient demographic, utilization, and comorbidity data from the National Patient Care Database. Pharmacy data were extracted from the Decision Support System National Data Extracts and Pharmacy Benefits Management. Vital status information was obtained from the Vital Status file, which incorporates data from Veterans' death benefits claims, inpatient deaths, Medicare Vital Status files, and the Social Security Administration's death master file. We used encrypted patient identifiers link to information across these databases.

We obtained demographic information (age, sex, race, marital status) from inpatient and outpatient data. We categorized race as white, black, Hispanic, and other/unknown. To infer active smoking and/or tobacco cessation attempts, we identified ICD-9 codes for tobacco use (305.1, V15.82), smoking cessation clinic use, and/or use of medications for the treatment of nicotine dependence (Zyban, nicotine replacement, or varenicline). We used VA Priority status as a proxy for socioeconomic and disability status. VA priority groups are a way for the VA to focus limited funds to those veterans most in need. The highest group (priority group 1) must have at least a 50% service-connected disability. Priority groups 2 through 6 includes veterans with up to 40% service-connected disability, former prisoners of war, those awarded certain honors, veterans with lower incomes, and the catastrophically disabled. The lowest groups (priority groups 7 and 8) include veterans with relatively higher incomes who agree to pay copayments [9].

We also obtained information on comorbid conditions from inpatient and outpatient administrative data. We defined alcohol abuse using ICD-9 codes 291, 303, 305.0, and illicit drug use with ICD-9 codes 292, 304, 305 excluding 305.0-1. We used the Charlson comorbidity index to quantify levels of preexisting comorbidity [10], adapted for administrative databases, using ICD-9 codes for 19 comorbid conditions from prior outpatient and inpatient encounters. We defined cardiovascular events and lung cancer that were diagnosed during the hospitalization as previously described. [11, 12]

Outcomes

Our primary study outcome was readmission within 30-days of hospital discharge for pneumonia from any VA acute care hospital only. Medicare has used a 30-day readmission as a quality indicator as readmissions that occur closer to discharge are believed to be more likely due to events during the index hospitalization [13].

Statistical Analyses

We randomly divided patients from our initial cohort into equal derivation or validation cohorts. We assessed differences between the two groups using Student's t-test for continuous variables and chi-square test for categorical variables. We performed univariate logistic regression analyses in the derivation cohort to examine the relationship between 30-day rehospitalization and each of our potential covariates. We entered covariates that were significant at $p < 0.10$ in the univariate analyses into a multiple regression model.

Significant covariates at $p < 0.10$ were then entered into the final model. For this model, we used bootstrapping with replacement in 1,000 replications to obtain standard errors of our coefficients and associated p-values. Since some subjects had more than one index admission, we used robust variance-covariance matrix estimators to compute standard errors for model coefficients.

We used the c-statistic to assess the discrimination of our model. Calibration of this model was measured using the Hosmer-Lemeshow chi-square goodness of fit test, using 10 quantiles to group the data. We evaluated differences in discrimination between the derivation and validation cohorts by comparing c-statistics.

Lastly, using the final model in the derivation cohort, five categories of severity were created based on subjects' predicted risk values for readmission. Severity categories in both the derivation and validation cohorts were then graphically evaluated.

We defined statistical significance as a two-tailed p value of < 0.05 . We used STATA 12 (College Station, Texas) for all analyses.

RESULTS

Patient Characteristics

Of the 45,134 eligible patients, 13.2% were rehospitalized within 30-days and rates by hospital ranged from 1.6% to 20.7%. Table 1 presents the comparison of baseline difference between the derivation and validation cohorts. There were no statistically significant differences between the two groups. Overall, the mean age was 77 years, 98% were male, and 54% were married. Over 90% of the patients received guideline concordant antibiotics and only 10% required stay in the intensive care unit. The average length of stay was 7 days (standard deviation of 11 days).

Predictors of 30-Day Readmission

All variables listed in Table 1 were evaluated. The derivation logistic regression model is shown in Table 2 and only include predictors associated with rehospitalization. Antibiotic

concordance and time in ICU were evaluated and did not predict readmission. Variables that were significantly associated with rehospitalization in the models included age, marital status, the number of emergency department clinic visits a year prior, prior admission, number of non-pharmacy clinic visits in 1 year prior, and hospital length of stay. The increasing age and number of ER and clinic visits were associated with higher odds of readmission. The longer the length of stay and those patients that were married had a lower odds of subsequent readmission. Other variables included were the presence of chronic renal disease, prior malignancy, nursing home residence, congestive heart failure, and prior use of oral corticosteroids. The c-statistics for the derivation and validation models were 0.615 and 0.613, respectively. There were no significant differences in any of the variables between the derivation and validation cohorts.

Figure 1 shows the derivation and validation models are similar in percent re-hospitalization group in all severity groups.

DISCUSSION

Sociodemographics and comorbidities, which are “host” factors, were significantly associated with rehospitalization of elderly patients admitted for pneumonia and length of hospital stay, the only pneumonia-specific variable associated with readmission [6, 14–16]. Higher rates of all types of prior utilization (inpatient, outpatient, emergency department visits) were also associated with higher risk of rehospitalization. Our findings highlight the role that “host” factors play in rehospitalization of elderly patients with pneumonia and that there are few potentially modifiable targets to help reduce readmissions after pneumonia.

These findings may indicate two things: patients in this cohort are sicker than the general population at baseline so require more medical care, and/or they are being evaluated by more providers increasing the chance they will be readmitted. For the former, an example is the association of prior use of oral corticosteroids with rehospitalization, which may demonstrate poorly controlled comorbidities (e.g., chronic obstructive lung disease with recent exacerbation) that require steroid use. Given the VA’s integrated healthcare system, greater access to primary care providers may lead to more opportunities for patients to voice their complaints, which can lead to more readmissions among severely ill patients. Weinberger et. al. had found an increase in hospital readmissions with increased access to primary care[17]. The association of readmission and length of stay is less likely related to pneumonia-related clinical instabilities on discharge, but rather to the ability to address other potential issues related to the patient’s underlying comorbidities [5].

Our study presents a model looking at rehospitalization of patients with pneumonia, including factors not addressed in a prior publication by Lindenauer et al., the model CMS uses to predict early rehospitalization [18]. Lindenauer’s model included 39 variables (of which 37 were clinical variables), including “vertebral fractures”, “other injuries”, “other gastrointestinal disorders”, “other urinary tract disorders”, as well as demographic and comorbidity factors that were similar to some of those included in our study. Their 30-day rehospitalization medical record and administrative models had a c-statistic of 0.59 and 0.63, respectively. A major difference between our presented model and the latter is the data on

utilization of medical care. Despite this additional data, the resulting c-statistics of 0.615 and 0.613, were not qualitatively better.

The similar c-statistics demonstrates that factors, which play a significant role in determining rehospitalization, have yet to be identified, and it is not clear if researchers will be able to identify potentially modifiable risk factors for rehospitalization. Examples of other potential factors that should be examined in future studies include family/community support, quality of the transition of care back to their residence, and quality of inpatient care. Calvillo-King et al. found a broad range of social factors that potentially affect the risk of post-discharge readmission for chronic heart failure [19]. For pneumonia, there were few studies that assessed the risk of social factors on readmission, and these were most commonly rather broad categories such as race/ethnicity or nursing home residence. In the publications that studied social factors and readmissions, there were inconsistent associations between hospital readmission with lower education, low income, and unemployment. Future papers may benefit from taking these additional factors into account.

Potential limitations to our study include the overwhelming majority of patients being male and in a VA population, which may limit generalizability. In addition, no data was available on the quality of the transition of care. A limitation to this study is the age of the data, evaluation of outpatient follow-up, and antibiotics on discharge. Though our data is up to 10 years old, this study evaluates risk factors that are still relevant. Finally some variables noted in the CMS model were not evaluated in our study, such as protein-calorie malnutrition, skin ulcers, and fracture history. Given these differences, we were unable to directly compare the performance of the two models.

Despite examining novel factors related to prior health care utilization our model still performed sub-optimally in identifying subjects at risk for rehospitalization after pneumonia. Future studies need to examine other social factors and patient frailty that may increase the risk of readmissions, as well as examining the impact of clinical stability on discharge. It remains to be determined if readmissions after pneumonia are truly preventable or mostly attributable to the patient's prior health status.

Acknowledgments

The project described was supported by Grant Number R01NR010828 from the National Institute of Nursing Research. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Nursing Research or the National Institutes of Health. This material is the result of work supported with resources and the use of facilities at the VA North Texas Health Care System. Funding agencies had no role in conducting the study, or role in the preparation, review, or approval of the manuscript.

References

1. Thomas CP, Ryan M, Chapman JD, et al. Incidence and cost of pneumonia in medicare beneficiaries. *Chest*. 2012; 142(4):973–81. [PubMed: 22406959]
2. Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program. *N Engl J Med*. 2009; 360(14):1418–28. [PubMed: 19339721]
3. Adamuz J, Viasus D, Camprecios-Rodriguez P, et al. A prospective cohort study of healthcare visits and rehospitalizations after discharge of patients with community-acquired pneumonia. *Respirology*. 2011; 16(7):1119–26. [PubMed: 21736665]

4. Capelastegui A, Espana PP, Bilbao A, et al. Pneumonia: criteria for patient instability on hospital discharge. *Chest*. 2008; 134(3):595–600. [PubMed: 18490403]
5. Halm EA, Fine MJ, Kapoor WN, et al. Instability on hospital discharge and the risk of adverse outcomes in patients with pneumonia. *Arch Intern Med*. 2002; 162(11):1278–84. [PubMed: 12038946]
6. Jasti H, Mortensen EM, Obrosky DS, et al. Causes and risk factors for rehospitalization of patients hospitalized with community-acquired pneumonia. *Clin Infect Dis*. 2008; 46(4):550–6. [PubMed: 18194099]
7. Mortensen EM, Nakashima B, Cornell J, et al. Population-based study of statins, angiotensin II receptor blockers, and angiotensin-converting enzyme inhibitors on pneumonia-related outcomes. *Clin Infect Dis*. 2012; 55(11):1466–73. [PubMed: 22918991]
8. Meehan TP, Fine MJ, Krumholz HM, et al. Quality of care, process, and outcomes in elderly patients with pneumonia. *JAMA*. 1997; 278(23):2080–4. [PubMed: 9403422]
9. Affairs, U.S.D.o.V. Health benefits priority groups table. [cited 2013 July 1]; Available from: http://www.va.gov/healthbenefits/resources/priority_groups.asp%5D
10. Charlson M, Szatrowski TP, Peterson J, et al. Validation of a combined comorbidity index. *J Clin Epidemiol*. 1994; 47(11):1245–1251. [PubMed: 7722560]
11. Perry TW, Pugh MJ, Waterer GW, et al. Incidence of cardiovascular events after hospital admission for pneumonia. *The American Journal of Medicine*. 2011; 124(3):244–51. [PubMed: 21396508]
12. Mortensen EM, Copeland LA, Pugh MJ, et al. Diagnosis of pulmonary malignancy after hospitalization for pneumonia. *Am J Med*. 2010; 123(1):66–71. [PubMed: 20102994]
13. MedPAC. 2013. [cited 2013 July 11]; Available from: <http://www.hospitalcompare.gov>
14. Epstein AM, Jha AK, Orav EJ. The relationship between hospital admission rates and rehospitalizations. *N Engl J Med*. 2011; 365(24):2287–95. [PubMed: 22168643]
15. Capelastegui A, Espana Yandiola PP, Quintana JM, et al. Predictors of short-term rehospitalization following discharge of patients hospitalized with community-acquired pneumonia. *Chest*. 2009; 136(4):1079–85. [PubMed: 19395580]
16. Herzog NS, Bratzler DW, Houck PM, et al. Effects of previous influenza vaccination on subsequent readmission and mortality in elderly patients hospitalized with pneumonia. *Am J Med*. 2003; 115(6):454–61. [PubMed: 14563502]
17. Weinberger M, Oddone EZ, Henderson WG. Does Increased Access to Primary Care Reduce Hospital Readmissions? *New England Journal of Medicine*. 1996; 334(22):1441–1447. [PubMed: 8618584]
18. Lindenauer PK, Normand SL, Drye EE, et al. Development, validation, and results of a measure of 30-day readmission following hospitalization for pneumonia. *J Hosp Med*. 2011; 6(3):142–50. [PubMed: 21387551]
19. Calvillo-King L, Arnold D, Eubank KJ, et al. Impact of social factors on risk of readmission or mortality in pneumonia and heart failure: systematic review. *J Gen Intern Med*. 2013; 28(2):269–82. [PubMed: 23054925]

Figure 1.
Risk of 30-day readmission in the Derivation versus Validation Cohort by Pneumonia Severity on Admission

Table 1

Comparison of Patients Hospitalized with Pneumonia in the Derivation and Validation Cohorts

Characteristics	Derivation Cohort N=22,567(%)	Validation Cohort N=22,567(%)	P-value
Sociodemographics			
Age, mean (SD)	77.3 (6.5)	77.2 (6.5)	0.30
Male	22,133 (98.1)	22,175 (98.3)	0.14
White	18,858 (83.6)	18,719 (83.0)	0.08
Black	2,562 (11.4)	2,583 (11.5)	0.76
Hispanic	1,337 (5.9)	1,414 (6.3)	0.13
Married	12,166 (53.9)	12,258 (54.3)	0.39
VA Priority Group*			0.28
1	4,286 (19.0)	4,281 (19.0)	
2–6	16,306 (72.3)	16,399 (72.8)	
7–8	1,954 (8.7)	1,860 (8.3)	
Nursing Home Residence	199 (0.9)	216 (1.0)	0.41
Smoker Status	8,579 (38.0)	8,677 (38.5)	0.34
Index Hospitalization Factors			
Myocardial Infarction During Hospitalization	469 (2.1)	452 (2.0)	0.57
Heart Failure During Hospitalization	4,817 (21.4)	4,772 (21.2)	0.61
Arrhythmia During Hospitalization	4,356 (19.3)	4,287 (19.0)	0.41
Length of Stay, mean (SD)	7.2 (10.4)	7.2 (11.7)	0.76
Lung Cancer Diagnosis During Hospitalization	31 (0.1)	31 (0.1)	1.00
ICU Admission	2,282 (10.1)	2,336 (10.4)	0.40
Guideline Concordant Antibiotics	20,724 (91.8)	20,643 (91.5)	0.17
Invasive Mechanical Ventilation	793 (3.5)	832 (3.7)	0.32
Vasopressor Use	495 (2.2)	489 (2.2)	0.85
Hospital Complications			
Renal Organ Failure	2,671 (11.8)	2,640 (11.7)	0.65
Cardiac Organ Failure	1,004 (4.5)	959 (4.3)	0.30
Hepatic Organ Failure	33 (0.2)	40 (0.2)	0.41
Respiratory Organ Failure	1,623 (7.2)	1,583 (7.0)	0.46
Comorbid Illnesses			
Medical			
Myocardial Infarction	1,528 (6.8)	1,542 (6.8)	0.79
Congestive Heart Failure	5,815 (25.8)	5,697 (25.2)	0.20
Peripheral Vascular Disease	3,413 (15.1)	3,454 (15.3)	0.59
Cerebrovascular Disease	3,993 (17.7)	4,078 (18.1)	0.30
Dementia	1,141 (5.1)	1,110 (4.9)	0.50
COPD	12,168 (53.9)	12,076 (53.5)	0.39

Characteristics	Derivation Cohort N=22,567(%)	Validation Cohort N=22,567(%)	P-value
Rheumatologic Disease	646 (2.9)	660 (2.9)	0.69
Peptic Ulcer Disease	795 (3.5)	723 (3.2)	0.06
Severe Liver Disease	169 (0.8)	169 (0.8)	1.00
Mild Liver Disease	78 (0.4)	80 (0.4)	0.87
Diabetes	7,310 (32.4)	7,410 (32.8)	0.32
Diabetes with Complications	2,229 (9.9)	2,248 (10.0)	0.77
Chronic Renal Disease	2,836 (12.6)	2,745 (12.2)	0.19
Hemi/paraplegia	319 (1.4)	338 (1.5)	0.46
Any Prior Malignancy	5,226 (23.2)	5,269 (23.4)	0.63
Metastatic Solid Tumor	748 (3.3)	795 (3.5)	0.22
AIDS	56 (0.3)	49 (0.2)	0.49
HIV	23 (0.1)	13 (0.1)	0.10
Alcohol Abuse	897 (4.0)	934 (4.1)	0.38
Drug Abuse	254 (1.1)	255 (1.1)	0.96
Psychiatric			
Anxiety Disorder	1,692 (7.5)	1,722 (7.6)	0.59
Depression Indicator	3,655 (16.2)	3,718 (16.5)	0.42
Bipolar Disorder	430 (1.9)	433 (1.9)	0.92
Cataract Indicator	6,106 (27.1)	6,180 (27.4)	0.43
Prostatitis Indicator	4,680 (20.7)	4,593 (20.4)	0.31
Schizophrenia Indicator	685 (3.0)	676 (3.0)	0.80
Posttraumatic Stress Disorder	844 (3.7)	862 (3.8)	0.66
Medication History Within 90-days			
Cardiovascular Drugs, mean (SD)	1.8 (1.6)	1.8 (1.6)	0.65
Diabetes Drugs, mean (SD)	0.3 (0.7)	0.3 (0.7)	0.60
Inhaled Corticosteroids, mean (SD)	0.4 (0.9)	0.4 (0.9)	0.28
Pulmonary Drug, mean (SD)	1.3 (2.0)	1.3 (2.0)	0.36
Oral Corticosteroids	5,363 (23.8)	5,505 (24.4)	0.12
Prior Medical Utilization			
Number of Primary Care Clinic Visits Within 1 Year, mean (SD)	4.9 (4.2)	4.9 (4.2)	0.67
Number of Emergency Department Within 1 Year, mean (SD)	1.2 (2.0)	1.2 (2.0)	0.86
Number of Outpatient Clinic Visits Within 1 Year, mean (SD)	14.7 (13.1)	14.6 (13.0)	0.33
Number of Specialty Clinic Visits Within 1 Year, mean (SD)	3.4 (5.5)	3.3 (5.4)	0.50
Prior Hospital Admission Within 90- days	5,062 (22.4)	5,141 (22.8)	0.37

Table 2

Multivariable Predictors of 30-day readmission (Derivation Cohort)*

		95% CI		
Predictors	OR	Lower	Upper	P-value
Sociodemographics				
Age at Admission	1.01	1.00	1.02	<0.01
White	1.14	1.02	1.28	0.02
Non-white	1.00			
Hispanic	1.26	1.07	1.47	0.01
Not Hispanic	1.00			
Married	0.89	0.82	0.96	<0.01
Not married	1.00			
Nursing Home Residence	1.68	1.18	2.39	<0.01
Comorbid Illnesses				
Peptic Ulcer Disease	1.24	1.02	1.50	0.03
Chronic Renal Disease	1.18	1.06	1.33	<0.01
Any Prior Malignancy	1.17	1.06	1.28	<0.01
Metastatic solid tumor	1.20	0.98	1.47	0.08
AIDS	1.99	1.06	3.74	0.03
Index Admission Factors				
Heart Failure on Initial Admission	1.18	1.07	1.29	<0.01
Length of Stay (per day)	0.97	0.97	0.98	<0.01
Medication History				
Cardiovascular Drug Count 90 days Prior	1.03	1.00	1.06	0.02
Oral Corticosteroids 90 days Prior	1.16	1.06	1.27	<0.01
Prior Medical Utilization				
Number of Emergency Department Visits Within 1 Year	1.04	1.02	1.06	<0.01
Number of Specialty Clinic Visits Within 1 Year	1.01	1.00	1.02	0.01
Prior Hospital Admission within 90-days	1.60	1.46	1.75	<0.01
Number of Non-Pharmacy Clinic Visits Within 1 Year	1.00	1.00	1.01	<0.01

* includes only predictors from Table 1 that predicted readmission.

† reference group includes native Hawaiian, American Indian, black, Asian, and unknown

‡ reference group includes non-Hispanic white, non-Hispanic black, Asian, American Indian, and unknown

§ reference group includes never married, separated/divorced, widowed, and unknown