First Report of *Anaplasma phagocytophilum* and *Babesia microti* in Rodents in Finland

Eva R. Kallio,^{1,2} Michael Begon,² Richard J. Birtles,³ Kevin J. Bown,³
Esa Koskela,¹ Tapio Mappes,¹ and Phillip C. Watts^{1,2}

Abstract

Tick-borne diseases pose an increasingly important public health problem in Europe. Rodents are the reservoir host for many tick-transmitted pathogens, including *Anaplasma phagocytophilum* and *Babesia microti*, which can cause human granulocytic anaplasmosis and babesiosis, respectively. To estimate the presence of these pathogens in rodents in Finland, we examined blood samples from 151 bank voles (*Myodes glareolus*) and demonstrate, for the first time, that *A. phagocytophilum* and *B. microti* commonly infect bank voles (in 22% and 40% of animals, respectively) in Finland. Sequence analysis of a fragment of 18S rRNA showed that the *B. microti* strain isolated was identical to the Munich strain, which is considered to be nonzoonotic. The *A. phagocytophilum* strain (based on a fragment of the *msh4* gene) was identical to one found earlier in rodents in the United Kingdom that is transmitted by the tick *Ixodes trianguliceps*, all the life stages of which feed on small mammals. The infection probability of *B. microti* in the bank voles was the greater the older the individual was, and males were more often infected than females. *A. phagocytophilum* infection probability first increased and then decreased with the age of individual without any difference between sexes. While these pathogens presumably pose a limited zoonotic risk to humans in Finland, they might have important interactions with other rodent pathogens and therefore affect infection dynamics of, for example, zoonotic pathogens.

Key Words: *Anaplasma phagocytophilum*—*Babesia microti*—Bank voles—*Myodes glareolus*—Anaplasmosis—Babesiosis—Finland.

Introduction

TICK-BORNE ZOO NOTIC DISEASES are an increasingly important health problem in Europe for a number of reasons. For example, following the expansion in the geographic range of the tick *Ixodes ricinus* (Jaenson et al. 2012a, Medlock et al. 2013), tick-borne diseases such as tick-borne encephalitis (TBE) and Lyme borreliosis are being found in new areas (Rizzoli et al. 2011, Jaenson et al. 2012b). Moreover, different interactions between people and the environment increase the numbers of humans exposed to ticks and tick-borne pathogens (Sumilo et al. 2008). In addition, several emerging zoonoses, such as human granulocytic anaplasmosis (HGA) and babesiosis, that are caused by the obligate intracellular bacterium *Anaplasma phagocytophilum* and members of the protozoan hemoparasitic genus *Babesia*, respectively, are posing risks to humans in Europe (Doudier et al. 2010, Hildebrandt et al. 2011).

Babesia microti is the main causative agent for human babesiosis in the United States (Gray et al. 2010). Although there is only one confirmed human infection by *B. microti* in Europe (Hildebrandt et al. 2007), because most cases are caused by *B. divergens* (Gray et al. 2010), serological studies indicate that human *B. microti* infections are more common than previously recognized (Hunfeldt et al. 2002). *B. microti* is widely distributed across the Northern Hemisphere (Goethert and Telford 2003), and at least two *B. microti* strains (Munich and Jena), which may differ in terms of their association with human disease, are found in Europe (Pieniazek et al. 2006, Siński et al. 2006, Bown et al. 2008, Beck et al. 2011, Katargina et al. 2011).

In Europe, some 70 confirmed HGA cases have been reported to date (Dumler et al. 2005, Edouard et al. 2012). However, seroepidemiological studies have revealed high (up to 21%) *A. phagocytophilum* antibody prevalences in some areas in Europe (Dumler et al. 2005), suggesting that asymptomatic

¹Department of Biological and Environmental Science, University of Jyväskylä, Jyväskylä, Finland.

²Department of Evolution, Ecology and Behaviour, Institute of Integrative Biology, University of Liverpool, Liverpool, United Kingdom.

³School of Environment and Life Sciences, University of Salford, Salford, United Kingdom.

human infections might be common (Doudier et al. 2010). *A. phagocytophilum* also has indirect implications for humans because it causes, for example, a tick-borne fever in livestock and febrile fevers in companion animals (cats and dogs) (Rar and Golovljova 2011). Many wildlife mammalian species, such as rodents, shrews, and cervids, are naturally infected with *A. phagocytophilum* (Rar and Golovljova 2011). Different genetic variants of *A. phagocytophilum* circulate in different tick and mammalian species (Bown et al. 2008), and these strains may differ in their pathogenicity to humans (Doudier et al. 2010). The distribution of *A. phagocytophilum* is worldwide (Rar and Golovljova 2011).

Wild rodents and shrews are important reservoir hosts for many tick-borne pathogens, and therefore they have a key role in the natural circulation of tick-borne infections, including *A. phagocytophilum* and *B. microti* (Goethert and Telford 2003, Bown et al. 2008, Bown et al. 2011). In the rodent reservoirs, *A. phagocytophilum* infection is transient, lasting for 1–2 months, whereas *B. microti* infection is chronic (Bown et al. 2003, Bown et al. 2008). The transmission of both *A. phagocytophilum* and *B. microti* is transstadial, i.e., an uninfected tick (larvae or nymph) becomes infected while feeding on an infectious reservoir host that itself had become infected while feeding on an infected tick (nymph or adult) (Gray et al. 2002, Woldehivet 2006, Hunfeld et al. 2008). The primary vectors of *A. phagocytophilum* and *B. microti* are ticks from the genus *Ixodes* (Gray et al. 2002, Woldehivet 2006, Hunfeld et al. 2008).

Here we quantify the presence and prevalence of *A. phagocytophilum* and *B. microti* in Finland in the bank vole (*Myodes glareolus*) and identify the pathogen strains that are circulating in the host population. We find that *A. phagocytophilum* and *B. microti* infection prevalence in bank voles is high (22% and 40%, respectively), and we provide the first evidence that these pathogens are circulating in Finnish rodents.

Materials and Methods

Rodent trapping was carried out in July, 2008, as a part of an ongoing long-term study on bank vole (*M. glareolus*) populations in the Konnevesi area in central Finland (62°83'79" N, 26°82'09" E) (for details of the study sites and trapping procedures, see Kallio et al. 2009). The bank vole dominates the small rodent community in the study sites (our unpublished data). All captured bank voles (63 females and 88 males) were taken to the laboratory where their body mass, head width, and sex were recorded and a blood sample was taken from the retro-orbital sinus with capillary tubes (Haematocrit Capillaries, Hirschmann Laborgeräte, Germany). The presence of ticks was not examined. The samples were placed on dry ice immediately after collecting and stored at –80°C until use.

Presence or absence of pathogens was assessed using PCR-based assays. Briefly, total DNA was extracted from bank vole blood using alkaline digestion (for details, see Bown et al. 2003). Negative controls (one per every four samples) were included in the DNA extraction and in the PCR assays, and none of the negative controls were found positive in the PCR assays. Detection of *A. phagocytophilum* used a nested PCR-assay that targets the 16S rRNA gene (primers in Massung et al. 1998; PCR conditions are described by Bown et al. 2006). The positive control was a DNA extract derived from a strain of *A. phagocytophilum* isolated from a sheep (*Ovis aries*) in the

United Kingdom. To quantify the strain variation, six randomly chosen positive samples were screened using a separate nested PCR that targets a fragment of the *msp4* gene, with thermal cycling conditions and primers described by Bown et al. (2007); amplicons were sequenced. *B. microti* was detected using an Apicomplexa-specific nested PCR that targets a fragment of the 18S rRNA gene (Simpson et al. 2005). The positive control used was *B. microti* Munich strain. Because this method is not *B. microti* specific, all putative positive samples were sequenced to separate *B. microti* from *Hepatozoon* sp. infections. Sequence identity was determined by BLAST search (Altschul et al. 1990) against the National Center for Biotechnology Information (NCBI) Nucleotide database. Only samples that were confirmed to be infected by *B. microti* on the basis of the sequence data were considered as positive results.

We analyzed separately the probability of being infected (binary response variable) with *A. phagocytophilum* and *B. microti* using a generalized linear mixed model (GLMM) approach with a logit link function and binomial distribution. The explanatory variables assessed in the full model were gender, head width, and head width squared and whether the individual was co-infected with the other parasite. Body size is often used as a proxy for age in wild rodents. In central Finland, for example, bank vole head width is approximately on average 8.1–8.4 mm at birth, 11.6–12.0 mm at weaning (~3 weeks age), 12.7 mm and 13.4 mm at the age of ~2 and ~5 months, respectively, and 13.8–14.0 at the age of 10 months (Koskela et al. 1998, Oksanen et al. 2001, Kallio et al. 2006, our unpublished data). Models were fitted using the Laplace approximation method (lmer function in lme4 package in R software, available under GNU license at www.r-project.org). To control for the potential correlation amongst individuals that were captured at the same location, the trapping site was included as a random effect in the models. Starting from the full model, terms were omitted if they did not reduce the Akaike information criterion (AIC) by more than 2 units when included (Burnham and Anderson 2002).

Results

Out of the 151 bank voles captured in July, 2008, 33 individuals (21.9%; 95% confidence interval [CI] 15.3–28.5) were infected with *A. phagocytophilum*, 60 individuals (39.7%; 95% CI 31.9–47.5) were infected with *B. microti*, and 11 individuals (7.3%; 95% CI 3.1–11.4) were infected with both pathogens. The probability of a bank vole being infected with *A. phagocytophilum* peaked at intermediate ages (i.e., head width of ~12.5–13.5 mm, which corresponds approximately the age of 2–5 months; see Materials and Methods) (Fig. 1, Table 1). Neither host sex nor *B. microti* infection status had a significant effect on *A. phagocytophilum* infection probability. *B. microti* infection probability increased with age (increasing head width) in both sexes with males being more likely to be infected than females (Fig. 2, Table 2). *A. phagocytophilum* infection did not influence the probability of being infected with *B. microti*.

Analysis of the *msp4* fragment showed the strain of *A. phagocytophilum* to be identical to that previously reported in field voles in the United Kingdom (GenBank accession no. FJ469653) (Bown et al. 2007). All 60 sequences of 18S rRNA fragment from putative positive *B. microti*-infected animals were identical to the Munich strain of *B. microti* (GenBank accession no. AB071177).

FIG. 1. Predicted probability of being infected with *A. phagocytophilum* in relation to individual's head width (mm).

Discussion

In this study we present the first evidence that *B. microti* and *A. phagocytophilum* pathogens are present in wild rodents in Finland. *A. phagocytophilum* is known to cause tick-borne fever in cattle in Finland (Tuomi 1966), and, recently, a domestic cat was diagnosed with *A. phagocytophilum* infection (Heikkilä et al. 2010). *B. microti*, in turn, has been reported in a single *Ixodes persulcatus* tick in western Finland (Aleksiev et al. 2007). However, this is the first study examining the presence of *A. phagocytophilum* and *B. microti* in wildlife reservoir host species.

The *B. microti* Munich strain found in this study has been suggested not to be pathogenic to humans (Gray 2006, Siński et al. 2006), although it may be transmitted by *I. ricinus* ticks (Welc-Fałęciak et al. 2012), which is the most important tick species in transmitting tick-borne pathogens to humans in Europe (Jaenson et al. 2012a). In the United Kingdom, however, *B. microti* Munich is transmitted by the nidicolous tick *Ixodes trianguliceps*, all life stages of which feed on rodents (Randolph 1995, Bown et al. 2008). Similarly, the *A. phagocytophilum* strain found here is identical to the one found in the field vole (*Microtus agrestis*) in the United Kingdom, which was apparently restricted to an enzootic small mammal–*I. trianguliceps* cycle (Bown et al. 2006, Bown et al. 2008, Bown et al. 2009). Therefore, it is not likely that

FIG. 2. Predicted probability of being infected with *B. microti* in relation to individual's head width (mm) and sex. (Black line, top) male; (grey line, bottom) female.

the *B. microti* or *A. phagocytophilum* strains found here pose any direct risk to humans in Finland. However, this study was carried out in only one location, and only bank voles were examined. Because different strains of *A. phagocytophilum* may exploit different tick and mammal species supporting separate enzootic cycles (Bown et al. 2008), it is possible that other genetic variants are circulating in Finland. Moreover, *B. microti* strain Jena, which is considered to be zoonotic, has been found in neighboring Estonia (Katargina et al. 2011), which may suggest that this strain could be also found in Finland. Hence, further work has to widen the scope of sampling and, especially, focus on questing *Ixodes* ticks, which pose a risk to humans, to determine whether zoonotic *A. phagocytophilum* or *B. microti* strains exist in Finland.

Even if the *A. phagocytophilum* and *B. microti* found in bank voles do not pose a direct risk to humans in Finland, these pathogens may interact with other pathogens, within their rodent hosts, because these pathogens have significant interactions with other infections in field voles in the United Kingdom (Telfer et al. 2010). For instance, *A. phagocytophilum* infection increases cowpox virus and decreases *Bartonella* sp. and *B. microti* infection probabilities, whereas *B. microti* decreases the risk of *Bartonella* sp. infection while increasing the

TABLE 1. BANK VOLE INDIVIDUAL'S LIKELIHOOD OF BEING *A. PHAGOCYTOPHILUM* INFECTED (IN LOGIT SCALE) IN RELATION TO ITS HEAD WIDTH

Source of variation	Estimate (SE)	z value	P value
Intercept	−165.464 (85.30)	−1.932	0.053
Head width	24.450 (13.011)	1.956	0.051
Head width squared	−0.984 (0.494)	−1.994	0.046
σ^2	0.637		

σ^2 is the variance attributable to random effect.
sd, standard deviation of σ^2 ; SE, standard error.

TABLE 2. BANK VOLE INDIVIDUAL'S LIKELIHOOD OF BEING *B. MICROTI* INFECTED (IN LOGIT SCALE) IN RELATION TO ITS SEX AND HEAD WIDTH

Source of variation	Estimate (SE)	z value	P value
Intercept	−19.667 (4.181)	−4.703	<0.001
Head width	1.370 (0.308)	4.443	<0.001
Sex Male	1.373 (0.414)	3.316	<0.001
σ^2	0.225		
sd	0.474		

σ^2 , the variance attributable to random effect.
sd, standard deviation of σ^2 ; SE, standard error.

risk of *A. phagocytophilum* infection. These strong individual level effects are likely to be translated to infection dynamics at the population level (Telfer et al. 2010), and for zoonotic pathogens may be further reflected in human epidemics.

We do not know yet which tick species is responsible for the circulation of these pathogens in our study area, where both *I. ricinus* and *I. trianguliceps* are found (our unpublished data). In the United Kingdom, the same strains of *A. phagocytophilum* and *B. microti* are transmitted primarily by *I. trianguliceps* (Bown et al. 2008). The distribution of *I. trianguliceps* covers most of Finland (Ulmanen 1972), thus the pathogens transmitted by *I. trianguliceps* might also be found across Finland. Moreover, *I. trianguliceps* infests many small mammal species, such as bank voles (*Myodes glareolus*), field voles (*Microtus agrestis*), root voles (*Microtus oeconomus*), common shrews (*Sorex araneus*), and pygmy shrews (*S. minutus*) (Ulmanen 1972, Nilsson 1974, Randolph 1975, Bown et al. 2003, Bown et al. 2011), many of which are competent reservoir hosts species for a variety of tick-borne pathogens, including *A. phagocytophilum* and *B. microti* (Randolph 1995, Liz et al. 2000, Bown et al. 2003, Bown et al. 2006). Therefore, it is likely that *A. phagocytophilum* and *B. microti* are widely distributed in small mammal species in Finland.

The infection prevalences reported here are in line with earlier studies on *A. phagocytophilum* and *B. microti* in rodents from elsewhere in Europe (e.g., Liz 2002, Duh et al. 2003, Siński et al. 2006, Bown et al. 2008, Bown et al. 2011, Beck et al. 2011, Rar et al. 2011). The peak in *A. phagocytophilum* infection likelihood at intermediate individual age (i.e., head width) is likely to reflect the transient nature of this infection (Bown et al. 2003), whereas the constantly increasing likelihood of *B. microti* infection with age reflects a chronic infection (Bown et al. 2008). Although both pathogens are likely to be transmitted by the same vector species, only *B. microti* showed some difference in infection probability between sexes (more common among males).

Conclusion

To summarize, *A. phagocytophilum* and *B. microti* are prevalent in bank voles in central Finland. Because they are transmitted by *I. trianguliceps* ticks (Bown et al. 2008), whose distribution range covers most of Finland (Ulmanen 1972), we predict that these pathogens might be widespread in Finland. Although it is unlikely that *B. microti* or *A. phagocytophilum* strains found here cause any direct risk to humans in Finland, these pathogens may interact with other pathogens (Telfer et al. 2010) and therefore affect infection dynamics of, for example, zoonotic pathogens. Therefore, further studies are needed to examine the role of these *B. microti* and *A. phagocytophilum* strains in the pathogen community of Finnish rodents and to examine the presence of other, potentially zoonotic strains in Finland.

Acknowledgments

This research adhered to the Association for the Study of Animal Behaviour/Animal Behaviour Society Guidelines for the Use of Animals in Research, the legal requirements in Finland, and all institutional guidelines. We wish to thank H. Helle for help with fieldwork and sampling. This study was financially supported by Academy of Finland (grant nos. 250524 to E.R.K.; 218107 and 257340 to E.K.; 132190 to T.M.).

We are also grateful for support provided by the Konnevesi Research Station.

Author Disclosure Statement

No competing financial interests exist.

References

- Alekseev AN, Dubinina HV, Jääskeläinen AE, Vapalahti O, et al. First report on tick-borne pathogens and exoskeletal anomalies in *Ixodes persulcatus* Schulze (*Acari: Ixodidae*) collected in Kokkola coastal region, Finland. *Int J Acarol* 2007; 33:253–258.
- Altschul SF, Gish W, Miller W, Myers EW, et al. Basic Local Alignment Search Tool. *J Mol Biol* 1990; 215:403–410.
- Beck R, Vojta L, Curkovic S, Mrljak V, et al. Molecular survey of *Babesia microti* in wild rodents in central Croatia. *Vector Borne Zoonotic Dis* 2011; 11:81–83.
- Bown KJ, Begon M, Bennett M, Woldehiwet Z, et al. Seasonal dynamics of *Anaplasma phagocytophilum* in a rodent-tick (*Ixodes trianguliceps*) system, United Kingdom. *Emerg Infect Dis* 2003; 9:63–70.
- Bown KJ, Begon M, Bennett M, Birtles RJ, et al. Sympatric *Ixodes trianguliceps* and *Ixodes ricinus* ticks feeding on field voles (*Microtus agrestis*): Potential for increased risk of *Anaplasma phagocytophilum* in the United Kingdom? *Vector Borne Zoonotic Dis* 2006; 6:404–410.
- Bown KJ, Lambin X, Ogden NH, Petrovec M, et al. High-resolution genetic fingerprinting of European strains of *Anaplasma phagocytophilum* by use of multilocus variable-number tandem-repeat analysis. *J Clin Microbiol* 2007; 45:1771–1776.
- Bown KJ, Lambin X, Telford GR, Ogden NH, et al. Relative importance of *Ixodes ricinus* and *Ixodes trianguliceps* as vectors for *Anaplasma phagocytophilum* and *Babesia microti* in field vole (*Microtus agrestis*) populations. *Appl Environ Microbiol* 2008; 74:7118–7125.
- Bown KJ, Lambin X, Ogden NH, Begon M, et al. Delineating *Anaplasma phagocytophilum* ecotypes in coexisting, discrete enzootic cycles. *Emerg Infect Dis* 2009; 15:1948–1954.
- Bown KJ, Lambin X, Telford G, Heyder-Bruckner D, et al. The common shrew (*Sorex araneus*): A neglected host of tick-borne infections? *Vector Borne Zoonotic Dis* 2011; 11:947–953.
- Burnham KP, Anderson DR. *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*. Springer, 2002.
- Doudier B, Olano J, Parola P, Brouqui P. Factors contributing to emergence of *Ehrlichia* and *Anaplasma* spp. as human pathogens. *Veterinary Parasitol* 2010; 167:149–154.
- Duh D, Petrovec M, Trilar T, Avsic-Zupanc T. The molecular evidence of *Babesia microti* infection in small mammals collected in Slovenia. *Parasitology* 2003; 126:113–117.
- Dumler JS, Choi KY, Garcia-Garcia JC, Barat NS, et al. Human granulocytic anaplasmosis and *Anaplasma phagocytophilum*. *Emerg Infect Dis* 2005; 12:1828–1834.
- Edouard S, Koebel C, Goehring F, Socolovschi C, et al. Emergence of human granulocytic anaplasmosis in France. *Ticks Tick Borne Dis* 2012; 3:403–405.
- Goethert HK, Telford SR 3rd. What is *Babesia microti*? *Parasitology* 2003; 127:301–309.
- Gray JS. Identity of the causal agents of human babesiosis in Europe. *Int J Med Microbiol* 2006; S1:131–136.
- Gray JS, von Stedingk LV, Gürtelschmid M, Granström M. Transmission studies of *Babesia microti* in *Ixodes ricinus* ticks and gerbils. *J Clin Microbiol* 2002; 40:1259–1263.

- Gray JS, Zintl A, Hildebrandt A, Hunfeld KP, et al. Zoonotic babesiosis: Overview of the disease and novel aspects of pathogen identity. *Ticks Tick Borne Dis* 2010; 1:3–10.
- Heikkilä HM, Bondarenko A, Mihalkov A, Pfister K, et al. *Anaplasma phagocytophilum* infection in a domestic cat in Finland: Case report. *Acta Vet Scand* 2010; 52:62.
- Hildebrandt A, Hunfeld KP, Baier M, Krumbholz A, et al. First confirmed autochthonous case of human *Babesia microti* infection in Europe. *Eur J Clin Microbiol Infect Dis* 2007; 26:595–601.
- Hildebrandt A, Fritsch J, Franke J, Sachse S, Dorn W, Straube E. Co-circulation of emerging tick-borne pathogens in Middle Germany. *Vector Borne Zoonotic Dis* 2011; 11: 533–537.
- Hunfeld KP, Lambert A, Kampen H, Albert S, et al. Seroprevalence of *Babesia* infections in humans exposed to ticks in midwestern Germany. *J Clin Microbiol* 2002; 40:2431–2436.
- Hunfeld KP, Hildebrandt A, Gray JS. Babesiosis: Recent insights into an ancient disease. *Int J Parasitol* 2008; 38:1219–1237.
- Jaenson TG, Jaenson DG, Eisen L, Peterson E, et al. Changes in the geographic distribution and abundance of tick *Ixodes ricinus* during the past 30 years in Sweden. *Parasit Vectors* 2012a; 5:8.
- Jaenson TG, Hjertqvist M, Bergstrom T, Lundkvist A. Why is tick-borne encephalitis increasing? A review of the key factors causing the increasing incidence of human TBE in Sweden. *Parasit Vectors* 2012b; 5:184.
- Kallio ER, Poikonen A, Vaheri A, Vapalahti T, et al. Maternal antibodies postpone hantavirus infection and enhance individual breeding success. *Proc R Soc B* 2006; 273:2771–2776.
- Kallio ER, Begon M, Henttonen H, Koskela E, et al. Cyclic hantavirus epidemics in humans—Predicted by rodent host dynamics. *Epidemics* 2009; 1:101–107.
- Katargina O, Geller J, Vasilenko V, Kuznetsova T, et al. Detection and characterization of *Babesia* species in *Ixodes* ticks in Estonia. *Vector Borne Zoonotic Dis* 2011; 11:923–928.
- Koskela E, Jonsson P, Hartikainen T, Mappes T. Maternal antibodies postpone hantavirus infection and enhance individual breeding success. *Proc R Soc B* 1998; 265:1129–1134.
- Liz JS. Ehrlichiosis in *Ixodes ricinus* and wild mammals. *Int J Med Microbiol* 2002; 291(Suppl 33):104–105.
- Liz JS, Andres L, Sumner JW, Massung RF, et al. PCR detection of granulocytic Ehrlichiae in *Ixodes ricinus* ticks and wild small mammals in western Switzerland. *J Clin Microbiol* 2000; 38:1002–1007.
- Massung R, Slater K, Owens J, Nicholson W, et al. Nested PCR assay for detection of granulocytic Ehrlichiae. *J Clin Microbiol* 1998; 36:1090–1095.
- Medlock JM, Hansford KM, Bormane A, Derdakova M, et al. Driving forces for changes in geographical distribution of *Ixodes ricinus* ticks in Europe. *Parasit Vectors* 2013; 6:1.
- Nilsson A. Host relations and population changes of *Ixodes trianguliceps* (Acari) in northern Scandinavia. *Oikos* 1974; 25: 315–320.
- Oksanen T, Jonsson P, Koskela E, Mappes T. Optimal allocation of reproductive effort: Manipulation of offspring number and size in the bank vole. *Proc R Soc B* 2001; 268:661–666.
- Pieniazek N, Sawczuk M, Skotarczak B. Molecular identification of *Babesia* parasites isolated from *Ixodes ricinus* ticks collected in northwestern Poland. *J Parasitol* 2006; 92:32–35.
- Randolph SE. Patterns of distribution of the tick *Ixodes trianguliceps* Birula on its hosts. *J Anim Ecol* 1975; 44:451–474.
- Randolph SE. Quantifying parameters in the transmission of *Babesia microti* by the tick *Ixodes trianguliceps* amongst voles (*Clethrionomys glareolus*). *Parasitology* 1995; 110:287–295.
- Rar V, Golovljova I. *Anaplasma*, *Ehrlichia*, and “*Candidatus* Neoehrlichia” bacteria: Pathogenicity, biodiversity, and molecular genetic characteristics, a review. *Infect Genet Evol* 2011; 11:1842–1861.
- Rar VA, Epikhina TI, Livanova NN, Panov VV. Genetic diversity of *Babesia* in *Ixodes persulcatus* and small mammals from North Ural and West Siberia, Russia. *Parasitology* 2011; 138:175–182.
- Rizzoli A, Haufler H, Carpi G, Vourc HG, et al. Lyme borreliosis in Europe. *Euro Surveill* 2011; 16:19906.
- Simpson VR, Panciera RJ, Hargreaves J, McGarry JW, et al. Myocarditis and myositis due to infection with *Hepatozoon* species in pine martens (*Martes martes*) in Scotland. *Vet Rec* 2005; 156:442–446.
- Siński E, Bajer A, Welc R, Pawelczyk A, et al. *Babesia microti*: Prevalence in wild rodents and *Ixodes ricinus* ticks from the Mazury Lakes District of North-Eastern Poland. *Int J Med Microbiol* 2006; 296(Suppl 40):137–143.
- Sumilo D, Asokliene L, Avsic-Zupanc T, Bormane A, et al. Behavioural responses to perceived risk of tick-borne encephalitis: Vaccination and avoidance in the Baltics and Slovenia. *Vaccine* 2008; 26:2580–2588.
- Telfer S, Lambin X, Birtles R, Beldomenico P, et al. Species interactions in a parasite community drive infection risk in a wildlife population. *Science* 2010; 330:243–246.
- Tuomi J. Studies in epidemiology of bovine tick-borne fever in Finland and a clinical description of field cases. *Ann Med Exp Biol Fenn* 1966; 44(Suppl 6):1–62.
- Ulmann I. Distribution and ecology of *Ixodes trianguliceps* Birula (Acarine, Ixodidae) in Finland. *Ann Zool Fennici* 1972; 9:111–115.
- Welc-Fałęciak R, Bajer A, Paziawska-Harris A, Baumann-Popczyk A, et al. Diversity of *Babesia* in *Ixodes ricinus* ticks in Poland. *Adv Medic Sci* 2012; 57:364–369.
- Woldehivet Z. *Anaplasma phagocytophilum* in ruminants in Europe. *Ann NY Acad Sci* 2006; 1078:446–460.

Address correspondence to:

Eva R. Kallio

Department of Biological and Environmental Sciences

University of Jyväskylä

P.O.Box 35 (YAC)

FIN-40014 University of Jyväskylä

Jyväskylä

Finland

E-mail: eva.kallio@ju.fi