Molecular and Serological Study of Rickettsial Infection in Humans, and in Wild and Farm Animals, in the Province of Burgos, Spain

Lourdes Lledó,¹ Gerardo Domínguez-Peñafiel,² Consuelo Giménez-Pardo,¹ Isabel Gegúndez,¹ Rosario González,¹ and José Vicente Saz¹

Abstract

Limited information is available on the presence of rickettsial infection in humans and animal reservoirs in Spain. Exposure to spotted fever group rickettsia in healthy humans and in farm and wild animals in the Province of Burgos, Spain, was examined by serological methods. Rickettsial DNA was also sought by PCR in animal samples. Of 102 human serum samples examined by indirect immunofluorescence assays (IFA), 5.88% were positive for antibodies against *Rickettsia conorii* (titers 1/128–1/512). Significant differences were detected in human seroprevalence with respect to age. In further IFAs, 102 out of 375 (27.2%) serum samples from the wild animals reacted with *R. conorii* antigens (titers 1/64–1/1024); 32 out of 281 (11.38%) samples from farm animals were also positive for *R. conorii* (titers 1/64–1/2048). The prevalence detected among total wild animals was significantly higher than among total farm animals. No rickettsial DNA was found by PCR in any farm or wild animal sample.

Key Words: Epidemiology—Rickettsia—Vector-borne—Reservoir host.

Introduction

IN RECENT YEARS, EUROPE HAS EXPERIENCED an increase in the incidence of certain zoonoses, particularly those transmitted by arthropod vectors, such as rickettsiosis (Blanco and Oteo 2006). The number of methods available for detecting and identifying rickettsia has increased and allowed the detection of new rickettsial pathogens as well as already-known types in new locations. Increasing interest in rickettsial diseases now demands that the epidemiology of their causal agents be reassessed (Richards 2012, Santibáñez et al. 2013).

Recent epidemiological studies undertaken in Spain have detected emerging species of spotted fever group *Rickettsiae* (SFGR) such as *R. slovaca*, *R. massiliae*, *R. raoultii*, *R. felis*, and *R. monacensis* in arthropods (Márquez et al. 2006, Márquez 2008, Lledó et al. 2010). Seroprevalence studies in humans and animals have detected antibodies to *R. slovaca* and *R. felis* as well as reemerging *R. typhi* and *R. conorii*; indeed, some clinical cases by emerging species (like *R. sibirica mongolitimonae*) have also been reported by serological or molecular methods (Bernabeu-Wittel et al. 2006, Lledó et al. 2006, Amusatogui et al. 2008, Antón et al. 2008, Nogueras et al. 2013, Ramos et al 2013).

Mediterranean spotted fever (MSF), caused by *R. conorii*, is an endemic rickettsiosis of southern and eastern Europe and parts of Africa and Asia (Brouqui et al. 2007). It is mainly transmitted by the tick *Rhipicephalus sanguineus*. Most clinical cases are therefore noted in rural areas. Our ecological and epidemiological comprehension of MSF has experienced important advances in the last 10 years as the disease has emerged and reemerged in different countries (Brouqui et al. 2007, Rovey and Raoult 2008). However, many questions remain unanswered, including the identities of all the vectors and reservoirs of *R. conorii*, and whether other risk factors exist for the onset of the severe form of MSF (Rovey and Raoult 2008, Schex et al. 2011). Although MSF is regarded as a historically endemic disease in Spain, with the highest incidence usually seen in the summer months and mainly in rural areas, new epidemiological studies are needed to update our knowledge of its distribution and epidemiology.

The aim of the present study was to determine the prevalence of antibodies to SFGR in humans and animals in a region of the Province of Burgos in northern Spain, where human activities commonly involve continued contact with animals. No previous studies have been performed on the distribution and prevalence of *R. conorii* infection in this area.

¹Departamento de Biomedicina y Biotecnología, Universidad de Alcalá, Alcalá de Henares, Spain.

²Consejería de Sanidad y Bienestar Social de la Junta de Castilla y León, Burgos, Spain.

Materials and Methods

Study area

This study was performed in the Merindades area of the Province of Burgos in northern Spain (central point 42°55'52"N, 3°29'2"W). Here, mean daily summer temperatures range between 16°C and 20°C, whereas those for winter range between 2°C and 5°C. Rainfall ranges from 900 to 1100 mm/year and is usually high in winter. The area is mainly rural, but recreational activities attracting nonresidents have increased in recent years.

Human serum samples

A total of 102 samples of human serum (male/female ratio 83.3/16.7%; age 55 ± 11.4 year, range 24–88 years) were collected from visitors to health centers run by the Spanish

National Institute of Health over 2006–2007. (Each subject had either attended a health care center for a reason unrelated to infectious disease and were volunteers.) All serum samples were maintained at -20°C until analysis. The following information was recorded for each donor: Age, sex, occupation, area of residence, travel history, and history of contact with animals and arthropod bites. All subjects gave their informed consent to be included in the study in compliance with the ethical standards of the Human Experimentation Committee of the University of Alcalá de Henares and the Helsinki Declaration of 1964 (as revised in 2004).

Animal serum samples

Blood, serum, and tissue (lung, kidney, liver, and brain [the latter only for rodents]) samples from 375 wild animals belonging to 19 species (Table 1) were collected between June,

TABLE 1. WILD ANIMALS STUDIED: PREVALENCE RESULTS FOR SERUM, BLOOD, AND TISSUE SAMPLES

Species examined	Serum samples studied/ prevalence	Blood samples studied/ positives	Tissue samples studied/ positives
<i>Arvicola terrestris</i> (Water vole)	2/0%	—	—
<i>Apodemus flavicollis</i> (Yellow-necked mouse)	9/33.3%	—	9/0
<i>Apodemus sylvaticus</i> (Wood mouse)	24/33.3%	—	24/0
<i>Canis lupus</i> (Wolf)	3/0%	2/0	—
<i>Genetta genetta</i> (Genet)	2/50%	1/0	—
<i>Lepus europaeus</i> (European hare)	16/18.7%	12/0	1/0
<i>Martes foina</i> (Beech marten)	2/0%	1/0	2/0
<i>Martes martes</i> (Pine marten)	2/0%	—	1/0
<i>Meles meles</i> (Badger)	7/28.57%	2/0	2/0
<i>Mus musculus</i> (House mouse)	22/4.5%	22/0	—
<i>Mustela erminea</i> (Stoat)	1/0%	—	—
<i>Mustela nivalis</i> (Least weasel)	2/0%	—	1/0
<i>Mustela putorius</i> (Polecat)	2/0%	—	1/0
<i>Myodes glareolus</i> (Bank vole)	5/20%	—	6/0
<i>Neovison vison</i> (American mink)	121/54.5%	—	25/0
<i>Sciurus vulgaris</i> (Red squirrel)	4/25%	2/0	3/0
<i>Sus scrofa</i> (European wild boar)	103/14.5%	38/0	1/0
<i>Talpa occidentalis</i> (Iberian mole)	3/33.3%	1/0	3/0
<i>Vulpes vulpes</i> (Fox)	45/0%	3/0	3/0
Total	375/27.2%	84/0	82/0

TABLE 2. FARM ANIMALS STUDIED: PREVALENCE RESULTS FOR SERUM AND BLOOD SAMPLES

Mammal species	Serum samples studied/prevalence	Blood samples studied/positive
Cattle	105/21.9%	31/0
Horses	23/8.7%	9/0
Pigs	50/14%	—
Sheep	103/0%	31/0
Total	281/11.38%	71/0

2006, and September, 2007. Wild animals were captured alive or found dead (victims of road accidents). Blood and serum samples from farm animals (cattle, horses, pigs, and sheep, all in extensive or semiextensive farming systems) were collected over 2006 and 2007 (Table 2). All samples were stored at -20°C until analyses. Permission to take and study these animal samples was obtained from the regional government of Castilla y León in compliance with current legislation and in keeping with the ethical guidelines of the Committee on Animal Experimentation of the University of Alcalá de Henares.

Serology

Sera were tested by the indirect immunofluorescence assay (IFA), as described by Phillip et al. (1976). The rickettsia providing the antigen was *R. conorii* (strain Malish 7). Rickettsia were propagated in Vero E6 cells (ATCC CRL 1586) and fixed on spot slides. The fluorescein-labeled conjugate used was adapted according to the species studied. All antibodies (immunoglobulin G [IgG]) were from Sigma (St. Louis, MO) (Table 3). Positive and negative control sera were included (provided by Dr. Fatima Alves, Centro de Estudos de Vectores e Doenças Infecciosas, Portugal-CEVDI). Titers of $\geq 1/128$ were considered positive for humans, and $\geq 1/64$ were considered positive for animals.

DNA extraction and PCR

Total DNA was extracted from wild animal blood and tissues and from farm animal blood using the COBAS AmpliPrep® system (Roche Diagnostics, GmbH, Mannheim, Germany) according to the manufacturer's recommendations. DNA of *Rickettsia* genera was detected by PCR-

amplification of: (1) The citrate synthase gene *gltA*, using primers RpCs 1258 and RpCs 877, which amplify a 381-bp fragment; (2) the gene coding for protein 190-kDa *OmpA* using the primers Rr 190.70p and Rr 190.602n, which amplify a 532-bp fragment; and (3) the gene *ompB*, using the primers 120-M59' and 120-807, which amplify a 833-bp fragment. The first two amplifications were performed following the protocol of Regnery et al. (1991); the third was performed following that of Roux and Raoult (2000). To prevent DNA contamination and the carryover of amplified products, sterile tools were used at all times in each step of the analysis (DNA extraction, preparation of the reaction mixture, and amplification and analysis of the PCR products). Each step was performed in separate work areas. A negative control (Milli-Q water) was included in all amplifications.

Data analyses

The chi-squared or Fisher exact test was used to compare differences in proportions employing two-way tables with a StatView® software in Apple™ format. Significance was set at $p \leq 0.05$.

Results

IgG antibodies to *R. conorii* were found in six human serum samples (total 5.8%; females 5.8%, males 5.8%). The age of the seropositive subjects ranged from 68 to 82 years (76 ± 4.8 years). No significant differences in seroprevalence were seen with respect to sex, but significantly more positive results were recorded among the those over 60 years ($p \leq 0.001$; $\chi^2 = 7.35$). The IgG antibody titers for seropositive subjects ranged between 1/128 and 1/512 (Table 4). All seropositive human subjects had contact with animals; three had occupational risks. One of the subjects referred to a history of arthropod bites.

A total of 102 sera with antibodies specific to *R. conorii* were detected among the wild animals (prevalence 27.2%), while 32 were detected among the farm animals (prevalence 11.47%; $p \leq 0.001$ compared to the wild animals; $\chi^2 = 24.7$). Titers for these positive animals ranged from 1/64 to 1/2048 (Table 4). Among the wild animals, mink showed the highest prevalence ($p \leq 0.001$ compared to the remaining species; $\chi^2 = 67.46$) (Table 1). Among the farm animals, cattle showed the highest prevalence ($P \leq 0.001$ compared to the remaining farm species; $\chi^2 = 18.37$) (Table 2).

No rickettsial DNA was detected by PCR in any animal sample.

Discussion

Defining the vector species of a disease is of the utmost importance for its control, yet in some cases it remains unclear which rickettsial species are carried by which arthropod vectors, nor is it certain which mammalian species serve as their reservoirs (Schex et al. 2011). In many cases, the natural epidemiological cycle is well known, but in recent years there have been changes in these classic epidemiological cycles (Lledó et al. 2003). After many years of falling prevalence, there would now seem to be a reemergence of rickettsial disease in Spain, perhaps associated with changes in recreational habits that bring people into closer contact with the reservoir hosts and arthropod vectors.

The present serological results confirm the presence of antibodies to *R. conorii* in humans and most of the animal

TABLE 3. THE FLUORESCCEIN-LABELED CONJUGATE USED ACCORDING TO HUMAN AND ANIMAL SPECIES STUDIED

Human and animals	Fluorescein-labeled conjugate
Human	Rabbit anti-human IgG
Canidae	Rabbit anti-dog IgG
Felidae and Mustelidae	Goat anti-cat IgG
Rat, squirrel, and mole	Rabbit anti-rat IgG
Mouses	Rabbit anti-mouse IgG
Leporidae	Anti-rabbit IgG
Suidae	Anti-pig IgG
Horses	Goat anti-horse IgG
Cattle	Goat anti-bovine IgG
Sheep	Rabbit anti-sheep IgG

IgG, immunoglobulin G.

TABLE 4. SERUM TITERS FOR ANTI-*R. CONORII* ANTIBODIES (IGG)

Groups	Titer 1/64	Titer 1/128	Titer 1/256	Titer 1/512	Titer 1/1024	Titer 1/2048
Human samples		2 (1.9%)	3 (2.9%)	1 (0.9%)	—	—
Wild animal samples	62 (16.5%)	24 (6.4%)	8 (2.1%)	4 (1%)	1 (0.2%)	—
Farm animal samples	3 (1%)	4 (1.4%)	3 (1%)	7 (2.5%)	9 (3.2%)	6 (2.3%)

IgG, immunoglobulin G.

species examined in the study area. However, because of serological cross-reactivity between rickettsiae of the spotted fevers group (Santibañez et al. 2006), we consider the results as preliminary, while another assay can be used to validate IFA results by future studies using more antigens and serological methods that are more specific, such as western blot. Therefore, it is possible that some of the seropositive cases observed in the present study were (or had been) infected with SFGR rickettsiae other than *R. conorii*.

Seroprevalence in humans (5.8%) was much lower than that recorded in the 1980s for the Spanish provinces of Seville (26.3%) (García and Najera 1984) and Salamanca (73.5%) (Herrero et al. 1989), but is reminiscent of more recent figures reported for the provinces of Soria (5%) (Saz et al. 1993), León (1%) (Rojo-Vázquez 1997), and the Spanish northwest (8%) (Segura-Porta et al. 1998) and south (8.7%) (Bernabeu-Wittel et al. 2006). Compared to Portugal (7.6%) (Bacellar et al. 1991), Croatia (43.7%) (Punda-Polic et al. 2003), Greece (7.9%) (Daniel et al. 2002), southern France (18%) (Raoult et al. 1987), Slovakia (32%) (Sekeyova et al. 2012), and Israel (10%) (Harrus et al. 2007), the present figure is low.

The present seroprevalence results were also similar to those reported in the above studies with respect to age, sex, and contact with animals and ticks. In the present work, seroprevalence was significantly higher in those over 60 years old, as reported for the Spanish province of Albacete (Bartolomé et al. 2005). Several clinical studies (Bartolomé et al. 2005, Aliaga et al. 2009, Baltadzhiev et al. 2012) have reported local reemergence of the disease in Spain and Bulgaria; adults over 40 years of age have been those most affected, with a peak seen in patients over 60 years, many of whom have suffered severe illness. The present study, and those mentioned above, all seem to suggest that MSF continues to be most common among people living or working in rural areas, as suggested by Segura-Porta et al. (1998) and Podsiadly et al. (2010), although other authors have found higher prevalence figures in urban and suburban areas than in semirural areas (Raoult et al. 1987, Bolaños-Riveros et al. 2011).

Of the 19 species of wild animals studied, 11 had members seropositive for *R. conorii*. In general, the prevalences detected in the wild species were significantly higher than those recorded for farm species, as reported by other authors (Jilintai et al. 2008). The prevalence of seropositive wild animals ranged from 4.5% for the house mouse *Mus musculus* to 54.5% for the American mink (*Neovison vison*). (This is the first study to report antibodies to *R. conorii* in *N. vison* in Spain.) American mink found their way into the wild in Europe after escaping from captivity. They are known to approach farms, and they may therefore play an important role in both the maintenance of *R. conorii* in nature and its transmission to farm animals.

All of the wild small mammal species examined had antibodies to *R. conorii*, with the wood mouse *Apodemus sylvaticus* returning the highest seroprevalence. Even in places as distant as Japan, the prevalence of antibodies against SFGR in mice (mainly wood mice) is high (Morita et al. 1990, Okabayashi et al. 1996). In California too, a seroprevalence for SFGR of 14% has been reported for rodents (Adjemian et al. 2008). However, in a recent molecular study conducted in the Basque Country (northern Spain), all of the small mammals examined were negative for SFGR (Barandika et al. 2007), although in Germany SFGR DNA was detected in seven rodents out of 124 studied (Schex et al. 2011).

Some authors report other wild animals may play a host reservoir role for SFGR, such as wild boar (*Sus scrofa*) (Ortuño et al. 2007, Selmi et al. 2009), rabbit (Ruiz-Beltran et al. 1992), deer (Inokuma et al. 2008, Jilintai et al. 2008), and jackal (Waner et al. 1999). In the present work, antibodies were detected in wild boar, European hare, and several carnivorous animals. However, unlike that reported by Boretta et al. (2009), no infections were detected in foxes.

The domestic animals most studied to date have been pet dogs and cats; indeed, the dog is the main domestic animal reservoir of *R. conorii* worldwide. In the present study, however, farm animals were studied (which in rural areas have much contact with humans), which may maintain tick populations and serve as pathogen reservoirs (Ruiz-Fons et al. 2006). As a group, the farm animals had higher titers than humans or wild animals. The seroprevalence detected in cattle (21.9%) was comparable to that reported for seroprevalence to SFGR from Croatia (27.3%) (Punda-Polic et al. 1995) but higher than that observed in Kenya (16.3%), Sri Lanka (15%) (Kováčová et al. 1996), and Japan (9.6%) (Jilintai et al. 2008). In a study performed in Salamanca (Spain), a seroprevalence figure of 78.1% was reported for cattle (Herrero et al. 1989). The seroprevalence recorded in the literature for *R. conorii* in horses is very variable, ranging from 0% in Sicily (Torina et al. 2007) to 100% in the Spanish province of Salamanca (Herrero et al. 1989) (although here only three horses were studied). For other SFGR species outside Europe, high prevalences have been reported in Brazil (Horta et al. 2004) and Panama (65%) (Bermúdez et al. 2011).

In the present work, a seroprevalence of 0% was recorded for sheep. This was somewhat unexpected because other authors have reported values of 38.9% in Salamanca (Herrero et al. 1989), 9.4% in Croatia (Punda-Polic et al. 1995), and 15% in Kenya (Mutai et al. 2013) for antibodies against *R. conorii* and other SFGR species. However, in Sicily, Torina et al. (2007) also found no seropositive sheep.

The literature contains little information on seroprevalence in pigs. In the studies that have been performed, the number

of specimens analyzed has been very low, and the results were highly variable (Herrero et al. 1989, Torina et al. 2007). Studies on larger numbers of animals are required to examine the role of this animal in SFGR infection. In the present study, a prevalence of 14% was recorded.

Finally, PCR detected no *Rickettsia* spp. DNA in any of the animal samples tested, even though in some animal species the prevalence of antibodies to *R. conorii* and antibody titers, were high. Other authors have reported similar situations, finding no bacterial DNA in tissue or blood samples even though the donor animals had high antibody titers (Boretti et al. 2009, Ortuño et al. 2012).

Conclusions

In summary, the present results expand our knowledge regarding the seroprevalence of SFGR in the Province of Burgos and shed light on the epidemiology of tick-borne rickettsial disease. Some infections of SFGR have a re-emergence of cases involving serious disease. This may due to a change in natural epidemiological cycles, with other wild and peridomestic reservoirs becoming available. The large number of seropositive animals (wild and farm) detected suggests that human contact with animals is a risk factor for the acquisition of *R. conorii* infection or other species of SFGR. Further research should attempt to determine the role of different species studied in the epidemiology and ecology of these infections and involve other areas of the country.

Author Disclosure Statement

No competing financial interests exist.

References

- Adjemian JZ, Adjemian MK, Foley P, Chomel BB, et al. Evidence of multiple zoonotic agents in a wild rodent community in the eastern Sierra Nevada. *J Wild Dis* 2008; 44:737–742.
- Aliaga L, Sánchez-Blázquez P, Rodríguez-Granger J, Sampedro A, et al. Mediterranean spotted fever with encephalitis. *J Med Microbiol* 2009; 58:521–525.
- Amusatogui I, Tesouro MA, Kakoma I, Sainz A. Serological reactivity to *Ehrlichia canis*, *Anaplasma phagocytophilum*, *Neorickettsia risticii*, *Borrelia burgdorferi* and *Rickettsia conorii* in dogs from northwestern Spain. *Vector Borne Zoonotic Dis* 2008; 8:797–803.
- Antón E, Nogueras MM, Pons I, Font B, et al. *Rickettsia slovaca* infection in humans in the northeast of Spain: Seroprevalence study. *Vector Borne Zoonotic Dis* 2008; 8:689–694.
- Bacellar F, Nuncio MS, Reháček J, Filipe AR. Rickettsiae and rickettsioses in Portugal. *Eur J Epidemiol* 1991; 7:291–293.
- Baltadzhiev IG, Popivanova NI, Stoilova YM, Kevorkian AK. Mediterranean spotted fever classification by disease course and criteria for determining the disease severity. *Folia Med (Plovdiv)* 2012; 54:53–61.
- Barandika JF, Hurtado A, García-Esteban C, Gil H, et al. Tick-borne zoonotic bacteria in wild and domestic small mammals in northern Spain. *Appl Environ Microbiol* 2007; 73:6166–6171.
- Bartolomé J, Lorente S, Hernández-Pérez N, Martínez-Alfaro E, et al. Estudio clínico y epidemiológico de las rickettsiosis del Grupo de las Fiebres Manchadas en Albacete, España. *Enferm Infecc Microbiol Clin* 2005; 23:194–196.
- Bermúdez CS, Zaldivar AY, Spolidorio MG, Moraes-Filho J, et al. Rickettsial infection in domestic mammals and their ectoparasites in El Valle de Antón, Coclé, Panamá. *Vet Parasitol* 2011; 177:134–138.
- Bernabeu-Wittel M, del Toro MD, Nogueras MM, Muniain MA, et al. Seroepidemiological study of *Rickettsia felis*, *Rickettsia typhi*, and *Rickettsia conorii* infection among the population of southern Spain. *Eur J Clin Microbiol Infect Dis* 2006; 25:375–381.
- Blanco JR, Oteo JA. Rickettsiosis in Europe. *Ann NY Acad Sci* 2006; 1078:26–36.
- Boretti FS, Perreteri A, Meli ML, Cattori V, et al. Molecular Investigations of *Rickettsia helvetica* infections in dogs, foxes, humans, and *Ixodes* ticks. *Appl Environ Microbiol* 2009; 75:3230–3237.
- Brouqui P, Parola P, Fournier PE, Raoult D. Spotted fever rickettsioses in southern and eastern Europe. *FEMS Immunol Med Microbiol* 2007; 49:2–12.
- Daniel SA, Manika K, Arvanmdou M, Antoniadis A. prevalence of *Rickettsia conorii* and *Rickettsia typhi* infections in the population of northern Greece. *Am J Trop Med Hyg* 2002; 66:76–79.
- García A, Nájera E. Estudio epidemiológico de las rickettsiosis en la provincia de Sevilla, basado en las reacciones serológicas de inmunofluorescencia indirecta. *Rev Sanid Hig Pública* 1984; 58:83–98.
- Harrus S, Lioe Y, Perros M, Grisaru-Spen G, et al. *Rickettsia conorii* in humans and dogs: A seroepidemiologic survey of two rural villages in Israel. *Am J Trop Med Hyg* 2007; 77:133–135.
- Herrero J, Ruiz-Beltrán R, Martín AM, García EJ. Mediterranean spotted fever in Salamanca, Spain. Epidemiological study in patients and serosurvey in animals and healthy human population. *Acta Trop* 1989; 46:335–350.
- Horta MC, Labruna MB, Sangioni LA, Vianna MC, et al. Prevalence of antibodies to spotted fever group rickettsiae in humans and domestic animals in a Brazilian spotted fever-endemic area in the state of São Paulo, Brazil: Serologic evidence for infection by *Rickettsia rickettsii* and another spotted fever group Rickettsia. *Am J Trop Med Hyg* 2004; 71:93–97.
- Inokuma H, Seino N, Suzuki M, Kaji K, et al. Detection of *Rickettsia helvetica* DNA from peripheral blood of sika deer (*Cervus nippon yezoensis*) in Japan. *J Wild Dis* 2008; 44:164–167.
- Jilintai, Seino N, Matsumoto K, Hayakawa D, et al. Serological and molecular survey of Rickettsial infection in cattle and sika deer in a pastureland in Hidaka district, Hokkaido, Japan. *Jpn J Infect Dis* 2008; 61:315–317.
- Kováčová E, Sixl W, Stünzner D, Urvölgyi J, et al. Serological examination of human and animal sera from six countries of three continents for the presence of rickettsial antibodies. *Eur J Epidemiol* 1996; 12:85–89.
- Lledó L, Gegúndez MI, Serrano JL, Saz JV, et al. A seroepidemiological study of *Rickettsia typhi* infection in dogs from Soria province, central Spain. *Ann Trop Med Parasitol* 2003; 97:861–864.
- Lledó L, Gegúndez MI, Fernandes N, Sousa R, et al. The seroprevalence of human infection with *Rickettsia slovaca*, in an area of northern Spain. *Ann Trop Med Parasitol* 2006; 100:337–343.
- Lledó L, Giménez-Pardo C, Domínguez-Peñafiel G, Sousa R, et al. Molecular detection of hemoprotozoa and Rickettsia species in arthropods collected from wild animals in the Burgos Province, Spain. *Vector Borne Zoonotic Dis* 2010; 10:735–738.

- Márquez FJ. Spotted fever group *Rickettsia* in ticks from southeastern Spain natural parks. *Exp Appl Acarol* 2008; 45:185–194.
- Márquez FJ, Muniaín MA, Rodríguez-Liebana JJ, Toro MD, et al. Incidence and distribution pattern of *Rickettsia felis* in peridomestic fleas from Andalusia, Southeast Spain. *Ann NY Acad Sci* 2006; 1078:344–346.
- Morita C, Yamamoto S, Tsuchiya K, Yoshida Y, et al. Prevalence of spotted fever group rickettsia antibody in *Apodemus speciosus* captured in an endemic focus in Miyazaki Prefecture, Japan. *Jpn J Med Sci Biol* 1990; 43:15–18.
- Mutai BK, Wainaina JM, Magiri CG, Nganga JK, et al. Zoonotic surveillance for rickettsiae in domestic animals in Kenya. *Vector Borne Zoonotic Dis* 2013; 13:360–366.
- Nogueras MM, Pons I, Pla J, Ortuño A, et al. The role of dogs in the eco-epidemiology of *Rickettsia typhi*, etiological agent of Murine typhus. *Vet Microbiol* 2013; 12:97–102.
- Okabayashi T, Miura N, Miyazaki S, Nigo T, et al. Epidemiological survey of small rodents for spotted fever rickettsial antibody in Hokkaido, Japan. *Jpn J Med Sci Biol* 1996; 49:63–68.
- Ortuño A, Quesada M, López-Claessens S, Castellá J, et al. The role of wild boar (*Sus crofa*) in the eco-epidemiology of *R. slovaca* in northeastern Spain. *Vector Borne Zoonotic Dis* 2007; 7:59–64.
- Ortuño A, Pons I, Quesada M, Lario S, et al. Evaluation of the presence of *Rickettsia slovaca* infection in domestic ruminants in Catalonia, Northeastern Spain 2012; 12:1019–1022.
- Philip RN, Casper EA, Ormsbee RA, Peacock MG, et al. Microimmunofluorescence test for the serological study of Rocky Mountain spotted fever and typhus. *J Clin Microbiol* 1976; 3:51–61.
- Podsiadły E, Chmielewski T, Karbowiak G, Kędra E, et al. The occurrence of spotted fever rickettsioses and other tick-borne infections in forest workers in Poland. *Vector Borne Zoonotic Dis* 2011; 11:985–989.
- Punda-Polic V, Poljak S, Bubic A, Bradaric N, et al. Antibodies to spotted fever group rickettsiae and *Coxiella burnetii* among domestic animals in southern Croatia. *Acta Microbiol Immunol Hung* 1995; 42:339–344.
- Punda-Polic V, Klismanic Z, Capkun V. Prevalence of antibodies to spotted fever group rickettsiae in the region of Split (southern Croatia). *European J Epidemiol* 2003; 18:451–455.
- Ramos JM, Jado I, Padilla S, Masiá M, et al. Human infection with *Rickettsia sibirica mongolitimonae*, Spain. *Emerg Infect Dis* 2013; 19:267–269.
- Raoult D, Toga B, Chaudet H, Chiche-Portiche C. Rickettsial antibody in southern France: antibodies to *Rickettsia conorii* and *Coxiella burnetii* among urban, sub-urban and semi-rural blood donors. *Trans R Soc Trop Med Hyg* 1987; 81:80–81.
- Regnery RL, Spruill CL, Plykaitis BD. Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J Bacteriol* 1991; 173:1576–1589.
- Richards AL. Worldwide detection and identification of new and old rickettsiae and rickettsial diseases. *FEMS Immunol Med Microbiol* 2012; 64:107–110.
- Rojo-Vázquez J. Seroprevalencia de las infecciones causadas por *Borrelia burgdorferi* y *Rickettsia conorii* en humanos y perros en el área primaria de salud de San Andreas de Rabanedo, León, España. *Rev Esp Salud Pública* 1997; 71:173–180.
- Roux V, Raoult D. Phylogenetic analysis of members of the genus *Rickettsia* using the gene encoding the outer-membrane protein rOmpB (ompB). *Int J Syst Evol Microbiol* 2000; 50:1449–1455.
- Rovero C, Raoult D. Mediterranean spotted fever. *Infect Dis Clin North Am* 2008; 22:515–530.
- Ruiz-Beltrán R, Herrero JJ, Martín AM, Criado LA. Role of *Lagomorpha* in the wild cycle of *Rickettsia conorii* in Salamanca (Spain). *Eur J Epidemiol* 1992; 8:136–139.
- Ruiz-Fons F, Fernández de Mera IG, Acevedo P, Höfle L, et al. *Ixodes* ticks parasitizing Iberian red deer (*Cervus elaphus hispanicus*) and European wild boar (*Sus scrofa*) from Spain: Geographical and temporal distribution. *Vet Parasitol* 2006; 140:133–142.
- Santibáñez S, Ibarra V, Portillo A, Blanco JR, et al. Evaluation of IgG antibody response against *Rickettsia conorii* and *Rickettsia slovaca* in patients with DEBONEL/TIBOLA. *Ann NY Acad Sci* 2006; 1078:570–572.
- Santibáñez S, Portillo A, Santibáñez P, Palomar AM, et al. Usefulness of rickettsial PCR assays for the molecular diagnosis of human rickettsioses. *Enferm Infecc Microbiol Clin* 2013; 3:283–288.
- Saz JV, Bacellar F, Merino FJ, Filipe A. Seroprevalencia de la infección por *Coxiella burnetii* y *Rickettsia conorii* en la provincia de Soria. *Enferm Infecc Microbiol Clin* 1993; 11:469–473.
- Schex S, Dobler G, Riehm J, Müller J, et al. *Rickettsia spp.* in wild small mammals in lower Babaria, south-eastern Germany. *Vector Borne Zoonotic Dis* 2011; 11:493–502.
- Segura-Porta F, Diestre G, Ortuño A, Sanfeliu I, et al. Prevalence of antibodies to spotted fever group rickettsiae in human beings and dogs from an endemic area of Mediterranean Spotted Fever in Catalonia, Spain. *Eur J Epidemiol* 1998; 14:395–398.
- Sekeyova Z, Subramanian G, Mediannikov O, Diaz MQ, et al. Evaluation of clinical specimens for *Rickettsia*, *Bartonella*, *Borrelia*, *Coxiella*, *Anaplasma*, *Francisella* and *Diplorickettsia* positivity using serological and molecular biology methods. *FEMS Immunol Med Microbiol* 2012; 64:82–91.
- Selmi M, Martello E, Bertolotti L, Bisanzio D, et al. *Rickettsia slovaca* and *Rickettsia raoultii* in *Dermacentor marginatus* ticks collected on wild boars in Tuscany, Italy. *J Med Entomol* 2009; 46:1490–1493.
- Torina A, Vicente J, Alongi A, Scimeca S, et al. Observed prevalence of tick-borne pathogens in domestic animals in Sicily, Italy during 2003–2005. *Zoon Pub Health* 2007; 54:8–15.
- Waner T, Baneth G, Strenger C, Keysary A, et al. Antibodies reactive with *Ehrlichia canis*, *Ehrlichia phagocytophila* genogroup antigens and the spotted fever group rickettsial antigens, in free-ranging jackals (*Canis aureus s yriacus*) from Israel. *Vet Parasitol* 1999; 82:121–128.

Address correspondence to:

Lourdes Lledó

Departamento de Biomedicina y Biotecnología

Universidad de Alcalá

Ctra. Madrid-Barcelona

km 33.6, 28871 Alcalá de Henares

Madrid

Spain

E-mail: lourdes.lledo@uah.es