Published in final edited form as:

Spine J. 2014 September 1; 14(9): 2000–2007. doi:10.1016/j.spinee.2013.11.052.

INSOMNIA IN CHRONIC DISABLING MUSCULOSKELETAL PAIN DISORDERS IS INDEPENDENT OF PAIN AND DEPRESSION

Sali Asih, M.S*, Randy Neblett, M.A., L.P.C., BCB*, Tom G. Mayer, M.D**, Emily Brede, Ph.D., RN*, and Robert J. Gatchel, Ph.D., ABPP***

*PRIDE Research Foundation, 5701 Maple Ave. #100, Dallas, TX 75235

**Department of Orthopedic Surgery, University of Texas Southwestern Medical Center, 5323 Harry Hines Blvd., Dallas, TX 75235

***Department of Psychology, College of Science. The University of Texas at Arlington, 313 Life Science Building, Arlington, Texas 76019

Abstract

BACKGROUND CONTEXT—Insomnia is frequently experienced by patients suffering from chronic musculoskeletal disorders, but is often seen as simply a symptom of pain or depression, and not as an independent disorder. Compared to those who experience only chronic pain, patients with both chronic pain and insomnia report higher pain intensity, more depressive symptoms, and greater distress. However, insomnia has not yet been systematically studied in a chronic musculoskeletal pain with disability population.

PURPOSES—This study assessed the prevalence and severity of patient-reported insomnia, as well as the relationship among insomnia, pain intensity, and depressive symptoms, in a chronic musculoskeletal pain with disability population.

STUDY DESIGN/SETTING—A retrospective study of prospectively captured data.

PATIENT SAMPLE—A consecutive cohort of 326 chronic musculoskeletal pain with disability patients (85% with spinal injuries) entered a functional restoration treatment program. All patients signed a consent form to participate in this protocol.

OUTCOME MEASURES—Insomnia was assessed with the Insomnia Severity Index (ISI), a validated patient-report measure of insomnia symptoms. Four patient groups were formed: No Clinically Significant Insomnia (score 0–7); Sub-Threshold Insomnia (score 8–14); Moderate Clinical Insomnia (score 15–21); and Severe Clinical Insomnia (score 22–28). Three patterns of sleep disturbance were also evaluated: Early, Middle, and Late Insomnia. Additional validated

© 2013 Elsevier Inc. All rights reserved.

Corresponding Author: Tom G. Mayer, M.D., 5701 Maple Ave. #100, Dallas, TX 75235, Phone: (214) 351-6600, Fax: (214) 351-3026, tgmayer@pridedallas.com.

The Abstract of this study was accepted as a podium presentation at the 27th Annual Meeting of NASS in Dallas, TX

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

psychosocial patient-report data were collected, including the Pain Visual Analog Scale (PVAS), the Beck Depression Inventory (BDI), and the Pain Disability Questionnaire (PDQ).

METHODS—Patients completed a standard psychosocial assessment battery upon admission to the functional restoration program. The program included a quantitatively-directed exercise process in conjunction with a multimodal disability management approach. The four insomnia groups were compared on demographic and psychosocial variables. The shared variances among insomnia, depression, and pain were determined by partial correlational analyses. The writing of this article was supported in part by Grant 1K05 MH 71892 from NIH, focusing on evidence-based assessment and treatment approaches to musculoskeletal pain and the monitoring of valid outcomes. None of the authors involved in this study had a conflict of interest.

RESULTS—The presence of No Clinically Significant Insomnia, Sub-threshold Insomnia, Moderate Clinical Insomnia and Severe Clinical Insomnia was found in 5.5%, 21.2%, 39.6%, and 33.7% of the cohort, respectively. More than 70% of patients reported moderate to severe insomnia symptoms, which is a considerably higher prevalence than found in most patient cohorts studied previously. A step-wise pattern was found, in which Severe Clinical Insomnia patients reported the highest pain, the most severe depressive symptoms, and the greatest disability. The Severe Clinical Insomnia patients also reported a higher number of sleep disturbance types (Early, Middle, and Late insomnia) than the other 3 groups. In fact, 62.9% of them reported all 3 disturbance types. Although correlations were found between insomnia and depressive symptoms, and between insomnia and pain, the shared variances were small (12.9% and 3.6%, respectively), indicating that depression and pain are separate constructs from insomnia.

CONCLUSION—This research indicates that insomnia is a significant and pervasive problem in a chronic musculoskeletal pain with disability population. Most importantly, although insomnia has traditionally been assumed to be simply a symptom of pain or depression, the findings of the present study reveal that it is a construct relatively independent from both pain and depression. Specific insomnia assessment and treatment is therefore recommended for this chronic musculoskeletal pain with disability population.

INTRODUCTION

Insomnia is highly prevalent in the general population, ranging from 19% to 27% of adults over the age of 18 [1–4]. Insomnia includes difficulty initiating sleep, disrupted sleep, and early morning awakening [5]. People with insomnia report greater health problems, more limited physical activity, more interference with daily living, and greater emotional difficulty compared to healthy people without insomnia [2, 6]. Poor sleep quality and insufficient sleep is associated with decreased pain thresholds in normal, healthy subjects [6, 7], and hyperalgesia in chronic pain subjects [8–10]. Moreover, insomnia is frequently reported by patients with chronic pain, with about 88% reporting at least one type of sleep disturbance [8, 11]. Many studies have found that increases in insomnia severity are significantly correlated with increases in pain intensity, using both subjective measures, such as sleep diaries and patient-report instruments [8, 12–13], and objective measures, such as polysomnography, which uses EEG recordings [12, 14–15], and actigraphy, which measures the frequency of movements during sleep [13, 16]. Furthermore, the combination of insomnia and chronic pain has been found to be associated with a variety of negative outcomes,

including anxiety, depression, and suicidal ideation [7–9, 17–18]. A cycle of sleep deprivation and increased pain, due to hyperalgesia and/or decreased pain thresholds, can severely limit the daily functioning of chronic pain patients [19, 20].

Insomnia is a symptom of major depressive disorder, according to the DSM-IV-TR classification system [21], and is a significant predictor of people developing depression, based on longitudinal studies [22]. When insomnia occurs comorbidly with depression, it is commonly mis-diagnosed as simply a symptom of the depressive disorder, and not recognized as a separate disorder [5, 23]. A similar phenomenon has also been observed in relation to pain and insomnia. Pain causes insomnia, which in turn, causes increased pain intensity, resulting in more disrupted sleep. Due to the high prevalence of insomnia in chronic pain populations, insomnia is often seen as a secondary symptom of chronic pain [24]. This view of insomnia may be especially common in assessing and treating chronic pain patients who have both depression and insomnia [3]. However, this view has shifted gradually as new evidence has emerged, pointing toward insomnia as a primary disorder for many chronic pain patients [18, 25].

Although insomnia has been studied in diverse chronic pain populations, it has not been studied in chronic musculoskeletal pain with disability patients undergoing functional restoration treatment. This patient population typically exhibits severe physical and psychosocial deterioration, often accompanying a workers' compensation claim. This patient population is highly associated with various psychiatric disorders, including pain disorder, major depressive disorder, and opioid dependence, that often have post-injury onset [26]. Based on the findings that other pain conditions are associated with insomnia, it is highly probable that insomnia also plays a role in the course of chronic musculoskeletal pain with disability. However, prior research has not made a distinction between patients with high- versus low-frequency or severity of insomnia symptoms, with only a few studies categorizing participants based on whether they had symptoms severe enough and frequent enough to be classified as having "clinical insomnia" [9, 27]. In addition, little prior research has differentiated between participants with moderate to severe insomnia, sub-threshold insomnia, and minimal insomnia [18]. Thus, in order to more systematically address this issue, the present study was designed to evaluate patient-reported insomnia, including prevalence, pattern (early, middle, and late insomnia), and severity, and its association with patient-reported pain intensity, depressive symptoms, and perceived disability.

In addition, the present study was intended to identify differences in pain, depression, and disability among patients with different levels of insomnia severity. It was hypothesized that insomnia would be highly prevalent in chronic musculoskeletal pain with disability patients, and that associations would be found between insomnia and other psychosocial distress measures. Identifying the role of insomnia in this population may lead to a greater recognition of sleep disturbances and better treatments for patients suffering from both chronic musculoskeletal pain with disability and insomnia.

METHODS

Participants

A retrospective study of prospectively captured data that consisted of a consecutive cohort of 326 chronic musculoskeletal pain with disability patients were admitted to a functional restoration (FR) program at a regional rehabilitation center. All patients signed a consent form to participate in this protocol. The majority of patients had spinal disorders (85%), of which most were compensable under state or federal workers' compensation. The inclusion criteria for participation in the treatment program were: (1) at least four months of disability post-injury; (2) primary and/or secondary care was unsuccessful in improving pain and function; (3) the patient failed to improve despite previous surgical treatment or was not a surgical candidate; and (4) severe pain and functional limitations remained. The participants in this study formed a consecutive cohort that was admitted between June 2010 and December 2011. Of the initial sample of 420 patients, 94 patients were excluded because they did not complete all of the items on the Insomnia Severity Index (ISI), the primary sleep measure in this study, leaving 326 patients with complete data for subsequent analyses. Patients who did, and those who did not complete the ISI, were not significantly different in age, gender proportion, or length of disability. Figure 1 depicts the flow of patients through the study.

Measures

The ISI is a brief patient-report measure that assesses the severity of both night-time and daytime components of insomnia, using a 5-point Likert scale (0 = "Not at all" to 4 = "Extremely"), generating a total score ranging from 0 to 28 [27]. A cut-off score of 15 has been shown to have good sensitivity (94%) and specificity (94%) in identifying clinical insomnia [28]. Four groups were formed based on the ISI total score collected upon admission to the treatment program: No Clinically Significant Insomnia (ISI score: 0–7, n=18), Sub-threshold Insomnia (ISI score: 8–14, n=69), Moderate Clinical Insomnia (ISI score: 15–21, n=129), and Severe Clinical Insomnia (ISI score: 22–28, n=110) [27]. In addition, the four groups were evaluated for three patterns of sleep disturbance: Early (difficulty initiating sleep), Middle (difficulty staying sleep) and Late Insomnia (early morning awakening). The ISI asks subjects to rate the level of problems for each insomnia type (from 0 = "None" to 4 = "Very Severe"). A score of 3 (indicating a severe disturbance) was chosen to indicate the presence of early, middle, or late insomnia [29]. Demographic comparisons among the No Clinically Significant Insomnia, Sub-threshold Insomnia, the Moderate Clinical Insomnia and the Severe Clinical Insomnia showed no significant differences in age, gender, ethnicity, length of disability, or pre-treatment surgery (see Table 1).

In addition to the ISI, patient-report measures evaluating pain intensity, depressive symptoms, and disability, were also administered. The Pain Visual Analog Scale (PVAS) was used to measure pain intensity, with scores ranging from 0–10 [30]. The Beck Depression Inventory (BDI) [31] was used to assess depressive symptoms, with scores ranging from 0–63. The Pain Disability Questionnaire (PDQ, with scores ranging from 0–150) [32], and the Oswestry Disability Index (ODI, with scores ranging from 0–100) [33],

were used to evaluate perceived disability. For all measures, a higher score indicates more severe symptoms, and all measures have been shown to be reliable and valid for use in chronic pain populations.

Procedure

All patients participated in an interdisciplinary FR treatment program, consisting of a quantitatively-directed exercise progression, supervised by physical and occupational therapists, and aimed at restoring muscular strength, flexibility, endurance, and cardiovascular fitness. The exercise program was administered in conjunction with multimodal disability management, which included cognitive behavioral therapy, biofeedback, stress-management training, educational classes, vocational reintegration, and future fitness management [34–36]. All patients received a comprehensive evaluation prior to the start of the program, which included a physical examination, a medical history, a medical case management/disability assessment, a quantitative functional capacity evaluation, and a psychosocial intake interview. Demographic data and socioeconomic information, such as occupational characteristics and work status, were collected as part of the intake interview. Patients completed a battery of patient-report instruments upon admission to the FR program, including the ISI, PVAS, BDI, ODI, and the PDQ. Because all assessments were part of the standard medical record for patients participating in the interdisciplinary FR program, the study was granted an exemption from review by the Institutional Review Board of the University of Texas at Arlington.

Statistical Methods

An *a priori* power analysis was conducted prior to data collection to determine an appropriate sample size [37, 38]. In order to detect a medium effect size of 0.5 on the basis of Cohen's f [39], with power $\beta = 0.8$ and $\alpha = 0.05$, a total of 180 subjects was required. Thus, this study had a more than sufficient sample size ($n = 326$). The insomnia groups were compared on demographic and psychosocial variables using one-way ANOVAs for continuous variables, and chi-square tests for categorical variables. Finally, the shared variances of insomnia, depression, and pain were determined by partial correlation analyses.

The writing of this article was supported in part by Grant 1K05 MH 71892 from the National Institutes of Health, which focuses on evidence-based assessment and treatment approaches to musculoskeletal pain and the monitoring of valid outcomes. None of the authors involved in this study had a conflict of interest.

RESULTS

The Pattern of Insomnia and Psychosocial Patient-Report Measures

Of the total sample, 36.4% of patients reported severe problems with all 3 types of sleep disturbance patterns (Early, Middle, and Late Insomnia); 40.5% reported severe problems with 2 types; and 23.1% reported severe problems with a single sleep disturbance. Of those patients who reported severe problems with two or more sleep disturbance patterns, 59.0% reported difficulty falling asleep, 91.6% reported difficulty staying asleep, and 49.4% reported early-morning waking. Patients reporting a severe problem with a single sleep

pattern most frequently reported difficulty staying asleep (41.5%). The patterns of severe sleep disturbance for each of the insomnia groups are presented in Figure 2. The prevalence of patients in the Sub-threshold Insomnia group who reported severe problems with a single sleep disturbance, and two sleep disturbance patterns, was 77.3% and 22.7% respectively. The prevalence of patients in the Moderate Clinical Insomnia group who reported a single severe sleep disturbance, two severe sleep disturbances, and three severe sleep disturbance patterns, was 38.8%, 47.2%, and 14%, respectively. The prevalence of patients in the Severe Clinical Insomnia group who reported two severe sleep disturbances and three severe sleep disturbances was 37.1% and 62.9%. No patients in the No Clinically Significant Insomnia group reported a severe sleep disturbance pattern.

The psychosocial patient-report measures for the insomnia groups are presented in Table 2. As can be seen, the psychosocial measures were significantly different among the four insomnia groups ($p < .001$) with a step-wise pattern. The patients in the No Clinically Significant Insomnia group reported the lowest level of pain intensity, depressive symptoms, and pain-related disability. In contrast, the Severe Clinical Insomnia patients reported the highest level of pain intensity, depressive symptoms, and pain-related disability. The psychosocial self-report outcomes for Sub-threshold Insomnia and Moderate Clinical Insomnia groups lay in between, with poorer outcomes associated with the more severe group. Further, the Severe Clinical Insomnia group had the highest percentage of patients with moderate to severe depressive symptoms on the BDI (54.1%), and severe pain-related disability on the PDQ (92.7%), compared to other insomnia groups.

The Relationship among Insomnia, Pain, and Depression

The correlations among insomnia, pain, and depression are presented in Table 3. Insomnia was moderately correlated with pain ($r = .29, p < .001$). After controlling for depression, the correlation between pain and insomnia was weak ($r = .19, p = .001$) with a shared variance of only 3.6%. The correlation between depressive symptoms and insomnia was moderate ($r = .41, p < .001$). After controlling for pain, the correlation between insomnia and depressive symptoms was moderate ($r = .36, p < .001$) with a shared variance of 12.9%. The small proportion of shared variance between insomnia and pain, and between insomnia and depressive symptoms, indicated that insomnia is only a very small part of pain and depression constructs.

DISCUSSION

The purpose of this present study was to evaluate the prevalence and severity of insomnia in a chronic musculoskeletal pain with disability population. Of 326 patients assessed, only 6% reported no clinically significant insomnia symptoms. Further, 73% met criteria for Clinical Insomnia (moderate to severe symptoms), and 21% were classified as having Sub-threshold Insomnia. The prevalence of clinical insomnia in this population is therefore extremely high, and is more than double the rate of insomnia found in the general population [1–4]. The mean ISI score for those patients classified with Clinical Insomnia (21.2) was slightly higher in this sample of chronic musculoskeletal pain with disability patients than mean scores reported for a separate cohort of insomniacs with chronic low back pain (19.1) [9]. In

addition, the presence of clinical insomnia, particularly severe clinical insomnia, was associated with greater patient-reported psychosocial distress, including pain, depressive symptoms, and perceived disability.

This study also evaluated the relationship among insomnia, pain, and depressive symptoms in much greater detail. As noted previously, insomnia is often viewed as simply a symptom of depression, and not as an independent disorder [23]. Indeed, we did find that insomnia was significantly related to depressive symptoms. However, the shared variance between insomnia and depressive symptoms was relatively small (12.9%). Insomnia was also significantly correlated with pain intensity, although the shared variance between insomnia and pain was negligible (3.6%). Taken together, these results clearly demonstrate that insomnia can occur independently of depressive symptoms and pain within a chronic musculoskeletal pain with disability population. These findings support previous research results suggesting that insomnia, pain, and depressive symptoms are related yet independent constructs. For example, Wilson and colleagues [40] found that 40% of subjects in a chronic musculoskeletal pain population had insomnia without comorbid major depression, 27% had major depression with comorbid insomnia, and 33% did not have either insomnia or depression. A study of psychiatric patients with major depressive disorder found that insomnia was a significant predictor of pain, even after controlling for depression. As noted by Wickwire and Smith [5], insomnia can originate from a painful medical condition, but in time, may evolve into its own disorder, related to, but separate from, the precipitating condition. Dysfunctional sleep and sleep habits can become primary factors that perpetuate and maintain the insomnia, so that sleep disturbance is no longer directly caused by pain intensity [41]. When this happens, insomnia can evolve from a secondary symptom of chronic pain to a primary, comorbid condition, with similar characteristics to primary insomnia [42]. This appears to be the case with the chronic musculoskeletal pain with disability patients in the present study.

In analyzing types of sleep disturbance patterns (Early, Middle, and Late Insomnia), 36.4% of the patients reported severe problems with all 3 types, 40.5% reported severe problems with 2 types, and 23.1% reported severe problems with a single type of sleep disturbance. Severe problems with all 3 sleep disturbance patterns was reported only by patients meeting criteria for clinical insomnia (moderate and severe) patients. The majority of Sub-threshold Insomnia subjects (77.3%) reported severe problems with only 1 sleep disturbance. Compared to Late Insomnia, the prevalence of severe problems with Early and Middle Insomnia combined was somewhat higher in this population. High rates of severe problems with Early Insomnia may be related to the tendency of chronic musculoskeletal pain with disability patients who are not working to spend the majority of their day reclining or napping, which may interfere with their normal sleep patterns. Education about proper sleep hygiene may be beneficial for these patients with Early Insomnia.

Some limitations pertaining to this study were identified. A relatively large percentage of patients failed to complete all items on the ISI, so those data were unavailable for analysis. As a result, there is a possibility that a selection bias may have occurred. Another limitation was the absence of a control group due to the chosen study design. It should also be noted that other sleep disorders, such as sleep apnea, can also affect the prevalence and severity of

patient-reported insomnia in chronic musculoskeletal pain with disability patients. Unfortunately, information on other sleep disorders was not included in the standard medical record. In addition, data on the onset (pre-injury or post-injury) or duration of insomnia were not available in the present study. Furthermore, pain levels in this study were assessed only during treatment intake and not while patients were experiencing insomnia. Therefore, future clinical research studies should examine other characteristics of sleep disturbance, and assess pain ratings, and other psychosocial measures, at specific times when patients are unable to sleep, in order to provide even more insight into the relationships among insomnia, pain, and depression in chronic musculoskeletal pain with disability patients.

In conclusion, we found a high prevalence of clinical insomnia within a chronic musculoskeletal pain with disability population, and chronic musculoskeletal pain with disability was almost universally associated with some level of insomnia. Clinical Insomnia was associated with increased patient-reported pain, depressive symptoms, and perceived disability. Although Clinical Insomnia was associated with depressive symptoms and pain, we found a significant amount of variance that was not related to these two variables, indicating that Clinical Insomnia can be a distinct and independent disorder within this population. The significant clinical implications of these findings are that, in order to maximize treatment responsiveness, the specific evaluation and treatment of clinical insomnia within a chronic musculoskeletal pain with disability population appears warranted.

Acknowledgments

The writing of this article was supported in part by Grant 1K05 MH 71892 from the National Institutes of Health, which focuses on evidence-based assessment and treatment approaches to musculoskeletal pain and the monitoring of valid outcomes.

References

1. Kim K, Uchiyama M, Okawa M, Liu X, Ogihara R. An epidemiological study of insomnia among the Japanese general population. *Sleep*. 2000; 23(1):41–7. [PubMed: 10678464]
2. Léger D, Scheuermaier K, Philip P, Paillard M, Guilleminault C. SF-36: evaluation of quality of life in severe and mild insomniacs compared with good sleepers. *Psychosom Med*. 2001; 63(1):49–54. [PubMed: 11211064]
3. National Sleep Foundation. Adult Sleep Habits and Styles [Internet]. Arlington: National Sleep Foundation; 2011. cited 2012 November 19. Available from: <http://www.sleepfoundation.org/article/sleep-america-polls/2005-adult-sleep-habits-and-styles>
4. Ohayon MM. Prevalence and Correlates of Nonrestorative Sleep Complaints. *Arch Intern Med*. 2005; 165(1):35–41. [PubMed: 15642872]
5. Wickwire, EM.; Smith, MT. Insomnia in chronic pain. In: Sateia, MJ.; Buysse, DJ., editors. *Insomnia: Diagnosis and Treatment*. London: Informa Healthcare; 2010. p. 139–52.
6. Zammit GK, Weiner J, Daamto N, Sillup GP, McMillan CA. Quality of life in people with insomnia. *Sleep*. 1999; 22(Suppl 2):S379–45. [PubMed: 10394611]
7. Smith MT, Haythornthwaite JA. How do sleep disturbance and chronic pain inter-relate? insights from the longitudinal and cognitive-behavioral clinical trials literature. *Sleep Med Rev*. 2004; 8(2): 119–32. [PubMed: 15033151]
8. McCracken LM, Iverson GL. Disrupted sleep patterns and daily functioning in patients with chronic pain. *Pain Res Manag*. 2002; 7(2):75–9. [PubMed: 12185371]

9. Tang NKY, Wright KJ, Salkovskis PM. Prevalence and correlates of clinical insomnia co-occurring with chronic back pain. *J Sleep Res.* 2007; 16(1):85–95. [PubMed: 17309767]
10. Marty M, Rozenberg S, Duplan B, Thomas P, Duquesnoy B, Allaert F. Quality of sleep in patients with chronic low back pain: a case-control study. *Eur Spine J.* 2008; 17(6):839–44. [PubMed: 18389288]
11. Smith MT, Perlis ML, Smith MS, Giles DE, Carmody TP. Sleep quality and presleep arousal in chronic pain. *J Behav Med.* 2000; 23(1):1–13. [PubMed: 10749008]
12. Atkinson JH, Ancoli-Israel S, Slater MA, Garfin SR, Gillin C. Subjective sleep disturbance in chronic back pain. *Clin J Pain.* 1988; 4(4):225–32.
13. Wilson KG, Watson ST, Currie SR. Daily diary and ambulatory activity monitoring of sleep in patients with insomnia associated with chronic musculoskeletal pain. *Pain.* 1998; 75(1):75–84. [PubMed: 9539676]
14. Rains JC, Penzien DB. Sleep and chronic pain: challenges to the α -EEG sleep pattern as a pain specific sleep anomaly. *J Psychosom Res.* 2003; 54(1):77–83. [PubMed: 12505558]
15. Wittig RM, Zorick FJ, Blumer D, Heilbronn M, Roth T. Disturbed sleep in patients complaining of chronic pain. *J Nerv Ment Dis.* 1982; 170(7):429–31. [PubMed: 7086403]
16. Lavie P, Epstein R, Tzischinsky O, Gilad D, Nahir M, Lorber M, Scharf Y. Actigraphic measurements of sleep in rheumatoid arthritis: comparison of patients with low back pain and healthy controls. *J Rheumatol.* 1992; 19(3):362–5. [PubMed: 1533678]
17. Nicassio PM, Moxham EG, Schuman CE, Gevirtz RN. The contribution of pain, reported sleep quality, and depressive symptoms to fatigue in fibromyalgia. *Pain.* 2002; 100(3):271–9. [PubMed: 12467998]
18. Kelly GA, Blake C, Power C, O’Keeffe D, Fullen B. The association between chronic low back pain and sleep: a systematic review. *Clin J Pain.* 2011; 27(2):169–81. [PubMed: 20842008]
19. Lautenbacher S, Kundermann B, Krieg JC. Sleep deprivation and pain perception. *Sleep Med Rev.* 2006; 10(5):357–69. [PubMed: 16386930]
20. Marin R, Cyhan T, Miklos W. Sleep disturbance in patients with chronic low back pain. *Am J Phys Med Rehabil.* 2006; 85(5):430–5. [PubMed: 16628150]
21. American Psychiatric Association. *Diagnostics and statistical manual of mental disorders IV text revised (DSM-IV TR).* Washington, DC: American Psychiatric Association; 2000.
22. Baglioni C, Battagliese G, Feige B, Spiegelhalder K, Nissen C, Voderholzer U, Lombardo C, Riemann D. Insomnia as a predictor of depression: a meta-analytic evaluation of longitudinal epidemiological studies. *J Affect Disord.* 2011; 135(1):10–9. [PubMed: 21300408]
23. Riemann D, Voderholzer U. Primary insomnia: a risk factor to develop depression? *J Affect Disord.* 2003; 76(1):255–9. [PubMed: 12943956]
24. Léger, D. Sleep and quality of life in insomnia. In: Verster, LC.; Pandi-Perumal, SR.; Streiner, DL., editors. *Sleep and Quality of Life in Clinical Medicine.* Totowa, NJ: Humana Press; 2008. p. 47-52.
25. Chung KF, Tso KC. Relationship between insomnia and pain in major depressive disorder: a sleep diary and actigraphy study. *Sleep Med.* 2010; 11(8):752–8. [PubMed: 20133187]
26. Dersh J, Mayer T, Theodore BR, Polatin P, Gathel RJ. Do psychiatric disorders first appear preinjury or postinjury in chronic disabling occupational spinal disorders? *Spine (Phila Pa 1976).* 2007; 32(9):1045–51. [PubMed: 17450081]
27. Bastien CH, Vallières A, Morin CM. Validation of the insomnia severity index as an outcome measure for insomnia research. *Sleep Med.* 2001; 2(4):297–307. [PubMed: 11438246]
28. Smith S, Trinder J. Detecting insomnia: comparison of four self-report measures of sleep in a young adult population. *J Sleep Res.* 2001; 10(3):229–35. [PubMed: 11696076]
29. Morin CM, Bélanger L, LeBlanc M, Ivers H, Savard J, Espie CA, Mérette C, Baillargeon L, Grégoire JP. The natural history of insomnia: a population-based 3-year longitudinal study. *Arch Intern Med.* 2009; 169(5):447–53. [PubMed: 19273774]
30. McGeary DD, Mayer TG, Gatchel RJ. High pain ratings predict treatment failure in chronic occupational musculoskeletal disorders. *J Bone Joint Surg Am.* 2006; 88(2):317–25. [PubMed: 16452743]

31. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry*. 1961; 4(6):561–71. [PubMed: 13688369]
32. Anagnostis C, Gatchel RJ, Mayer TG. The pain disability questionnaire: a new psychometrically sound measure for chronic musculoskeletal disorders. *Spine (Phila Pa 1976)*. 2004; 29(20):2290–302. [PubMed: 15480144]
33. Fairbank JC, Couper J, Davies JB, O'Brien JP. The oswestry low back pain disability questionnaire. *Physiotherapy*. 1980; 66(8):271–3. [PubMed: 6450426]
34. Mayer T, Gatchel R, Kishino N, Keeley J, Capra P, Mayer H, Barnett J, Mooney V. Objective assessment of spine function following industrial accident: a prospective study with comparison group and one-year follow-up; Volvo Award in Clinical Sciences. *Spine (Phila Pa 1976)*. 1985; 10(6):482–93. [PubMed: 2934829]
35. Mayer TG, Gatchel RJ, Kishino N, Keeley J, Mooney V. A prospective two-year study of functional restoration in industrial low back injury: an objective assessment procedure. *JAMA*. 1987; 258(13):1763–7. [PubMed: 2957520]
36. Mayer, TG.; Gatchel, RJ. *Functional restoration for spinal disorders: The sports medicine approach*. Philadelphia, PA: Lea & Febiger; 1988.
37. Faul F, Erdfelder E, Lang A, Buchner A. G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*. 2007; 05;39(2):175–91.
38. Faul F, Erdfelder E, Buchner A, Lang A. Statistical power analyses using G*Power 3.1: Tests for correlation and regression analyses. *Behav Res Methods*. 2009; 11;41(4):1149–60.
39. Cohen J. A power primer. *Psychol Bull*. 1992; 07;112(1):155–9.
40. Wilson KG, Eriksson MY, D'Eon JL, Mikail S, Emery P. Major depression and insomnia in chronic pain. *Clin J Pain*. 2002; 18(2):77–83. [PubMed: 11882770]
41. Currie SR, Wilson KG, Curran D. Clinical significance and predictors of treatment response to cognitive-behavior therapy for insomnia secondary to chronic pain. *J Behav Med*. 2002; 25(2): 135–53. [PubMed: 11977435]
42. Tang NKY, Goodchild CE, Hester J, Salkovskis PM. Pain-related insomnia versus primary insomnia: a comparison study of sleep pattern, psychological characteristics, and cognitive-behavioral Processes. *Clin J Pain*. 2012; 28(5):428–6. [PubMed: 22179550]

Figure 1.
Flow of participants through the program

Figure 2.**Insomnia Pattern**

*Early, middle or late insomnia would be assigned when participant scored 3 and above in item 1a, 1b or 1c No participant in the No Clinically Significant Insomnia group scored 3 or 4 in either item

Table 1

Demographic variables according to Insomnia Severity Index (N = 326)

Variable	Insomnia Severity Index score			F/ χ^2 value	p-value
	No Clinically Significant Insomnia n = 18 (5.5%)	Sub-Threshold Insomnia n = 69 (21.2%)	Moderate Clinical Insomnia n = 129 (39.6%)	Severe Clinical Insomnia n = 110 (33.7%)	
Gender, n (% male)	12 (66.7%)	46 (66.7%)	87 (67.4%)	75 (68.2%)	.997
Age, mean (SD)	48.0 (10.7)	45.3 (12.8)	45.9 (9.8)	44.6 (10.7)	.602
Area of Injury, n (%)					.270
Cervical only	0%	1 (1.4%)	1 (0.8%)	6 (5.5%)	
Thoracic/Lumbar only	4 (22.2%)	28 (40.6%)	35 (27.1%)	38 (34.5%)	
Extremity only	5 (27.8%)	9 (13.0%)	15 (11.6%)	14 (12.7%)	
Multiple spinal	2 (11.1%)	5 (7.2%)	15 (11.6%)	6 (5.5%)	
Spinal + additional injury	7 (38.9%)	26 (37.7%)	59 (45.7%)	44 (40%)	
Other	0%	0%	4 (3.1%)	2 (1.8%)	
Race, n (%)					.148
Caucasian	12 (70.6%)	42 (61.8%)	68 (54.0%)	69 (63.9%)	
African-American	2 (11.8%)	12 (17.6%)	40 (31.7%)	30 (27.8%)	
Hispanic or Latino	3 (17.6%)	13 (19.1%)	15 (11.9%)	9 (8.3%)	
Other	0%	1 (1.5%)	3 (2.4%)	0%	
Pre-treatment surgeries, n (%)	10 (58.8%)	35 (56.5%)	68 (55.7%)	60 (55.6%)	.995
Length of disability in months, mean (SD)	10.9 (10.4)	28.2 (46.5)	35.1 (54.0)	28.0 (45.1)	.238

Table 2

Patient-reported psychosocial variables (N = 326)

Variable	Insomnia Severity Index score			F	p-value	ES ¹
	No Clinically Significant Insomnia n = 18 (5.5%)	Sub-Threshold Insomnia n = 69 (21.2%)	Moderate Clinical Insomnia n = 129 (39.6%)	Severe Clinical Insomnia n = 110 (33.7%)		
Insomnia Severity Index (ISI), mean (SD) Valid n = 326	3.9 (2.5)	11.6 (2.1)	18.3 (1.8)	24.6 (2.1)	940.53	<.001
Pain Visual Analog Scale (PVAS), mean (SD)	6.2 (2.1)	6.8 (2.1)	7.2 (1.9)	7.9 (1.5)	8.34	<.001
% PVAS 6 (n, %) ² Valid n = 324	15 (83.3%)	56 (82.4%)	105 (81.4%)	103 (94.5%)	9.73	.021
Beck Depression Inventory (BDI), mean (SD)	12.2 (8.4)	13.7 (8.0)	18.8 (9.8)	23.4 (11.8)	15.6	<.001
% BDI 20 (n, %) ² Valid n = 324	2 (11.1%)	14 (20.6%)	52 (40.3%)	59 (54.1%)	26.10	<.001
Pain Disability Questionnaire (PDQ), mean (SD)	85.5 (21.2)	88.7 (25.9)	98.8 (27.2)	109.0 (21.6)	11.51	<.001
% PDQT 71 (n, %) ² Valid n = 321	14 (77.8%)	53 (79.1%)	110 (85.9%)	101 (92.7%)	7.80	.05
Oswestry Disability Index (ODI), mean (SD)	30.7 (12.1)	37.5 (15.9)	41.3 (17.4)	50.1 (17.1)	12.53	<.001
% ODI 41 (n, %) ² Valid n = 312	3 (16.7%)	28 (41.8%)	61 (50.4%)	75 (70.8%)	26.66	<.001

¹ Effect Size (η^2): Small = 0.20, Medium = 0.50, Large = 0.80

Effect size (Cohen W) is defined 0.10 as small, 0.30 as medium, and 0.50 as large

² The cut-off scores were based on: J Bone Joint Surg Am. 2006;88(2):317–25 (PVAS); J Am Acad Orthop Surg. 2008;16(2):107–12 (BDI-II); J Occup Rehabil. 2006;16(1):75–94 (PDQ); Physiotherapy. 1980;66(8):271–3 (ODI).

Table 3

Correlation between Insomnia, Pain and Depression

Variable ¹	Insomnia	Pain	Depressive Symptoms
Insomnia	-	0.19 ²	0.36 ³
Pain		-	0.20 ⁴
Depressive Symptoms			-

¹ Insomnia was assessed with the ISI; Pain Intensity was assessed with the PVAS; Depressive symptoms were assessed with the BDI.

² Correlation after controlling for depressive symptoms; $p = 0.001$

³ Correlation after controlling for pain intensity; $p < 0.001$

⁴ Correlation after controlling for insomnia; $p < 0.001$