Published in final edited form as:

Psychon Bull Rev. 2014 August ; 21(4): 969–975. doi:10.3758/s13423-013-0562-5.

Visual adaptation of the perception of “life”: animacy is a basic perceptual dimension of faces

Kami Koldewyn¹, Patricia Hanus¹, and Benjamin Balas²

¹Department of Brain and Cognitive Science and McGovern Institute for Brain Research, Massachusetts Institute of Technology, Cambridge, MA, USA

²Department of Psychology, North Dakota State University, Fargo, ND, USA

Abstract

One critical component of understanding another’s mind is the perception of “life” in a face. However, little is known about the cognitive and neural mechanisms underlying this perception of animacy. Here, using a visual adaptation paradigm, we ask whether face animacy is (1) a basic dimension of face perception and (2) supported by a common neural mechanism across distinct face categories defined by age and species. Observers rated the perceived animacy of adult human faces before and after adaptation to (1) adult faces, (2) child faces, and (3) dog faces. When testing the perception of animacy in human faces, we found significant adaptation to both adult and child faces, but not dog faces. We did, however, find significant adaptation when morphed dog images and dog adaptors were used. Thus, animacy perception in faces appears to be a basic dimension of face perception that is species-specific, but not constrained by age categories.

1. Introduction

Quick, accurate decisions about which objects in the visual field are alive and capable of action are critical for survival. Visual cues that contribute to perceiving an object as “biological” also carry significant social information. The human visual system is tuned to social objects, including faces (Kanwisher & Yovel, 2006), bodies (Peelen & Downing, 2007) and biological motion (Pelphrey & Carter, 2008). Faces in particular capture and hold our attention (e.g. Langton, Law, Burton, & Schweinberger, 2008; Ro, Russell, & Lavie, 2001), whether they are real people, dynamic movies, veridical representations or schematic drawings. This privileged response to socially relevant stimuli persists even when it is suboptimal. For instance, the addition of eyes to otherwise non-biological objects creates a profound sense of “animacy” that can interfere with task-relevant cues (Gao, McCarthy, & Scholl, 2010). At some point, however, basic detection of a biological agent must yield to an assessment of whether that agent is capable/worthy of social engagement. Distinguishing social from non-social objects is fundamentally important to our ability to function in the social world.

Faces provide salient and informative cues for determining animacy and sociability (Balas & Horksi, 2012; Looser & Wheatley, 2010). Face animacy is perceived categorically - a steep shift in perception results from gradual morphing between real and artificial face images (Looser & Wheatley, 2010). Further, people are more sensitive to appearance changes near the point of subjective equality (the point of maximal animate/inanimate ambiguity) and animacy judgments track with judgments about whether faces have other socially relevant characteristics like having a “mind”, the ability to plan and the ability to engage in social interactions (Gao et al., 2010; Gao & Scholl, 2011; Looser & Wheatley, 2010). These findings suggest two important things: First, the perception of “animacy” is closely tied to the perception of others as socially capable beings. Second, animacy may be a separable dimension of faces – like gender and age. If so, animacy may be a property of faces coded for by a separable neural population.

Face after-effects are an effective way to investigate the processes underlying face perception and recognition (Webster & MacLeod, 2011). High-level visual adaptation has helped characterize many dimensions of faces, including identity (G. Rhodes & Jeffery, 2006), gender (Webster, Kaping, Mizokami, & Duhamel, 2004), and age (O’Neil & Webster, 2011; Schweinberger et al., 2010). These aftereffects are thought to primarily reflect neural changes at face-specific levels of visual processing since adaptation effects transfer across changes in image position, size and orientation (Leopold, O’Toole, Vetter, & Blanz, 2001; G. Rhodes, Jeffery, Watson, Clifford, & Nakayama, 2003; Watson & Clifford, 2003; Zhao & Chubb, 2001). Adaptation aftereffects result from a reduction in the response of particular sub-populations of neurons tuned to the properties of the adapting stimulus. In the case of faces, adaptation aftereffects can demonstrate that the population of neurons coding facial characteristics is sensitive to a particular dimension, especially if adaptation effects can be transferred across identities or face categories.

We used a high-level adaptation paradigm to ask two questions regarding how face animacy is coded in the visual system: Is animacy a dimension of face perception? Is animacy represented in a category-specific fashion or is animacy perception across face categories supported by a common neural mechanism?

2. Experiment 1: Will animacy show adaptation that transfers across individuals?

2.1. Methods

2.1.1. Participants—Twenty-four young adults (9 female, mean age: 21.9) from the MIT and NDSU communities participated in experiment 1. All participants had normal or corrected to normal vision.

2.1.2. Stimuli: animacy morphs—Four grey-scale images of adult female Caucasian faces with neutral expressions were morphed with visually similar images of doll faces using FantaMorph software (Version 4; Abrosoft Co., Beijing, China). Fantamorph interpolates two original images to create morph continua of images that range from fully human to fully

inanimate (Figure 1). Each continuum was comprised of eleven images, spanning real to artificial appearance in 10% increments.

2.1.3. Procedure—Experiment one consisted of two parts: A baseline phase, followed by an adaptation phase. In the baseline phase, participants rated the animacy of 220 face images (two continua from the full set of four) presented in a randomized order. Each stimulus was presented for 1000 msec and participants rated each image 10 times over the course of the baseline testing using a 1-7 scale, where 1 was “completely inanimate” and 7 was “completely human”. We counterbalanced morph pair combinations across participants.

In the adaptation phase, participants were first presented with either a fully human or fully artificial face for 30 seconds. We selected this adapting face from one of the two continua not seen during the participants’ baseline phase. After this extended adaptation period, participants made animacy judgments using the same procedure used during the baseline phase, with the exception that each trial was preceded by a 2-second “top-up” adapting stimulus. We used the same adapting face for the entire adaptation phase. Test stimuli belonged to the morph continua not presented during the baseline phase. To minimize the contribution of image-level adaptation effects, we presented adapting stimuli at a larger size ($12^\circ \times 12^\circ$ visual angle) than the test stimuli ($4.85^\circ \times 4.85^\circ$).

2.2. Results

We transformed participants’ 1-7 rating to a 0-1 scale (without normalizing their range of responses) so that we could estimate the point of subjective equality (PSE) by fitting a psychometric function to each participant’s data. We fit logistic functions to each participants’ normalized ratings using the MATLAB Palamedes toolbox (Prins & Kingdom, 2009) and calculated each participant’s aftereffect size by subtracting the adapted PSE from the baseline PSE. The adapted PSE should move toward the adaptor; e.g., if the adapting image was a doll, doll faces should look more human post-adaptation. Thus, if face animacy is “adaptable”, PSE shifts should be in the opposite direction for adaptation to a human face vs. a doll face. PSE shifts in the expected direction were expressed as positive numbers while shifts in the opposite direction were expressed as negative numbers. This allowed us to collapse across doll and human adaptors by comparing the measured PSE values to zero in all cases. Because we have an a priori expectation for the direction of adaptation aftereffects, one-sample t-tests examining these aftereffects were done using a one-tailed t-test.

Adaptation aftereffects were significant when collapsed across human and doll adaptation trials ($t(23) = 4.04$, $p = .001$, one-sample t-test; $d = 1.27$) and the strength of adaptation did not differ for human or doll adapting images ($t(22) = .73$, $p = .468$, two-sample t-test; $d = 0.302$). Additionally, since the adapting face belonged to one of the two morph continua tested during the adaptation phase, we also asked if adaptation was stronger when adaptor and test image were identity matched. Both continua showed significant adaptation effects that did not differ from one another ($t(22) = .668$, $p = .51$, paired-sample t-test; $d = .227$) suggesting that animacy adaptation is neither image-specific nor identity-specific.

2.3. Discussion

In Experiment 1, we observed animacy aftereffects obtained for morphed images of adult faces and dolls. In Experiment 2, we examined the transfer of animacy adaptation aftereffects between child faces and adult faces. The “other age” effect (M. G. Rhodes & Anastasi, 2012) demonstrates that child and adult faces are processed differentially and may be represented by distinct neural sub-populations, as suggested by previous adaptation studies (O’Neil & Webster, 2011; Schweinberger et al., 2010). If animacy perception is category-specific, adapting to young children’s faces or dolls that look like children should not affect the perceived animacy of adult faces. If, however, animacy is a face characteristic that adapts within face space regardless of age category, aftereffects from child faces should be as strong as those seen to adult faces.

3. Experiment 2: Will animacy aftereffects transfer between different categories of human faces?

3.1. Methods

3.1.1. Participants—Sixteen young adults (9 female, mean age: 26.6) from the MIT community participated in experiment 2. None of these participants were subjects in Experiment 1.

3.1.2. Stimuli: animacy morphs—Stimuli were created as described in Experiment 1. In addition to the four adult morph continua, four grey-scale images of young Caucasian children’s faces with neutral expressions were morphed with visually similar images of child-like doll faces.

3.1.3. Procedure—We used the same testing procedure described in Experiment 1, except that, after baseline animacy ratings were obtained using the adult human/doll morphs, the adaptor used during the adaptation phase was either a real child’s face, or a child-like doll. As in Experiment 1, the adult morph pairs used during the baseline phase were different than those used during the adaptation phase and pairs were counterbalanced across participants. In addition, the morph pair from which the adaptation image was drawn was counterbalanced across participants.

3.2. Results

Adaptation aftereffects were significant when collapsed across child and doll adaptation trials ($t(15) = 6.96, p < .001$, one-sample t-test; $d = 1.76$) and were also significant for adaptation to both human child faces ($t(7) = 5.5, p = .001$, one-sample t-test; $d = 1.94$) and doll faces ($t(7) = 5.6, p = .001$, one-sample t-test; $d = 1.96$) separately. Animacy adaptation thus exhibits transfer across age categories, suggesting a shared neural mechanism supporting perceived animacy.

3.3. Discussion

Our first two experiments revealed that animacy “adapts” and transfers across age categories. Next, we examined the transfer of animacy aftereffects between dog faces and

adult human faces to determine if animacy perception is species-specific. If animacy aftereffects are driven by how generally “alive” a face appears to be, adaptation to a dog’s face or the face of a stuffed dog toy may also show transfer to adult human faces. If, however, animacy adapts within a species-specific face space, as might be expected from the “other-species effect” (Pascalis, De Haan, & Nelson, 2002; Sugita, 2008), animacy aftereffects may not cross the species barrier. In separate tasks, we examined the extent to which animacy aftereffects transferred between dog adaptors and human test images (Experiment 3a) and also whether or not animacy aftereffects were evident using dog faces as adapting *and* test stimuli (Experiment 3b).

4. Experiment 3: Will animacy aftereffects transfer across species?

4.1. Methods

4.1.1. Participants—Sixteen young adults (8 female; mean age: 19.2) from the NDSU community participated in Experiment 3a and an additional sixteen (7 female; mean age: 25.2) from the MIT community in Experiment 3b. None of these participants were tested in either previous task.

4.1.2. Stimuli: animacy morphs—We created morph continua as described in Experiment 1. In addition to the other morph continua already described, four grey-scale images of dog faces were morphed with visually similar images of plush dog faces.

4.1.3. Procedure for Experiment 3a—We adopted the same procedure described in Experiment 1, except that after baseline animacy ratings were obtained using the adult human/doll morphs, the adaptor image used during the adaptation phase was from either of the extreme ends of the dog/stuffed-toy continua. As in Experiment 1, the adult morph pairs used during the baseline phase were different than those used during the adaptation phase and pairs were counterbalanced across participants. In addition, the morph pair from which the adaptation image was drawn was counterbalanced across participants.

4.1.4. Procedure for Experiment 3b—Experiment 3b: When testing within-species adaptation using the dog/toy morphs, the procedure was exactly as described in Experiment 1, except using the dog/toy morphs instead of the human/doll morphs. In addition, participants were asked to rate dog/toy morphs as either “fully dog-like” or “fully toy-like”.

4.2 Results and Discussion

Adaptation to dog/toy dog faces did not produce a significant adaption aftereffect when human/doll morphs were the test items ($t(15) = -.48$, $p = .64$, one-sample t-test; $d = .12$). However, animacy aftereffects were obtained when both the adaptor and test images were dogs ($t(15) = 2.15$, $p = .025$, one-sample t-test; $d = .54$). Thus, while facial animacy *is* adaptable within non-human species, animacy aftereffects do not transfer between different species.

5. Experiment 5: Will animacy adaptation aftereffects be seen in inversion?

Experiment 4 examined whether animacy adaptation aftereffects would still be seen if the adapting stimulus was inverted. If animacy adaptation aftereffects rely on face-specific mechanisms in face space, they should be weaker when the adapting stimulus is inverted.

5.1. Methods

5.1.1. Participants—Twenty-four young adults from the MIT community (13 female, mean age: 24.3) participated in experiment 4. None of the participants in experiment 4 were tested in experiments 1, 2 or 3.

5.1.2. Stimuli: animacy morphs—Stimuli were the same as those described in Experiment 1.

5.1.3. Procedure—The testing procedure was exactly as described in Experiment 1 except that, after baseline animacy ratings were obtained, the adaptation image used during the adaptation phase was inverted.

5.2. Results and Discussion

Animacy ratings were transformed as described for experiment 1 to obtain each participant's PSE for both baseline and adaptation conditions. Adaptation to inverted faces did produce an aftereffect that was very close to significant ($t(23) = 1.6, p = .06$, one-sample t-test; $d = .327$). Despite the fact that we do see near-significant adaptation aftereffects to an inverted face, the adaptation aftereffect seen to the inverted faces was significantly weaker than that seen to upright faces ($t(46) = 2.1, p = .04$, two-sample t-test, two-tailed; $d = .61$).

6. General Discussion

Our results revealed reliable adaptation aftereffects in the perception of “animacy” in human faces. Following adaptation to either a wholly inanimate doll face or a fully human face, the perceived animacy of subsequently presented faces was shifted away from the adapting stimulus, consistent with other examples of face adaptation. These animacy aftereffects were significantly weaker when the adapting stimulus was inverted, demonstrating that at least part of this aftereffect may be face-specific. These results demonstrate that the perception of animacy in faces is flexible and can be tuned, even over a very short time scale, by experience – thus suggesting that animacy is a psychologically real dimension of face variability.

Our data further demonstrate significant adaptation across age groups (child vs. adult), suggesting that animacy adaptation transfers not only across different identities (as demonstrated in experiment 1) but also across different face *categories* (demonstrated in experiment 2). The transfer of adaptation effects across categories has a limit, however, as the results of experiment 3 show that adaptation effects do not transfer between faces of different species.

What cues may be driving these animacy adaptation aftereffects? Although our data suggest that animacy adaptation is a high-level aftereffect, it is difficult to disentangle the visual cues that may contribute to the effect. Two cues that could drive part of this adaptation effect are skin-texture and the reflective qualities of the eyes. Adaptation aftereffects to manipulations of surface texture/reflectance have previously been demonstrated in non-face objects (Motoyoshi, 2007) and could explain some of the adaptation effect seen in our data. The clear difference in adaptation strength between upright and inverted adaptors, however, suggests that surface-based cues alone cannot explain the entire effect. Shape-based cues may also drive part of this effect. Although doll and human faces did not differ from each other systematically across morph pairs, certain differences remained relatively stable – for example dolls’ eyes tended to be slightly bigger and lower in the face than those of real human faces. These sorts of shape-based cues may be similar to previously reported shape-based manipulations of faces (e.g. expansion/contraction of face features) that have shown adaptation after-effects on the perception of facial characteristics like “normalcy” and “attractiveness” (e.g., (G. Rhodes et al., 2003). It is important to note, here, that we tested the transfer of adaptation effects across age and species categories only in one direction (child to adult, dog to human), leaving open the possibility that adaptation transfer in the other directions might differ. Although the one-way nature of these tests does not undermine our conclusions, investigating adaptation effect transfer in the other direction may provide clues to the source of these effects, especially if they prove to be asymmetrical. More generally, a systematic investigation of the contribution of both surface/texture and shape-based cues to animacy aftereffects will be important in future research investigating the perception of face animacy.

What do these results tell us about face perception itself? First, our data suggest that animacy is a *perceptual* dimension of faces, susceptible to high-level adaptation, thus lending additional support to earlier evidence that the human face processing system includes sensitivity to the “animacy” of faces (Balas & Horski, 2012; Looser & Wheatley, 2010). Second, the observed transfer between child faces and adult faces suggests that, while child faces are farther from the norm for adult observers than adult faces, they are still represented in the same face space as adult faces and that changes in animacy are perceived similarly across the two age categories. Animacy is not, however, perceptually monolithic. The lack of transfer between human and dog faces both reassures us that the effect is unlikely to be the result of response-bias and reinforces the perceptual nature of the effect – if the effect relied on the “concept” of animacy, we might expect to see transfer of adaptation from dogs to humans. Perceptually, animacy in human faces and dog faces may be supported by different visual cues – texture, surface reflectance and shape cues may still be relevant to the perception of animacy in dog faces, but in a quite different way than in human faces. In addition, our results are consistent with a model in which dog faces are not represented in the same face space as human faces but may be represented in a face space of their own.

What does animacy adaptation tell us about the neural architecture of the face-processing system? Previous research demonstrated that differences in the processing of human and doll faces did not emerge until relatively late in perceptual processing (~400ms) (Wheatley,

Weinberg, Looser, Moran, & Hajcak, 2011) and may take place in face-sensitive regions of the superior temporal sulcus (STS) (Looser, Guntupalli, & Wheatley, 2012). Our data cannot directly address the question of where animacy may be coded in the brain but, in the context of these previous results, our data suggest that processing differences between animate and inanimate faces may be driven by face-selective neural populations within the STS that drive animacy perception. If so, animacy perception in faces may also be related to the perception of animacy in other social stimuli that engage the pSTS, including both point-light human figures engaged in “biological motion” (e.g. Grossman, Jardine, & Pyles, 2010) and simple animated shapes whose movements are perceived to have intention (Gao, Scholl, & McCarthy, 2012; Lee, Gao, & McCarthy, 2012). Further research is needed to clarify if animacy perception in faces and biological motion perception engage similar brain regions and whether adaptation effects using biological motion stimuli (Jordan, Fallah, & Stoner, 2006; Troje, Sadr, Geyer, & Nakayama, 2006) can also reflect differences in animacy (for instance contrasting adaptation effects to the movements of a humanoid robot against typical human movements). Regardless of its neural substrate, our results demonstrate that changes in animacy activate and adapt the tuning of the neurons that underlie face perception - at the very least suggesting that animacy perception is a fundamental part of the face processing system.

Acknowledgments

This study was supported, in part, by a grant from the Simons Foundation to the Simons Center for the Social Brain at MIT. BB was supported by NIGMS #P20 GM103505 and ND EPSCoR NSF #EPS-0814442.

References

- Balas B, Horki J. You can take the eyes out of the doll, but.... Perception. 2012; 41(3):361–364. [PubMed: 22808589]
- Gao T, McCarthy G, Scholl BJ. The wolfpack effect. Perception of animacy irresistibly influences interactive behavior. Psychological science. 2010; 21(12):1845–1853. doi: 10.1177/0956797610388814. [PubMed: 21078895]
- Gao T, Scholl BJ. Chasing vs. stalking: interrupting the perception of animacy. Journal of experimental psychology. Human perception and performance. 2011; 37(3):669–684. doi:10.1037/a0020735. [PubMed: 21639674]
- Gao T, Scholl BJ, McCarthy G. Dissociating the detection of intentionality from animacy in the right posterior superior temporal sulcus. The Journal of neuroscience: the official journal of the Society for Neuroscience. 2012; 32(41):14276–14280. doi:10.1523/JNEUROSCI.0562-12.2012. [PubMed: 23055497]
- Grossman ED, Jardine NL, Pyles JA. fMR-Adaptation Reveals Invariant Coding of Biological Motion on the Human STS. Frontiers in human neuroscience. 2010; 4:15. doi:10.3389/neuro.09.015.2010. [PubMed: 20431723]
- Jordan H, Fallah M, Stoner GR. Adaptation of gender derived from biological motion. Nature neuroscience. 2006; 9(6):738–739. doi:10.1038/nn1710.
- Kanwisher N, Yovel G. The fusiform face area: a cortical region specialized for the perception of faces. Philosophical transactions of the Royal Society of London. Series B, Biological sciences. 2006; 361(1476):2109–2128. doi:10.1098/rstb.2006.1934.
- Langton SRH, Law AS, Burton AM, Schweinberger SR. Attention capture by faces. Cognition. 2008; 107(1):330–342. doi:10.1016/j.cognition.2007.07.012. [PubMed: 17767926]
- Lee SM, Gao T, McCarthy G. Attributing intentions to random motion engages the posterior superior temporal sulcus. Social cognitive and affective neuroscience. 2012 doi:10.1093/scan/nss110.

- Leopold DA, O'Toole AJ, Vetter T, Blanz V. Prototype-referenced shape encoding revealed by high-level aftereffects. *Nature neuroscience*. 2001; 4(1):89–94. doi:10.1038/82947.
- Looser CE, Guntupalli JS, Wheatley T. Multivoxel patterns in face-sensitive temporal regions reveal an encoding schema based on detecting life in a face. *Social cognitive and affective neuroscience*. 2012 doi:10.1093/scan/nss078.
- Looser CE, Wheatley T. The tipping point of animacy. How, when, and where we perceive life in a face. *Psychological science*. 2010; 21(12):1854–1862. doi:10.1177/0956797610388044. [PubMed: 21097720]
- O'Neil SF, Webster MA. Adaptation and the perception of facial age. *Visual cognition*. 2011; 19(4): 534–550. doi:10.1080/13506285.2011.561262. [PubMed: 22215952]
- Pascalis O, De Haan M, Nelson CA. Is face processing species-specific during the first year of life? *Science (New York, N.Y.)*. 2002; 296(5571):1321–1323. doi:10.1126/science.1070223.
- Peelen MV, Downing PE. The neural basis of visual body perception. *Nature reviews. Neuroscience*. 2007; 8(8):636–648. doi:10.1038/nrn2195.
- Pelphrey KA, Carter EJ. Brain mechanisms for social perception: lessons from autism and typical development. *Annals of the New York Academy of Sciences*. 2008; 1145:283–299. doi:10.1196/annals.1416.007. [PubMed: 19076404]
- Prins, N.; Kingdom, FA. Palamedes: Matlab routines for analyzing psychophysical data. 2009. Retrieved from <http://www.palamedestoolbox.org>
- Rhodes G, Jeffery L. Adaptive norm-based coding of facial identity. *Vision research*. 2006; 46(18): 2977–2987. doi:10.1016/j.visres.2006.03.002. [PubMed: 16647736]
- Rhodes G, Jeffery L, Watson TL, Clifford CWG, Nakayama K. Fitting the mind to the world: face adaptation and attractiveness aftereffects. *Psychological science*. 2003; 14(6):558–566. [PubMed: 14629686]
- Rhodes MG, Anastasi JS. The own-age bias in face recognition: a meta-analytic and theoretical review. *Psychological bulletin*. 2012; 138(1):146–174. doi:10.1037/a0025750. [PubMed: 22061689]
- Ro T, Russell C, Lavie N. Changing faces: a detection advantage in the flicker paradigm. *Psychological science*. 2001; 12(1):94–99. [PubMed: 11294237]
- Schweinberger SR, Zäske R, Walther C, Golle J, Kovács G, Wiese H. Young without plastic surgery: perceptual adaptation to the age of female and male faces. *Vision research*. 2010; 50(23):2570–2576. doi:10.1016/j.visres.2010.08.017. [PubMed: 20800608]
- Sugita Y. Face perception in monkeys reared with no exposure to faces. *Proceedings of the National Academy of Sciences of the United States of America*. 2008; 105(1):394–398. doi:10.1073/pnas.0706079105. [PubMed: 18172214]
- Troje NF, Sadr J, Geyer H, Nakayama K. Adaptation aftereffects in the perception of gender from biological motion. *Journal of vision*. 2006; 6(8):850–857. doi:10.1167/6.8.7. [PubMed: 16895463]
- Watson TL, Clifford CWG. Pulling faces: an investigation of the face-distortion aftereffect. *Perception*. 2003; 32(9):1109–1116. [PubMed: 14651323]
- Webster MA, Kaping D, Mizokami Y, Duhamel P. Adaptation to natural facial categories. *Nature*. 2004; 428(6982):557–561. doi:10.1038/nature02420. [PubMed: 15058304]
- Webster MA, MacLeod DIA. Visual adaptation and face perception. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*. 2011; 366(1571):1702–1725. doi: 10.1098/rstb.2010.0360.
- Wheatley T, Weinberg A, Looser C, Moran T, Hajcak G. Mind perception: real but not artificial faces sustain neural activity beyond the N170/VPP. *PloS one*. 2011; 6(3):e17960. doi:10.1371/journal.pone.0017960. [PubMed: 21483856]
- Zhao L, Chubb C. The size-tuning of the face-distortion after-effect. *Vision research*. 2001; 41(23): 2979–2994. [PubMed: 11704237]

Figure 1.

A and B) Examples of adult and dog face pairs used in experiments 1-3 to create morph sequences. Eleven morph levels evenly spaced along the morph sequence for each face pair were used in the experiments. C) Examples of adaptors used in experiments 1-3.

Figure 2.

Average psychometric curves for animacy ratings in each experiment (A-D). In each case, black lines indicate the average baseline animacy rating, blue and magenta lines represent post-adaptation response following adaptation to a real or toy face, respectively. Error bars = ± 1 SEM.

Figure 3.

Average aftereffect size for all 4 experiments, error bars = ± 1 SEM. We observe significant animacy aftereffects in all tasks, with the exception of transfer between dog and human faces.