Published in final edited form as:

Prev Med. 2013 June ; 56(6): 410–412. doi:10.1016/j.ypmed.2013.02.013.

Leisure-time Physical Activity in Relation to Depressive Symptoms in African-Americans: Results from the National Survey of American Life

Elisa R. Torres, PhD^a, Carolyn M. Sampselle, PhD^b, David L. Ronis, PhD^{c,d}, Harold W. Neighbors, PhD^e, and Kimberlee A. Gretebeck, PhD^f

Elisa R. Torres: ertorres@wisc.edu; Carolyn M. Sampselle: csampsll@umich.edu; David L. Ronis: dronis@umich.edu; Harold W. Neighbors: woodyn@umich.edu; Kimberlee A. Gretebeck: kgretebeck@wisc.edu

^aSchool of Nursing, University of Wisconsin, 600 Highland Ave., Clinical Science Center K6/326, Madison, WI 53792-2455, United States, Phone: 1 (608) 263-5268, Fax: 1 (608) 263-5458

^bSchool of Nursing, University of Michigan, Director, Community Engagement Program, Michigan Institute for Clinical and Health Research, 400 North Ingalls Building, Ann Arbor, MI 48109-5482

^cSchool of Nursing, University of Michigan, Director of Statistical Consulting Team, Room 4330, 400 North Ingalls Building, Ann Arbor, MI 48109-5482

^dAnn Arbor VA, 2215 Fuller Road, Ann Arbor, MI 48105

^eSchool of Public Health, University of Michigan, Associate Director, Center for Research on Ethnicity, Culture and Health, 1415 Washington Heights, Ann Arbor, Michigan 48104-3028

^fSchool of Nursing, University of Wisconsin, 600 Highland Ave., Madison, WI 53792-2455

Abstract

Objective—To examine the frequency of leisure-time physical activity (LTPA) in relation to depressive symptoms in a nationally representative sample of African American (AA) women and AA men with guidance by Stokols' Social Ecological Framework.

Method—A secondary analysis of AA women (n=1,811) and AA men (n=1,038) was performed on the National Survey of American Life, where a four stage national area probability sampling was conducted. Interviews were conducted 2001-2003. Clinically depressed AA were excluded from the current study. LTPA was measured by self-report frequency (never, rarely, sometimes, often) of participation in sports/exercise. Depressive symptoms were measured with the Center for Epidemiologic Studies Depression scale. Logistic regression for complex samples was used to examine the relationship between LTPA and depressive symptoms, adjusting for biopsychobehavioral and sociophysical environmental factors.

Results—Compared with AA women and AA men who reported never participating in LTPA, the multivariate OR for depressive symptoms in AA women and AA men who reported

---

Corresponding author: Elisa R. Torres, PhD, School of Nursing, University of Wisconsin, 600 Highland Ave., Clinical Science Center K6/326, Madison, WI 53792-2455, United States Phone: 1 (608) 263-5268 Fax: 1 (608) 263-5458, ertorres@wisc.edu.

Conflict of Interest Statement: The authors declare that there are no conflicts of interest.

participating in LTPA often was 0.42 (95% CI=0.24-0.72) and 0.41 (95% CI=0.25-0.69) respectively.

Conclusion—Increased frequency of LTPA was associated with fewer depressive symptoms in a nationally representative sample of non-clinically depressed AAs.

Keywords

exercise; sport; depression; preventive medicine

Introduction

Leisure-time physical activity (LTPA) is exercise, sports and/or recreation that are not associated with occupation, household, or transportation. Higher frequencies of LTPA are associated with less depressive symptoms in African-Americans (AAs) (Farmer et al., 1988; Wise et al., 2006). Farmer et al. examined recreational physical activity in the U.S. population as a whole, and stratified results by race and sex. Due to the small number of AAs in the sample, sample weights were not used (Farmer et al., 1988). Wise et al. examined exercise in exclusively AA women across the U.S. with convenience sampling. Neither study provided nationally representative estimates or reported guidance by theory. Since LTPA is hypothesized to relieve depressive symptoms through biological, psychological and social mechanisms (Paluska and Schwenk, 2000), theories that consider multiple levels of influence are recommended (Torres et al., 2010).

The Social Ecology of Health Promotion offers a broad understanding of how biological, psychological, sociocultural and physical environmental factors affect well-being (Stokols, 1992) and is applied to LTPA (Linde, et al., 2012) and depressive symptoms (Katz, et al., 2008) in AAs to describe biopsychobehavioral (age, body mass index [BMI], disability, family history of depression, discrimination) and sociophysical environmental (household income, national region and neighborhood safety) factors (Torres et al., 2010). LTPA (Wise, et al., 2006) and depression (Williams et al., 2007) both decrease with age. LTPA is inversely associated with depressive symptoms among the obese but not non-obese (Wise et al., 2006). Disability is associated with less LTPA (National Center for Health Statistics [NCHS], 2011) and more depression (Williams et al., 2007). Family history of depression is a proxy for genetics, which modifies the relation between physical activity and depressive symptoms (Mata et al., 2010). Perceived discrimination is predictive of depressive symptoms (Banks et al., 2006). Higher household income is associated with higher LTPA (NCHS, 2011) and lower depressive symptoms (Banks et al., 2006). LTPA is associated with living in the West (Wise et al., 2006), while living in the South and West have a lower prevalence of depression than the Northeast and Midwest (Williams et al., 2007). The purpose of this study was to examine the frequency of LTPA in relation to depressive symptoms in a nationally representative sample of AAs while controlling for biopsychobehavioral and sociophysical environmental factors.

Method

A secondary analysis was performed on the National Survey of American Life (NSAL). Heeringa et al. (2004) described a four stage national area probability sampling. Primary stage comprised a stratified probability sample of 1990 census block households reported to be <10% or 10% AA (Heeringa et al., 2004). Second stage included area segments formed by linking geographically continuous census blocks (Heeringa et al., 2004). Third stage included a systematic random sample of housing units which were contacted in person by an interviewer (Heeringa et al., 2004). If the interviewer reported that one or more eligible adults lived at the sample housing unit address, the interviewer prepared a complete list of household members and randomly selected a respondent for the study interview, comprising the fourth stage (Heeringa et al., 2004). Pretesting of questionnaires and training of interviewers was described by Jackson et al. (2004). Oral consent was obtained prior to initiating the interview. There was racial matching of interviewers and respondents, with face-to-face interviews lasting on average 2 hours and 20 minutes (Jackson et al., 2004). The response rate was 70.7% (Jackson et al., 2004). Of the 3,570 AAs in the NSAL, individuals who met the DSM-IV-TR criteria for MDD (n=75), Dysthymia (n=23) or Bipolar Disorder (n=13) were excluded since clinical depression may influence frequency of LTPA (Wise, et al., 2006). Cases missing data on depressive symptoms, LTPA or covariates were excluded, resulting in a sample size of 2,849. The Institutional Review Board at the University of Michigan granted approval for the NSAL (Jackson et al., 2004) and the current study.

Depressive symptoms were measured with the Center for Epidemiologic Studies Depression (CESD) Scale (Radloff, 1977). The original 20-item CESD was reduced to 12-items in the NSAL. Eleven items in AA women ($\alpha=0.80$) and 10 items in AA men ($\alpha=0.73$) demonstrated validity and reliability (Torres, 2012). A cutoff score of nine for the 11-item CESD and eight for the 10-item CESD was used, with higher scores indicating more symptoms (Torres, 2012).

LTPA was measured with one question asking how often the respondent engaged in sports/exercise; never, rarely, sometimes, often. Scores were coded with higher scores indicating more LTPA. The question came from the Americans' Changing Lives questionnaire (Lantz et al., 1998).

BMI was calculated by self-reported weight(lbs)/(703)/height(in²), which approximates BMI in kg/m²(U.S. Department of Health & Human Services, 2000). Disability was measured with three items from the WHO's Disability Assessment Schedule II mobility domain (Rehm et al., 2006), $\alpha=0.68-0.76$. Respondents reported whether they had a family history of depression. The Everyday Discrimination Scale measures chronic, routine, and relatively minor experiences of unfair treatment (Williams et al., 1997), consisting of ten items, the averaged higher scores indicating more discrimination, $\alpha=0.88-0.89$. National region was based on the four Census regions. Neighborhood safety was measured by two questions from the National Survey of Black Americans, with averaged higher scores indicating a safer neighborhood, $\alpha=0.72-0.73$.

Analyses were performed in STATA 10.0. Weights constructed specifically for the study design and methodology were used (Heeringa and Berglund, 2013). T-tests and Chi-square for complex samples determined whether the sex difference for each descriptive characteristic was statistically significant. Logistic regression for complex samples was used to compute odds ratios (ORs) and 95% confidence intervals.

Results

Compared to AA men, AA women reported higher BMIs, discrimination, disability, family histories of depression, and lower household incomes (Table 1). There were statistically significant sex differences in LTPA. More AA women reported never participating in LTPA than AA men ($X^2=14.8$, $p=.0005$) and more AA men reported participating in LTPA often than AA women ($X^2=35.0$, $p<.0001$).

Compared with AA women who reported never participating in LTPA, the OR for depressive symptoms in AA women who reported participating in LTPA rarely and often were 0.59 and 0.42, respectively (Table 2). AA women had higher OR for depressive symptoms when they reported more disability, a family history of depression, and lower household incomes and discrimination.

Compared with AA men who reported never participating in LTPA, the OR for depressive symptoms in AA men who reported participating in LTPA sometimes and often were 0.49 and 0.41, respectively (Table 2). AA men had higher OR for depressive symptoms when they reported more disability and less discrimination.

Discussion

The biggest decrease in the odds of depressive symptoms was observed among AAs who reported participating in LTPA often compared to never. These results are consistent with the idea that LTPA may prevent depressive symptoms in AAs. It is important to stratify results by sex since AA women report more depressive symptoms (Torres, 2012) and less frequency in LTPA than AA men. AA women also reported more family histories of depression and lower household incomes, which were associated with higher OR for depressive symptoms in AA women but not men. The Social Ecology of Health Promotion was beneficial in selecting potential confounders a priori.

Generalization of results is limited to non-clinically depressed AAs. It is not possible to determine whether LTPA was an antecedent to, or a consequence of, depressive symptoms. However, by excluding clinically-depressed individuals, the possibility that depressive symptoms influenced LTPA was minimized. The association between LTPA and depressive symptoms may have been confounded by unmeasured factors. Effort was made to include the most relevant factors by use of a theory. All measures were self-report. LTPA in the NSAL was measured with only one question without 1) established psychometrics; 2) definition of never, rarely, sometimes, often; 3) reference to a time frame; 4) or differentiation between sports and exercise.

Increased frequency of LTPA was associated with fewer depressive symptoms in a nationally representative sample of non-depressed AAs. Future research should include valid and reliable measures of LTPA, validation of self-report measures, and address intensity, duration, and types of LTPA.

Acknowledgments

The work of this study was supported by grants from the National Institutes for Health (U01MH057716, 1F31NR010669, T32NR07110, and R25GM058641).

References

- Banks KH, Kohn-Wood LP, Spencer M. An examination of the African American experience of everyday discrimination and symptoms of psychological distress. *Community Ment Health J.* 2006; 42:555–570. [PubMed: 16897412]
- Farmer ME, Locke BZ, Mo cicki EK, Dannenberg AL, Larson DB, Radloff LS. Physical activity and depressive symptoms: the NHANES I Epidemiologic Follow-up Study. *Am J Epidemiol.* 1988; 128:1340–1351. [PubMed: 3264110]
- Heeringa, SG.; Berglund, P. National Institutes of Mental Health (NIMH) Collaborative Psychiatric Epidemiology Survey Program (CPES) Data Set. [Accessed January 31, 2013] Integrated weights and sampling error codes for design-based analysis. Available at http://www.icpsr.umich.edu/icpsrweb/CPES/about_cpes/weighting/index.jsp
- Heeringa SG, Wagner J, Torres M, Duan N, Adams T, Berglund P. Sample designs and sampling methods for the Collaborative Psychiatric Epidemiology Studies (CPES). *Int J Methods Psychiatr Res.* 2004; 13:221–240. [PubMed: 15719530]
- Jackson JS, Torres M, Caldwell CH, Neighbors HW, Nesse RM, Taylor RJ, Trierweiler SJ, Williams DR. The National Survey of American Life: a study of racial, ethnic and cultural influences on mental disorders and mental health. *Int J Methods in Psychiatr Res.* 2004; 13:196–207. [PubMed: 15719528]
- Katz K, Blake S, Milligan R, Sharps P, White D, Rodan M, Rossi M, Murray K. The design, implementation and acceptability of an integrated intervention to address multiple behavioral and psychosocial risk factors among pregnant African American women. *BMC Pregnancy Childbirth.* 2008; 8:22–22. [PubMed: 18578875]
- Lantz PM, House JS, Lepkowski JM, Williams DR, Mero RP, Chen J. Socioeconomic factors, health behaviors, and mortality: results from a nationally representative prospective study of US adults. *JAMA.* 1998; 279:1703–1708. [PubMed: 9624022]
- Linde J, Nygaard K, MacLehose R, Mitchell N, Harnack L, Cousins J, Graham D, Jeffery R. HealthWorks: results of a multi-component group-randomized worksite environmental intervention trial for weight gain prevention. *Int J Behav Nutr Phys Act.* 2012; 9:14–14. [PubMed: 22340088]
- Mata J, Thompson RJ, Gotlib IH. BDNF genotype moderates the relation between physical activity and depressive symptoms. *Health Psychol.* 2010; 29:130–133. [PubMed: 20230085]
- National Center for Health Statistics (NCHS). [Accessed January 31, 2013] Health, United States, 2011. 2011. Available at <http://www.cdc.gov/nchs/data/hs/hs11.pdf>
- Paluska SA, Schwenk TL. Physical activity and mental health: current concepts. *Sports Med.* 2000; 29:167–180. [PubMed: 10739267]
- Radloff LS. The CES-D Scale: a self-report depression scale for research in the general population. *Appl Psychol Meas.* 1977; 1:385–401.
- Rehm J, Üstün TB, Saxena S, Nelson CB, Chatterji S, Ivis F, Adlart E. On the development and psychometric testing of the WHO screening instrument to assess disablement in the general population. *Int J Methods Psychiatr Res.* 2006; 8:110–122.
- Stokols D. Establishing and maintaining healthy environments. Toward a social ecology of health promotion. *Am Psychol.* 1992; 47:6–22. [PubMed: 1539925]

- Torres ER. Psychometric properties of the Center for Epidemiologic Studies Depression Scale in African-American and Black Caribbean Adults. *Issues Ment Health Nurs*. 2012; 33:687–696. [PubMed: 23017046]
- Torres ER, Sampselle C, Gretebeck KA, Ronis DL, Neighbors HW. Physical activity effects on depressive symptoms in Black adults. *J Health Dispar Res Pract*. 2010; 4:71–88.
- U.S. Department of Health & Human Services. [Accessed January 31, 2013] The practical guide: identification, evaluation, and treatment of overweight and obesity in adults. 2000. Available at http://www.nhlbi.nih.gov/guidelines/obesity/prctgd_c.pdf
- Williams DR, Yu Y, Jackson JS, Anderson NB. Racial differences in physical and mental health: socioeconomic status, stress and discrimination. *J Health Psychol*. 1997; 2:335–351. [PubMed: 22013026]
- Williams DR, Gonzalez HM, Neighbors H, Nesse R, Abelson JM, Sweetman J, Jackson JS. Prevalence and Distribution of Major Depressive Disorder in African Americans, Caribbean Blacks, and Non-Hispanic Whites: Results From the National Survey of American Life. *Arch Gen Psychiatry*. 2007; 64:305–315. [PubMed: 17339519]
- Wise LA, Adams-Campbell LL, Palmer JR, Rosenberg L. Leisure time physical activity in relation to depressive symptoms in the Black Women's Health Study. *Ann Behav Med*. 2006; 32:68–76. [PubMed: 16827631]

Table 1
Sample Characteristics of African-Americans: The National Survey of American Life, 2001-2003

	Mean (Confidence Interval)		t	p
	African-American			
	Women (n=1,811)	Men (n=1,038)		
Age	42 (41-43)	41 (40-43)	0.2	.67
Body Mass Index	29 (29-30)	28 (28-29)	28.5	<.0001
Disability	4.7 (4.0-5.4)	3.6 (2.8-4.3)	5.7	.0229
Discrimination	78 (78-79)	76 (75-77)	17.0	.0004
Household Income ^a	32,540 (30,261-34,819)	42,612 (38,739-46,485)	39.5	<.0001
Neighborhood safety	3.1 (3.0-3.2)	3.1 (3.0-3.2)	1.3	.26
	No. (%)		X ²	p
Family History of Depression	544 (61.5)	243 (38.5)	8.3	.0067
National region			1.4	.26
West	112 (4.7)	86 (5.1)		
Northeast	204 (8.7)	122 (6.9)		
Midwest	288 (10.2)	156 (7.4)		
South	1207 (31.0)	674 (26.0)		
Leisure-time Physical Activity			14.4	<.0001
Never	447 (13)	180 (7)		
Rarely	331 (10)	146 (7)		
Sometimes	524 (16)	276 (12)		
Often	509 (16)	436 (19)		

^a top coded at \$200,000

Table 2
Odds Ratios for Depressive Symptoms in Relation to Leisure-Time Physical Activity in African-Americans: The National Survey of American Life, 2001-2003

Variable	African-American women (n=1,811)			African-American men (n=1,038)		
	No.	CESD11 9(%)	OR 95% CI	No.	CESD10 8(%)	OR 95% CI
LTPA						
Never	447	32	1.00	180	25	1.00
Rarely	331	16	0.59*	146	18	0.80 0.45-1.42
Sometimes	524	32	0.76 0.50-1.14	276	24	0.49* 0.28-0.88
Often	509	20	0.42* 0.24-0.72	436	33	0.41* 0.25-0.69
Biopsychobehavioral factors						
Age			0.99 0.98-1.00			0.99 0.98-1.00
BMI			1.01 0.98-1.03			1.01 0.97-1.04
Disability			1.02* 1.01-1.03			1.02 1.00-1.03
Family history of depression			1.69* 1.12-2.56			1.40 0.93-2.11
Perceived discrimination			0.93* 0.91-0.95			0.96* 0.94-0.98
Sociophysical environmental factors						
Household income			a*	b	c*	d
National region						
Northeast			1.00	1.00	1.00	
Midwest			1.67 0.69-4.04		1.95 0.78-4.89	
South			1.43 0.61-3.36		1.76 0.74-4.18	
West			1.33 0.55-3.27		0.99 0.32-3.11	
Neighborhood safety			0.87 0.75-1.01		0.85 0.69-1.04	
			p<.0001		p<.0001	

Note: CESD11=Center for Epidemiologic Studies Depression Scale, 11 items in National Survey of American Life; CESD10=Center for Epidemiologic Studies Depression Scale, 10 items in National Survey of American Life; OR=odds ratio; CI=95% Confidence Interval;

* p<.05;

a = .9999771 OR,

d =.9999804-.999975 CI. OR greater than 1 indicates greater odds of more depressive symptoms, while OR less than 1 indicates lesser odds of less depressive symptoms.

b =.9999697-.9999845 CI,

c =.9999889 OR,