Comprehensive Evaluation of Culture- Negative Endocarditis

with Use of Cardiac and 4-Dimensional-Flow
Magnetic Resonance Imaging

Samir R. Thadani, MD, MEng
Petter Dyverfeldt, PhD
Amy Gin, MD
Sam Chitsaz, MD
Rajni K. Rao, MD
Michael D. Hope, MD

Section Editor:

Raymond F. Stainback, MD,
Department of Adult
Cardiology, Texas Heart
Institute, 6624 Fannin St.,
Suite 2480, Houston, TX
77030

From: Division of Cardiology,
Department of Medicine
(Drs. Gin, Rao, and Thadani),
Department of Radiology
(Drs. Dyverfeldt and Hope),
and Cardiac Biomechanics
Laboratory (Dr. Chitsaz),
University of California,
San Francisco, San Fran-
cisco, California 94143;
and Division of Cardiology,
Department of Medicine
(Dr. Thadani), California
Pacific Medical Center,
San Francisco, California
94115

Dr. Thadani's secondary
affiliation is now the Division
of Cardiology, Department
of Medicine, Kaiser Perma-
nente, South San Francisco,
California.

Address for reprints:

Samir R. Thadani, MD,
Division of Cardiology,
University of California, San
Francisco, 505 Parnassus
Ave., Box 0214, San Fran-
cisco, CA 94143-0214

E-mail:

samir.thadani@ucsf.edu

© 2014 by the Texas Heart®
Institute, Houston

A 42-year-old, otherwise healthy man with exposure to domesticated buffalo presented with a 3-day history of intermittent headaches and word-finding difficulties. Magnetic resonance images (MRI) revealed an ischemic infarct in the left temporal lobe. A high-pitched, intense, pandsystolic murmur was detected at the right upper sternal border. A transthoracic echocardiogram showed moderate-to-severe aortic insufficiency. A transesophageal echocardiogram (TEE) showed a bicuspid aortic valve with possible leaflet perforation. Cardiac magnetic resonance (CMR) images confirmed regurgitation through a perforated bicuspid leaflet, excluded an abscess, and revealed a dilated ascending thoracic aorta (Fig. 1). The eccentric regurgitant jet and asymmetric aortic dilation were best seen by means of time-resolved, 3-dimensional, phase-contrast MRI (4-dimensional [4D]-flow MRI), with use of instantaneous streamline and dynamic pathline views of blood flow (Fig. 2). The

Fig. 1 Cardiac magnetic resonance image shows the dilated ascending aorta, and significant aortic regurgitation (arrow) through the perforated bicuspid aortic valve leaflet.

[Supplemental motion image is available for Figure 1.](#)

Fig. 2 Four-dimensional-flow magnetic resonance imaging of the thoracic aorta. Streamlines are color-coded for velocity and reveal instantaneous flow across the aortic valve (arrow) and in the aorta during **A)** systole and **B)** diastole.

[Supplemental motion image is available for Figure 2.](#)

Fig. 3 Photograph shows surgical specimen of the infected, bicuspid aortic valve with leaflet perforation (arrow).

patient's blood cultures remained negative; however, serologic studies subsequently were positive for *Coxiella burnetii*, suggesting a chronic, active infection. The patient was treated with doxycycline and hydroxychloroquine and underwent successful surgical aortic root and valve replacement (Fig. 3).

Comment

Coxiella burnetii (Q Fever) endocarditis is a rare form of endocarditis that can sometimes present as a stroke.¹ Most cases of Q fever occur in individuals who have been exposed to farm animals, including buffalo. Bicuspid aortic valve, the most common congenital heart defect, is often associated with substantial morbidity, including endocarditis and aortic dilation.² Although echocardiography, particularly TEE, typically yields adequate views of a bicuspid aortic valve and aortic root

dilation, CMR affords better spatial and contrast resolution and the ability to obtain images of the entire aorta. In conventional CMR, blood flow is measured by using velocities in 2 spatial dimensions, in individual slices at specific anatomic locations.³ Using the emerging technique of 4D-flow MRI, one can view the entire 3-dimensional volume of the aorta and evaluate the complex forward and regurgitant flow jets, color-coded by velocity, over time.^{3,4} High-velocity, eccentric blood flow in an area of asymmetric aortic dilation or aneurysm suggests increased wall shear stress with high potential for rupture. This combination of complete anatomic visualization and evaluation of pathologic hemodynamic conditions is not feasible with echocardiography or conventional CMR. Accordingly, 4D-flow MRI might best identify those patients who are in need of urgent intervention.⁴ As in our patient's case, difficult clinical presentations often require multiple imaging methods for effective management.

References

1. Raoult D, Marrie T, Mege J. Natural history and pathophysiology of Q fever. *Lancet Infect Dis* 2005;5(4):219-26.
2. Siu SC, Silversides CK. Bicuspid aortic valve disease. *J Am Coll Cardiol* 2010;55(25):2789-800.
3. Markl M, Chan FP, Alley MT, Wedding KL, Draney MT, Elkins CJ, et al. Time-resolved three-dimensional phase-contrast MRI [published erratum appears in *J Magn Reson Imaging* 2003;18(3):396]. *J Magn Reson Imaging* 2003;17(4):499-506.
4. Hope MD, Hope TA, Crook SE, Ordovas KG, Urbania TH, Alley MT, Higgins CB. 4D flow CMR in assessment of valve-related ascending aortic disease. *JACC Cardiovasc Imaging* 2011;4(7):781-7.