Published in final edited form as:

Tetrahedron Lett. 2014 May 1; 55(22): 3328–3331. doi:10.1016/j.tetlet.2014.04.043.

Exploiting the Divalent Nature of Isonitriles: a novel Pictet–Spengler Amidination process

Federico Medda^{a,b} and Christopher Hulme^{a,b,c,*}

^aUniversity of Arizona, College of Pharmacy, 1703 E. Mabel St., Tucson, AZ 85721, USA

^bBIO5 Oro Valley, 1580 E. Hanley Blvd., Oro Valley, AZ 85737, USA

^cUniversity of Arizona, Department of Chemistry and Biochemistry, Tucson, AZ 85721, USA

Abstract

An isocyanide-based multicomponent reaction (IMCR) utilized for the rapid assembly of novel, biologically relevant dihydropyrrolo[1,2-*a*]quinoxalines-amidines is herein presented. Starting from 1-(2-aminophenyl)pyrroles, aldehydes, and isonitriles, the target heterocyclic scaffold is assembled in a one-pot, operationally friendly process. With three points of diversity and formation of three chemical bonds in one step, this strategy proves to be very general. Novel, mild methodology for the generation of amidines from secondary amine anilines and isonitriles is also introduced.

Keywords

Isonitrile; Multicomponent reaction; Amidine; Pictet–Spengler

Multicomponent reactions (MCRs) are powerful transformations where three or more starting materials react in one pot, forming a final product incorporating all atoms of starting materials, with the exception of condensation byproducts, (i.e., H₂O, MeOH, or HCl).¹ The main advantage of MCRs thus resides in their capacity to generate complex chemotypes from simple starting materials, via atom-economic and convergent processes, with high bond-forming efficiency.¹ As such, the last two decades have witnessed renewed interest in MCRs as the need for facile access to libraries of pharmacologically relevant molecules has evolved.^{1,2}

Within the MCR arena, isocyanide-based MCRs (IMCRs) are particularly attractive, due in large part to the chemical versatility of the isocyanide functional group, which is capable of behaving as both a nucleophile and, when converted into a nitrilium ion, as an electrophile. As such, isocyanides are divalent in nature, a property that can be exploited to give rise to a

© 2014 Elsevier Ltd. All rights reserved.

*Corresponding Author Telephone: (520) 626-5322, Fax: (520) 626-0794, hulme@pharmacy.arizona.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

wide range of molecular scaffolds.^{1–3} Traditionally involved in the generation of novel biologically appealing chemotypes via post-condensation modifications of IMCRs,⁴ we recently turned our attention toward study of the Pictet–Spengler reaction, discovered in 1911, and classically defined as the condensation of a β -arylethylamine with a carbonyl compound in the presence of a protic or Lewis acid that gives rise to a substituted tetrahydroisoquinoline.⁵ Subsequently, the scope and utility of this transformation was improved through the employment of novel aryl-amines linked to electron-rich aromatic heterocycles,⁶ resulting in the synthesis of natural products and small molecules of biological interest.⁷

We were specifically drawn to the possibility of disrupting the Pictet–Spengler reaction of 1-(2-aminophenyl)pyrroles **1**, aldehydes **2**, and isonitriles **3** through enticing the pyrrole moiety to trap the intermediate nitrilium ion **4**, to afford heterocycles of structure **5**. However, studies showed the predominant pathway proceeded through the Pictet–Spengler reaction via **6**, which intriguingly was highly susceptible to reaction with protonated isonitriles **7**, affording products of generic structure **8** in a single step. Enticingly, these new products contained three points of diversity, being derived from a novel, albeit unplanned, MCR containing an unexpected amidine functional group (**8**, Scheme 1). Interestingly, reactions of isonitriles with carboxylic acids have been recently reported by Danishefsky and others under mild conditions.⁸ However, reactions of isonitriles with amines affording amidines are known to proceed only via relatively harsh metal catalysis,⁹ and only one report from 1962¹⁰ details a hydrochloric acid-mediated transformation of secondary anilinic amines with isonitriles to amidines which was not exemplified in any great detail. To the best of our knowledge, *para*-toluene sulfonic acid-promoted amidine formation of the intermediate aniline **6** represents a far milder example of this significantly under-utilized functional group transformation. Initial optimization of the IMCR focused on the one-pot reaction between amine **1a**, aldehyde **2a**, and isonitrile **3a** (Table 1) utilizing toluene as a solvent and *para*-toluenesulfonic acid as a catalyst, which resulted in acceptable yields of **8a**, as judged by LC/MS (entries 1–4). Conversely, use of protic solvents saw yields drop significantly (entries 5–6). A similar outcome was also seen using stronger acidic catalysis, including previously reported HCl¹⁰ (entries 7–10), probably due to heightened instability of the isonitrile **3a** in these media, which is prone to formamide formation if adventitious water is present. Indeed, the latter transformation is an excellent way of destroying low molecular weight pungent isonitriles during reaction work-up, again exploiting the inherent “divalent nature” of the isonitrile functional group.

Upon silica-based purification of **8a**, pure product was recovered in only 10% yield (entry 2), suggesting a somewhat surprising instability on prolonged exposure to silica. Encouragingly, reverse-phase (C18) column chromatography enabled a satisfactory increase in yield (**8a**, 62%), and definitive structural confirmation was obtained with a small-molecule X-ray crystal structure (**8a**, Figure 1).

Isolated yields were further improved via microwave irradiation in DCE (Table 1), coupled with shorter reaction times (entries 11 and 12). Evaluation of various acid catalysts under these improved conditions (entries 13–17) showed none to be as effective as *para*-toluene sulfonic acid. Scheme 2 depicts a proposed mechanism where amine **1** and aldehyde **2** react

to generate the expected Pictet–Spengler product **6** via iminium species **9**. Evidence for the intermediary formation of **6** was observed through LC/MS monitoring, and reaction with the protonated isonitrile **7** affords the final exocyclic amidine moiety **8**.

To the best of our knowledge, this is the first example of a Pictet–Spengler reaction embedded in a one-pot IMCR process that is capable of forming three new chemical bonds in one operation with the generation of an exocyclic amidine moiety. These novel scaffolds are likely to possess a range of biological activities.¹¹

With an optimized procedure in hand, reaction scope was explored and twelve analogs (**8a–l**, Figure 2) prepared using three amines (**1a–c**), seven aldehydes (**2a–g**), and seven isonitriles (**3a–g**) as reagents [Note: only three 1-(2-aminophenyl)pyrroles were found to be commercially available]. To our delight, the one pot procedure proved to be general, tolerating the full range of starting materials in high yield (39%–87%). As noted, the Pictet–Spengler intermediate was observed during reaction monitoring, and therefore two Pictet–Spengler intermediates, **6a** and **6b**, were isolated and treated with an isonitrile in the presence of *para*-toluene sulfonic acid at elevated temperatures in toluene. Final products **8a–b** were obtained in 52% and 61% isolated yields respectively (Scheme 3). Moreover, in a preliminary scope study of the amidination process, the indoline **10a** and 1,2,3,4-tetrahydroquinoline **10b** both afforded amidine products **11a–b** in moderate to good yields (Scheme 4). However, non-anilinic amines (piperidine and pyrrolidine) failed to produce the desired amidine, presumably due to their inherent higher pK_a values leading to competitive amine protonation and deactivation.

In summary, we have herein reported a novel, one-pot IMCR that enables the assembly of biologically relevant amidine-containing dihydropyrrolo[1,2-*a*]quinoxalines and additionally shown new utility and method improvement of a rarely used amidination reaction of anilines and isonitriles. Due to the facile and practical production protocol of the IMCR, it is envisioned that access to larger libraries of diverse analogs will be feasible for further biological annotation.

Acknowledgments

Financial support from the National Institutes of Health (Grant P41GM086190) is gratefully acknowledged. Dr. Sue A. Roberts and Mr. Gabriel B. Hall are thanked for small molecule crystal structure determination.

References

1. (a) Zhu, J.; Bienaymé, H., editors. Multicomponent Reactions. Wiley-VCH; Weinheim, Germany: 2005. (b) Dömling A. Chem Rev. 2006; 106:17. [PubMed: 16402771] (c) Dömling A, Wang W, Wang K. Chem Rev. 2012; 112:3083. [PubMed: 22435608] (d) Ganem B. Acc Chem Res. 2009; 42:463–472. [PubMed: 19175315] (e) El Kaim L, Grimaud. Tetrahedron. 2009; 65:2153–2171. (f) Banfi L, Riva R, Basso A. Synlett. 2010; 1:23–41. (g) Van der Heijden G, Ruijter E, Orru RVA. Synlett. 2013; 24:666–685. (h) Brauch S, van Berkel Sander S, Westermann B. Chem Soc Rev. 2013; 42:4948–4962. [PubMed: 23426583]
2. (a) Bienaymé H, Hulme C, Odon G, Schmitt P. Chem-Eur J. 2000; 6:3321. [PubMed: 11039522] (b) Hulme C, Gore V. Curr Med Chem. 2003; 10:51. [PubMed: 12570721] (c) Akritopolou-Zanze I. Curr Opin Chem Biol. 2008; 12:324. [PubMed: 18312861]

3. (a) Ugi I, Werner B, Dömling A. *Molecules*. 2003; 8:53.(b) Dömling A, Ugi I. *Angew Chem Int Ed*. 2000; 39:3168.
4. For recent examples of post-condensation modification of either Ugi or Passerini products, see: Gunawan S, Petit J, Hulme C. *ACS Comb Sci*. 2012; 14:160. [PubMed: 22330239] Gunawan S, Nichol G, Hulme C. *Tetrahedron Lett*. 2012; 53:1664. [PubMed: 22707799] Shaw AY, Medda F, Hulme C. *Tetrahedron Lett*. 2012; 53:1313. [PubMed: 22368307] Xu Z, Shaw AY, Dietrich J, Cappelli AP, Nichol G, Hulme C. *Mol Divers*. 2012; 16:73. [PubMed: 22237832] Shaw AY, McLaren JA, Nichol GS, Hulme C. *Tetrahedron Lett*. 2012; 53:2592. [PubMed: 22544981] Shaw AY, Xu Z, Hulme C. *Tetrahedron Lett*. 2012; 53:1998. [PubMed: 23559684] Medda F, Hulme C. *Tetrahedron Lett*. 2012; 53:5593.
5. (a) Pictet A, Spengler T. *Chem Ber*. 1911; 44:2030.(b) Raines S, Chai S-Y, Palopoli FP. *J Heterocycl Chem*. 1976; 13:711–16.(c) Verma AK, Jha RR, Sankar VK, Aggarwal T, Singh RP, Chandra R. *Eur J Org Chem*. 2011:6998–7010.
6. (a) Kundu B, Sawant D, Chhabra R. *J Comb Chem*. 2005; 7:317. [PubMed: 15762762] (b) Kundu B, Sawant D, Partani P, Kesarwani AP. *J Org Chem*. 2005; 70:4889. [PubMed: 15932339] (c) Duggineni S, Sawant D, Saha B, Kundu B. *Tetrahedron*. 2006; 62:3228.(d) Sharma S, Saha B, Sawant D, Kundu B. *J Comb Chem*. 2007; 9:783. [PubMed: 17569562] (e) Saha B, Sharma D, Sawant D, Kundu B. *Tetrahedron*. 2008; 64:8676.
7. For reviews, see: Royer J, Bonin M, Micouin L. *Chem Rev*. 2004; 104:2311. [PubMed: 15137793] Nielsen TE, Diness F, Meldal M. *Curr Opin Drug Discovery Dev*. 2003; 6:801.Cox ED, Cook J. *Chem Rev*. 1995; 95:1797.Larghi EL, Amongero M, Bracca ABJ, Kauffman TS. *ARKIVOC*. 2005; xii:98.Stöckigt J, Antonchick AP, Wu F, Waldmann E. *Angew Chem Int Ed*. 2011; 50:8558.
8. For recent examples, see: Wilson RM, Stockdill JL, Wu X, Li X, Vadola PA, Park PK, Wang P, Danishefsky SJ. *Angew Chem Int Ed*. 2012; 51:2.Yuan Y, Zhu J, Li X, Wu X, Danishefsky SJ. *Tetrahedron Lett*. 2009; 50:2329. [PubMed: 20161291] Jones GO, Li X, Hayden AE, Houk KN, Danishefsky SJ. *Org Lett*. 2008; 10:4093. [PubMed: 18707109] Li X, Danishefsky SJ. *Nat Protoc*. 2008; 3:1666. [PubMed: 18833204] Hou J, Ajami D, Rebek J Jr. *J Am Chem Soc*. 2008; 130:7810. [PubMed: 18507459] Li X, Yuan Y, Berkowitz WF, Todaro LJ, Danishefsky SJ. *J Am Chem Soc*. 2008; 130:13222. [PubMed: 18783224] Shaabani A, Rezayan AH, Sarvary A, Keshipour S, Khavasi HR. *Tetrahedron Lett*. 2010; 51:4091.
9. For representative examples, see: Lygin AV, de Meijere A. *Eur J Org Chem*. 2009:5138.Saegusa T, Ito K, Kobayashi K, Hirota H, Yoshioka H. *Tetrahedron Lett*. 1966; 7:6121.Saegusa T, Ito K, Hirota K, Yoshioka H. *Bull Chem Soc Jpn*. 1969; 42:3310.Saegusa T, Murase I, Ito K. *J Org Chem*. 1971; 36:2876.Hegarty AF, Chandler A. *Tetrahedron Lett*. 1980; 21:885.Smith R, Livinghouse T. *Synthetic Commun*. 1984; 14:639.Sawai H, Takizawa T. *J Organomet Chem*. 1975; 94:333.
10. (a) Mimin YV. *J Gen Chem USSR (Engl Transl)*. 1962; 32:3867.(b) Mimin YV. *J Gen Chem USSR (Engl Transl)*. 1962; 32:3795.
11. (a) Desplat V, Geneste A, Begorre M, Fabre SB, Brajot S, Massip S, Thiolat D, Mossalayi D, Jarry C, Guillon J. *J Enzym Inhib Med Chem*. 2008; 23:648.(b) Xu H, Fan L. *Eur J Med Chem*. 2011; 46:1919. [PubMed: 21388718] (c) Szabó G, Kiss R, Páyer-Lengyel D, Vukics K, Szikra J, Baki A, Molnár L, Fisher J, Keserü GM. *Bioorg Med Chem Lett*. 2009; 19:3471. [PubMed: 19457667] (g) Albright, JD. US Patent. US 5733905 A 19980331. 1998. (e) Albright, JD. US Patent. US 5736540 A 19980407. 1998. (f) Sum F, Dusza J, Santos ED, Grosu G, Reich M, Du X, Albright JD, Chan P, Coupet J, Ru X. *Bioorg Med Chem Lett*. 2003; 13:2195. [PubMed: 12798333] (g) Albright, JD. US Pat. US 5753648 A 19980519. 1998. (h) Albright, JD. PCT Int Appl. WO 9749707 A1 19971231. 1997. (i) Albright, JD. US Patent. US 5700796 A 19971223. 1997. (j) Bissantz, C. US Pat Appl Publ. US 20100105906 A1 20100429. 2010. (k) Sabatucci, JP. PCT Int Appl. WO 2006068928 A1 20060629. 2006. (l) Ho, GD. PCT Int Appl. WO 2010151595 A1 20101229. 2006. (m) Gardiner, EM. PCT Int Appl. WO 2007008541 A2 20070118. 2007.

Figure 1.
Crystal structure of compound **8a**.

Figure 2.
Structures of congeners **8a-l**.

Scheme 1.

Synthesis of *N*-((4-methylpyrrolo[1,2-*a*]quinoxalin-5(4*H*)-yl)methylene)methanamines **8** via a Pictet–Spengler amidination sequence.

Scheme 2.
Proposed mechanism for the formation of **8**.

Scheme 3.

Two-step preparation of dihydropyrrolo quinoxaline amidines **8a-b**.

Scheme 4.

Reaction of isonitrile **3a** and indolines **10a–b** to afford corresponding amidines **11a–b**.

Table 1

Optimization of the One-pot Model Reaction.^a

entry	solvent	acid (eq.)	conditions	conversion ^b
1	toluene	TsOH (0.2)	rt, 18 h	56
2	toluene	TsOH (0.4)	rt, 18 h	70
3	toluene	TsOH (0.2)	reflux, 18 h	63
4	toluene	TsOH (0.4)	reflux, 18 h	45
5	MeOH	TsOH (0.4)	rt, 18 h	24
6	MeOH	TsOH (0.4)	reflux, 18 h	–
7	AcOH	cHCl ^c	rt, 18 h	21
8	AcOH	–	rt, 18 h	35
9	AcOH	cHCl ^c	reflux, 18 h	–
10	AcOH	–	reflux, 18 h	–
11	DCE	TsOH (0.4)	MW, 120 °C, 30'	97
12	DCE	TsOH (0.4)	MW, 120 °C, 60'	87
13	DCE	CF ₃ CH ₂ OH ^d	MW, 120 °C, 30'	65
14	DCE	AcOH (0.4)	MW, 120 °C, 30'	27
15	DCE	cHCl (0.4)	MW, 120 °C, 30'	53
16	DCE	TFA (0.4)	MW, 120 °C, 30'	46
17	DCE	HClO ₄ (0.4)	MW, 120 °C, 30'	29

^a Starting materials used in a 1:1:1 molar ratio;^b Conversion is expressed as % and was judged by means of LCMS analysis (see supplementary information);^c One drop of 12N HCl used;^d DCE:CF₃CH₂OH (2:1).