Published in final edited form as:

Curr Org Synth. 2013 June 1; 10(3): 366–383. doi:10.2174/1570179411310030003.

Synthetic Studies of Glycosylphosphatidylinositol (GPI) Anchors and GPI-Anchored Peptides, Glycopeptides, and Proteins

Zhongwu Guo

Department of Chemistry, Wayne State University, 5101 Cass Avenue, Detroit, Michigan 48202, USA Phone: +1-313-577-2557 Fax: +1-313-577-8822 zwguo@chem.wayne.edu

Abstract

Glycosylphosphatidylinositol (GPI) anchorage of proteins and glycoproteins onto the cell surface is ubiquitous in eukaryotes, and GPI-anchored proteins and glycoproteins play an important role in many biological processes. To study GPI anchorage and explore the functions of GPIs and GPI-anchored proteins and glycoproteins, it is essential to have access to these molecules in homogeneous and structurally defined forms. This review is focused on the progress that our laboratory has made towards the chemical and chemoenzymatic synthesis of structurally defined GPI anchors and GPI-anchored peptides, glycopeptides, and proteins. Briefly, highly convergent strategies were developed for GPI synthesis and were employed to successfully synthesize a number of GPIs, including those carrying unsaturated lipids and other useful functionalities such as the azido and alkynyl groups. The latter enabled further site-specific modification of GPIs by click chemistry. GPI-linked peptides, glycopeptides, and proteins were prepared by regioselective chemical coupling of properly protected GPIs and peptides/glycopeptides or through site-specific ligation of synthetic GPIs and peptides/glycopeptides/proteins under the influence of sortase A. The investigation of interactions between GPI anchors and pore-forming bacterial toxins by means of synthetic GPI anchors and GPI analogs is also discussed.

Keywords

Glycosylphosphatidylinositol; GPI; Carbohydrate; Peptide; Glycopeptide; Protein; Glycolipid; Sortase

Introduction

Glycosylphosphatidylinositols (GPIs) are a class of complex glycolipids, which anchor surface proteins and glycoproteins to the cell membrane [1,2]. A recent study estimated that 0.5-1.0% of the total proteins encoded in the eukaryotic genome are GPI-anchored [3]. Consequently, GPI attachment to proteins and glycoproteins is one of the most common posttranslational modifications [1,2,4,5]. GPI-anchored proteins and glycoproteins play an essential role in many biological events, such as cell recognition and interaction [6], biological signal transduction [7-10], cell activation [9,10], cell surface enzymatic reaction [11], and bacterial or viral infection [12]. GPI anchors have also been revealed to mediate protein and glycoprotein endocytosis and turnover [7,8,13], and the metabolites of GPIs have been shown to function as secondary messengers in certain hormonal pathways [14,15].

Since the first structural characterization of an intact GPI anchor by Ferguson and coworkers in 1988 [16], more than 30 GPI anchors and a larger number of GPI-anchored proteins and glycoproteins have been isolated and characterized from various eukaryotic species ranging from yeasts and parasites to human [1,2]. Although structurally diverse, all protein-anchoring GPIs identified so far share the highly conserved core structure, H₂Net-(P)-6Man α (1 $\rightarrow$ 2)Man α (1 $\rightarrow$ 6)Man α (1 $\rightarrow$ 4)GlcNH₂ α (1 $\rightarrow$ 6)-*myo*-inositol-1-(P)-lipid (**1**), whereas all GPI-anchored proteins and glycoproteins have their polypeptide C-terminus attached to the phosphoethanolamine [(P)-EtNH₂] moiety at the mannose-III (Man-III) 6-*O*-position of the GPI core structure (Figure 1) [1]. Proteins and glycoproteins are associated with cells through the anchorage of GPI lipids into the cell membrane as depicted in Figure 1. The structural diversity of GPIs is mainly composed of structural variations of the lipid chains and additional phosphoethanolamine moieties and carbohydrate chains attached to Man-I and Man-III residues, respectively.

To study GPIs and GPI-anchored proteins/glycoproteins at the molecular level, it is essential to have access to these biomolecules and related derivatives in homogeneous forms and sufficient quantities. However, this is difficult to achieve via isolation and purification of natural products from biological sources, due to the problem of structural microheterogeneity. As a result, much recent effort has been focused on the development of methodologies for the chemical and chemoenzymatic synthesis of GPIs and GPI-anchored proteins and glycoproteins [17,18].

Ogawa and coworkers reported the first total synthesis of an intact GPI anchor in 1991 [19-21]. Since then, several GPI anchors have been prepared by the Fraser-Reid [22-25], Ley [26,27], Martin-Lomas [28,29], Nikolaev [30], Schmidt [31-34], Seeberger [35-38], and Vishwakarma [39,40] groups, as well as ours [41-44]. Therefore, GPI chemical synthesis has witnessed great progresses in the past two decades. In contrast, the synthesis of GPI-anchored proteins/glycoproteins has made less advancement and remains a formidable challenge. To the best of our knowledge, there have been only a few reported attempts for the synthesis of GPI-linked peptides/proteins. Nakahara and coworkers [45] reported first the synthesis of GPI-peptide conjugates by coupling short peptides with simple GPI analogs through native chemical ligation (NCL), by which Bertozzi and coworkers [46] have successfully ligated a protein and a simple GPI analog. NCL was also used by Seeberger and coworkers [47] to prepare the first GPI-anchored protein with prion protein linked to a GPI anchor. GPI anchors were also coupled with proteins via unnatural linkers to obtain undefined neoglycoprotein mixtures that were studied as GPI-based vaccines [36]. Alternatively, we have [48,49] prepared GPI-linked peptides and glycopeptides by regiospecific coupling of extensively protected peptides/glycopeptides carrying free C-termini with GPIs carrying a free (P)-EtNH₂ moiety, followed by global deprotection. Recently, we [50-52] have developed a chemoenzymatic strategy for GPI-anchored peptide/glycopeptide synthesis, which may be elaborated into a generally useful strategy for accessing GPI-anchored proteins and glycoproteins.

This review focuses on the progress made in our laboratory towards the chemical and chemoenzymatic synthesis of GPI anchors and GPI conjugates. These studies have facilitated the access to GPIs, GPI-linked peptides/glycopeptides/proteins, and other GPI

conjugates to enable the investigation of biological issues related to GPI anchorage, which is also discussed in brief.

Chemical Synthesis of GPIs and Related Derivatives

GPI synthesis is a notable challenge, as it involves multiple branches of organic chemistry, including carbohydrate, lipid, phosphate, and inositol chemistries. To address any potential compatibility concerns, most reported GPI syntheses have adopted a strategy that started with the construction of the GPI skeleton, such as **2**, followed by manipulation of the protecting groups and final-stage installation of the phosphate and phospholipid functionalities at the proper positions (Scheme 1). This strategy has been employed to successfully synthesize a number of GPIs. However, its application to the synthesis of GPI anchor of the human sperm CD52 antigen (Figure 2) was unsuccessful [41], as the late-stage phospholipidation reaction was problematic. Compared to the GPI anchors previously synthesized by the strategy outlined in Scheme 1, the human CD52 GPI anchor contains a large acyl group attached to the inositol 2-*O*-position. This structure not only required the redesign of GPI synthetic strategy but also necessitated the preparation of a unique inositol derivative, which has to have the 1,2,6-*O*-positions differentiated from each other and from the remaining positions of inositol as well.

As depicted in Scheme 2, our first synthesis of an inositol derivative **6** with differentiated 1,2,6-*O*-positions commenced with racemic *myo*-inositol [53]. After *myo*-inositol was transformed into optically pure (+)-**7** and (–)-**7** by a reported procedure involving lipase-mediated enantiomeric resolution [54], the two free hydroxyl groups of (+)-**7** were differentiated upon stannylene acetal-directed regioselective 6-*O*-allylation followed by 1-*O*-*para*-methoxybenzylation. When the resulting product **8** was treated with HCl briefly, the strained *trans*-ketal was selectively removed to produce **9**, which was followed by 4,5-di-*O*-benzylation and removal of the remaining ketal to give **10**. Then, stannylene acetal-directed regioselective 3-*O*-benzylation of **10** afforded **6**, which was obtained from (+)-**7** in 6 separate steps and a 32% overall yield. Meanwhile, (–)-**7** was also utilized to prepare **6** through a series of similar reactions that were implemented in an essentially reversed order. As outlined in Scheme 2, after (–)-**7** was di-*O*-benzylated, the inositol 6-*O*-position was protected with an allyl group upon selective removal of the *trans*-ketal and stannylene acetal-directed regioselective allylation. Thereafter, the remaining ketal was removed, and the 1,2-*O*-positions of **13** were differentiated, again by means of stannylene acetal-directed regioselective alkylation. Thus, **6** was prepared from (–)-**7** in 6 separate steps and a 41% overall yield. The synthetic strategy outlined here is highlighted by the use of both enantiomers of **7** for the preparation of the same synthetic target, which had significantly improved the overall synthetic efficiency, and by repeated application of stannylene acetal-directed regioselective alkylation reactions. A burdensome step of this synthesis is the preparation of (+)-**7** and (–)-**7** from *myo*-inositol.

Recently, we developed another synthetic method for 1,2,6-*O*-differentiated inositol derivatives, starting with methyl α -D-glucopyranoside **14** (Scheme 3) [55], in which a key step was the stereospecific Ferrier rearrangement of **15** to produce **16** [56,57]. Thereafter, the carbonyl group of **16** was stereospecifically reduced with NaBH(OAc)₃, followed by 1-

O-deacetylation to afford **18**. Subsequently, two routes were established for the differentiation of the free hydroxyl groups in **18**. One route was to protect the 1,2-*cis*-diol with an isopropylidene ketal, followed by 6-*O*-alkylation and deketalation to give **13**. Then, the 1,2-*O*-positions were distinguished by stannylene acetal-directed regioselective alkylation to give **6**. Alternatively, the 1,2-*cis*-diol functionality was protected with a *p*-methoxybenzylidene group, which gave diastereomers **21** and **22**. After the free hydroxyl group in **21** and **22** was allylated, the acetal ring of **23** and **24** was regioselectively opened with NaBH₃CN under acidic condition to afford **25** and **6**, respectively. Both **6** and **25**, which were prepared from **14** in 10-12 steps and 13-19% overall yields, are useful for the synthesis of GPI anchors containing a 2-*O*-acylated inositol residue.

In all GPIs, the glucosamine (GlcNH₂) residue is linked to the inositol residue through an α -glycosidic linkage, which is relatively difficult to form. To get α -glycosylation, the glucosamine 2-*N*-position must be protected by a non-participating protecting group; otherwise, the glycosylation reaction would result in β -glycosides exclusively. Therefore, in GPI synthesis the most commonly used protecting tactic for GlcNH₂ is to convert the 2-amino group to an azido group, which can be readily and selectively converted to a free amino group under mild reductive conditions later on. Our group has developed effective procedures for the synthesis of 2-azido-2-deoxy-glucose derivatives as glycosyl donors.

Our first synthesis of a glucosamine derivative **31** (Scheme 4) useful for GPI synthesis started from commercially available D-glucal [53]. After D-glucal was transformed into Cerny epoxide **26** via a one-pot protocol [58], its epoxide ring was opened with sodium azide, which was followed by conventional *O*-benzylation of the resulting alcohol to give **27** in excellent overall yields. Regioselective opening of the strained 5-membered ring in **27** through acetolysis under acidic conditions afforded diacetate **28** as an anomeric mixture (α : β = 3.5:1), of which the anomeric position was regioselectively deprotected with benzylamine to yield hemiacetal **29**. Treatment of **29** with DAST was followed by replacement of the acetyl group at the 6-*O*-position with a benzyl group to eventually produce glycosyl fluoride **31** as an inseparable anomeric mixture (α : β = 1:6), which was directly used as a glycosyl donor.

For the synthesis of GPI anchors having a palmitoyl group at the inositol 2-*O*-position, the traditional strategy outlined in Scheme 1 was problematic [41]. To deal with this issue, we developed a new strategy with phospholipidated pseudodisaccharide **32** as the key intermediate. The synthesis of **32** is outlined in Scheme 5 [41]. After a palmitoyl group was attached to the 2-*O*-position of **6** to form **33**, the 6-*O*-position was deprotected on PdCl₂-catalyzed deallylation. Glycosylation of **34** by **31** with dichlorohafnocene/silver perchlorate as the promoter gave an anomeric mixture (α : β = 4:3), slightly in favor of the desired α -anomer **35**, which was readily separable by flash column chromatography. The removal of the PMB group in **35** by cerium ammonium nitrate (CAN) oxidation was followed by phospholipidation to produce an inseparable diastereoisomeric mixture **38**. Finally, the allyl group of **38** was removed to afford **32**, which was utilized to assemble GPI anchors.

For the final assembly of sperm CD52 GPI anchor (Scheme 6) [41], **32** was glycosylated by trimannosyl donor **39** with *N*-iodosuccinimide (NIS)/triflic acid (TfOH) as the promoter.

After selective deprotection of the product to expose the nonreducing end 6-*O*-position, the diastereoisomeric phosphates were separable by column chromatography. Next, a phosphoethanolamine moiety was attached to the free hydroxyl group of **40** to produce a 1:1 diastereoisomeric mixture of phosphate **42** in a good yield (74%). Finally, global deprotection of **42** was achieved in two steps, including removal of the Fmoc and cyanoethyl groups by DBU and removal of all benzyl groups with the concomitant reduction of the azido group by hydrogenation using 10% Pd/C as the catalyst, to afford sperm CD52 GPI anchor **43**. This work represented the first chemical synthesis of a GPI anchor having a large acyl chain linked to the inositol 2-*O*-position.

Evidently, elongating the carbohydrate chain of **32** with **39** led to the GPI backbone **40** in one step. We expect this new and highly convergent strategy to be widely useful for the synthesis of various GPIs and GPI derivatives. As a proof, the strategy was successfully utilized to rapidly assemble a number of GPIs and GPI analogs, such as **45-49** shown in Scheme 7 [59], including a fully phosphorylated GPI anchor **49** of the sperm CD52 antigen [42]. Another advantage of this new synthetic strategy is that the relatively difficult reactions involved in GPI synthesis, such as the glycosylation reaction to install the α -linked glucosamine unit and the reaction to introduce the phospholipid moiety, are realized at an early stage of the synthesis, which has improved the overall synthetic efficiency.

Chemical Synthesis of Functionalized GPIs and GPI Derivatives

To probe the biological relevance of the structural variants of GPI anchors and to better understand the functions of GPI anchorage, it is necessary to have access to functionalized GPIs with homogeneous and defined structures. Although strategies have been developed for the synthesis of GPI anchors as mentioned above, limitations associated with these strategies prevent the inclusion of certain combinations of useful functionalities. For example, most of the reported GPI syntheses utilized the benzyl group as a permanent hydroxyl protection, which prevents the incorporation of alkene, alkyne, azide, sulfide, thiol, and other functionalities intolerant to Pd-catalyzed hydrogenolysis. To address this issue, Nikolaev and coworkers [30] employed the benzoyl group as a permanent hydroxyl protection in the synthesis of GPIs containing unsaturated lipids. However, this strategy may not be applicable to target molecules that contain other acyl groups or peptides/glycopeptides due to their sensitivity to sodium methoxide employed for global deprotection. For example, the reported final deacylation step gave only 40% yield [30].

We established an alternative strategy for functionalized GPI synthesis employing the PMB group for permanent hydroxyl protection [43]. PMB ethers can be cleaved under mild acidic or oxidative conditions, which we anticipated would provide the flexibility for the incorporation of a range of functional groups in the target molecule. This methodology was applied to the synthesis of GPI anchors bearing unsaturated lipid chains and other functionalities.

In this process, we developed methods for the synthesis of involved key intermediates, such as 1,6-*O*-differentiated inositol derivative **50** (Scheme 8) [43]. After the free hydroxyl groups of racemic ($\pm$)-**7** were protected as allyl ethers, the ketals in resultant ($\pm$)-**51** were

methanolized with HCl as catalyst, which was followed by methoxybenzylation. Deallylation of resultant ($\pm$)-**53** by Cha's method [60] gave diol ($\pm$)-**54**, of which the 1-*O*- and 6-*O*-positions were differentiated by the stannylene acetal-directed regioselective 1-*O*-allylation. The resulting enantiomers ($\pm$)-**50** were resolved with its free 6-*O*-position as an anchoring point for the resolution reagent. Consequently, ($\pm$)-**50** was esterified with (1*S*)-(-)-camphanic chloride, and the diastereomeric products were separated by HPLC to afford optically pure (-)-**55** that was saponified to yield **50** (44%) in two steps, including the enantiomeric resolution, of which the maximum yield is 50%.

PMB-protected trimannose fragment **63** was assembled from monosaccharide building blocks **56**, **57**, and **59** (Scheme 9) [43]. Glycosylation of **57** by imidate **56** with TMSOTf as the promoter generated only the α -disaccharide, which was deacetylated with $K_2CO_3/MeOH$ to afford **58**. This product was glycosylated by **59** under the same condition to give trimannose **60** in a 5:1 α/β ratio, which was improved to α -only (76%) after the switch of solvent to Et_2O . The allyl group of **60** was then replaced with a triethylsilyl (TES) group upon Cha's deallylation and silylation, to avoid potential interference with unsaturated lipids introduced later on. Finally, **62** was converted to imidate **63** upon NIS/AgOTf-promoted hydrolysis of the thioacetal followed by treatment of the resulting hemiacetal with Cl_3CCN and DBU.

The target GPI anchor **72** carrying unsaturated lipids was finally assembled by the highly convergent strategy that was discussed above, and again the azido group was used to protect the glucosamine 2-amino group (Scheme 10) [43]. Glycosylation of **50** by imidate **64** with TMSOTf as the promoter afforded an anomeric pseudodisaccharide mixture (1.2:1.0), slightly in favor of the desired α -isomer **65**. Deallylation of **65** with the Ir-complex/ $HgCl_2$ protocol was carried out to expose the inositol 1-*O*-position, to which the unsaturated phospholipid was installed via reaction with freshly prepared phosphoramidite **67** under the influence of tetrazole and *in situ* oxidation with *tert*-butyl hydroperoxide. The 4-*O*-*tert*-butyldimethylsilyl (TBS) group was removed with $Et_3N \cdot 3HF$ to give the key intermediate **68**. Pseudodisaccharide **68** reacted with trimannose **63** smoothly to afford the desired α -product, which was deprotected with $Et_3N \cdot 3HF$ to remove the TBS and TES groups. Regioselective introduction of the phosphoethanolamine group to the Man-III 6-*O*-position of **69** was realized by reaction with phosphoramidite **70** for a short period (1 h) followed by oxidation to give **71**. The target GPI **72** was obtained in an 81% yield upon global deprotection of **71** using a one-pot three-step protocol, that is, (i) Zn-mediated reduction of the azido group; (ii) removal of the base-labile Fmoc and cyanoethoxy groups with DBU; (iii) hydrolysis of all PMB ethers with 10% TFA.

We anticipated that the synthetic strategy using the PMB group for permanent hydroxyl protection should be generally applicable to other GPI anchor, carbohydrate, and glycoconjugate syntheses. This was demonstrated with synthesis of a natural GPI anchor **73** of the human lymphocyte CD52 antigen, which carries a sensitive polyunsaturated lipid chain, and functionalized GPIs **74** and **75**, which contain an alkyne and an azido group, respectively (Figure 3) [44,61]. Further elaboration of GPIs **74** and **75**, as well as their coupling with various biomolecules, can be readily achieved by click chemistry. For

example, fluorescein and biotin as imaging and affinity probes were effectively attached to **74** and **75** via Cu(I)-catalyzed and Cu-free strain-promoted click reactions to result in GPI-fluoro conjugate **76** and GPI-biotin conjugate **77**, respectively. Functionalized GPI anchors **74** and **75**, GPI conjugates **76** and **77**, and their derivatives form a set of useful tools for biological studies of GPI anchorage.

GPI **73** was prepared (Scheme 11) by the same strategy as that for **72**, with monosaccharides **78-81** and phospholipidated pseudodisaccharide **83** as the key building blocks. In brief, after the tetramannosyl donor **82** was assembled via a series of glycosylation reactions, it was coupled to **83** by Schmidt glycosylation to afford the GPI skeleton **84** in a good yield (>39%). Thereafter, the 6-*O*- and 2-*O*-positions of Man-III and Man-I, respectively, were simultaneously deprotected with HF and phosphorylated to yield **84**, which was globally deprotected by the three step-one pot protocol as shown in Scheme 10 to finally afford the target GPI **73** in an excellent yield (78%). The deprotection procedure did not affect the sensitive polyunsaturated lipid in **73**. Functionalized GPI **74** was prepared by the same scheme but with **86** and **87** (Figure 4) as the key intermediates.

For the synthesis of **75**, because of the presence of an azido group in the synthetic target, the azido sugar residue as a protected form of glucosamine cannot be carried out to the final stage of the synthesis. Therefore, after pseudodisaccharide **66** was obtained, its azido group was immediately transformed into an amine and protected with an Fmoc group that can be easily removed under mild basic conditions later on. The resultant **88** was then phospholipidated and regioselectively deprotected to produce **89** (Scheme 12). Glycosylation of **89** by trimannosyl donor **91** and deprotection of the product gave **92**. The target molecule **75** was finally obtained upon phosphorylation of **92** followed by global deprotection.

Chemical Synthesis of GPI-Anchored Peptides and Glycopeptides

As mentioned above, all of the identified GPI-anchored proteins/glycoproteins have the polypeptide C-terminus linked to the phosphoethanolamine moiety at the GPI core glycan non-reducing end. Therefore, it is theoretically possible to develop a broadly applicable method for GPI-anchored protein/glycoprotein synthesis through site-specific conjugation of synthetic GPIs and proteins/glycoproteins. In this regard, conventional chemical coupling reactions may be used, so long as GPIs and peptides/glycopeptides are properly protected to guarantee site-specific coupling [48,49]. For GPI conjugation with full-size proteins, however, native chemical ligation (NCL) may be more suitable [46,47]. Alternatively, proteins and glycoproteins may be coupled with GPIs through specific and flexible enzymatic reactions [50-52]. To develop a practical method for the synthesis of GPI-anchored proteins and glycoproteins, we have explored both chemical and enzymatic methods for the conjugation of GPIs and peptides or proteins.

First, we used a model GPI and a short peptide to probe the coupling reactions of protected GPIs and peptides (Scheme 13) [49]. It was shown that the reaction between **94** and activated peptide **95** was effective and racemization-free and that global deprotection of coupling product **96**, including treatment with base to remove the Fmoc group and Pd-catalyzed hydrogenolysis to remove the CBz group and all of the benzyl groups, was

straightforward and effective. This work has demonstrated that the conventional chemical methods may be used to synthesize GPI-anchored peptides and glycopeptides.

This strategy was then employed to synthesize a complex GPI-anchored glycopeptide **98** (Figure 5). An extensively protected glycopeptide **99** carrying the entire peptide sequence of the human CD52 antigen[48] and an acid-labile *N*-linked fucosylated hexasaccharide was the key synthetic intermediate. The peptide *C*-terminus of **99** was free to facilitate its regiospecific coupling with GPIs.

Glycopeptide **99** was prepared by solid-phase peptide synthesis (SPPS) with acid-sensitive 2-chlorotrityl resin as solid-phase support, which enabled the retrieval of the target glycopeptide from the resin under very mild acidic conditions that did not affect the fucosyl linkage or the peptide side chain protections [62]. SPPS was performed on an automatic peptide synthesizer using conventional Fmoc-amino acid chemistry with commercial Fmoc-amino acids and the fully synthetic hexasaccharide-Asn conjugate **101** as building blocks (Scheme 14). Glycopeptide **99** was successfully retrieved from the solid-phase support by 10% HOAc in trifluoroethanol (TFE) and dichloromethane (1:8).

The fucosylated hexasaccharide-Asn conjugate **101** was prepared according to a procedure outlined in Scheme 15 [63]. The difficult β -mannosylation reaction involved was achieved by the Crich protocol [64], which gave **105** in an excellent yield (92%) and stereoselectivity ($\beta/\alpha = 5/1$). Then, **105** was transformed into trichloroacetimidate **106** upon protecting group manipulations and anomeric trichloroacetimidation. The oligosaccharide was further elaborated with **107** and **109** as glycosyl donors to give **110**, the common core pentasaccharide of *N*-glycans. The subsequent α -fucosylation was achieved by the Lemieux protocol [65], which resulted in the α -anomer only, followed by protecting group manipulation to form **112** with an azido group attached to the reducing end to facilitate coupling with amino acids. After the azido group in **112** was selectively reduced under a hydrogen atmosphere in the presence of Lindlar catalyst and subsequent *in situ* coupling of the resultant glycosyl amine to aspartic acid via reacting with activated ester **113**, the amino acid carboxylic group was deprotected under mild acidic conditions to produce **101**.

GPI analog **114** used to couple with **99** was prepared by the same strategy as that of **43** (Schemes 5 and 6) [41]. The coupling reaction between **99** and **114** was achieved in DCM/*N*-methylpyrrolidinone (NMP) (2:1) under a N_2 atmosphere using 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (EDC) and HOBT as condensation reagents to get **115** (70% Scheme 16). The reaction did not result in observable racemization. Global deprotection of **115** was performed in two steps, i.e., hydrogenolytic debenzoylation in a H_2 atmosphere with 10% Pd/C as the catalyst and removal of the amino acid side chain protecting groups with 15% trifluoroacetic acid (TFA). This deprotection protocol offered an excellent overall yield (85%) of the target **98**. This work represented the first chemical synthesis of a complex GPI-linked glycopeptide.

Chemoenzymatic Synthesis of GPI-Anchored Peptides and Glycopeptides

Though preparing GPI-anchored peptides/glycopeptides via chemical ligation of properly protected GPIs and peptides/glycopeptides was proved feasible, its application to full-size proteins and glycoproteins can be difficult. To address this issue, we explored a novel strategy that is based on enzymatic coupling of peptides/proteins with GPIs for GPI-anchored peptide/protein synthesis. A chemoenzymatic strategy can be broadly useful because enzymes are especially suitable for dealing with large biomolecules, including full-size proteins and complex GPI anchors.

In nature, GPI attachment to proteins is a posttranslational modification, accomplished by enzymatic ligation of biosynthetic GPIs and proteins in the endoplasmic reticulum (ER) lumen via a process known as GPI transamidation (Figure 6) [66,67], and the enzyme involved is GPI transamidase (GPI-T) composed of 5 subunits. It was further revealed that GPI transamidation is regulated by a GPI attachment signal, a short peptide near the C-terminus of target proteins destined for GPI addition, and that GPI transamidation is independent of GPI and protein biosynthesis [68]. GPI-T recognizes the GPI attachment signal, cleaves the peptide bond between the target protein and the ω -amino acid of the signal peptide to form a reactive thioester with the target protein, and finally transfers the acyl group to the P-EtNH₂ at the non-reducing end of the GPI core to afford GPI-anchored protein (Figure 6) [68].

Theoretically, GPI-T may be utilized to perform GPI-protein ligation *in vitro* for the synthesis of GPI-anchored proteins. However, GPI-T is a membrane-associated enzyme and consisted of at least 5 different subunits, and several of the subunits have multiple transmembrane domains [69,70]. Therefore, it can be very challenging to obtain GPI-T by recombinant method. To identify a proper enzyme for GPI-protein ligation, we became interested in sortases, a family of transpeptidases found in gram-positive bacteria. Sortases are used by bacteria to anchor surface proteins to cell wall peptidoglycans [71]. Their functions and functional mechanisms are similar to that of GPI-T. As illustrated in Figure 7, sortase also recognizes a unique peptide sequence, known as the sorting signal, at the target protein C-terminus, cleaves a specific peptide bond of this signal, and then links the target protein to a nucleophilic amino group. For example, sortase A (SrtA) of the *Staphylococcus aureus* origin recognizes a pentapeptide sequence, LPXTG where X can be D, E, A, N, Q, or K [72], cleaves the peptide bond between T and G, and finally links the carboxyl group of T to the N-terminal glycine of bacterial cell wall peptidoglycans. Most importantly, compared to GPI-T, the structure of sortases is very simple. Sortases are single-unit proteins containing usually only one transmembrane domain. Therefore, recombinant sortases are much more easily available than GPI-T.

Moreover, literature results have shown that SrtA is quite promiscuous [73-77]. Therefore, we expected that SrtA might be able to accept GPIs as substrates and used to couple proteins and glycoproteins to GPIs for the development of a general strategy for GPI-anchored protein and glycoprotein synthesis, as outlined in Scheme 17 [50]. To be accepted by a sortase, peptides, glycopeptides, proteins or glycoproteins (**116**) used must contain the sorting signal, such as LPXTG, which can be readily achieved during their chemical

synthesis or recombinant expression. However, a worth-mentioning issue is that although the resulting GPI conjugates have GPIs linked to the polypeptide C-terminus via a native peptide linkage, they will differ from natural products in that a short peptide, namely the sorting signal, is inserted between the GPI anchor and the target protein. How this structural difference may affect the properties and bioactivities of GPI-anchored proteins and glycoproteins is an interesting question worthy of investigation.

To prove the synthetic strategy outlined in Scheme 17 and establish the appropriate enzymatic reaction conditions, we first tested the ligation of short peptides and GPI analogs [50]. The enzyme used in this work was a water-soluble variant of the wild-type SrtA, containing only the extracellular sequence, which was prepared by recombinant method using an *E. coli* expression vector. We found that GPI anchor itself was not a good SrtA substrate but by attaching one or two glycine residues to the phosphoethanolamine moiety, the resulting GPI analogs were good SrtA substrates. As a result, under the influence of SrtA, GPI analogs **120** and **122** were efficiently coupled to peptide **119** to produce conjugation products **121** and **122**, respectively. We further observed that the reaction yield was dependent upon the enzyme concentration. For example, in the presence of 6.8 μM and 30 μM of SrtA, the conjugation products were isolated in 45% and 95% yields, respectively (Scheme 18). In addition, we observed that GPI analogs **120** and **122** did not display significantly different reactivity as SrtA substrates for the reaction with **119**.

Next, we examined the SrtA-catalyzed coupling of **120** with peptides **124-126** having the sorting signal at different locations (Scheme 19) [51]. Both **125** and **126** were accepted by SrtA to yield coupling products **127** (61%) **128** (78%), respectively, but peptide **124** with the sorting signal directly exposed at the peptide C-terminus was not a good substrate. These results suggest that, for the sorting signal to be efficiently recognized by and react with SrtA, it should not be exposed at the peptide C-terminus.

SrtA was then used to prepare analogs of the human CD52 antigen (Scheme 20) [51]. For this purpose, **129** and **130** carrying one and two Gly residues at the nonreducing end, respectively, were compared for coupling with CD52 peptide **125**. Both reactions were fast and afforded **132** and **133** in similar yields (78% and 81%, respectively). These studies showed again that GPI analogs carrying one and two Gly residues were similar as peptide acceptors for SrtA-catalyzed transpeptidation. Interestingly, the more sterically hindered trisaccharide GPI analogs **129** and **130** showed more effective reactions than the monosaccharide analog **120**, suggesting that SrtA may accept more complex GPIs. Finally, **129** was effectively coupled to a small protein **133** to afford the human CD24 antigen analog **134** in a 73% yield (Scheme 20).

We found that SrtA could also accept sterically hindered glycopeptides for ligation with GPI analogs to yield GPI-linked glycopeptides (Scheme 21) [52]. Glycopeptide **118** contained a partial sequence of MUC1 and three O-linked glycans, one linked to Thr immediately next to the sorting signal. It was achieved by SPPS and standard Fmoc chemistry with commercial Fmoc amino acids and synthetic glycosyl amino acids Fmoc-Ser(Ac α GalNAc)-OH and Fmoc-Thr(Ac α GalNAc)-OH [78] as building blocks. After the synthetic glycopeptide was detached from the solid-phase support, its glycans were

deprotected with NaOMe to give **135**. The reactions of **135** with **120** and **129** in the presence of SrtA went smoothly to form **136** and **137** in good yields (76% and 84%, respectively). We also proved that the peptide chain of the resulting GPI-linked glycopeptide could be further elongated to generate more complex GPI-linked glycopeptides. For example, after **136** was treated with DBU to disclose its *N*-terminal Gly residue, it could be further elongated with **135** in the presence of SrtA to afford **138** in a 59% yield, as well as a small amount of the corresponding *N*-terminal deprotected product.

In summary, we have demonstrated that GPI derivatives with one or two Gly residues attached to the phosphoethanolamine moiety at the core glycan non-reducing end are excellent SrtA substrates and can be effectively ligated with peptides, small proteins, and complex glycopeptides under the influence of SrtA. This enzymatic ligation method has been used to successfully prepare several analogs of the human CD52 and CD24 antigens and a short analog of GPI-anchored MUC1. Our effort is currently focused on the study of this strategy for the synthesis of more complex structures, including SrtA-catalyzed ligation of intact GPI anchors with full-size proteins and glycoproteins. Based on our present results, as well as literature reports that SrtA accepts large proteins and various sterically hindered nucleophiles [73-77], we are confident that SrtA can be utilized to ligate more complex GPI anchors and proteins/glycoproteins, thus to establish a practical, efficient, and generally useful method for the synthesis of GPI-anchored peptides, glycopeptides, proteins, and glycoproteins.

Biological Studies Using Synthetic GPIs and GPI Derivatives

Many pathogenic bacteria secrete toxins as the major virulent factors, and a significant fraction of these toxins exhibit cytotoxicity via the formation of pores in the host cell membrane [79], which would break the cell permeability barrier to prompt harmful ion exchange and other lethal damages [80]. It has also been revealed that some pore-forming toxins need to bind to GPIs or GPI-anchored proteins/glycoproteins on the host cell surface to assist their pore formation and membrane insertion processes [79-82]. Among a number of established GPI-dependent pore-forming bacterial toxins, aerolysin secreted by *Aeromonas* [83] and CAMP factor secreted by *Streptococcus agalactiae* [84] are the most extensively studied.

Although it has been generally established that aerolysin-type toxins interact mainly with GPIs on the host cell surface, how these toxins bind to GPIs and what structural motifs of GPIs are necessary for the binding are still unclear. Clarification of these questions is important not only for gaining insights into the functional mechanisms of these toxins, but also for the design and development of novel antimicrobial strategies. For this purpose, it is necessary to have access to structurally defined GPIs and GPI analogs and systematic information about the structure-activity relationships of GPIs as the binding partners of pore-forming toxins. In this regard, our synthetic GPIs and GPI derivatives can be very useful [85].

To study the interactions between GPI anchors and toxins, we prepared a number of GPI derivatives (Scheme 7) and, in collaboration with Xiangqun Zeng, evaluated their ability to

bind to CAMP factor by means of quartz crystal microbalance (QCM) [59]. We found that CAMP factor exhibited specific and high-affinity binding to both intact GPI anchors and partial structures of the GPI core structure. Moreover, we found that the intact and fully phosphorylated GPI anchor **50** and the pseudodisaccharide fragment **46** of the GPI core structure had essentially the same bonding affinity to CAMP factor.

In collaboration with Michael Palmer, we also evaluated the influence of synthetic GPI analogs on the cytotoxicity of CAMP factor [86]. It was shown that **139**, a GPI analog without the phospholipid but with a palmitoyl group and a dipeptide linked to the inositol 2-*O*-position and the phosphoethanolamine moiety at the glycan non-reducing end, respectively, had obvious and concentration-dependent inhibition on CAMP factor-mediated lysis of sheep erythrocytes (Figure 8). At the concentration of 4 μ M, **139** could completely protect cells from CAMP factor-induced lysis. The results suggest specific binding of **139** to CAMP factor, which might have resulted in the inhibition of CAMP factor interaction with cell surface GPI anchors.

Both the binding assays [59] and the hemolysis inhibition study [86] have validated the strong and specific binding of CAMP factor to GPIs and GPI analogs. The observation that a GPI fragment **46** had essentially the same binding affinity as the intact GPI anchor **50** suggests that CAMP factor may interact mainly with the pseudodisaccharide and/or the phospholipid moieties of GPI anchors. The observation that GPI analog **139**, which does not contain the phospholipid moiety, could inhibit the interaction between CAMP factor and cells suggests that the phospholipid moiety may not be involved in the binding. Consequently, we proposed that the GlcNH₂ α (1 $\rightarrow$ 6)-inositol motif might be key for the GPI-CAMP factor interaction. These studies have laid the foundation for the design and development of novel inhibitors for CAMP factor and similar pore-forming toxins.

Conclusion

In the past decade, our laboratory has been focused on the development of synthetic methods for GPIs, functionalized GPIs and GPI conjugates, and GPI-anchored peptides, glycopeptides, and proteins. First, we have established a new and highly convergent strategy for GPI synthesis, which was utilized to chemically achieve for the first time a GPI anchor having a large acyl group linked to the inositol 2-*O*-position, as well as several other GPIs and GPI analogs. Second, we have established a method for the chemical synthesis of GPI anchors carrying unsaturated lipids and other functionalities incompatible with the conditions used to deprotect benzyl ethers and acyl esters by using the PMB group for global hydroxyl protection. Third, the PMB strategy was used to synthesize functionalized GPIs enable further modifications by click reaction for the easy installation of molecular tags, such as imaging and affinity probes, for biological studies. Forth, we have explored the synthesis of GPI-anchored peptides/glycopeptides by both chemical and enzymatic ligation. These studies have significantly facilitated the access to GPI anchors and related molecules, which enabled us to investigate some biological problems, such as the interaction between GPI anchors and pore-forming bacterial toxins.

Currently, our laboratory is focused on the synthesis of various biofunctionalized GPI anchors, as well as development and optimization of chemoenzymatic methods for the synthesis of GPI-anchored proteins and glycoproteins. In the meanwhile, we are also interested in the application of the synthetic GPIs and GPI conjugates to chemical biology, such as analyzing GPI-anchored proteomics, investigating the arrangement, shedding and turnover processes, and other properties of GPI anchors and GPI-anchored proteins in the cell membrane, etc. These studies will help gain insights into GPI anchorage and understand the functions and functional mechanisms of GPI anchors and GPI-anchored proteins and glycoproteins. They are not only our future research directions but also very important topics in GPI research that have attracted interests from many other research groups.

Acknowledgments

The author gratefully acknowledges his students and research associates, whose names are in the cited references, as well as his collaborators mentioned in the text, for their dedication and great contributions in various aspects to this research. The author also thanks National Science Foundation (CHE-0320878, 0715275, and 1053848), National Institutes of Health (R01 GM090270 and CA95142), Case Western Reserve University, and Wayne State University for generous financial supports.

Biography

Zhongwu Guo received his Ph.D. degree in organic chemistry from the Institute of Organic Chemistry, Polish Academy of Sciences with Prof. Aleksander Zamojski in 1991 and his postdoctoral training in glycoscience at Shanghai Institute of Organic Chemistry (SIOC), Chinese Academy of Sciences. He was appointed Assistant and Associate Professor at SIOC 1994-1996, RIKEN Fellow at RIKEN, Japan 1996-1997, Assistant Research Officer at National Research Council of Canada 1997-1999, and Assistant and Associate Professor at Case Western Reserve University 1999-2005. He has been Professor of Chemistry at Wayne State University since 2005. Currently, he is also serving as the Editor-in-Chief of the *Journal of Carbohydrate Chemistry*. His research interests are focused on carbohydrate chemistry, glycobiology, and medicinal chemistry, including the development of novel methodologies for complex carbohydrate and glycoconjugate synthesis and the application of synthetic glycoconjugates to the study of biological problems and to the development of new therapeutic strategies for various diseases.

REFERENCES

1. Ferguson MAJ, Williams AF. Cell-surface anchoring of proteins via glycosylphosphatidylinositol structure. *Ann. Rev. Biochem.* 1988; 57:285–320. [PubMed: 3052274]
2. Englund PT. The structure and biosynthesis of glycosyl phosphatidylinositol protein anchors. *Ann. Rev. Biochem.* 1993; 62:121–138. [PubMed: 8352586]
3. Eisenhaber B, Bork P, Eisenhaber F. Post-translational GPI lipid anchor modification of proteins in kingdoms of life: analysis of protein sequence data from complete genomes. *Protein Eng.* 2001; 14:17–25. [PubMed: 11287675]
4. Paulick MG, Bertozzi CR. The glycosylphosphatidylinositol anchor: A complex membrane-anchoring structure for proteins. *Biochemistry.* 2008; 47:6991–7000. [PubMed: 18557633]
5. Udenfriend S, Kodukula K. How glycosylphosphatidylinositol-anchored membrane protein are made. *Ann. Rev. Biochem.* 1995; 64:563–591. [PubMed: 7574493]

6. He H-T, Finne J, Goridis C. Biosynthesis, membrane association and release of N-CAM-120, a phosphatidylinositol-linked form of the neuron cell adhesion molecule. *J. Cell Biol.* 1987; 105:2489–2500. [PubMed: 3693391]
7. Robinson PJ. Signal transduction by GPI-anchored membrane proteins. *Cell Biol. Intern. Rep.* 1991; 15:761–767.
8. Eardley DD, Koshland ME. Glycosylphosphatidylinositol: a candidate system for IL-2 signal transduction. *Science.* 1991; 251:78–81. [PubMed: 1824727]
9. Robinson PJ, Millrain M, Antoniou J, Simpson E, Mellor AL. A glycosphospholipid anchor is required for Qa-2-mediated T cell activation. *Nature.* 1989; 342:85–87. [PubMed: 2530453]
10. Ilangumaran S, He H-T, Hoessli DC. Microdomains in lymphocyte signalling: beyond GPI-anchored proteins. *Immunol. Today.* 2000; 21:2–7. [PubMed: 10637551]
11. Hwa K. Glycosyl phosphatidylinositol-linked glycoconjugates: structure, biosynthesis and function. *Adv. Exp. Med. Biology.* 2001; 491:207–214.
12. Cross GAM. Cellular and genetic aspects of antigenic variation in trypanosomes. *Ann. Rev. Immunol.* 1990; 8:83–110. [PubMed: 2188680]
13. Thomas JR, Dwek RA, Rademacher TW. Structure, biosynthesis, and function of glycosylphosphatidylinositols. *Biochemistry.* 1990; 29:5413–5422. [PubMed: 2143679]
14. Field MC. Is there evidence for phospho-oligosaccharides as insulin mediators? *Glycobiology.* 1997; 7:161–168. [PubMed: 9134421]
15. Gaulton GN, Pratt JC. Glycosylated phosphatidylinositol molecules as second messengers. *Seminars Immunol.* 1994; 6:97–104.
16. Ferguson MAJ, Homans SW, Dwek RA, Rademacher TW. Glycosyl-phosphatidylinositol moiety that anchors *Trypanosoma brucei* variant surface glycoprotein to the membrane. *Science.* 1988; 239:753–759. [PubMed: 3340856]
17. Guo Z, Bishop L. Chemical synthesis of GPIs and GPI-anchored glycopeptides. *Euro. J. Org. Chem.* 2004;3585–3596.
18. Nikolaev AV, Al-Maharik N. Synthetic glycosylphosphatidylinositol (GPI) anchors: How these complex molecules have been made. *Nat. Prod. Rep.* 2011; 28:970–1020. [PubMed: 21448495]
19. Murakata C, Ogawa T. A total synthesis of GPI anchor of *Trypanosoma brucei*. *Tetrahedron Lett.* 1991; 32:671–674.
20. Murakata C, Ogawa T. Stereoselective total synthesis of the glycosylphosphatidylinositol (GPI) anchor of *Trypanosoma brucei*. *Carbohydr. Res.* 1992; 235:95–114. [PubMed: 1473115]
21. Murakata C, Ogawa T. Stereoselective synthesis of glycobiosyl phosphatidylinositol, a part structure of the glycosylphosphatidylinositol (GPI) anchor of *Trypanosoma brucei*. *Carbohydr. Res.* 1992; 234:75–91. [PubMed: 1468082]
22. Campbell AS, Fraser-Reid B. First synthesis of a fully phosphorylated GPI membrane anchor: Rat brain Thy-1. *J. Am. Chem. Soc.* 1995; 117:10387–10388.
23. Udodong UE, Madsen R, Roberts C, Fraser-Reid B. A ready, convergent synthesis of the heptasaccharide GPI membrane anchor of rat brain Thy-1. *J. Am. Chem. Soc.* 1993; 115:7886–7887.
24. Lu J, Jayaprakash KN, Fraser-Reid B. First synthesis of a malarial prototype: a fully lipidated and phosphorylated GPI membrane anchor. *Tetrahedron Lett.* 2004; 45:879–882.
25. Lu J, Jayaprakash KN, Schlueter U, Fraser-Reid B. Synthesis of a malaria candidate glycosylphosphatidylinositol (GPI) structure: A strategy for fully inositol acylated and phosphorylated GPIs. *J. Am. Chem. Soc.* 2004; 126:7540–7547. [PubMed: 15198601]
26. Baeschlin DK, Chaperon AR, Charbonneau V, Green LG, Ley SV, Lucking U, Walther E. Rapid assembly of oligosaccharides: total synthesis of a glycosylphosphatidylinositol anchor of *Trypanosoma brucei*. *Angew. Chem. Int. Ed.* 1998; 37:3423–3428.
27. Baeschlin DK, Chaperon AR, Green LG, Hahn MG, Ince SJ, Ley SV. 1,2-Diacetals in synthesis: total synthesis of a glycosylphosphatidylinositol anchor of *Trypanosoma brucei*. *Chem. Eur. J.* 2000; 6:172–186. [PubMed: 10747402]
28. Lopez-Prados J, Martin-Lomas M. Inositolphosphoglycan mediators: An effective synthesis of the conserved linear GPI anchor structure. *J. Carbohydr. Chem.* 2006; 24:393–414.

29. Reichardt N-C, Martin-Lomas M. A practical solid-phase synthesis of glycosylphosphatidylinositol precursors. *Angew. Chem. Int. Ed.* 2003; 42:4674–4677.
30. Yashunsky DV, Borodkin VS, Ferguson MAJ, Nikolaev AV. The chemical synthesis of bioactive glycosylphosphatidylinositols from *Trypanosoma cruzi* containing an unsaturated fatty acid in the lipid. *Angew. Chem. Int. Ed.* 2006; 45:468–474.
31. Pekari K, Schmidt RR. A variable concept for the preparation of branched glycosyl phosphatidyl inositol anchors. *J. Org. Chem.* 2003; 68:1295–1308. [PubMed: 12585868]
32. Mayer TG, Kratzer B, Schmidt RR. Synthesis of a GPI anchor of yeast (*Saccharomyces cerevisiae*). *Angew. Chem. Int. Ed.* 1994; 33:2177–2181.
33. Mayer TG, Schmidt RR. Glycosyl phosphatidylinositol (GPI) anchor synthesis based on versatile building blocks - Total synthesis of a GPI anchor of Yeast. *Eur. J. Org. Chem.* 1999:1153–1165.
34. Pekari K, Tailler D, Weingart R, Schmidt RR. Synthesis of the fully phosphorylated GPI anchor pseudohexasaccharide of *Toxoplasma gondii*. *J. Org. Chem.* 2001; 66:7432–7442. [PubMed: 11681958]
35. Hewitt MC, Snyder DA, Seeberger PH. Rapid synthesis of a glycosylphosphatidylinositol- based malaria vaccine using automated solid-phase oligosaccharide synthesis. *J. Am. Chem. Soc.* 2002; 124:13434–13436. [PubMed: 12418894]
36. Schofield L, Hewitt MC, Evans K, Siomos M-A, Seeberger PH. Synthetic GPI as a candidate antitoxic vaccine in a model of malaria. *Nature.* 2002; 418:785–789. [PubMed: 12181569]
37. Liu X, Kwon Y, Seeberger PH. Convergent synthesis of a fully lipidated glycosylphosphatidylinositol anchor of *plasmodium falciparum*. *J. Am. Chem. Soc.* 2005; 127:5004–5005. [PubMed: 15810819]
38. Tsai Y-H, Gtze S, Azzouz N, Hahm HK, Seeberger PH, Silva DV. A general method for synthesis of GPI anchors illustrated by the total synthesis of the low-molecular-weight antigen from *Toxoplasma gondii*. *Angew. Chem. Int. Ed.* 2011; 50:9961–9964.
39. Ali A, Gowda DC, Vishwakarma RA. A new approach to construct full-length glycosylphosphatidylinositols of parasitic protozoa and [4-deoxy-Man-III]-GPI analogues. *Chem. Commun.* 2005:519–521.
40. Ali A, Vishwakarma RA. Total synthesis of the fully lipidated glycosylphosphatidylinositol (GPI) anchor of malarial parasite *Plasmodium falciparum*. *Tetrahedron.* 2010; 66:4357–4369.
41. Xue J, Guo Z. Convergent synthesis of a GPI containing an acylated inositol. *J. Am. Chem. Soc.* 2003; 125:16334–16339. [PubMed: 14692775]
42. Wu X, Guo Z. Convergent synthesis of a fully phosphorylated GPI anchor of the CD52 antigen. *Org. Lett.* 2007; 9:4311–4313. [PubMed: 17877362]
43. Swarts BM, Guo Z. Synthesis of a glycosylphosphatidylinositol (GPI) anchor carrying unsaturated lipid chains. *J. Am. Chem. Soc.* 2010:6648–6650. [PubMed: 20423078]
44. Burgula S, Swarts BM, Guo Z. Total synthesis of a glycosylphosphatidylinositol anchor of the human lymphocyte CD52 antigen. *Chem. Eur. J.* 2012; 18:1194–1201. [PubMed: 22189835]
45. Tanaka Y, Nakahara Y, Hojo H, Nakahara Y. Studies directed toward the synthesis of protein-bound GPI anchor. *Tetrahedron.* 2003; 59:4059–4067.
46. Paulick MG, Wise AR, Forstner MB, Groves JT, Bertozzi CR. Synthetic analogues of glycosylphosphatidylinositol-anchored proteins and their behavior in supported lipid bilayers. *J. Am. Chem. Soc.* 2007; 129:11543–11550. [PubMed: 17715922]
47. Becker CFW, Liu X, Olschewski D, Castelli R, Seidel R, Seeberger PH. Semisynthesis of a glycosylphosphatidylinositol-anchored prion protein. *Angew. Chem., Int. Ed.* 2008; 47:8215–8219.
48. Xue J, Shao N, Guo Z. First total synthesis of a GPI-anchored peptide. *J. Org. Chem.* 2003; 68:4020–4029. [PubMed: 12737586]
49. Shao N, Xue J, Guo Z. Chemical synthesis of a skeleton structure of sperm CD52 - A GPI-anchored glycopeptide. *Angew. Chem. Int. Ed.* 2004; 43:1569–1573.
50. Guo X, Wang Q, Swarts BM, Guo Z. Sortase-catalyzed peptide-glycosylphosphatidylinositol analog ligation. *J. Am. Chem. Soc.* 2009; 131:9878–9879. [PubMed: 19583255]

51. Wu Z, Guo X, Wang Q, Swarts BM, Guo Z. Sortase A-catalyzed transpeptidation of glycosylphosphatidylinositol derivatives for chemoenzymatic synthesis of GPI-anchored proteins. *J. Am. Chem. Soc.* 2010; 132:1567–1571. [PubMed: 20078124]
52. Wu Z, Guo X, Guo Z. Chemoenzymatic synthesis of glycosylphosphatidylinositol-anchored glycopeptides. *Chem. Comm.* 2010:5773–5774. [PubMed: 20574591]
53. Xue J, Guo Z. Convergent synthesis of a GPI anchor of sperm CD52 antigens. *Bioorg. Med. Chem. Lett.* 2002; 12:2015–2018. [PubMed: 12113831]
54. Gou D, Liu Y, Chen C. An efficient chemoenzymatic access to optically active *myo*-inositol polyphosphates. *Carbohydr. Res.* 1992; 234:51.
55. Yu F, Guo Z. Efficient syntheses of chiral *myo*-inositol derivatives - Key intermediates in glycosylphosphatidylinositol (GPI) syntheses. *Bioorg. Med. Chem. Lett.* 2009; 19:3852–3855. [PubMed: 19380226]
56. Jia ZJ, Olsson L, Fraser-Reid B. Ready routes to key *myo*-inositol component of GPIs employing microbial arene oxidation or Ferrier reaction. *J. Chem. Soc., Perkin Trans.* 1998; 1:631–632.
57. Ferrier RJ, Middleton S. The conversion of carbohydrate derivatives into functionalized cyclohexanes and cyclopentanes. *Chem. Rev.*, 1993, 93 (8), 2779–2831. *Chem. Rev.* 1993; 93:2779–2831.
58. Xue J, Guo Z. A facile synthesis of Cerny epoxides and selectively blocked derivatives of 2-azido-2-deoxy- β -D-glucopyranose. *Tetrahedron Lett.* 2001; 42:6487–6489.
59. Wu X, Shen Z, Zeng X, Lang S, Palmer M, Guo Z. Synthesis and biological evaluation of sperm CD52 GPI anchor and related derivatives as binding receptors of pore-forming CAMP factor. *Carbohydr. Res.* 2008; 343:1718–1729. [PubMed: 18439573]
60. Lee J, Cha JK. Selective cleavage of allyl ethers. *Tetrahedron Lett.* 1996; 37:3663–3666.
61. Swarts BM, Guo Z. Chemical synthesis and functionalization of clickable glycosylphosphatidylinositol anchors. *Chem. Sci.* 2011; 2:2342–2352. [PubMed: 22163072]
62. Shao N, Xue J, Guo Z. Chemical synthesis of CD52 glycopeptides containing the acid-labile fucosyl linkage. *J. Org. Chem.* 2003; 68:9003–9011. [PubMed: 14604374]
63. Shao N, Guo Z. A facile synthesis of the fucosylated N-linked glycan core and its application to solid-phase synthesis of CD52 glycopeptide. *Pol. J. Chem.* 2005; 79:297–307.
64. Crich D, Sun S. Direct synthesis of β -mannopuranosides and other hindered glycosides from thioglycoside. *J. Am. Chem. Soc.* 1998; 120:435–436.
65. Lemieux RU, Hendriks KB, Stick RV, James K. Halide ion catalyzed glycosidation reactions. Syntheses of α -linked disaccharides. *J. Am. Chem. Soc.* 1975; 97:4056–4062.
66. Kinoshita T, Inoue N. Dissecting and manipulating the pathway for glycosylphosphatidylinositol-anchor biosynthesis. *Curr. Opin. Chem. Biol.* 2000; 4:632–638. [PubMed: 11102867]
67. Takeda J, Kinoshita T. GPI-anchor biosynthesis. *Trends Biochem. Sci.* 1995; 20:367–371. [PubMed: 7482705]
68. Amthauer R, Kodukula K, Gerber L, Udenfriend S. Evidence that the putative COOH-terminal signal transamidase involved in glycosylphosphatidylinositol protein synthesis is present in the endoplasmic reticulum. *PNAS.* 1993; 90:3973–3977. [PubMed: 8387204]
69. Nagamune K, Ohishi K, Ashida H, Hong Y, Hino J, Kangawa K, Inoue N, Maeda Y, Kinoshita T. GPI transamidase of *Trypanosoma brucei* has two previously uncharacterized (trypanosomatid transamidase 1 and 2) and three common subunits. *PNAS.* 2003; 100:10682–10687. [PubMed: 12958211]
70. Grimme SJ, Gao X, Martin PS, Tu K, Tcheperegine SE, Corrado K, Farewell AE, Orlean P, Bi E. Deficiencies in the Endoplasmic Reticulum (ER)-membrane protein Gab1p perturb transfer of glycosylphosphatidylinositol to proteins and cause perinuclear ER-associated actin bar formation. *Mol. Biol. Cell.* 2004; 15:2758–2770. [PubMed: 15075373]
71. Marraffini LA, DeDent AC, Schneewind O. Sortases and the art of anchoring proteins to the envelopes of Gram-positive bacteria. *Microbiol. Mol. Biol. Rev.* 2006; 70:192–221. [PubMed: 16524923]
72. Ton-That H, Liu G, Mazmanian SK, Faull KF, Schneewind O. Purification and characterization of sortase, the transpeptidase that cleaves surface proteins of *Staphylococcus aureus* at the LPXTG motif. *PNAS.* 1999; 96:12424–12429. [PubMed: 10535938]

73. Pritz S, Kraetke O, Klose A, Klose J, Rothmund S, Fechner K, Bienert M, Beyermann M. Synthesis of protein mimics with nonlinear backbone topology by a combined recombinant, enzymatic, and chemical synthesis strategy. *Angew. Chem. Int. Ed.* 2008; 47:3642–3645.
74. Samantaray S, Marathe U, Dasgupta S, Nandicoori VK, Roy RP. Peptide-sugar ligation catalyzed by transpeptidase sortase: A facile approach to neoglycoconjugate synthesis. *J. Am. Chem. Soc.* 2008; 130:2132–2133. [PubMed: 18229923]
75. Pritz S, Wolf Y, Kraetke O, Klose J, Bienert M, Beyermann M. Synthesis of biologically active peptide nucleic acid-peptide conjugates by sortase-mediated ligation. *J. Org. Chem.* 2007; 72:3909–3912. [PubMed: 17432905]
76. Parthasarathy R, Subramanian S, Boder ET. Sortase A as a novel molecular “stapler” for sequence-specific protein conjugation. *Bioconjugate Chem.* 2007; 18:469–476.
77. Mao H, Hart SA, Schink A, Pollok BA. Sortase-mediated protein ligation: A new method for protein engineering. *J. Am. Chem. Soc.* 2004; 126:2670–2671. [PubMed: 14995162]
78. Arsequell G, Valencia G. O-Glycosyl α -amino acids as building blocks for glycopeptide synthesis. *Tetrahedron Asym.* 1997; 8:2839–2876.
79. van der Goot, FG. Pore-Forming Toxins. Vol. 257. Springer-Verlag; Berlin: 2001.
80. Braun V, Focareta T. Pore-forming bacterial protein hemolysins (cytolysins). *Crit. Rev. Microbiol.* 1991; 18:115–158. [PubMed: 1930675]
81. Montecucco C, Papini E, Schiavo G. Bacterial protein toxins penetrate cells via a four-step mechanism. *FEBS Lett.* 1994; 346:92–98. [PubMed: 8206166]
82. Shier, WT.; Mebs, D. Handbook of Toxicology. Marcel Dekker, Inc.; New York: 1990.
83. Parker MW, Buckley JT, Postma JPM, Tucker AD, Leonard K, Pattus F, Tsernoglou D. Structure of the Aeromonas toxin proaerolysin in its water-soluble and membrane-channel states. *Nature.* 1994; 367:292–295. [PubMed: 7510043]
84. Lang S, Palmer M. Characterization of Streptococcus agalactiae CAMP factor as a pore-forming toxin. *J. Biol. Chem.* 2003; 278:38167–38173. [PubMed: 12835325]
85. Wu Q, Guo Z. Glycosylphosphatidylinositols (GPIs) are potential targets for the development of novel Inhibitors for aerolysin-type of pore-forming bacterial toxins. *Med. Res. Rev.* 2010:258–269. [PubMed: 19557762]
86. Lang S, Xue J, Guo Z, Palmer M. Streptococcus agalactiae CAMP factor binds to GPI-anchored proteins. *Med. Microbiol. Immunol.* 2007; 196:1–10. [PubMed: 16773378]

Figure 1.

The highly conserved core structure (1) of protein/glycoprotein-anchoring GPIs, as well as GPI anchorage of surface proteins and glycoproteins onto cell membranes through the insertion of GPI lipid chains into the cell membrane bilayer.

R = permanent protecting groups; R^1 , R^2 , R' , R'' = temporary protecting groups; R^3 , R^4 = lipids and/or permanent protecting groups.

Scheme 1.

A widely adopted strategy for the synthesis of GPI anchors

Figure 2.
Representative structure of the human CD52 antigen

Scheme 2.

Synthesis of the inositol derivative **6** suitable for the synthesis of CD52 GPI anchor

Bu = butyl; PMP = *para*-methoxyphenyl; TBAI = tetrabutylammonium iodide

Scheme 3.

Alternative synthesis of inositol derivatives having differentiated 1,2,6-*O*-positions

Scheme 4.
Synthesis of the glycosaminy donor **31**

CAN = cerium ammonium nitrate; Cbz = benzyloxycarbonyl; Cp = cyclopentadiene; DCC = dicyclohexylcarbodiimide; DMAP = 4-Dimethylaminopyridine; *i*Pr = *iso*-propyl; Tf = triflic

Scheme 5.

Synthesis of phospholipidated pseudodisaccharide **32**

*t*Bu = *tert*-butyl; DBU = 1,8-Diazabicycloundec-7-ene; Et = ethyl; Fmoc = fluorenylmethyloxycarbonyl; NIS = *N*-iodosuccinimide; TBS = *tert*-butyldimethylsilyl

Scheme 6.
Convergent assembly of CD52 GPI anchor **43**

R^P = protecting group; R^2 = protecting group or palmitoyl

Scheme 7.

Convergent synthesis of GPIs and GPI analogs with phospholipidated pseudodisaccharide **44** as key intermediate

Scheme 8.
Synthesis of optically pure inositol derivative **50**

Scheme 9. Synthesis of PMB-protected trimannosyl donor **63**

Scheme 10.

Assembly of GPI **72** carrying unsaturated lipid chains

Figure 3.

Other GPI anchors and GPI derivatives carrying Pd-catalyzed hydrogenolysis-incompatible functionalities synthesized by the convergent strategy using the PMB group for global and permanent hydroxyl protection

Scheme 11.

Synthesis of the natural GPI anchor **73** of the human lymphocyte CD 52 antigen

Figure 4.
Key intermediates involved in the synthesis of functionalized GPI anchor **74**

Scheme 12. Synthesis of **75** carrying a “clickable” azido group

DCM = dichloromethane; Pfp = pentafluorophenyl.

Scheme 13.
synthesis of GPI-anchored peptide **97**

Figure 5.

GPI-anchored glycopeptide **98** prepared by chemical coupling of glycopeptide **99** and GPI

Scheme 14.
Solid-phase synthesis of glycopeptide **99**

Scheme 15.
Synthesis of glycan-asparagine conjugate **101**

EDC = 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide; HOBT = 1-*N*-hydroxybenzotriazole; NMP = *N*-methylpyrrolidinone

Scheme 16.

Final assembly of GPI-anchored glycopeptide **98**

Figure 6.

The addition of GPIs to target proteins to form GPI-anchored proteins is achieved in the ER lumen via GPI transamidation catalyzed by GPI-T, with both the GPI anchor and the nascent protein, as well as the enzyme, attached to the ER membrane inner surface. This reaction is regulated by a peptide signal near the nascent protein C-terminus, which is cut off during the transamidation process. GPI-anchored proteins formed can be further processed and finally delivered onto the cell surface through the Golgi network.

Figure 7.

Sorting pathway of surface proteins in gram-positive bacteria. After surface proteins containing a sorting signal at the C-terminus are synthesized in the bacterial cytoplasm, they are translocated onto the cell surface. Sortase cleaves a peptide bond of the sorting signal to generate a reactive acyl intermediate and then transfers the acyl group to the N-terminus of the pentaglycine cross bridge of the cell wall peptidoglycan. As a result, surface proteins are anchored onto the bacterial cell wall.

Scheme 17.

A general strategy for GPI-anchored protein synthesis based on sortase-mediated GPI-protein ligation

Scheme 18.

SrtA-mediated coupling reactions of peptides and GPI analogs

Scheme 19.

SrtA-mediated chemoenzymatic synthesis of CD52 analogs

Scheme 20.

SrtA-mediated chemoenzymatic synthesis of CD52 and CD24 analogs

Scheme 21.

SrtA-mediated chemoenzymatic synthesis of GPI-anchored glycopeptides

Figure 8.

Inhibition of CAMP factor-induced sheep erythrocyte lysis by **139**. After CAMP factor was first incubated with 1,2-dipalmitoyl-*sn*-glycero-3-phosphatidylcholine (DPPC) alone or with DPPC and the indicated concentrations of **121** (see labels in the figure), its solution was added to the sphingomyelinase-sensitized erythrocytes. Then, hemolysis was observed by decreasing cell turbidity at OD₆₀₀.