The SKPO-1 Peroxidase Functions in the Hypodermis to Protect *Caenorhabditis elegans* From Bacterial Infection

George R. Tiller and Danielle A. Garsin¹

Department of Microbiology and Molecular Genetics, University of Texas Health Science Center and Graduate School of Biomedical Sciences, Houston, Texas 77030

ABSTRACT In recent years, the synergistic relationship between NADPH oxidase (NOX)/dual oxidase (DUOX) enzymes and peroxidases has received increased attention. Peroxidases utilize NOX/DUOX-generated H_2O_2 for a myriad of functions including, but not limited to, thyroid hormone biosynthesis, cross-linking extracellular matrices (ECM), and immune defense. We postulated that one or more peroxidases produced by *Caenorhabditis elegans* would act in host defense, possibly in conjunction with BLI-3, the only NOX/DUOX enzyme encoded by the genome that is expressed. Animals exposed to RNA interference (RNAi) of the putative peroxidase genes were screened for susceptibility to the human pathogen *Enterococcus faecalis*. One of three genes identified, *skpo-1* (ShkT-containing peroxidase), was studied in depth. Animals mutant for this gene were significantly more susceptible to *E. faecalis*, but not *Pseudomonas aeruginosa*. A slight decrease in longevity was also observed. The *skpo-1* mutant animals had a dumpy phenotype of incomplete penetrance; half the animals displayed a dumpy phenotype ranging from slight to severe, and half were morphologically wild type. The SKPO-1 protein contains the critical catalytic residues necessary for peroxidase activity, and in a whole animal assay, more H_2O_2 was detected from the mutant compared to the wild type, consistent with the loss of an H_2O_2 sink. By using tissue-specific *skpo-1* RNAi and immunohistochemical localization with an anti-SKPO-1 antibody, it was determined that the peroxidase is functionally and physically present in the hypodermis. In conclusion, these results characterize a peroxidase that functions protectively in the hypodermis during exposure to *E. faecalis*.

HEME-containing peroxidases play critical, wide-ranging roles in biological systems. Once called the “animal heme peroxidases,” members of the peroxidase–cyclooxygenase superfamily are actually found in all kingdoms of life (Zamocky *et al.* 2008). The roles of most are poorly characterized, but some are clearly involved in immune defense. For example, the most famous and best-studied member of this group, myeloperoxidase (MPO), is found in the granulocytes of neutrophils where it catalyzes the formation of the potent oxidant HOCl from H_2O_2 and Cl^- to kill invading microbes (reviewed by Klebanoff 2005). Another is lactoperoxidase (LPO), which is found on mucosal surfaces and generates

the protective oxidant hypothiocyanite ($OSCN^-$) from H_2O_2 and thiocyanate (SCN^-). This process is impaired in the lungs of patients with cystic fibrosis (CF), contributing to the poor clearance of pathogens such as *Pseudomonas aeruginosa* and *Staphylococcus aureus*, the most common causes of lung infection in patients with CF (Conner *et al.* 2002; Geiszt *et al.* 2003; Forteza *et al.* 2005; Moskwa *et al.* 2007).

Many of these peroxidases are functionally associated with members of the NADPH oxidase (NOX)/dual oxidase (DUOX) family of proteins. These enzymes generate the H_2O_2 required as substrate for the peroxidases. For example, Nox2 is the source of H_2O_2 for MPO. DUOXs differ from NOXs in that they encode a peroxidase domain in addition to the oxidant-generating NADPH oxidase domain (reviewed by Rada *et al.* 2008; Sumimoto 2008). Despite the fact that they have a peroxidase domain, DUOXs appear to still associate with separate peroxidases. For example, LPO utilizes H_2O_2 generated by Duox1 or Duox2 (Conner *et al.* 2002; Geiszt *et al.* 2003; Forteza *et al.* 2005). In fact, the peroxidase

Copyright © 2014 by the Genetics Society of America
doi: 10.1534/genetics.113.160606

Manuscript received December 10, 2013; accepted for publication March 9, 2014;
published Early Online March 12, 2014.

Supporting information is available online at <http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.113.160606/-/DC1>.

¹Corresponding author: University of Texas Health Science Center at Houston,
Department of Microbiology and Molecular Genetics, 6431 Fannin St., MSB 1.168,
Houston, TX 77030. E-mail: Danielle.A.Garsin@uth.tmc.edu

domains of human Duox1/2 are reported to lack peroxidase activity and may have other functions, such as serving as an interaction domain for separate, active peroxidases (Meitzler and Ortiz de Montellano 2009, 2011; Meitzler et al. 2013).

Caenorhabditis elegans has been used as a model host to study various aspects of the innate immune response, including the purposeful generation of reactive oxygen species (ROS) as a defense mechanism (Chavez et al. 2007, 2009). The animal encodes only one functional NADPH oxidase, a dual oxidase called **BLI-3** (Edens et al. 2001). When exposed to human pathogens such as the Gram-positive, opportunistic bacterium, *Enterococcus faecalis*, H₂O₂ is released (Chavez et al. 2007). The response appears protective, as its loss by reducing the expression of *bli-3* by RNA interference (RNAi) renders the animals more sensitive to killing by the pathogen (Chavez et al. 2009). Using indirect immunofluorescence, **BLI-3** was localized to the hypodermis, which is essentially the “skin” of *C. elegans* (Edens et al. 2001). There is also speculation that **BLI-3** may be present in the intestinal cells (Chavez et al. 2007, 2009). In addition to playing a role in innate immunity, **BLI-3** is essential to the normal development of the worm because it contributes to the generation of the tyrosine-linked collagen necessary for proper biogenesis of the cuticle, i.e., the exoskeleton of the animal. Its role appears twofold. First, it generates the necessary substrate H₂O₂, and then it oxidizes the tyrosines of the collagen proteins that then combine to form di- and trityrosines, thereby cross-linking the cuticle (Edens et al. 2001). Unlike human Duox1/2, the peroxidase domain of **BLI-3** has low levels of peroxidase activity that is essential to this process, and mutations in the peroxidase domain that disrupt this activity result in a “blistered” (*bli*) phenotype (Brenner 1974; Simmer et al. 2003; Meitzler and Ortiz de Montellano 2009, 2010, 2011). However, more recent work has demonstrated that the peroxidase domain of **BLI-3** is not the only peroxidase involved in this process. A separate peroxidase, **MLT-7**, also contributes to cuticle cross-linking and loss of this activity results in the same *bli* phenotype as loss of the **BLI-3** peroxidase domain. Additionally, loss of both peroxidase activities greatly increases the severity of cuticle blistering (Thein et al. 2009).

Because of the prevalent involvement of peroxidases in immune responses, we hypothesized that *C. elegans* might also utilize a peroxidase(s) in host defense, perhaps in conjunction with its NADPH oxidase, **BLI-3**. The results of an earlier investigation indicated that the peroxidase domain of **BLI-3** is not involved, as point mutants in this domain had wild-type resistance to *E. faecalis*, despite their blistered phenotype (Chavez et al. 2009). In this study, we characterized the effects of reducing the expression of putative peroxidases encoded by the *C. elegans* genome on susceptibility to *E. faecalis*. We identified three genes whose loss resulted in susceptibility. One gene, named *skpo-1*, for ShkT-containing peroxidase, was investigated in depth.

Materials and Methods

Creation of complementary DNA pL4440 constructs for double-stranded RNA production in *Escherichia coli*

A standard RNA isolation procedure was used to procure total RNA from healthy N2 adults grown on *Escherichia coli* OP50. We followed the first-strand complementary DNA (cDNA) synthesis protocol from the Life Technologies SuperScript III Synthesis system to produce first-strand total cDNA. We then followed a standard PCR protocol using primers specific to *F32A52.A*, *F32A52.B*, and *pxn-1*, respectively (Supporting Information, Table S5). PCR products were digested and ligated into pL4440 and transformed into chemically competent *E. coli* HT115 to produce double-stranded RNA (dsRNA) to the gene of interest, respectively.

C. elegans strains and growth conditions

C. elegans strains were grown and maintained as previously described (Hope 1999). The strains used for the hypodermal and intestinal RNAi studies are as follows: *rde-1(ne219)*; *Is[wrt-2prom::RDE-1::unc54 3'utr; myo2p::RFP3]* and *sid-1(qt9)*; *Is[vha-6::sid-1]*, respectively. The RB1437 [*skpo-1(ok1640)* II] strain is a partial deletion mutant for *skpo-1* and was verified via sequencing (www.wormbase.org). The RB1437 [*skpo-1(ok1640)* II] strain was created by the *C. elegans* Reverse Genetics Core Facility at the University of British Columbia, part of the International *C. elegans* Gene Knockout Consortium and obtained through the *Caenorhabditis* Genetics Center. The independent lines of *skpo-1* strains, GF89–GF91, were obtained through backcrossing the RB1437 [*skpo-1(ok1640)* II] strain with the wild-type N2 Bristol strain five times (backcrossing, BC ×5).

RNAi

RNAi was performed by exposing L1–L4 stage larvae to *E. coli* HT115-expressing dsRNA to target genes. RNAi clones were obtained from the *C. elegans* library (Fraser et al. 2000; Kamath et al. 2003). *skpo-1* worms exposed to pathogens were prone to a maternal bagging phenotype; therefore, *cdc-25.1* RNAi was used to induce sterility in some experiments (van der Hoeven et al. 2011).

Survival and longevity assays

Unless otherwise indicated, the following bacterial strains were used: OP50 (*E. coli*) (Brenner 1974), OG1RF (*E. faecalis*) (Dunny et al. 1978), and PA14 (*P. aeruginosa*) (Rahme et al. 1995). Exposure to RNAi, survival assays, and longevity assays were performed as previously described (Garsin et al. 2001, 2003; Kim et al. 2002; van der Hoeven et al. 2011). Briefly, for *E. faecalis* survival assays, *E. faecalis* grown in brain heart infusion (BHI) medium for 5 hr was seeded (10 µl) onto BHI plates (gentamycin 50 µg/ml) and incubated at 37° for 24 hr, whereas for *P. aeruginosa* survival assays, *P. aeruginosa* was cultured in Luria broth (LB) overnight at 37°, seeded (10 µl) onto slow-killing plates, and incubated first for 24 hr at 37° and then for the duration of

the experiment at 25°. For the *E. coli* longevity assays, 20× *E. coli* was seeded (100 µl) onto nematode growth (NG) medium plates supplemented with 5-Fluoro-2'-deoxyuridine (25 µg/ml) and streptomycin (25 µg/ml). Seeded NG plates were incubated at 25° for 24 hr. In all assays, a total of 90 L4 larvae were transferred to three replica plates and the assay was performed at 25°. Worms were scored as live and dead at various time points.

Amplex Red assay for H₂O₂ measurements

The Amplex Red hydrogen peroxide/peroxidase kit (Invitrogen Molecular Probes, Eugene, OR) was previously adapted to *C. elegans* to measure pathogen-stimulated hydrogen peroxide release (Chavez *et al.* 2007, 2009). The same protocol was followed with the following modifications: L4 worms were exposed to a bacterial strain for 12 hr, and the fluorescence of 30 worms per well was measured after 30 min of incubation with Amplex Red (540/590 nm excitation and emission, respectively). A total of 80 µM diphenyleneiodinium chloride (DPI) (TCI, Tokyo) was added to some wells and allowed to incubate for 15 min prior to addition of Amplex Red, HRP, and Na₂PO₄ (pH 7.4), and fluorescence was measured as above.

Indirect immunofluorescence

Immunofluorescence for *C. elegans* was performed according to Seydoux and Dunn (1997) on young adult animals. Custom peptide synthesis, rabbit immunization, and affinity purification of the SKPO-1 polyclonal antibody was performed by LifeTein (South Plainfield, NJ). The CRVGRRAFDIENGSR peptide corresponds to the C terminus of the *C. elegans* protein SKPO-1. Adult N2 and GF89 worms were imaged using an Olympus IX81 automated inverted microscope and Slidebook (version 5.0) software along with the SKPO-1 polyclonal primary antibody and Alexa Fluor 488 goat antirabbit IgG secondary antibody to localize SKPO-1 in *C. elegans*.

Immunogold labeling and transmission electron microscopy

Approximately 1000 N2 and GF89 animals were raised under standard conditions to the young adult stage. They were rinsed off of the propagation plates with M9 and spun down at 2000 rpm for 1 min. The worms were sequentially resuspended and pelleted three times in 1 ml M9 (Hope 1999). After the final wash, they were resuspended in 1 ml 3% formalin, 0.15% glutaraldehyde in Millonig's buffer, pH 7.45, and fixed for 3 days at 40°. Then the buffer was drawn off and the samples were incubated in 0.1% fresh made sodium borohydride in Millonig's buffer (room temperature, RT, for 10 min). This was followed by incubations in Millonig's buffer (RT, 10 min, repeated twice). The samples were then dehydrated at room temperature by incubating in 50% ethanol (10 min, twice) and 70% ethanol, (15 min, twice). The samples were then permeated with 50% LR White resin and 50% ethanol for 60 min, followed by an

overnight incubation at room temperature in 50% LR White resin sealed on a rotator. The samples were then incubated at room temperature for 4 hr in 70% LR White resin and then for 5 hr in 100% LR White resin, while on a rotator. The pellets were then sealed into BEEM capsules and allowed to polymerize in a 53° oven overnight. Thin sections (120 nm) of the LR White blocks were cut using a DiATOME diamond knife and a Leica Ultracut R microtome. Sections were floated onto 200 mesh nickel grids. The grids were then floated in 50 mM glycine (RT, 15 min) and then washed twice by floating in drops of PBS. The samples were blocked by floating them for 30 min in Aurion Blocking Solution followed by washes in incubation buffer (PBS with 0.1% Aurion BSA-c) (5 min, three times). The SKPO-1 antibody was diluted into the incubation buffer at 1:1000 and the grids were incubated for 1.5 hr at RT. This was followed by washes in incubation buffer (10 min, four times) and then incubation with the secondary antibody (EMS goat antirabbit IgG with 10 nm gold) diluted 1:20 in incubation buffer. The samples were washed again in incubation buffer (10 min, five times) followed by PBS (10 min, three times). A postfix treatment of 3% glutaraldehyde was applied for 10 min followed by a final wash with distilled water (5 min, two times). The grids were dried in a 70° oven and then imaged using a JEOL 1200 transmission electron microscope at 60 kV and captured with a 2k × 2k Gatan CCD camera.

Statistical analysis

Amplex Red, survival, and longevity assays were analyzed using GraphPad Prism version 5.0 (GraphPad Software, San Diego). Student's paired *t*-test was used to determine the statistical significance of the Amplex Red data. *P*-values of <0.05 were considered to be statistically significant. Statistically significant differences are denoted in the figures with asterisks. Kaplan–Meier log rank analysis was used to compare survival and longevity curves pairwise and to calculate the median survival. *P*-values of <0.05 were considered to be statistically significant.

Results

Host resistance to *E. faecalis* is affected by *skpo-1*

To investigate if *C. elegans* possesses any peroxidases involved in host defense, we utilized an RNAi-killing assay to screen candidate peroxidase genes. The candidate genes were found using the BLI-3 peroxidase domain as the BLAST query on www.wormbase.org. Twelve putative peroxidase-encoding genes were identified, 9 of which were available in the RNAi library (Fraser *et al.* 2000; Kamath *et al.* 2003), including the previously studied *mlt-7* (Thein *et al.* 2009). Standard molecular biology techniques were used to generate the three missing RNAi constructs (*Materials and Methods*). Using a background in which RNAi efficiency is increased (*eri-1*) (Kennedy *et al.* 2004), the expression of these genes was reduced in *C. elegans*. The animals were

Table 1 Susceptibility to *E. faecalis* following RNAi of putative peroxidase genes

Gene	Susceptibility phenotype	P-value	Median survival (days)
<i>mlt-7</i> control	Wild type	0.3070	6
<i>pxn-1</i> control	Wild type	0.9218	6
<i>pxn-2</i> control	Wild type	0.8233	4
<i>C16C8.2</i> control	Wild type	0.1067	4
<i>C46A5.4</i> control	Wild type	0.7206	6
<i>F09F3.5</i> control	Susceptible	0.0058	6
<i>F32A5.2a</i> control	Wild type	0.2499	7
<i>F32A5.2b</i> control	Wild type	0.4115	5
<i>F49E12.1 (skpo-1)</i> control	Susceptible	0.0053	5
<i>R08F11.7</i> control	Susceptible	0.0004	6
<i>K10B4.1</i> control	Wild type	0.7904	7
<i>T06D8.10</i> control	Wild type	0.6636	8
			8

then exposed to *E. faecalis* and survival was scored over time. By these means, the genes were screened for possible roles in innate immune function. The reduced expression of three genes—*F09F3.5*, *R08F11.7*, and *F49E12.1*—resulted in a statistically significant susceptibility phenotype (Table 1). We decided to focus on *F49E12.1*, as it possessed high homology to human myeloperoxidase (hMPO, 34% sequence identity) and a deletion mutant was available through the *Caenorhabditis* Genetics Center (CGC). As described in more detail below, the predicted protein associated with this gene contains an N-terminal metridin Shk toxin (ShkT)-like domain and a C-terminal peroxidase domain and was therefore named **SKPO-1** for ShkT-containing peroxidase.

An example of the survival of *skpo-1* RNAi *eri-1* mutant animals on *E. faecalis* compared to vector control (VC) RNAi *eri-1* mutant animals is shown in Figure 1A. *skpo-1* RNAi animals displayed an enhanced susceptibility phenotype relative to VC RNAi ($P = 0.0011$). The experiment was repeated five times and the data were tabulated in Table S1. We also examined the lifespan of these animals to see if *skpo-1* RNAi caused a general fitness defect. However, on *E. coli* OP50, lifespan of the *skpo-1* RNAi animals was not significantly different when compared to VC RNAi animals ($P = 0.3772$) (Figure 1B and Table S2).

To further examine the phenotype resulting from the loss of *skpo-1*, we obtained the partial deletion strain, RB1437 from the CGC. The deletion is between bases 827 and 3202 of the 5008 full-length transcript, which ablates a significant portion of the protein, including the critical amino acid res-

idues necessary for peroxidase activity in classical animal heme peroxidases (reviewed by Ortiz de Montellano 2008). Strain RB1437 was backcrossed five times into our wild-type N2 strain and three lines were generated: GF89, GF90, and GF91. As shown in Figure S1, these animals were very susceptible to *E. faecalis* compared to wild-type N2 animals. Because significant differences in susceptibility were not observed between the strains, we continued our studies using GF89, which is referred to as *skpo-1*.

We observed a significant “bagging” phenotype—hatching of embryos inside the hermaphrodite that had failed to be expelled—when the *skpo-1* mutant animals were exposed to *E. faecalis*, unlike the *skpo-1* RNAi animals. There was concern that this was artificially enhancing their sensitivity to *E. faecalis*, thus *cdc-25.1* RNAi was employed to induce sterility, as done in previous *C. elegans* pathogenesis studies (Shapira *et al.* 2006; Irazoqui *et al.* 2008; van der Hoeven *et al.* 2011) (Figure 1, C–E). *skpo-1* mutant animals grown on *cdc-25.1* dsRNA-expressing *E. coli* HT115 did not display this maternal bagging phenotype when they were subsequently exposed to *E. faecalis*. *cdc-25.1* RNAi-exposed *skpo-1* mutant worms retained a more pronounced susceptibility phenotype to *E. faecalis* relative to similarly exposed wild-type animals ($P < 0.0001$) (Figure 1C and Table S1), suggesting that the pathogen sensitivity is not completely explained by an increase in internal hatching of progeny.

Regardless of whether *skpo-1* mutant worms were sterile, *i.e.*, *cdc-25.1*-treated (Figure 1D and Table S2), or fecund (Figure S2 and Table S2), they displayed a slight reduction in lifespan, relative to wild type, when exposed to live *E. coli* OP50 ($P < 0.0001$, for both). Because live OP50 has been

Figure 1 SKPO-1 contributes to *C. elegans* resistance to *E. faecalis*. (A) Survival of *eri-1* mutant worms on *E. faecalis* OG1RF following exposure to vector control (VC) RNAi and *skpo-1* RNAi ($P = 0.0011$). (B) Longevity of *eri-1* mutant worms on *E. coli* OP50 following exposure to VC RNAi and *skpo-1* RNAi ($P = 0.3772$). (C) Survival of wild-type and *skpo-1* mutant worms on *E. faecalis* following exposure to *cdc-25.1* RNAi ($P < 0.0001$). (D) Longevity of wild-type and *skpo-1* mutant worms on *E. coli* OP50 following exposure to *cdc-25.1* RNAi ($P < 0.0001$). (E) The relative mortality of *skpo-1* mutant worms exposed to *E. faecalis* (EF) is expressed as a ratio of (LT_{50} of wild-type *cdc-25.1* RNAi on EF/ LT_{50} of *skpo-1* mutant *cdc-25.1* RNAi on EF) over (LT_{50} of wild-type *cdc-25.1* RNAi on *E. coli*/ LT_{50} of *skpo-1* mutant *cdc-25.1* RNAi on *E. coli*). The average of four independent experiments, with 90 animals each, was used to calculate the relative mortality. (F) Survival of wild-type and *skpo-1* mutant worms on *P. aeruginosa* following exposure to *cdc-25.1* RNAi ($P = 0.3783$). Error bars represent standard error of the mean (SEM) and the asterisk indicates a significant difference between wild-type and *skpo-1* mutant worms ($P = 0.0091$). The P -values were calculated using Student's paired t -test. The median survival for survival and longevity assays are listed in Table S1 and Table S2 along with replicates of the experiments.

shown to have slightly pathogenic effects (Garigan *et al.* 2002), we also examined lifespan on heat-killed OP50 and found that *cdc-25.1* RNAi-exposed *skpo-1* mutant worms still displayed a reduction in lifespan (Figure S3 and Table S2). To determine if the defect in lifespan completely accounted for the pathogen sensitivity or not, the relative mortality of *skpo-1* mutant worms, compared to wild type, was calculated (Figure 1F). Relative mortality is the ratio of the lethal time to kill 50% of the organisms (LT_{50}) of pathogen-infected animals to uninfected animals with the ratio of wild-type animals normalized to one and has been used in previous work to analyze susceptibility phenotypes (Tenor *et al.* 2004; Chavez *et al.* 2007). The survival defect for the *skpo-1* mutant animals on *E. faecalis* was more severe than on nonpathogenic *E. coli* compared to wild-type animals ($P = 0.0091$) (Figure 1E and Table S3), arguing against a general fitness defect.

To further examine the extent of the susceptibility-to-pathogen phenotype and determine if it was pathogen specific, we used *P. aeruginosa* instead of *E. faecalis* in the killing assay ($P = 0.3783$) (Figure 1F and Table S4). Interestingly, we observed no significant difference in survival between *cdc-25.1* RNAi *skpo-1* mutants and *cdc-25.1* RNAi wild-type animals when exposed to *P. aeruginosa*. These results further argue against a general fitness defect and suggest that the function of SKPO-1 may be beneficial during infection with some, but not all pathogens.

To determine if the susceptibility phenotype on *E. faecalis* was a result of changes in pathogen burden, we assessed how many colony-forming units (CFUs) were in the intestines of the infected worms. Wild-type and *skpo-1* mutant animals were raised on *E. coli* OP50 until L4 and then exposed to *E. faecalis* for 12 or 36 hr. The infected worms were washed to remove surface bacteria, homogenized, and then

Figure 2 *skpo-1* mutant displays multiple morphotypes. (A) Wild-type young adult representing the approximate average size and morphology of a typical *C. elegans*. (B) *skpo-1* RNAi young adult raised on *skpo-1* dsRNA-expressing *E. coli* HT115 from L1–L4 stages. (C–E) Young adult *skpo-1* mutants ranged from very dumpy to wild type in size. The $\times 10$ microscopy images are representative of >100 wild-type, *skpo-1* RNAi, and *skpo-1* worms observed, respectively.

serial dilutions of the homogenates were plated. Interestingly, we observed no significant difference in CFUs per worm between the wild-type and *skpo-1* mutant animals at either time point (Figure S4). The result suggests that the susceptibility phenotype of the *skpo-1* mutant cannot be explained by an increased load of bacteria in the intestine.

To test for alterations in the immune response between the *skpo-1* and wild-type animals a small panel of genes known to be upregulated in response to *E. faecalis* were examined, *clec-35*, *-42*, *-60*, and *-71*. These genes encode C-type lectins (*clec*), many of which are upregulated in response to pathogens (Irazoqui *et al.* 2010; Engelmann *et al.* 2011). Sterile (by exposure to *cdc-25* RNAi) wild-type and *skpo-1* mutant animals at the L4 stage were exposed to *E. faecalis* or *E. coli* for 18 hr, at which point the animals were lysed and RNA was extracted. By qRT-PCR, no significant difference in the expression of *clec-35* and *-71* was observed between the wild type and *skpo-1* mutants exposed to *E. faecalis*. However, we observed *clec-60*'s expression was significantly higher in the *skpo-1* mutant and there was a trend toward increased expression of *clec-42*, which was not statistically significant (Figure S5). These data suggest that the immune response is altered in the *skpo-1* mutant.

Morphological characterization of *skpo-1* mutant animals

Animals deficient in the peroxidase *mlt-7* displayed altered morphology, including a *bli* (blistered) phenotype, indicative of incomplete cross-linking of the cuticle (Thein *et al.* 2009). The phenotype strongly suggests that *MLT-7*'s peroxidase activity contributes to cuticle formation (Thein *et al.* 2009). Interestingly, despite this rather dramatic cuticle defect, an increase in susceptibility to *E. faecalis* was not observed (Table 1). To determine if *SKPO-1* is also involved in

cuticle biogenesis, we observed the morphology of *skpo-1* RNAi and mutant animals. RNAi of *skpo-1* did not result in any visible morphological change in the *eri-1* mutant animals (Figure 2B). However, the *skpo-1* mutant did display some morphological phenotypes. We observed young adult animals under the dissecting microscope ($N = 300$) and found that they ranged from very dumpy (16.5%), to slightly dumpy (32.9%), to wild type (50.6%) in appearance (Figure 2, C–E). No blistering of the cuticle was ever observed. Because a dumpy phenotype can be associated with cuticle defects (Page and Johnstone 2007), these results suggest that *SKPO-1* may have some role in cuticle biogenesis, but one that is different than *MLT-7*. Additionally, we observed no significant difference in susceptibility between *skpo-1* mutant morphotypes, *i.e.*, all were equally hypersusceptible to *E. faecalis* relative to wild-type animals (Figure S1).

SKPO-1 features and activity

As mentioned, *SKPO-1* contains a ShkT-like domain at its N terminus from residues ~ 21 –56 (www.wormbase.org) in addition to the predicted peroxidase domain (Figure 3A). It shares this feature with *MLT-7* (Thein *et al.* 2009), and putative peroxidases *C16C8.2* and *F32A5.2* (Table 1). For this reason we propose naming *F49E12.1*, *SKPO-1*, and *C16C8.2* and *F32A5.2*, *SKPO-2*, and *SKPO-3* for ShkT-containing peroxidase. In addition to the ShkT-like domain, the very 5' end contains a predicted signal sequence for secretion, and cleavage is predicted to occur at Ser19 (www.predisi.de).

To examine *SKPO-1*'s peroxidase domain, we aligned it with other well-characterized peroxidase domains of interest, those contained in hMPO, *BLI-3*, and *MLT-7*. Upon alignment, we observed *SKPO-1* possesses the catalytic triad (distal histidine, H²²²; arginine, R³³²; and proximal histidine, H⁴²⁸) necessary for peroxidase activity (reviewed by Ortiz de Montellano 2008). However, *SKPO-1* lacks the

Figure 3 Evidence that *skpo-1* is a potential peroxidase. (A) Peroxidase domain sequences were aligned against the putative peroxidase domain of SKPO-1. SKPO-1 possesses the distal histidine (H²²²), catalytic arginine (R³³²), and proximal histidine (H⁴²⁸), which are necessary for peroxidase activity. However, SKPO-1 lacks covalent heme-binding residues (S²²¹ and L³³⁵) that are characteristic of mammalian peroxidases (Ortiz de Montellano 2008). (B) *eri-1* mutant worms were grown on VC RNAi or *skpo-1* RNAi prior to exposure with either *E. coli* or *E. faecalis* for 12 hr at 25°. (C) Wild-type and *skpo-1* mutant worms were grown on *cdc-25.1* RNAi prior to exposure with either *E. coli* or *E. faecalis* for 12 hr at 25°. (B and C) Following exposure to *E. coli* or *E. faecalis*, the amount of H₂O₂ produced per minute was determined using the Amplex Red assay. Error bars represent the SEM and the asterisks indicate significant differences between *eri-1* worms exposed to VC RNAi or *skpo-1* RNAi that were infected with *E. faecalis* as well as between wild-type and *skpo-1* mutant worms exposed to *cdc-25.1* RNAi prior to infection with *E. faecalis* [$P = 0.0091$ (B) and $P < 0.0001$ (C)]. Additionally, wild-type and *skpo-1* mutant worms were exposed to 80 μ M diphenyleneiodinium chloride (DPI) and H₂O₂ levels were calculated for both *E. coli*- and *E. faecalis*-exposed animals (EC, $P = 0.0752$; EF, $P = 0.4161$, respectively) P -values were calculated via Student's paired t -test. Data in B and C are representative of at least two independent replicates.

residues necessary for covalent heme binding, characteristic of the animal heme peroxidase family (reviewed by Ortiz de Montellano 2008). Interestingly, human Duox1/2 (hDuox1/2) also lack the conserved aspartic and glutamic residues required for covalent linkage of the heme prosthetic group; however, both of these recombinant peroxidase domains still bind heme, albeit weakly (Meitzler and Ortiz de Montellano 2009, 2011). This noncovalent binding may explain MLT-7's peroxidase activity even though it lacks these covalent heme-binding residues (Thein *et al.* 2009).

By comparison, SKPO-1 possesses all of the highlighted residues in MLT-7 (Figure 3A) and shares significant identity with hMPO (34% identity). Thus, we decided to indirectly assay SKPO-1's potential peroxidase activity using an Amplex Red assay, modified for whole animals, that detects H₂O₂ (Chavez *et al.* 2007, 2009). Previously, we observed that exposure to *E. faecalis* causes a significant release in H₂O₂ dependent on the NADPH oxidase, BLI-3 (Chavez *et al.* 2007, 2009). We reasoned that loss of an important peroxidase during this release might increase the amount of H₂O₂ detected. After infecting VC and *skpo-1* RNAi *eri-1* mutant animals with *E. faecalis* for 12 hr, released H₂O₂ was measured ($P = 0.0091$) (Figure 3B). *skpo-1* RNAi *eri-1* mutant animals released significantly greater amounts of H₂O₂, relative to VC RNAi *eri-1* mutants, consistent with the loss of a predicted H₂O₂ sink. The same result was observed with the *skpo-1* mutants compared to wild-type animals ($P < 0.0001$) (Figure 3C). The difference required *E. faecalis*, and the concomitant release of H₂O₂, as no significant differences were observed between wild-type and *skpo-1*-deficient animals on *E. coli* (Figure 3, B and C).

To investigate if the increased amount of H₂O₂ produced by the *skpo-1* mutant was dependent on BLI-3 activity, we added the NADPH oxidase inhibitor, DPI, to the assay. We previously demonstrated that BLI-3 is the only expressed NADPH oxidase encoded by the genome (Chavez *et al.* 2009), and DPI can be utilized to reduce its activity in wild-type animals (Chavez *et al.* 2007). In Figure 3C, DPI also abrogated the enhanced H₂O₂ generation observed in *skpo-1* mutant animals. Taken together, Figure 3 supports the hypothesis that SKPO-1 is a peroxidase that utilizes H₂O₂ produced by BLI-3 during infection. Unfortunately, attempts to purify the protein to demonstrate definitive peroxidase activity *in vitro* were not successful.

SKPO-1 localizes to the hypodermis

C. elegans interfaces with *E. faecalis* at the cuticle, synthesized by the underlying hypodermis, as it crawls through the pathogen lawn, and in its intestine, due to ingestion of the bacteria (Garsin *et al.* 2001). It has previously been demonstrated that *C. elegans* mounts immune responses at these host-pathogen boundaries, depending on the pathogen and the nature of the infection (Wong *et al.* 2007; Pujol *et al.* 2008; Irazoqui *et al.* 2010). To address what tissue SKPO-1's activity is required for normal levels of resistance to *E. faecalis*, a tissue-specific RNAi approach was employed. RNAi-defective strains of *C. elegans* were used in which RNAi activity was genetically restored to specific tissues through intestinal or hypodermal-specific promoters (*vha-6::SID-1* and *wrt-2::RDE-1*, respectively) (Melo and Ruvkun 2012). In Figure 4, A and C (and Table S1), we tested the susceptibility of the hypodermal and intestinal-specific RNAi

Figure 4 SKPO-1 is necessary in the hypodermis for resistance to *E. faecalis* infection. (A) Survival of the hypodermal RNAi strain on *E. faecalis* following exposure to VC RNAi or *skpo-1* RNAi ($P = 0.0002$). (B) Longevity assay on *E. coli* OP50 of the hypodermal RNAi strain following exposure to VC RNAi or *skpo-1* RNAi ($P = 0.9997$). (C) Survival of the intestinal RNAi strain on *E. faecalis* following exposure to VC RNAi or *skpo-1* RNAi ($P = 0.9435$). (D) Longevity assay of the intestinal RNAi strain on *E. coli* OP50 following exposure to VC RNAi or *skpo-1* RNAi ($P = 0.6379$).

strains to *E. faecalis*. As in Figure 1, these strains were exposed to VC and *skpo-1* RNAi prior to infection. In the hypodermal RNAi strain, we observed an enhanced susceptibility phenotype to *E. faecalis* in the *skpo-1* RNAi animals relative to VC RNAi ($P = 0.0002$); however, in the intestinal RNAi strain, no significant difference between *skpo-1* and VC RNAi animals was observed when on *E. faecalis* ($P = 0.9435$). On *E. coli*, no significant difference was observed between VC and *skpo-1* RNAi for either the hypodermal or intestinal RNAi line ($P = 0.9997$ and $P = 0.6379$; Figure 4, B and D, respectively, and Table S2). From these experiments we conclude that SKPO-1's functional activity during pathogen exposure is required in the *C. elegans* hypodermis.

To determine in which tissue SKPO-1 is produced, we used indirect immunofluorescence to visualize SKPO-1 localization in young adult animals raised under standard conditions. Using rabbits, a polyclonal primary antibody was raised against a chemically synthesized SKPO-1 peptide (see Materials and Methods). A freeze-cracking methodology was used to disrupt the cuticle of the animal and allow for internal fixation and staining (Seydoux and Dunn 1997). Following fixation, the samples were double stained with the polyclonal primary antibody to SKPO-1 and an Alexa Fluor 488 secondary antibody. Localization was clearly observed in the hypodermis of wild-type animals, and an example is shown in Figure 5, A–C and D–F (40 times and 10 times, respectively). Note the strong fluorescent staining just under the outermost layer of the animal. No internal organs

showed evidence of staining, including the gonad and the intestine, which in this animal had become partially separated from the rest of the body. In contrast to wild type, *skpo-1* mutant animals showed no evidence of staining in any organ (Figure 5, G–I). We also examined wild-type and *skpo-1* mutant animals that had been exposed to *E. faecalis* for 24 hr prior to staining and did not observe any differences in the localization of SKPO-1 (data not shown).

By using a secondary antibody conjugated to 10 nm gold particles for immunogold labeling, we examined localization at higher resolution by transmission electron microscopy (TEM). As seen in Figure 5J, the black dots, indicating the gold particles, were located just under the cuticle layer in association with the hypodermal cells of wild-type animals. Hardly any particles were observed in the *skpo-1* mutant animals (Figure 5K). The few observed were randomly scattered. In total, these results demonstrate SKPO-1 is both physically present and functionally active in the *C. elegans* hypodermis.

Discussion

In this work, we demonstrated that a previously unstudied *C. elegans* protein with a peroxidase domain, F49E12.1, plays a protective role during infection with *E. faecalis*. We named this protein SKPO-1 for ShkT-containing peroxidase, because it contains an N-terminal ShkT-like domain. The ShkT domain family was originally defined as a potassium channel blocker in the sea anemone (*Metridium senile*).

Figure 5 SKPO-1 localizes to the *C. elegans* hypodermis. (A–F and J) Wild-type and (G–I and K) *skpo-1* mutant worms were immunostained with anti-SKPO-1 polyclonal antibodies and imaged using fluorescence (A–I) or transmission electron microscopy (J and K), respectively. (A–C) The $\times 40$ magnification of a wild-type worm shows hypodermal SKPO-1 localization. (D–F) The $\times 10$ magnification of a wild-type worm. (G–I) The $\times 10$ magnification of a *skpo-1* worm shows loss of SKPO-1 staining. (J) In wild-type worms, the black dots, indicative of immunogold labeling, are localized beneath the cuticle layer, but external to the outer hypodermal cell surface. (K) In *skpo-1* mutant worms, very few black dots are observed. White arrowhead, apical hypodermal surface; black arrowhead, 10 nm gold-labeled secondary to SKPO-1). Microscopy images are representative of >100 (fluorescent) or >10 (TEM) wild-type and *skpo-1* mutant worms observed.

Binding to the potassium channel requires two conserved residues that are not found in this particular ShkT-like domain of **SKPO-1** or others from *C. elegans* (data not shown). It is postulated that the more general function of ShkT-like domains are as contact surfaces for protein interactions (Tsang *et al.* 2007). For this reason, **SKPO-1** and other ShkT-containing peroxidases may be most closely related to the peroxidasins, subfamily 2 of the peroxidase-cyclooxygenase superfamily, and in fact, **SKPO-1** was placed in this family by phylogenetic analysis (Soudi *et al.* 2012). These peroxidase domain-containing proteins also have protein interaction domains, but they are typically type C-like immunoglobulin domains, leucine-rich repeats, or von Willebrand factor C modules. This is in contrast to the mammalian peroxidases, MPO, LPO, and eosinophil peroxidase that lack these extra domains and belong to subfamily 1 (Zamocky *et al.* 2008). Additionally,

it is thought that subfamily 1 evolved from subfamily 2 (Zamocky *et al.* 2008). Other *C. elegans* peroxidase domain-containing proteins that have a ShkT-like domain include **MLT-7** (Thein *et al.* 2009), **C16C8.2**, and **F32A5.2**, but no study has yet addressed this domain's function in the context of a peroxidase.

An unexpected finding was the discovery that **SKPO-1** is located in the hypodermis and is functionally protective in this tissue against *E. faecalis* infection. Infection of *C. elegans* with *E. faecalis* results in colonization of the gut, leading to distension of the intestinal lumen and clear signs of physical damage, such as effacement of the microvilli (Garsin *et al.* 2001; Cruz *et al.* 2013). Though the worm is exposed to *E. faecalis* on its outer surface as it moves through the pathogen lawn, there is no notable colonization or characterized physical damage to the cuticle or hypodermis. In contrast,

bacterial pathogens such as *Microbacterium nematophilum*, *Xenorhabdus nematophila*, and *Yersinia pestis* adhere to and colonize the cuticle surface (Hodgkin *et al.* 2000; Couillault and Ewbank 2002; Darby *et al.* 2002). The natural fungal pathogen *Drechmeria coniospora* initially adheres to the cuticle and then penetrates the hypodermis, whereas the human fungal pathogen *Candida albicans* first colonizes the intestine and eventually penetrates the cuticle from the inside out (Jansson *et al.* 1985; Breger *et al.* 2007). We previously showed that tissue-specific loss of *bli-3* in the hypodermis also increased susceptibility of *C. elegans* to *E. faecalis* (Chavez *et al.* 2009). Based on the protective effects of *SKPO-1* and *BLI-3* in this tissue, we postulate that a hypodermal immune response does play some role in protecting *C. elegans* during infection with *E. faecalis*. The question is by what mechanism?

Several models for how these proteins might exert their protective effects can be imagined. It could be that loss of *skpo-1* results in a weaker cuticle barrier that increases susceptibility. Alternatively, *SKPO-1* could use H₂O₂ produced by *BLI-3* to form more potent antioxidants, analogous to the human Duox/LPO system on mucosal surfaces (Conner *et al.* 2002; Geiszt *et al.* 2003; Forteza *et al.* 2005). Or the H₂O₂ produced by *BLI-3* could act as a signaling molecule in either an autonomous or noncell-autonomous fashion. Unlike many signaling molecules, H₂O₂ is highly diffusible, and it could directly move into other tissues, such as the intestine, to signal responses. Interestingly, in zebrafish, H₂O₂ produced by a Duox in response to wounding acts as an attractant in a diffusion gradient to draw leukocytes to the area of damage (Niethammer *et al.* 2009). Clearly, the amount of H₂O₂ is important. We previously showed that loss of *BLI-3* and the resulting decrease in H₂O₂ production caused an increase in susceptibility to infection (Chavez *et al.* 2009), whereas in this work we show that loss of a peroxidase and a concurrent increase in H₂O₂ also increases susceptibility. A similar situation is apparent from studies of infection using the model host *Drosophila melanogaster* (Ha *et al.* 2005a,b). Loss of an intestinal DUOX enzyme or an intestinal catalase both increase susceptibility to infection, but with opposite effects on ROS levels in this tissue. It is possible that *SKPO-1* is catalyzing the degradation of excess H₂O₂ to prevent host damage. We formerly demonstrated, by a variety of means, that infection causes oxidative stress in *C. elegans* (Chavez *et al.* 2007; Mohri-Shiomi and Garsin 2008; van der Hoeven *et al.* 2011), much of it dependent on *BLI-3* activity (van der Hoeven *et al.* 2011).

Interestingly, there is evidence for a hypodermal response to several pathogens that are thought to mainly cause infection in the intestine. Microarray studies that examined the transcriptional responses of *C. elegans* to *E. faecalis*, *Serratia marcescens*, *Erwinia carotovora*, *Photorhabdus luminescens*, *S. aureus*, and *P. aeruginosa* all noted a dramatic downregulation in the expression of genes related to cuticle biosynthesis, such as those encoding collagens (Wong *et al.* 2007; Irazoqui *et al.* 2010). The response is not thought to

be due to a general reduction of gene transcription in this tissue and may be indicative of several possibilities (Wong *et al.* 2007). The changes in transcription of the genes could be part of a protective response that is occurring in the hypodermis to protect against pathogens. Or the changes could be purposely caused by the pathogens as part of their virulence programs to damage the host. Alternatively, the changes in expression of the cuticle biosynthetic genes could be a neutral side effect resulting from alterations in signaling due to pathogen exposure. Another microarray study observed an increase in the expression of a large number of genes related to cuticle biosynthesis in an *arr-1* mutant that was found to have altered susceptibility to pathogens (Singh and Aballay 2012). *arr-1* encodes for arrestin-1, a G-protein coupled receptor (GPCR) adaptor protein that is necessary for GPCR signaling in several neurons of *C. elegans*. Whether these changes in resistance had anything to do with alterations in the amount of cuticle biosynthetic proteins being produced was not investigated (Singh and Aballay 2012). Overall, these studies suggest that pathogen exposure, even to those pathogens not thought to directly affect the cuticle, cause major changes in the expression of the cuticle biosynthetic genes that might be indicative of a response to the infection in the hypodermis.

In conclusion, we uncovered putative peroxidases that affected susceptibility to *E. faecalis* in *C. elegans*. We focused our characterization on *F49E12.1*, named *SKPO-1* because it contains a ShkT-like domain and a peroxidase domain. In addition to the pathogen susceptibility phenotype, loss of *skpo-1* resulted in a dumpy phenotype of uneven penetration, suggestive of some role in cuticle biosynthesis (Page and Johnstone 2007). In support of a functional peroxidase domain, we noted that *SKPO-1* shares the same critical residues as *MLT-7* in its active site (Thein *et al.* 2009), and animals mutant for *skpo-1* produce significantly more H₂O₂ during infection. The functional and physical location for the protein was shown to be the hypodermis, which was surprising since *E. faecalis* has been characterized as infecting the intestine of *C. elegans* (Garsin *et al.* 2001; Cruz *et al.* 2013). This result, along with other evidence from the literature (Wong *et al.* 2007; Irazoqui *et al.* 2010), suggests that the hypodermis plays an important role during exposure of *C. elegans* to many human pathogens that do not obviously colonize or damage the cuticle, warranting further investigation.

Acknowledgments

For the TEM imaging, we thank S. Kolodziej and P. Navarro (Department of Pathology and Laboratory Medicine EM Laboratory, University of Texas Health Science Center at Houston). We thank G. Ruvkun for the tissue-specific RNAi *C. elegans* strains and K. McCallum, S. Arur, K. Park, T. Furata, T. Leto, A. Page, M. Cruz, and W. Nauseef for helpful input on this project. We acknowledge the *Caenorhabditis* Genetics Center for providing us with many of the *C. elegans* strains

used in the study, J. Ahringer and Geneservice for the bacterial strains for RNAi, and WormBase for their help and advice in naming genes. This work was supported by Public Health Service grants R01AI076406 to D.A.G. and T32AI55449 to G.R.T. from the National Institute of Allergy and Infectious Diseases.

Literature Cited

- Breger, J., B. B. Fuchs, G. Aperis, T. I. Moy, F. M. Ausubel *et al.*, 2007 Antifungal chemical compounds identified using a *C. elegans* pathogenicity assay. *PLoS Pathog.* 3: e18.
- Brenner, S., 1974 The genetics of *Caenorhabditis elegans*. *Genetics* 77: 71–94.
- Chavez, V., A. Mohri-Shiomi, A. Maadani, L. A. Vega, and D. A. Garsin, 2007 Oxidative stress enzymes are required for DAF-16-mediated immunity due to generation of reactive oxygen species by *Caenorhabditis elegans*. *Genetics* 176: 1567–1577.
- Chavez, V., A. Mohri-Shiomi, and D. A. Garsin, 2009 Ce-Duox1/BLI-3 generates reactive oxygen species as a protective innate immune mechanism in *Caenorhabditis elegans*. *Infect. Immun.* 77: 4983–4989.
- Conner, G. E., M. Salathe, and R. Forteza, 2002 Lactoperoxidase and hydrogen peroxide metabolism in the airway. *Am. J. Respir. Crit. Care Med.* 166: S57–S61.
- Couillault, C., and J. J. Ewbank, 2002 Diverse bacteria are pathogens of *Caenorhabditis elegans*. *Infect. Immun.* 70: 4705–4707.
- Cruz, M. R., C. E. Graham, B. C. Gagliano, M. C. Lorenz, and D. A. Garsin, 2013 *Enterococcus faecalis* inhibits hyphal morphogenesis and virulence of *Candida albicans*. *Infect. Immun.* 81: 189–200.
- Darby, C., J. W. Hsu, N. Ghorri, and S. Falkow, 2002 *Caenorhabditis elegans*: plague bacteria biofilm blocks food intake. *Nature* 417: 243–244.
- Dunny, G. M., B. L. Brown, and D. B. Clewell, 1978 Induced cell aggregation and mating in *Streptococcus faecalis*: evidence for a bacterial sex pheromone. *Proc. Natl. Acad. Sci. USA* 75: 3479–3483.
- Edens, W. A., L. Sharling, G. Cheng, R. Shapira, J. M. Kinkade *et al.*, 2001 Tyrosine cross-linking of extracellular matrix is catalyzed by Duox, a multidomain oxidase/peroxidase with homology to the phagocyte oxidase subunit gp91phox. *J. Cell Biol.* 154: 879–891.
- Engelmann, I., A. Griffon, L. Tichit, F. Montanana-Sanchis, G. Wang *et al.*, 2011 A comprehensive analysis of gene expression changes provoked by bacterial and fungal infection in *C. elegans*. *PLoS ONE* 6: e19055.
- Forteza, R., M. Salathe, F. Miot, R. Forteza, and G. E. Conner, 2005 Regulated hydrogen peroxide production by Duox in human airway epithelial cells. *Am. J. Respir. Cell Mol. Biol.* 32: 462–469.
- Fraser, A. G., R. S. Kamath, P. Zipperlen, M. Martinez-Campos, M. Sohrmann *et al.*, 2000 Functional genomic analysis of *C. elegans* chromosome I by systematic RNA interference. *Nature* 408: 325–330.
- Garigan, D., A. L. Hsu, A. G. Fraser, R. S. Kamath, J. Ahringer *et al.*, 2002 Genetic analysis of tissue aging in *Caenorhabditis elegans*: a role for heat-shock factor and bacterial proliferation. *Genetics* 161: 1101–1112.
- Garsin, D. A., C. D. Sifri, E. Mylonakis, X. Qin, K. V. Singh *et al.*, 2001 A simple model host for identifying Gram-positive virulence factors. *Proc. Natl. Acad. Sci. USA* 98: 10892–10897.
- Garsin, D. A., J. M. Villanueva, J. Begun, D. H. Kim, C. D. Sifri *et al.*, 2003 Long-lived *C. elegans daf-2* mutants are resistant to bacterial pathogens. *Science* 300: 1921.
- Geiszt, M., K. Lekstrom, S. Brenner, S. M. Hewitt, R. Dana *et al.*, 2003 NAD(P)H oxidase 1, a product of differentiated colon epithelial cells, can partially replace glycoprotein 91phox in the regulated production of superoxide by phagocytes. *J. Immunol.* 171: 299–306.
- Ha, E. M., C. T. Oh, Y. S. Bae, and W. J. Lee, 2005a A direct role for dual oxidase in *Drosophila* gut immunity. *Science* 310: 847–850.
- Ha, E. M., C. T. Oh, J. H. Ryu, Y. S. Bae, S. W. Kang *et al.*, 2005b An antioxidant system required for host protection against gut infection in *Drosophila*. *Dev. Cell* 8: 125–132.
- Hodgkin, J., P. E. Kuwabara, and B. Corneliussen, 2000 A novel bacterial pathogen, *Microbacterium nematophilum*, induces morphological change in the nematode *C. elegans*. *Curr. Biol.* 10: 1615–1618.
- Hope, I. A., 1999 *C. elegans: A Practical Approach*, Oxford University Press, Oxford.
- Irazoqui, J. E., A. Ng, R. J. Xavier, and F. M. Ausubel, 2008 Role for beta-catenin and HOX transcription factors in *Caenorhabditis elegans* and mammalian host epithelial-pathogen interactions. *Proc. Natl. Acad. Sci. USA* 105: 17469–17474.
- Irazoqui, J. E., E. R. Troemel, R. L. Feinbaum, L. G. Luhachack, B. O. Cezairliyan *et al.*, 2010 Distinct pathogenesis and host responses during infection of *C. elegans* by *P. aeruginosa* and *S. aureus*. *PLoS Pathog.* 6: e1000982.
- Jansson, H. B., A. Jeyaprakash, and B. M. Zuckerman, 1985 Differential adhesion and infection of nematodes by the endoparasitic fungus *Meria coniospora* (Deuteromycetes). *Appl. Environ. Microbiol.* 49: 552–555.
- Kamath, R. S., A. G. Fraser, Y. Dong, G. Poulin, R. Durbin *et al.*, 2003 Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature* 421: 231–237.
- Kennedy, S., D. Wang, and G. Ruvkun, 2004 A conserved siRNA-degrading RNase negatively regulates RNA interference in *C. elegans*. *Nature* 427: 645–649.
- Kim, D. H., R. Feinbaum, G. Allosing, F. E. Emerson, D. A. Garsin *et al.*, 2002 A conserved p38 MAP kinase pathway in *Caenorhabditis elegans* innate immunity. *Science* 297: 623–626.
- Klebanoff, S. J., 2005 Myeloperoxidase: friend and foe. *J. Leukoc. Biol.* 77: 598–625.
- Meitzler, J. L., and P. R. Ortiz de Montellano, 2009 *Caenorhabditis elegans* and human dual oxidase 1 (DUOX1) “peroxidase” domains: insights into heme binding and catalytic activity. *J. Biol. Chem.* 284: 18634–18643.
- Meitzler, J. L., and P. R. Ortiz de Montellano, 2011 Structural stability and heme binding potential of the truncated human dual oxidase 2 (DUOX2) peroxidase domain. *Arch. Biochem. Biophys.* 512: 197–203.
- Meitzler, J. L., R. Brandman, and P. R. Ortiz de Montellano, 2010 Perturbed heme binding is responsible for the blistering phenotype associated with mutations in the *Caenorhabditis elegans* dual oxidase 1 (DUOX1) peroxidase domain. *J. Biol. Chem.* 285: 40991–41000.
- Meitzler, J. L., S. Hinde, B. Banfi, W. M. Nauseef, and P. R. Ortiz de Montellano, 2013 Conserved cysteine residues provide a protein-protein interaction surface in dual oxidase (DUOX) proteins. *J. Biol. Chem.* 288: 7147–7157.
- Melo, J. A., and G. Ruvkun, 2012 Inactivation of conserved *C. elegans* genes engages pathogen- and xenobiotic-associated defenses. *Cell* 149: 452–466.
- Mohri-Shiomi, A., and D. A. Garsin, 2008 Insulin signaling and the heat shock response modulate protein homeostasis in the *Caenorhabditis elegans* intestine during infection. *J. Biol. Chem.* 283: 194–201.
- Moskwa, P., D. Lorentzen, K. J. Excoffon, J. Zabner, P. B. McCray, Jr. *et al.*, 2007 A novel host defense system of airways is defective in cystic fibrosis. *Am. J. Respir. Crit. Care Med.* 175: 174–183.

- Niethammer, P., C. Grabher, A. T. Look, and T. J. Mitchison, 2009 A tissue-scale gradient of hydrogen peroxide mediates rapid wound detection in zebrafish. *Nature* 459: 996–999.
- Ortiz de Montellano, P. R., 2008 Mechanism and role of covalent heme binding in the CYP4 family of P450 enzymes and the mammalian peroxidases. *Drug Metab. Rev.* 40: 405–426.
- Page, A. P., and I. L. Johnstone, 2007 The cuticle (March 19, 2007), *WormBook*, ed. The *C. elegans* Research Community, *WormBook*, doi/10.1895/wormbook.1.138.1, <http://www.wormbook.org>.
- Pujol, N., O. Zugasti, D. Wong, C. Couillault, C. L. Kurz *et al.*, 2008 Anti-fungal innate immunity in *C. elegans* is enhanced by evolutionary diversification of antimicrobial peptides. *PLoS Pathog.* 4: e1000105.
- Rada, B., K. Lekstrom, S. Damian, C. Dupuy, and T. L. Leto, 2008 The *Pseudomonas* toxin pyocyanin inhibits the dual oxidase-based antimicrobial system as it imposes oxidative stress on airway epithelial cells. *J. Immunol.* 181: 4883–4893.
- Rahme, L. G., E. J. Stevens, S. F. Wolfort, J. Shao, R. G. Tompkins *et al.*, 1995 Common virulence factors for bacterial pathogenicity in plants and animals. [see comments] *Science* 268: 1899–1902.
- Seydoux, G., and M. A. Dunn, 1997 Transcriptionally repressed germ cells lack a subpopulation of phosphorylated RNA polymerase II in early embryos of *Caenorhabditis elegans* and *Drosophila melanogaster*. *Development* 124: 2191–2201.
- Shapira, M., B. J. Hamlin, J. Rong, K. Chen, M. Ronen *et al.*, 2006 A conserved role for a GATA transcription factor in regulating epithelial innate immune responses. *Proc. Natl. Acad. Sci. USA* 103: 14086–14091.
- Simmer, F., C. Moorman, A. M. van der Linden, E. Kuijk, P. V. van den Berghe *et al.*, 2003 Genome-wide RNAi of *C. elegans* using the hypersensitive rrf-3 strain reveals novel gene functions. *PLoS Biol.* 1: E12.
- Singh, V., and A. Aballay, 2012 Endoplasmic reticulum stress pathway required for immune homeostasis is neurally controlled by arrestin-1. *J. Biol. Chem.* 287: 33191–33197.
- Soudi, M., M. Zamocky, C. Jakopitsch, P. G. Furtmuller, and C. Obinger, 2012 Molecular evolution, structure, and function of peroxidasins. *Chem. Biodivers.* 9: 1776–1793.
- Sumimoto, H., 2008 Structure, regulation and evolution of Nox-family NADPH oxidases that produce reactive oxygen species. *FEBS J.* 275: 3249–3277.
- Tenor, J. L., B. A. McCormick, F. M. Ausubel, and A. Aballay, 2004 *Caenorhabditis elegans*-based screen identifies *Salmonella* virulence factors required for conserved host-pathogen interactions. *Curr. Biol.* 14: 1018–1024.
- Thein, M. C., A. D. Winter, G. Stepek, G. McCormack, G. Stapleton *et al.*, 2009 Combined extracellular matrix cross-linking activity of the peroxidase MLT-7 and the dual oxidase BLI-3 is critical for post-embryonic viability in *Caenorhabditis elegans*. *J. Biol. Chem.* 284: 17549–17563.
- Tsang, S. W., C. Q. Nguyen, D. H. Hall, and K. L. Chow, 2007 mab-7 encodes a novel transmembrane protein that orchestrates sensory ray morphogenesis in *C. elegans*. *Dev. Biol.* 312: 353–366.
- van der Hoeven, R., K. C. McCallum, M. R. Cruz, and D. A. Garsin, 2011 Ce-Duox1/BLI-3 generated reactive oxygen species trigger protective SKN-1 activity via p38 MAPK signaling during infection in *C. elegans*. *PLoS Pathog.* 7: e1002453.
- Wong, D., D. Bazopoulou, N. Pujol, N. Tavernarakis, and J. J. Ewbank, 2007 Genome-wide investigation reveals pathogen-specific and shared signatures in the response of *Caenorhabditis elegans* to infection. *Genome Biol.* 8: R194.
- Zamocky, M., C. Jakopitsch, P. G. Furtmuller, C. Dunand, and C. Obinger, 2008 The peroxidase-cyclooxygenase superfamily: reconstructed evolution of critical enzymes of the innate immune system. *Proteins* 72: 589–605.

Communicating editor: B. Lazzaro

GENETICS

Supporting Information

<http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.113.160606/-/DC1>

The SKPO-1 Peroxidase Functions in the Hypodermis to Protect *Caenorhabditis elegans* From Bacterial Infection

George R. Tiller and Danielle A. Garsin

SUPPLEMENTARY MATERIALS AND METHODS

RNA isolation and qRT-PCR

Prior to RNA isolation, wild type and *skpo-1* mutant animals were grown on *cdc-25.1* dsRNA expressing *E. coli* HT115 from the egg to L4 larval stage. L4 animals were then exposed to either *cdc-25.1* RNAi or *E. faecalis* OG1RF for 18 hours. The RNA was isolated using Trizol (Invitrogen) according to the manufacturer. RNA samples were treated with DNase I to eliminate contaminating DNA by the Turbo DNA free kit (Applied Biosystem) according to the manufacturer. qRT-PCR was performed as described (VAN DER HOEVEN *et al.* 2011). Primers used are listed in Table S6 (*act-1* served as the reference gene).

Bacterial Colonization

The CFU analysis was conducted in a manner similar to past work (GARSIN *et al.* 2001). Briefly, L4 wild type and *skpo-1* mutant animals grown on *E. coli* OP50 were exposed on 100 mm plates containing BHI-agar with gentamycin (10 µg/mL) seeded with 100 µl of *E. faecalis* OG1RF for either 12 or 36 hours at 25° C. Worms were washed 3 times with M9 buffer at 1.4 rpm. Worms were then washed twice with 25 mM tetramisole hydrochloride to prevent ingestion of the antibiotic treatment. Worms were incubated at room temperature for 60 minutes in 25 mM tetramisole hydrochloride supplemented with ampicillin and kanamycin, both at 1 mg/mL, to kill surface-attached *E. faecalis*. Worms were collected at 1.4 rpm and washed twice with 25 mM tetramisole hydrochloride prior to grinding. Ten worms in 10 µl were transferred to 200 µl of M9 and ground for 1 minute using a motorized pestle (Kontes cordless cat# K749540-0000 and pestles cat# K749521-1590). Serial dilutions were performed and 100 µl of each dilution plated onto 100 mm BHI gentamycin 10 µg/mL plates for 24 hours at 37° C.

Figure S1 Survival of *cdc-25.1* RNAi-treated *skpo-1* animal lines on *E. faecalis*. Wild type and *skpo-1* worms were grown on *cdc-25.1* dsRNA expressing *E. coli* HT115 from L1 through L4 larval stage prior to exposure to *E. faecalis* OG1RF. The *skpo-1* worms (GF89, GF90 and GF91) were more susceptible to *E. faecalis* relative to wild type (*P*-values < 0.0001, 0.0110, and < 0.0001, respectively).

Figure S2 Lifespan defect of *skpo-1* mutants on *E. coli* OP50 without prior *cdc-25.1* exposure. Wild type and *skpo-1* worms were grown on *E. coli* OP50 from L1 through L4 larval stage prior to the longevity assay in which they were transferred to NGM plates with FUDR (100 µg/ml) seeded with concentrated (20X) *E. coli* OP50. The average lifespan of the *skpo-1* worms was significantly shorter than their wild type, counterparts (P -value < 0.0001).

Figure S3 *skpo-1* mutants display a lifespan defect on heat-killed *E. coli* OP50. Wild type and *skpo-1* worms were grown on *cdc-25.1* dsRNA expressing *E. coli* HT115 from L1 through L4 larval stage prior to the longevity assay. The worms were transferred to NGM plates like in Figure S2; however, the *E. coli* OP50 had been heat-killed. We observed that the *skpo-1* worms displayed a longevity defect on heat-killed OP50, relative to wild type worms (P -value <0.0001).

Figure S4 The *skpo-1* mutant does not have increased intestinal bacterial load during *E. faecalis* infection. L4 wild type and *skpo-1* mutant animals were exposed to *E. faecalis* at 25° C for 12 or 36 hours, respectively. CFU values for both wild type and *skpo-1* animals indicated that there is no significant difference in intestinal colonization by *E. faecalis*. Three biological replicates for each strain were used per time point and the experiment repeated twice—i.e a total n = 60 per strain. *P*-values were 0.3913 and 0.3592 for 12 and 36 hours, respectively.

Figure S5 *clec-60* expression is elevated in *skpo-1* mutant animals, relative to wild type, in response to *E. faecalis* infection. L4 wild type and *skpo-1* mutant animals were exposed to *E. faecalis* for 18 hours at 25° C. Three technical replicates per strain per gene were assayed for relative expression of innate immune response genes and the experiment was repeated, independently, twice. These genes were previously found to display increased expression in wild type animals in response to *E. faecalis* (ENGELMANN *et al.* 2011). Of the genes surveyed, *clec-60* was the only gene to display a significant increase in expression during infection, relative to *act-1*. *P*-values were 0.1912, 0.1768, 0.04432, and 0.4684 for *clec-35*, 42, 60 and 71, respectively.

Table S1 Median survival and *P*-values of *E. faecalis* OG1RF killing assays

Figure number	Experiment number	Strain, exposure conditions during development	Median survival (days)	<i>P</i> -value
1A	1	wild type, VC RNAi	8	C
		wild type, <i>skpo-1</i> RNAi	7	=0.0404
	2	wild type, VC RNAi	7	C
		wild type, <i>skpo-1</i> RNAi	6	=0.0331
	3	<i>eri-1</i> , VC RNAi	8	C
		<i>eri-1</i> , <i>skpo-1</i> RNAi	7	=0.001
	4	<i>eri-1</i> , VC RNAi	7	C
		<i>eri-1</i> , <i>skpo-1</i> RNAi	5	=0.0053
	5	<i>eri-1</i> , VC RNAi	9	C
		<i>eri-1</i> , <i>skpo-1</i> RNAi	8	=0.0037
1C	1	wild type, <i>cdc-25.1</i> RNAi	10	C
		<i>skpo-1</i> , <i>cdc-25.1</i> RNAi	6	<0.0001
	2	wild type, <i>cdc-25.1</i> RNAi	10	C
		<i>skpo-1</i> , <i>cdc-25.1</i> RNAi	8	<0.0001
	3	wild type, <i>cdc-25.1</i> RNAi	13	C
		<i>skpo-1</i> , <i>cdc-25.1</i> RNAi	8	=0.0016
	4	wild type, <i>cdc-25.1</i> RNAi	13	C
		<i>skpo-1</i> , <i>cdc-25.1</i> RNAi	8	=0.0005
	5	wild type, <i>cdc-25.1</i> RNAi	13	C
		<i>skpo-1</i> , <i>cdc-25.1</i> RNAi	7	<0.0001
4A	1	<i>wrt-2</i> , VC RNAi	5	C
		<i>wrt-2</i> , <i>skpo-1</i> RNAi	4	=0.0002
	2	<i>wrt-2</i> , VC RNAi	4	C
		<i>wrt-2</i> , <i>skpo-1</i> RNAi	3	=0.0176
	3	<i>wrt-2</i> , VC RNAi	5	C
		<i>wrt-2</i> , <i>skpo-1</i> RNAi	4	=0.0376
	4	<i>wrt-2</i> , VC RNAi	4	C
		<i>wrt-2</i> , <i>skpo-1</i> RNAi	4	=0.0058
4C	1	<i>vha-6</i> , VC RNAi	11	C
		<i>vha-6</i> , <i>skpo-1</i> RNAi	11	=0.6379
	2	<i>vha-6</i> , VC RNAi	11	C
		<i>vha-6</i> , <i>skpo-1</i> RNAi	11	=0.735
S1	1	wild type, OP50	8	C
		<i>skpo-1</i> *, OP50	3	<0.0001
	2	wild type, OP50	8	C
		<i>skpo-1</i> *, OP50	3	<0.0001

C = control, *Data only shown for GF89 *skpo-1* strain.

Table S2 Median survival and *P*-values of *E. coli* OP50 longevity assays

Figure number	Experiment number	Strain, exposure conditions during development	Median survival (Days)	<i>P</i> -value
1B	1	<i>eri-1</i> , VC RNAi	10	C
		<i>eri-1, skpo-1</i> RNAi	10	=0.3772
	2	<i>eri-1</i> , VC RNAi	10	C
		<i>eri-1, skpo-1</i> RNAi	10	=0.8948
1D	1	wild type, <i>cdc-25.1</i> RNAi	10	C
		<i>skpo-1, cdc-25.1</i> RNAi	8	<0.0001
	2	wild type, <i>cdc-25.1</i> RNAi	14	C
		<i>skpo-1, cdc-25.1</i> RNAi	11	<0.0001
	3	wild type, <i>cdc-25.1</i> RNAi	14	C
		<i>skpo-1, cdc-25.1</i> RNAi	11	<0.0001
4B	1	<i>vha-6</i> , VC RNAi	10	C
		<i>vha-6, skpo-1</i> RNAi	10	=0.9997
	2	<i>vha-6</i> , VC RNAi	11	C
		<i>vha-6, skpo-1</i> RNAi	10	=0.4484
4D	1	<i>wrt-2</i> , VC RNAi	11	C
		<i>wrt-2, skpo-1</i> RNAi	11	=0.6379
	2	<i>wrt-2</i> , VC RNAi	11	C
		<i>wrt-2, skpo-1</i> RNAi	11	=0.7235
S2	1	wild type, OP50	14	C
		<i>skpo-1</i> , OP50	11	<0.0001
	2	wild type, OP50	14	C
		<i>skpo-1</i> , OP50	11	<0.0001
S3*	1	N2, <i>cdc-25.1</i> RNAi*	16	C
		<i>skpo-1, cdc-25.1</i> RNAi*	13	<0.0001
	2	N2, <i>cdc-25.1</i> RNAi*	14	C
		<i>skpo-1, cdc-25.1</i> RNAi*	12	<0.0001

C = control, *Assayed on heat-killed OP50

Table S3 Data for Relative Mortality Calculation

Figure number	Experiment number	Strain	Relative mortality	P-value
1F	Average of experiments 1-4	N2	1	C
			1	
			1	
			<u>1</u>	
			Avg = 1	
		<i>skpo-1</i>	1.312	=0.0091
			1.452	
			1.654	
			1.253	
			Avg = 1.418	

C = control

Table S4 Median survival and *P*-values of *P. aeruginosa* PA14 killing assays

Figure Number	Experiment number	Strain, exposure conditions during development	Median survival (Hours)	<i>P</i> -value
1E	1	N2, <i>cdc-25.1</i> RNAi	91	C
		<i>skpo-1</i> , <i>cdc-25.1</i> RNAi	88	=0.3783
	2	N2, <i>cdc-25.1</i> RNAi	89	C
		<i>skpo-1</i> , <i>cdc-25.1</i> RNAi	69	=0.0639
	3	N2, <i>cdc-25.1</i> RNAi	72	C
		<i>skpo-1</i> , <i>cdc-25.1</i> RNAi	72	=0.2613

Table S5 Primers for cDNA inserts ligated into pL4440 construct for dsRNA production in *E. coli*

Oligo name	Sequence
<i>F32A52.A</i> 5F	5' TTG CGG CCG CAA TTC ACC AGG GAA TTT ACA CC 3'
<i>F32A52.A</i> 3R	5' CCT CGA GGG TGG TCT TTT CAC ATT CTG G 3'
<i>F32A52.B</i> 5F	5' TTG CGG CCG CAA CAA CCA CTC ATT TCA CCA GG 3'
<i>F32A52.B</i> 3R	5' CCT CGA GGC TGT AGT TGT TAC TCT TTG TGG 3'
<i>pxn-1</i> 5F	5' TTG CGG CCG CAA GGA CTC TGG AAG GTA CAC 3'
<i>pxn-1</i> 3R	5' CCT CGA GGA TAC TTG GCA CTT CCA CTC T 3'

Table S6 qRT-PCR primers for assessing expression of innate immune response genes in response to *E. faecalis*

Oligo name	Sequence
<i>clec-35</i> 5F	5' AGA TGC TGG ACA GTG GAA AAG 3'
<i>clec-35</i> 3R	5' GTG CGG AGT ATT GTA GCG TAG 3'
<i>clec-42</i> 5F	5' GTA ACT CCG TAT TGG CTG GG 3'
<i>clec-42</i> 3R	5' GTA AAC GCA GCT TCC AAT CTC 3'
<i>clec-60</i> 5F	5' TGT AAG AGA ACA GTT GGA ACC C 3'
<i>clec-60</i> 3R	5' TAT GTG CAT GGG TAC TGA TCG 3'
<i>clec-71</i> 5F	5' ACG ACA GGA AGT GAT GTA TTG G 3'
<i>clec-71</i> 3R	5' TTG ACG GAC TTT AGC CAC TG 3'
<i>act-1</i> 5F	5' ACC ATG TAC CCA GGA ATT GC 3'
<i>act-1</i> 3R	5' TGG AAG GTG GAG AGG GAA G 3'

SUPPLEMENTARY LITERATURE CITED

Engelmann, I., A. Griffon, L. Tichit, F. Montanana-Sanchis, G. Wang *et al.*, 2011 A comprehensive analysis of gene expression changes provoked by bacterial and fungal infection in *C. elegans*. PLoS One 6: e19055.

Garsin, D. A., C. D. Sifri, E. Mylonakis, X. Qin, K. V. Singh *et al.*, 2001 A simple model host for identifying Gram-positive virulence factors. Proceedings of the National Academy of Sciences of the United States of America 98: 10892-10897.

van der Hoeven, R., K. C. McCallum, M. R. Cruz and D. A. Garsin, 2011 Ce-Duox1/BLI-3 Generated Reactive Oxygen Species Trigger Protective SKN-1 Activity via p38 MAPK Signaling during Infection in *C. elegans*. PLoS Pathog 7: e1002453.