Modification of spasticity by transcutaneous spinal cord stimulation in individuals with incomplete spinal cord injury

Ursula S. Hofstoetter¹, William B. McKay², Keith E. Tansey^{2,3,4}, Winfried Mayr¹, Helmut Kern^{5,6}, Karen Minassian¹

¹Center for Medical Physics and Biomedical Engineering, Medical University of Vienna, Vienna, Austria, ²Hulse Spinal Cord Injury Lab, Crawford Research Institute, Shepherd Center, Atlanta, GA, USA, ³Departments of Neurology and Physiology, Emory University School of Medicine, Atlanta, GA, USA, ⁴Spinal Cord Injury Clinic, Atlanta Veterans Administration Medical Center, Atlanta, GA, USA, ⁵Ludwig Boltzmann Institute of Electrical Stimulation and Physical Rehabilitation, Vienna, Austria, ⁶Department of Physical Medicine and Rehabilitation, Wilhelminenspital Wien, Vienna, Austria

Context/objective: To examine the effects of transcutaneous spinal cord stimulation (tSCS) on lower-limb spasticity.

Design: Interventional pilot study to produce preliminary data.

Setting: Department of Physical Medicine and Rehabilitation, Wilhelminenspital, Vienna, Austria.

Participants: Three subjects with chronic motor-incomplete spinal cord injury (SCI) who could walk ≥ 10 m.

Interventions: Two interconnected stimulating skin electrodes ($\varnothing$ 5 cm) were placed paraspinally at the T11/T12 vertebral levels, and two rectangular electrodes (8×13 cm) on the abdomen for the reference. Biphasic 2 ms-width pulses were delivered at 50 Hz for 30 minutes at intensities producing paraesthesias but no motor responses in the lower limbs.

Outcome measures: The Wartenberg pendulum test and neurological recordings of surface-electromyography (EMG) were used to assess effects on exaggerated reflex excitability. Non-functional co-activation during volitional movement was evaluated. The timed 10-m walk test provided measures of clinical function.

Results: The index of spasticity derived from the pendulum test changed from 0.8 ± 0.4 pre- to 0.9 ± 0.3 post-stimulation, with an improvement in the subject with the lowest pre-stimulation index. Exaggerated reflex responsiveness was decreased after tSCS across all subjects, with the most profound effect on passive lower-limb movement (pre- to post-tSCS EMG ratio: 0.2 ± 0.1), as was non-functional co-activation during voluntary movement. Gait speed values increased in two subjects by 39%.

Conclusion: These preliminary results suggest that tSCS, similar to epidurally delivered stimulation, may be used for spasticity control, without negatively impacting residual motor control in incomplete SCI. Further study in a larger population is warranted.

Keywords: Humans, Movement, Muscle spasticity, Spinal cord injuries, Spinal cord stimulation, Paraplegia, Tetraplegia, Rehabilitation

Introduction

Partial injury of descending spinal cord tracts alters the supraspinal motor control over the lumbosacral neural circuitry that is normally preserved caudal to the lesion. The consequences of such injuries are usually paresis or paralysis and spasticity of the lower

extremities.^{1,2} Also, when present, spasticity can negatively impact otherwise sufficient residual motor control.^{3,4}

Treatment modalities used for spasticity target the segmental pathophysiological changes within the spinal cord after the injury as well as changes in the muscle-tendon complex.⁵ They range from oral medication and physical interventions to motor nerve block injections and intrathecal drug delivery.^{5,6} Yet, therapeutic control of severe forms of spasticity, while

Correspondence to: Ursula S. Hofstoetter, Center for Medical Physics and Biomedical Engineering, Medical University of Vienna, Waehringer Guertel 18-20/4L, A-1090 Vienna, Austria. Email: ursula.hofstoetter@meduniwien.ac.at

maintaining or even enhancing residual motor capacities, has remained difficult.⁷ Surgical treatment of spasticity is based on the concept that spinal reflex circuitry becomes hyperactive after being deprived of supraspinal inhibitory control. By cutting the posterior roots, Otfried Foerster, in 1909, interrupted afferent inputs to the spinal cord to reduce the excitability of segmental neural circuits.⁸ More selective posterior root entry zone procedures were developed to decrease lower limb hypertonia and improve ambulation.⁹ However, such surgical approaches create permanent lesions within the neural tissue and have the potential to damage surviving neural function and a high incidence of recurrence of spasticity has been reported.¹⁰

The emergence of epidurally delivered electrical spinal cord stimulation (SCS) provided an alternative way to reversibly modify motor control in patients with upper motor neuron disorders.^{11–15} Richardson and McLone¹⁶ and Richardson *et al.*¹⁷ described the alleviation of severe lower limb spasticity in clinically complete thoracic spinal cord injury (SCI) by epidural stimulation applied over the lumbosacral spinal cord and cauda equina. In patients with motor-complete and incomplete SCI, epidural SCS was applied below and above the lesion at vertebral levels ranging from C2 to T10 and at stimulation frequencies of 30–1500 Hz, with variable results.^{18–22} Dimitrijevic *et al.*^{23,24} suggested that the diversity of physiological conditions after SCI and the placement of the epidural electrodes determined the efficacy of SCS in spasticity control. While interest in this technique declined in the 1990s, Pinter *et al.*²⁵ revisited its use in individuals with chronic SCI. They showed a significant suppression of severe lower limb spasticity when the epidural electrodes were placed over the lumbar posterior roots and the stimulation frequency was in a range of 50–100 Hz. The suggested mechanism was the modification of the excitability of neural circuits within the lumbar spinal cord through continuous posterior-root activation.^{25–27}

We previously described a transcutaneous stimulation method to depolarize large-diameter afferent fibers of

the L2–S2 posterior roots in humans.^{28–30} These input structures to the segmental neural circuits are hence (at least a sub-population of) the same neural targets as electrically stimulated by epidural implants over the lumbar spinal cord.^{22,30} In this exploratory work, we hypothesized that transcutaneous SCS (tSCS), when applied in a continuous mode at 50 Hz, can be used to modify the central state of excitability of lumbar neural circuits, and reduce spasticity.

Methods

Subjects and clinical data

After informed consent was obtained, three subjects (mean age 32.7 ± 4.1 years; one female) with chronic (10.7 ± 1.2 years post-onset) sensory- and motor-incomplete SCI were studied (Table 1). The participants were otherwise healthy adults. All subjects had spastic muscle hypertonia in the lower limbs and exhibited clonus but were able to complete the 10-m walk test without braces and were not taking anti-spasticity medications. The study was approved by the Ethics Committee of the City of Vienna, Austria, and conforms to the Helsinki Declaration of 1975, as revised in 2000.

Transcutaneous spinal cord stimulation

tSCS was applied using self-adhesive electrodes (Schwamedico GmbH, Ehringshausen, Germany). A pair of stimulating disk electrodes ($\varnothing = 5$ cm) was placed over the T11 and T12 spinous processes, manually identified by palpation, with one electrode on either side of the spine (Fig. 1A). A pair of rectangular reference electrodes ($8\text{ cm} \times 13\text{ cm}$ each) was placed over the lower anterior abdomen, in symmetry to the umbilicus. The two electrodes of each pair were connected together so as to function as single, larger electrodes. A constant-voltage stimulator delivered charge-balanced, symmetric, biphasic rectangular pulses of 2 ms width (1 ms per phase). Electrode placement was confirmed by eliciting posterior root-muscle reflexes (PRM reflexes) in lower limb muscles with the subjects relaxed in the supine position (Fig. 1B).^{29–31} The requirement was

Table 1 Subject characteristics

Subject no.	Sex	Age	Years post-injury	Level of SCI	AIS	Lower limb motor scores		WISCI	Device
						Left	Right		
1	M	38	9	C5	D	13	12	13	Walker
2	M	32	12	C5	D	12	17	16	Crutches
3	F	28	11	T9	D	11	13	16	Crutches

ASIA Impairment Scale (AIS) category, Walking Index for Spinal Cord Injury (WISCI), assistive devices needed for gait (Device). M, male; F, female.

Figure 1 Transcutaneous spinal cord stimulation. (A) Sketch of the placement of the stimulating and reference skin electrodes over the back and the lower abdomen, respectively, relative to the spine and the lumbosacral spinal cord. (B) The elicitation of posterior root-muscle reflexes in left (L) and right (R) quadriceps (Q), hamstrings (Ham), tibialis anterior (TA), and triceps surae (TS) was used as electrophysiological criterion to confirm the position of the stimulating paravertebral electrodes over the lumbar spinal cord. (C) The stimulation of posterior root afferents was verified by testing the recovery cycle of the evoked responses using double-stimuli at interstimulus intervals of 30, 50, and 100 ms. Electromyographic data derived from subject 2 in the supine position.

that single stimuli evoke PRM reflexes in the L2–L4 innervated quadriceps (Q) at lower thresholds than in the L5–S2 innervated triceps surae (TS).^{25,26,32} Paraspinal electrodes were relocated, up to 3 cm in either the rostral or caudal direction when needed to achieve this required PRM response distribution. Double-stimuli at interstimulus intervals of 30, 50, and 100 ms were applied to test the presence of post-stimulation attenuation of the detected responses to verify the stimulation of afferent fibers (Fig. 1C).^{29,33}

To induce the hypothesized neuromodulative effect on lumbar spinal neural circuits, tSCS was applied at a frequency of 50 Hz.²⁵ With the subject lying comfortably supine on an examination bed, the stimulation intensity was slowly increased starting from 0 V to allow for adaptation to the stimulation-induced effects. These effects included, in sequence of their occurrence with increasing stimulation: sensory perception of the stimulation under the paraspinal and subsequently the abdominal electrodes, contraction of the paraspinal and abdominal muscles, and paraesthesias in the lower limbs. tSCS was applied for 30 minutes at intensities producing paraesthesias in most of the lower limb dermatomes, yet subthreshold for generating lower limb

motor activity. Particularly, the average intensity applied amounted to $20 \text{ V} \pm 2 \text{ V}$ (per phase of the biphasic stimulation pulse), corresponding to $68\% \pm 16\%$ of individual lower limb PRM reflex threshold.^{19,24} Discomfort during the stimulation was not reported by any subject.

Assessments

All assessments were performed prior to and immediately following the 30-minute session of 50-Hz tSCS.

The Wartenberg pendulum test (WPT) is a quantitative method for the assessment of spastic hypertonia of the Q muscle.³⁴ With the subject in a sitting position, reclined at approximately 45° , the examiner lifted the relaxed test limb to a horizontal position, then released it and let it oscillate passively until it stopped in a gravity-neutral position. The test was repeated three times on each side.

Electromyographic (EMG) recordings from Q, hamstrings (Ham), tibialis anterior (TA), and TS muscles bilaterally were acquired using pairs of silver–silver chloride recording electrodes (Intec Medizintechnik GmbH, Klagenfurt, Austria), each placed centrally over the muscle bellies and oriented along the long

axis of the muscles with an inter-electrode distance of 3 cm.³⁵ Inter-electrode impedance was reduced to below 5 k Ω using abrasive paste. EMG signals were amplified (EMS-Handels GmbH, Korneuburg, Austria) with a gain of 502, filtered to a bandwidth of 10–500 Hz and digitized at 2048 samples per second per channel. Electro-goniometers (Penny & Giles Biometrics, Ltd., Gwent, UK) were used to measure knee and ankle movements. The goniometric data were synchronized to the EMG, bandpass-filtered between 0 and 100 Hz and sampled at 2048 Hz.

Reflex activities and motor tasks tested included, in the supine position, passive unilateral hip and knee flexion and extension movements, the manual elicitation of ankle clonus by rapid stretch of the Achilles tendon, and foot withdrawal by non-noxious mechanical plantar stimulation. Each of the maneuvers was repeated three times on each side, separated by phases of relaxation. In the sitting position, rhythmic voluntary unilateral ankle dorsiflexion and plantar flexion attempted at a self-selected speed was performed five times on each side.

The 10-m walking speed was acquired with subjects using walking aids (Table 1) as required for safety.³⁶ Each test was videotaped for offline visual examination.

In subjects 1 and 2, all segments of the assessment were completed in a single recording session. In subject 3, passive lower limb movements and plantar stimulation were performed in one session, and ankle clonus, volitional ankle movement, and the timed 10-m walk test were accomplished in a second session. The WPT was repeated in both sessions, and the respective measures were averaged, hence contributing single values per leg to the group results.

Data analysis

Data were analyzed offline using WinDaq Waveform Browser playback software (DATAQ Instruments, Inc., Akron, OH, USA) and Matlab R2012a (The MathWorks, Inc., Natick, MA, USA).

Wartenberg pendulum test

The index of spasticity, R_{2n} , was calculated as described by Bajd and Vodovnik³⁷:

$$R_{2n} = \frac{(\text{peak knee flexion angle of first swing}) - (\text{starting knee angle})}{1.6[(\text{final resting knee angle}) - (\text{starting knee angle})]}$$

Scores ≥ 1 signify non-spastic conditions, while $R_{2n} = 0$ is obtained in case of extreme spasticity. The index correlates well with the clinical evaluation of muscle

hypertonia graded on the Ashworth scale, and provides a more objective and finely graded measurement.³⁸ Here, mean R_{2n} values were averaged over the three repetitions on each side.

In addition, the magnitude of total ipsilateral lower limb EMG activity generated during the pendulum test was calculated as the sum of the root mean square (RMS) of the EMG activity across the four ipsilateral muscle groups, between the onset of the maneuver, drop of leg from horizontal position, and the time when the resting angle was reached. Data for each muscle were baseline-corrected by subtracting the RMS calculated from a 1-second period preceding the maneuver. Mean values of the three repetitions for each limb were evaluated. The corresponding value after tSCS was then individually normalized to the value before stimulation and normalized mean group results $\bar{A}_{\text{total}}$ of these ratios (derived from $n = 6$ lower limbs) were calculated.

Volitional ankle dorsiflexion and plantar flexion

The total maneuver-related magnitudes of ipsilateral muscle activation during five consecutive flexion and extension phases (identified by ankle goniometric data) were calculated based on the RMS values as described above and averaged. Agonist muscle activity during each movement phase was normalized to the total activity recorded across all ipsilateral muscles. Thus, the amount of TA activity (RMS_{TA}) relative to the total magnitude of ipsilateral muscle activation ($\text{RMS}_{\text{total}}$) during dorsiflexion was individually calculated as $\text{RMS}_{\text{TA}}/\text{RMS}_{\text{total}}$, and group results were obtained across the subjects. Accordingly, the relative TS activity during plantar flexion was computed as $\text{RMS}_{\text{TS}}/\text{RMS}_{\text{total}}$. Mean peak angles of the ankle movement for the five repetitions were also obtained.

Passive lower limb flexion-extension, ankle clonus, and plantar stimulation

In case of passive lower limb flexion and extension movements, base-line corrected RMS from ipsilateral muscles were calculated between the onsets and offsets plus 3 seconds for the movement identified by knee goniometric data. For ankle clonus and plantar stimulation, RMS of ipsilateral lower limb EMG activity was computed for 5-second time windows following the initiation of the maneuvers. Mean values for the maneuver-related magnitude of ipsilateral muscle activation were obtained from the three repetitions of each task, and normalized group mean results $\bar{A}_{\text{total}}$ were calculated from the available data sets.

10-m walk test

The time required to walk a distance of 10 m before and after stimulation with the same walking aids was measured.

Results

Pendulum test

The average spasticity index R_{2n} was 0.8 ± 0.4 before and 0.9 ± 0.3 after tSCS. The group result $\bar{A}_{\text{total}}$ of the magnitude of ipsilateral muscle activation after stimulation normalized to the corresponding value before stimulation was 0.9 ± 0.0 . The EMG activities produced during the WPT were decreased evenly across all subjects after stimulation, while the R_{2n} score improved in subject 3 only, who had the lowest pre-stimulation scores, with no changes in the R_{2n} in subjects 1 and 2. The R_{2n} scores obtained in subject 3 in the two sessions were reproducible for both lower limbs and the pre- and post-tSCS assessments, respectively. Before stimulation, the pendulum test documented restraints in the free-swinging movements of the lower leg in subject 3 (Fig. 2, left). The passive, gravity-driven movement toward the gravity-neutral end-point position was interrupted by numerous short bursts of stretch-elicited Q activity, accompanied by activation of Ham and, to some extent, of TA. After stimulation, activation of the Q only occurred during the initial two flexion–extension oscillations of the leg and the knee movements were a pseudo-sinusoidal, damped motion (Fig. 2, right).

Effects of tSCS on exaggerated reflex responses

Passive hip and knee flexion and extension movements (Fig. 3A) before tSCS induced responses with a spread of EMG activity to all ipsilateral lower limb muscles. These responses were almost completely absent in the post-tSCS assessment. Pre-tSCS, a brisk manual

stretch of the calf muscle elicited clonus-like activity in TS that spread to Ham and TA in the example given in Fig. 3B. After tSCS, no muscle activities except for a brief stretch-related reflex response in TS were detected when the same maneuver was performed. Non-noxious, cutaneous stimulation to the plantar surface of the foot with a blunt probe (Fig. 3C) resulted in withdrawal-reflex EMG activity in all ipsilateral lower limb muscles but after tSCS, was largely diminished.

The general decline in exaggerated reflex activities was reflected by the group results $\bar{A}_{\text{total}}$ of the maneuver-related magnitudes of ipsilateral lower limb EMG activity obtained for the post-tSCS assessment normalized to the corresponding values before tSCS (Fig. 3D). The post-tSCS RMS values were considerably reduced across all maneuvers and lower limbs, with the exception of the values derived from the ankle clonus tested in the respective left lower limbs of subjects 1 and 3. Specifically, $\bar{A}_{\text{total}}$ amounted to: passive lower limb movement, 0.2 ± 0.1 ; ankle clonus, 0.7 ± 0.3 ; and plantar stimulation, 0.4 ± 0.1 .

Effects of tSCS on voluntary movements

The EMG activity during volitional ankle dorsiflexion and plantar flexion performed before stimulation was generally characterized by co-activation patterns in the TA and TS muscles during both phases of movement (Fig. 4A, left). After stimulation, this co-activation was decreased (Fig. 4A, right). Furthermore, active range of motion for dorsiflexion was increased. The mean total range of ankle movement of the different subjects was increased after stimulation by a factor of 1.2 ± 0.1 . In addition, clonus-like activity in TA and TS that occurred in subjects 1 and 2 during the execution of the voluntary motor task was eliminated after tSCS (Fig. 4A and B).

Figure 2 Wartenberg pendulum test. Electromyographic (EMG) activity from quadriceps (Q), hamstrings (Ham), tibialis anterior (TA), and triceps surae (TS) along with knee goniometric data before and after transcutaneous spinal cord stimulation in subject 3. Dotted lines mark the drop of leg from the horizontal position and the final resting knee angle reached. After stimulation, EMG activity is clearly reduced and the knee angle oscillations are a damped, pseudo-sinusoidal motion.

Figure 3 Assessment of different manifestations of spasticity before and after transcutaneous spinal cord stimulation (tSCS). (A) Manually performed hip and knee flexion and extension movements; (B) the attempt to elicit an ankle clonus; and (C) a light stroke of the plantar surface with a blunt rod. Arrow heads in (B) and (C) depict onsets of the respective mechanical stimuli. Exemplary recordings were derived from subject 2. (D) Group results ($n = 6$) of maneuver-induced magnitudes of ipsilateral lower limb muscle activity after tSCS relative to corresponding values before tSCS. Note the general decline in exaggerated reflex activity in the post-tSCS assessment.

Group results (Fig. 4C) revealed that the magnitude of ipsilateral EMG activity during dorsiflexion remained almost unchanged compared to that from before stimulation ($\bar{A}_{total} = 0.9 \pm 0.4$). Yet, a more selective activation of the ankle flexor TA during dorsiflexion was observed in the post-tSCS assessment, as reflected by the group mean results of RMS_{TA}/RMS_{total} (0.6 ± 0.1 before stimulation and 0.8 ± 0.2 afterwards; Fig. 4D, left). During plantar flexion, the magnitude of ipsilateral EMG activity was reduced after stimulation ($\bar{A}_{total} = 0.5 \pm 0.2$). There was no increase in selectivity in agonist muscle activation, with the relative TS activity RMS_{TS}/RMS_{total} remaining at a low level (0.3 ± 0.2 before and 0.2 ± 0.2 after stimulation; Fig. 4D, right).

Effects of tSCS on the walking ability

Two of the three participants increased their over-ground walking speed after stimulation. Times required to walk 10 m at a self-selected fast, yet secure speed were 54 before and 34 seconds after stimulation for subject 1,

and 54 vs. 32 seconds for subject 2. Clonus-like activity interfering with the subjects' ability to walk before stimulation was visibly reduced after stimulation in these two subjects. Subject 3 had the fastest 10-m time prior to stimulation and did not show any increase after stimulation (23 vs. 23.5 seconds).

Subjects' reports

Apart from the anti-spastic effect, the participants described a feeling of lightness of the lower limbs, and increased sensation, especially of the foot sole during ground contact. The anti-spastic effects were subjectively reported to have persisted for 2–6 hours after tSCS. No negative effects were reported.

Discussion

We described the effects of a 30-minute session of 50-Hz transcutaneous lumbar SCS in three motor-incomplete SCI subjects with clinically evident spasticity. We observed that the responsiveness to muscle stretch, both tonic and phasic was decreased, as was the

Figure 4 Volitional unilateral dorsiflexion and plantar flexion in sitting position. **(A)** EMG patterns showing co-activation and ankle clonus (left) are reduced post-stimulation (right). Dashed lines superimposed on the ankle goniograms show the range between the neutral extended position of the ankle at rest and maximum dorsiflexion before stimulation, and a flexion bias of the rhythmic movement after stimulation is documented by the shift of the goniometric trace. Boxes are displayed with enlarged scales in **(B)** depicting the interference of clonus-like activity before stimulation that is eliminated after stimulation. **(C)** Group mean magnitudes of ipsilateral EMG activity during dorsiflexion and plantar flexion after stimulation normalized to the corresponding values before stimulation. **(D)** The contribution of TA to the total magnitude of ipsilateral muscle activation during dorsiflexion was increased after stimulation. TS activity during plantar flexion was at a low level before and after stimulation.

response to non-noxious, cutaneous stimulation of the plantar surface of the foot. Also, TA was more selectively activated during volitional dorsiflexion movements after tSCS. Finally, two of the three subjects increased their walking speeds considerably after tSCS.

To understand these observations, we must first look to the neuroanatomy stimulated by tSCS. The electrode set-up and stimulation frequency of tSCS resemble that of transcutaneous electrical nerve stimulation (TENS). However, clinical TENS therapies normally excite more superficially located neural structures, and the major application is in the treatment of localized pain conditions. On the other hand, tSCS targets deeply located neural structures within the vertebral canal with the aim to provide bilateral multi-segmental input to the spinal cord and subsequently modify the excitability of its circuits.²² Within the vertebral canal, large diameter afferent fibers of the L2–S2 posterior rootlets

to spinal cord junctions have the lowest thresholds for direct, electrical depolarization.²⁹ The ligaments and discs between the vertebral bones reduce the transverse electrical resistance of the thoraco-lumbar spine and allow some current to flow through the vertebral canal.^{30,39,40} The excitation thresholds for the posterior root fibers are considerably reduced at their point of entry into the spinal cord due to non-uniformities of the anatomy and its electrical conductivities along the fiber paths and changes of the fiber path direction with respect to the generated field.^{30,39,40} Among the fibers within the posterior roots, the Ia afferents have the largest fiber diameters. It is known that the Ia afferents, presumably among the fibers depolarized during tSCS, make strong synaptic connections to Ia inhibitory interneurons.⁴¹ Independently, there is indirect evidence in humans that epidural SCS can activate spinal circuitries mediating reciprocal inhibition.^{32,42} Reciprocal

inhibition might be one of the mechanisms sub-serving an inhibitory plurisegmental network situated within the lumbar spinal cord that was thought to be involved in the amelioration of severe lower-limb spasticity in a group of clinically complete mid-thoracic SCI people when epidural electrodes were placed over the lumbar posterior roots.²⁵

Another potential source of the effects observed here can be drawn from the reports that stimulation of the long tracts of the spinal cord reduces responsiveness to muscle spindle afferent input after spinal cord lesioning in cat models and SCI in humans.^{23,24,43,44} Here, we applied tSCS with adequate intensity to produce paresthesias that are generally associated with posterior-column depolarization during epidural SCS.⁴⁵⁻⁴⁷ Modeling of tSCS-induced currents, however, has shown that posterior column fibers have considerably higher thresholds than the posterior root fibers.⁴⁰ Thus, at the intensities applied here, tSCS was probably not depolarizing long-tract fibers directly. Posterior roots contain fibers from cutaneous mechanoreceptors and their depolarization can also produce such sensations.⁴⁸ Therefore, the report of paresthesias experienced here and the incomplete nature of the lesions in this study support the possibility that long loop, spinal-brainstem-spinal mechanisms might be involved in the suppression of segmental excitability.²³ Such combined posterior roots activation and increased descending inhibition via brainstem loops in the control of spinal spasticity was suggested by other human studies.^{17,23,25,49} Therefore, the most likely explanation for the effects seen in the three subjects reported here is a combination of both direct activation of local inhibitory circuits by posterior root fibers and the potentially more complex involvement of descending activation of presynaptic inhibition brought by long-loop mechanisms. This would also explain the observed reduction in the excessive withdrawal-like response elicited by non-noxious stimulation of the plantar surface of the foot, interpreted as a sign of decreased descending inhibitory control and altered intraspinal processing due to SCI.⁵⁰ Finally, a generalized reduction of motor outputs by post-synaptic inhibition only is less probable since there was no deterioration of the voluntary activation of motor neuron pools during single ankle joint movements and walking. The improvements in the 10-m walk test in two of the three incomplete SCI participants after tSCS were partially attributable to the reduced clonus-like activity interfering with gait. The influence of a practice effect was less probable since the increase in gait speed of the two subjects was definitely beyond the normal variability of the subjects' condition.

To what extent the above-discussed mechanisms contribute to the temporarily persisting modifications of the spinal cord neural circuits' excitability as assessed after tSCS was ended remains to be elucidated. Yet, this carry-over effect and its subjectively reported persistence for 2 to several hours after cessation of stimulation were not unexpected. In epidural SCS in motor disorders, it was repetitively observed that degradation of the improved function after withdrawal of stimulation could take hours or days.^{13,19,23,24} Temporary carry-over effects were also described in functional electrical stimulation of peripheral nerves.⁵¹⁻⁵³ However, proposed physiological mechanisms underlying these carry-over effects are speculative and include activity-dependent, temporarily persisting changes of synaptic transmission such as long-term potentiation and long-term depression behavior of spinal motor neurons.⁵⁴

The results were derived from few individuals and thus, generalization to a population with a variety of SCI profiles should be cautioned.⁵⁵ Further, the obvious sensory effects of the active stimulation limits the applicability of more conclusive study designs such as crossover studies with sham stimulation. Yet, we conducted thorough and quantitative assessment of many parameters of a highly integrative sensory-motor system.⁵⁵ The antispastic effect of tSCS was apparent from complementary biomechanical, neurophysiological, and clinical evaluations and there were no discrepancies in the modifications across the parameters after stimulation. The promising results justify a clinical study with a statistically appropriate subject population with different severities of SCI. The close investigation of the carry-over effects as well as methods for prolonging this beneficial period, e.g. by several sessions of tSCS over a longer period of time, should be also addressed.

Conclusion

Independent from the specific mechanisms, the present results suggest that 50-Hz tSCS for 30 minutes leads to the diminution of pathophysiology (spasticity) and the improvement of normal physiology (voluntary motor control). If shown to be of wide prevalence, it is possible that this therapeutic approach could be used instead of systemically applied medications or ablative treatments with their unfortunate and long-lasting side effects. The temporary relief of spasticity with tSCS could be a non-invasive treatment of use for even short periods, perhaps prior to a physical therapy session where decreased co-contractions and dyssynergias and improved voluntary control could be better trained.

Thus, this report serves as evidence that more study designed to characterize the extent and potential

mechanisms of tSCS effect on motor neuron excitability is needed. Taken together, these results suggest that tSCS may be able to modify spinal motor responsiveness to external inputs from muscle, tendon, joint and cutaneous receptors while maintaining its responsiveness to input from spinal motor processing generated from within the CNS.

Acknowledgments

Special thanks to Milan R. Dimitrijevic, Department of Physical Medicine and Rehabilitation, Baylor College of Medicine, Houston, TX, USA, and Foundation for Movement Recovery, Oslo, Norway. We wish to acknowledge the support of the Austrian Science Fund (FWF), Proj.Nr. L512-N13, the Vienna Science and Technology Fund (WWTF), Proj.Nr. LS11-057, and the Wings for Life Spinal Cord Research Foundation (WfL), Proj.Nr. WFL-AT-007/11.

References

- Sheean G. The pathophysiology of spasticity. *Eur J Neurol* 2002; 9(S1):3–9.
- Biering-Sørensen F, Nielsen JB, Klinge K. Spasticity-assessment: a review. *Spinal Cord* 2006;44(12):708–22.
- Krawetz P, Nance P. Gait analysis of spinal cord injured subjects: effects of injury level and spasticity. *Arch Phys Med Rehabil* 1996; 77(7):635–8.
- Adams MM, Hicks AL. Spasticity after spinal cord injury. *Spinal Cord* 2005;43(10):577–86.
- Tansey K, Dimitrijevic MM, Mayr W, Bijak M, Dimitrijevic MR. Restorative neurology of motor control after spinal cord injury. In: Dimitrijevic MR, Byron A, Vrbova G, McKay WB, (eds.) *Restorative neurology of spinal cord injury*. New York: Oxford University Press; 2011. p. 43–64.
- Elbasiouny SM, Moroz D, Bakr MM, Mushahwar VK. Management of spasticity after spinal cord injury: current techniques and future directions. *Neurorehabil Neural Repair* 2010; 24(1):23–33.
- Smyth MD, Peacock WJ. The surgical treatment of spasticity. *Muscle Nerve* 2000;23(2):153–63.
- Foerster O. Über eine neue operative Methode der Behandlung spastischer Lähmung mittels Resektion hinterer Rückenmarkswurzeln. *Z Orthop Chir* 1909;22:203–22.
- Nielsen JB, Crone C, Hultborn H. The spinal pathophysiology of spasticity – from a basic science point of view. *Acta Physiol (Oxf)* 2007;189(2):171–80.
- Sindou M, Pregel J, Boisson D, Eyssette M, Goutelle A. Surgical selective lesions of nerve fibers and myelotomies for modifying muscle hypertonia. In: Eccles J, Dimitrijevic MR, (eds.) *Upper motor neuron functions and dysfunctions. Recent achievements in restorative neurology*, Vol. 1. Basel: S Karger AG; 1985. p. 10–26.
- Shealy CN, Mortimer JT, Reswick JB. Electrical inhibition of pain by stimulation of the dorsal columns: preliminary clinical report. *Anesth Analg* 1967;46(4):489–91.
- Cook AW, Weinstein SP. Chronic dorsal column stimulation in multiple sclerosis. Preliminary report. *N Y State J Med* 1973; 73(24):2868–72.
- Cook AW. Electrical stimulation in multiple sclerosis. *Hosp Pract* 1976;11(4):51–8.
- Dooley DM. Percutaneous electrical stimulation of the spinal cord. 43rd Annual Meeting of the American Association of Neurological Surgeons. Miami Beach, Florida, April 1975.
- Illis LS, Oygur AE, Sedgwick EM, Awadalla MA. Dorsal-column stimulation in the rehabilitation of patients with multiple sclerosis. *Lancet* 1976;1(7974):1383–6.
- Richardson RR, McLone DG. Percutaneous epidural neurostimulation for paraplegic spasticity. *Surg Neurol* 1978;9(3):153–5.
- Richardson RR, Cerullo LJ, McLone DG, Gutierrez FA, Lewis V. Percutaneous epidural neurostimulation in modulation of paraplegic spasticity. Six case reports. *Acta Neurochir* 1979;49(3–4): 235–43.
- Siegfried J, Lazorthes Y, Broggi G. Electrical spinal cord stimulation for spastic movement disorders. *Appl Neurophysiol* 1981; 44(1–3):77–92.
- Barolat G, Myklebust JB, Wenninger W. Effects of spinal cord stimulation on spasticity and spasms secondary to myelopathy. *Appl Neurophysiol* 1988;51(1):29–44.
- Barolat G, Singh-Sahni K, Staas WE, Shatin D, Ketcik B, Allen K. Epidural spinal cord stimulation in the management of spasms in spinal cord injury: a prospective study. *Stereotact Funct Neurosurg* 1995;64(3):153–64.
- Waltz JM. Chronic stimulation for motor disorders. In: Gindenberg PL, Tasker RR, (eds.) *Textbook for stereotactic and functional neurosurgery*. New York: McGraw-Hill; 1998. p. 1087–99.
- Minassian K, Hofstoetter U, Tansey K, Mayr W. Neuromodulation of lower limb motor control in restorative neurology. *Clin Neurol Neurosurg* 2012;114(5):489–97.
- Dimitrijevic MM, Dimitrijevic MR, Illis LS, Nakajima K, Sharkey PC, Sherwood AM. Spinal cord stimulation for the control of spasticity in patients with chronic spinal cord injury: I. Clinical observations. *Cent Nerv Syst Trauma* 1986;3(2):129–44.
- Dimitrijevic MR, Illis LS, Nakajima K, Sharkey PC, Sherwood AM. Spinal cord stimulation for the control of spasticity in patients with chronic spinal cord injury: II. Neurophysiologic observations. *Cent Nerv Syst Trauma* 1986;3(2):145–52.
- Pinter MM, Gerstenbrand F, Dimitrijevic MR. Epidural electrical stimulation of posterior structures of the human lumbosacral cord: 3. Control of spasticity. *Spinal Cord* 2000;38(9):524–31.
- Murg M, Binder H, Dimitrijevic MR. Epidural electric stimulation of posterior structures of the human lumbar spinal cord: 1. muscle twitches – a functional method to define the site of stimulation. *Spinal Cord* 2000;38(7):394–402.
- Rattay F, Minassian K, Dimitrijevic MR. Epidural electrical stimulation of posterior structures of the human lumbosacral cord: 2. quantitative analysis by computer modeling. *Spinal Cord* 2000;38(8):473–89.
- Dimitrijevic MR, Dimitrijevic MM, Kern H, Minassian K, Rattay F. Electrophysiological characteristics of H-reflexes elicited by percutaneous stimulation of the cauda equina. Program No. 417.11. Abstract viewer/itinerary planner. Washington, DC: Society for Neuroscience; 2004.
- Minassian K, Persy I, Rattay F, Dimitrijevic MR, Hofer C, Kern H. Posterior root-muscle reflexes elicited by transcutaneous stimulation of the human lumbosacral cord. *Muscle Nerve* 2007;35(3): 327–36.
- Minassian K, Hofstoetter U, Rattay F. Transcutaneous lumbar posterior root stimulation for motor control studies and modification of motor activity after spinal cord injury. In: Dimitrijevic MR, Byron A, Vrbova G, McKay WB, (eds.) *Restorative neurology of spinal cord injury*. New York: Oxford University Press; 2011. p. 226–55.
- Hofstoetter US, Minassian K, Hofer C, Mayr W, Rattay F, Dimitrijevic MR. Modification of reflex responses to lumbar posterior root stimulation by motor tasks in healthy subjects. *Artif Organs* 2008;32(8):644–8.
- Minassian K, Persy I, Rattay F, Pinter MM, Kern H, Dimitrijevic MR. Human lumbar cord circuitries can be activated by extrinsic tonic input to generate locomotor-like activity. *Hum Mov Sci* 2007; 26(2):275–95.
- Roy FD, Gibson G, Stein RB. Effect of percutaneous stimulation at different spinal levels on the activation of sensory and motor roots. *Exp Brain Res* 2012;223(2):281–9.
- Wartenberg R. Pendulousness of the legs as a diagnostic test. *Neurology* 1951;1(1):18–24.
- Sherwood AM, McKay WB, Dimitrijevic MR. Motor control after spinal cord injury: assessment using surface EMG. *Muscle Nerve* 1996;19(8):966–79.
- Scivoletto G, Tamburella F, Laurenza L, Foti C, Ditunno JF, Molinari M. Validity and reliability of the 10-m walk test and

- the 6-min walk test in spinal cord injury patients. *Spinal Cord* 2011;49(6):736–40.
- 37 Bajd T, Vodovnik L. Pendulum testing of spasticity. *J Biomed Eng* 1984;6(1):9–16.
 - 38 Leslie GC, Muir C, Part NJ, Roberts RC. A comparison of the assessment of spasticity by the Wartenberg pendulum test and the Ashworth grading scale in patients with multiple sclerosis. *Clin Rehabil* 1992;6(1):41–8.
 - 39 Ladenbauer J, Minassian K, Hofstoetter US, Dimitrijevic MR, Rattay F. Stimulation of the human lumbar spinal cord with implanted and surface electrodes: a computer simulation study. *IEEE Trans Neural Syst Rehabil Eng* 2010;18(6):637–45.
 - 40 Danner SM, Hofstoetter US, Ladenbauer J, Rattay F, Minassian M. Can the human lumbar posterior columns be stimulated by transcutaneous spinal cord stimulation? A modeling study. *Artif Organs* 2011;35(3):257–62.
 - 41 Jankowska E. Interneuronal relay in spinal pathways from proprioceptors. *Prog Neurobiol* 1992;38(4):335–78.
 - 42 Hofstoetter US. Model of spinal cord reflex circuits in humans: stimulation frequency-dependence of segmental activities and their interactions [PhD dissertation]. Vienna University of Technology, 2009.
 - 43 Chapman CE, Ruegg DG, Wiesendanger M. Effects of dorsal cord stimulation on stretch reflexes. *Brain Res* 1983;258(2):211–5.
 - 44 Siegfried J, Krainick JU, Haas H, Adorjani C, Meyer M, Thoden U. Electrical spinal cord stimulation for spastic movement disorders. *Appl Neurophysiol* 1978;41(1-4):134–41.
 - 45 Barolat G, Massaro F, He J, Zeme S, Ketcik B. Mapping of sensory responses to epidural stimulation of the intraspinal neural structures in man. *J Neurosurg* 1993;78(2):233–9.
 - 46 Holsheimer J. Computer modelling of spinal cord stimulation and its contribution to therapeutic efficacy. *Spinal Cord* 1998;36(8):531–40.
 - 47 Holsheimer J. Which neuronal elements are activated directly by spinal cord stimulation. *Neuromodulation* 2002;5(1):25–31.
 - 48 Hunter JP, Ashby P. Segmental effects of epidural spinal cord stimulation in humans. *J Physiol* 1994;474(3):407–19.
 - 49 Gybels J, Van Roost D. Spinal cord stimulation for the modification of dystonic and hyperkinetic conditions: a critical review. In: Eccles J, Dimitrijevic MR, (eds.) *Recent achievements in restorative neurology*. Basel: Karger; 1985. p. 56–70.
 - 50 Sandrini G, Serrao M, Rossi P, Romaniello A, Cruccu G, Willer JC. The lower limb flexion reflex in humans. *Prog Neurobiol* 2005;77(6):353–95.
 - 51 Vodovnik L, Bowman BR, Hufford P. Effects of electrical stimulation on spinal spasticity. *Scand J Rehabil Med* 1984;16(1):29–34.
 - 52 Bajd T, Gregoric M, Vodovnik L, Benko H. Electrical stimulation in treating spasticity resulting from spinal cord injury. *Arch Phys Med Rehabil* 1985;66(8):515–7.
 - 53 Stein RB. Functional electrical stimulation after spinal cord injury. *J Neurotrauma* 1999;16(8):713–7.
 - 54 Nielsen JF, Sinkjaer T. Long-lasting depression of soleus motoneurons excitability following repetitive magnetic stimuli of the spinal cord in multiple sclerosis patients. *Mult Scler* 1997;3(1):18–30.
 - 55 Edgerton VR, Harkema S. Epidural stimulation of the spinal cord in spinal cord injury: current status and future challenges. *Expert Rev Neurother* 2011;11(10):1351–3.