Appropriate Head Position for Nasotracheal Intubation by Using Lightwand Device (Trachlight)

Yozo Manabe, DDS, PhD,* Shigeru Iwamoto, DDS,[†] Mika Seto, DDS, PhD,[‡] and Kazuna Sugiyama, DDS, PhD[§]

*Department of Systemic Management for Dentistry, Kagoshima University Medical and Dental Hospital, Kagoshima, Japan, [†]Section of Anesthesiology, Department of Diagnostics and General Care, Fukuoka Dental College, Fukuoka, Japan, [‡]Department of Dentistry and Oral Surgery, Fukuoka University, Fukuoka, Japan, and [§]Department of Dental Anesthesiology, Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima, Japan

The purpose of this study was to determine the relationship between the head position and the subsequent ease of nasotracheal intubation by using the lightwand device Trachlight (TL). Patients requiring nasotracheal intubation were subdivided into 3 groups according to the intubated head position (group S: sniffing position; group E: extension position; and group N: neutral position). The number of attempts, the total intubation time, and the failures of the TL intubation were recorded. Intubation difficulty by means of TL was assessed by the ordinal 6-point scale. Of the 300 patients enrolled in the study, TL intubation was successful in 91.3% of them. There was no significant difference in the success rate of the first attempt between the groups. No correlation between the ordinal scale and the head position was observed. The total intubation time and the ratio of “unsuccessful” cases were not significantly different among the 3 groups. TL is an effective alternative for patients who require nasotracheal intubation. Our study did not determine the most favorable head position for nasotracheal intubation with the TL, so we recommend that nasotracheal intubation with TL be started with the head in the neutral position and then changed to a more appropriate position, if necessary, on an individual basis.

Key Words: Nasotracheal intubation; Lightwand device (Trachlight); Appropriate head position.

Nasotracheal intubation is often indicated in oral and maxillofacial surgery. An alternative orotracheal intubation technique using the lightwand device Trachlight (TL) (Laerdal Medical, Armonk, NY) has been reported to facilitate tracheal intubation in patients with difficult airways.¹ Some reports discussing the validity of using TL for nasotracheal intubation are available.^{2,3} In addition, the incidence of complications with light-guided intubation is reported to be low because elevation of the epiglottis by the laryngoscope blade is not required.^{4,5}

The sniffing position is widely considered essential to the performance of orotracheal intubation for direct laryngoscopy. In most cases, fiberoptic nasotracheal intubation would be started with neutral head position. But, there is no report about the difficulty of nasotracheal intubation using TL in relation to the head position.

This study was designed to determine the relationship between the head position and the subsequent ease of nasotracheal intubation by TL.

METHODS

This study was approved by the Institutional Review Board at the Fukuoka Dental College (Fukuoka Dental College, Fukuoka, Japan; approved April 4, 2005;

Received July 17, 2012; accepted for publication March 3, 2014.
Address correspondence to Dr Yozo Manabe, 8-35-1, Sakuragaoka, Kagoshima 890-8544, Japan; manabey@dent.kagoshima-u.ac.jp.

Anesth Prog 61:47–52 2014

© 2014 by the American Dental Society of Anesthesiology

ISSN 0003-3006/14
SSDI 0003-3006(14)

Figure. Head position in intubation. The patients were randomly subdivided into 3 groups according to their head position in intubation. Group S: intubated in the “sniffing” position. Group E: intubated in the “extension” position. Group N: intubated in the “neutral” position.

Approval 66), and written informed consent was obtained from each patient.

Patients (American Society of Anesthesiologists [ASA] 1 or 2) scheduled for elective oral and maxillofacial surgery under general anesthesia that required nasotracheal intubation were included in this study. Patients who had a past history of upper airway surgery or a body mass index (BMI) $> 30 \text{ kg/m}^2$ were excluded from the study. Approved patients were randomly assigned to 3 equal-sized groups ($n = 100$) according to their head position for lightwand nasotracheal intubation: group S were intubated in the sniffing position; group E were intubated in the extension position; and group N were intubated in the neutral position. The sniffing position was obtained by placement of a 9-cm cushion under the head of the patient.⁶ The extension position was obtained by bending the head part of the bed at an angle of 30° posteriorly, and the neutral position was obtained by just lying on the flat bed without a cushion (Figure). They underwent a preoperative airway assessment that included a Mallampati score^{7,8} and mouth opening with the mouth fully opened (the interincisor gap was measured in millimeters) by an attending anesthesiologist. Demographics such as age, sex, weight, and height were also recorded.

Each patient was routinely monitored during the entire procedure. A rapid-acting anesthetic (propofol, 1–1.5 mg/kg, or thiamylal, 3–5 mg/kg) was administered intravenously, mask ventilation was ensured, and vecuronium bromide (0.1 mg/kg) was administered. The nasal mucosa of both nostrils was disinfected and probed with absorbent cotton immersed in 10% povidone-iodine with 1 : 100,000 epinephrine, and the wider, patent nostril was chosen for intubation. Then, lidocaine jelly was used as a lubricant for the nasopharyngeal airway tubes, which had increasing caliber sizes to dilate the meatus of the selected nasal cavity. Direct laryngoscopy

was performed in all cases by using a size 3 Macintosh laryngoscope blade for topical anesthesia on the supraglottic region and the vocal cords with 8% lidocaine spray. This process is standardized in our department.⁹ Visualization of the glottis during laryngoscopy was assessed by using the Cormack and Lehane classification¹⁰ by an attending anesthesiologist with at least 3 years of anesthesia experience. All direct laryngoscopy procedures were performed in the sniffing position.

Nasal RAE endotracheal tubes (NTT) (Mallinckrodt, Glens Falls, NY) were used in this study. Each NTT was cut to the appropriate length so that the tube tip could be adjusted to the light source. The NTT was inserted with the TL after removal of the stylet (TL-NTT) and soaking it in warm sterilized normal saline to reduce epistaxis and nasal damage.¹¹ The TL-NTT was inserted into the nostril and advanced until light was seen in the oropharynx. After further advancing, a blurred light was seen in the submandibular area. The TL-NTT tip was maneuvered anteromedially until a bright spot of light was seen at the cricothyroid membrane; the TL was then withdrawn and the NTT advanced into the trachea. The cuff of the tracheal tube was inflated with air at the peak airway pressure of 25 cm H_2O immediately after intubation to prevent leaks. The location of the NTT tip was confirmed by palpation of the inflated balloon cuff at the level of the suprasternal notch and by capnography. If the intubation was not performed within 60 seconds or a decrease in arterial oxygen saturation by pulse oximetry below 98% was observed, the TL was withdrawn, the opposite nostril and mouth were covered by hand, and ventilation through the NTT was resumed. Three attempts were permitted with ventilation interposed. If the TL intubation was unsuccessful after 3 consecutive attempts, intubation was performed using direct laryngoscopy. Only 1 anesthesiologist with 20 years of experience, who was blinded to the preopera-

Table 1. Ordinal 6-Point Scale of Nasotracheal Intubation Difficulty

Score	Description	Classification Criterion
1	Very easy	Successfully intubated within 15 seconds at the first attempt
2	Easy	Successfully intubated between 15 and 30 seconds at the first attempt
3	Moderate	Successfully intubated more than 30 seconds at the first attempt
4	Difficult	Successfully intubated at the second attempt
5	Very difficult	Successfully intubated at the third attempt
6	Unsuccessful	Intubation could not be achieved with 3 attempts of using the lightwand device

tive assessment and Cormack and Lehane classification results, performed all of the intubations using TL. Intubation difficulty was assessed by the ordinal 6-point scale⁹ (Table 1). The number of attempts, the total intubation time, and the failures of the TL intubation were recorded by another anesthesiologist. The total intubation time was defined as the sum of all intubation attempt durations, but the total time of interposed ventilation was excluded. The duration of each attempt was defined as the time from inserting the TL-NTT into the nostril to the time when the TL was removed from the NTT.

Statistics

Ratio of sex, nostril side, distribution of Mallampati class, and Cormack and Lehane classification between groups were analyzed using χ^2 test. Intubation difficulty between groups was analyzed using Kruskal-Wallis test. For the comparison of other data, analysis of variance with Sheffe post-hoc test was performed. All statistical analyses were performed using Stat View version 5.0 (SAS Institute Inc, Cary, NC). Data are presented as number of patients and mean (SD). $P < .05$ was considered statistically significant.

RESULTS

A total of 300 patients were enrolled into the study—100 patients in each group (S, E, and N). There were no

significant differences among the 3 groups in terms of height, weight, or the ratio by sex (Table 2). There were no significant differences in the ratio of the nostril side, cuff volume, interincisor gap, and the tube size (Table 3). TL intubation was successful in 274 of the 300 patients (91.3%), and it failed in 26 (8.7%). Two hundred forty patients (80.0%) were intubated at the first attempt, 19 (6.3%) required a second attempt, and 15 (5.0%) required a third attempt. There was no significant difference in the success rate of the first attempt among the groups. The total intubation time was not significantly different among the 3 groups (Table 3). Distributions of Mallampati score and Cormack and Lehane grade in each group were not significantly different (Tables 4 and 5). There were 9 unsuccessful cases in group S, 7 unsuccessful cases in group E, and 10 unsuccessful cases in group N. No significant differences in the ratio of unsuccessful cases were found among the groups (Table 6). In all cases, tracheal intubation was performed without severe hemodynamic instability. There were no adverse outcomes.

DISCUSSION

Nasotracheal intubation using a light guide has been reported since 1959¹²; however, no report has focused on nasotracheal intubation difficulty using TL in relation to the head position.

Nasotracheal intubation using the TL has been performed in several cases, and a number of institutions use the TL as the first choice of orotracheal intubation

Table 2. Patient Demographics*†

	Group S	Group E	Group N
No. of Patients	100	100	100
Sex, M/F	44/56	38/62	39/61
Age, y	35.5 ± 15.3	36.0 ± 16.1	33.0 ± 17.4
Height, cm	163.1 ± 7.8	162.5 ± 8.5	162.6 ± 8.0
Weight, kg	56.4 ± 9.5	57.2 ± 10.7	56.4 ± 9.3

* Group S, intubated in the “sniffing” position; group E, intubated in the “extension” position; and group N, intubated in the “neutral” position.

† Age, height, and weight are presented as mean ± SD.

Table 3. Result of Measurement*

	Group S	Group E	Group N
No. of patients	100	100	100
Interincisor gap, cm†	44.6 ± 7.1	43.9 ± 7.7	45.2 ± 5.8
Tube size, cm†	7.6 ± 0.4	7.5 ± 0.4	7.6 ± 0.4
Nostril side, right/left	62/38	66/34	75/25
Cuff volume, mL†	3.6 ± 1.2	3.9 ± 1.6	4.0 ± 1.5
Total time to intubate, s†	17.4 ± 29.0	16.9 ± 25.6	20.7 ± 35.7
Patients			
intubated at the first attempt, %	77.0	83.0	80.0
Patients			
intubated at the second attempt, %	4.0	8.0	7.0
Patients			
intubated at the third attempt, %	10.0	2.0	3.0
Success rate, %	91.0	93.0	90.0

* Group S, intubated in the “sniffing” position; group E, intubated in the “extension” position; and group N, intubated in the “neutral” position.

† Data are presented as mean ± SD.

Table 4. Distribution of Mallampati Score in Each Head position*

	Group S	Group E	Group N
No. of patients	100	100	100
Mallampati score class 1 (n = 228)	79	77	72
Mallampati score class 2 (n = 54)	15	16	23
Mallampati score class 3 (n = 9)	3	2	4
Mallampati score class 4 (n = 9)	3	5	1

* Group S, intubated in the “sniffing” position; group E, intubated in the “extension” position; and group N, intubated in the “neutral” position.

Table 5. Distribution of Cormack-Lehane Grade in Each Head Position*

	Group S	Group E	Group N
No. of patients	100	100	100
Cormack-Lehane score grade 1 (n = 247)	85	85	77
Cormack-Lehane score grade 2 (n = 32)	10	9	13
Cormack-Lehane score grade 3 (n = 21)	5	6	10
Cormack-Lehane score grade 4 (n = 0)	0	0	0

* Group S, intubated in the “sniffing” position; group E, intubated in the “extension” position; and group N, intubated in the “neutral” position.

Table 6. Relation Between Ordinal 6-Point Scale and Head Position

	Group S	Group E	Group N
No. of patients	100	100	100
Score 1 “Very easy” (n = 216)	71	75	70
Score 2 “Easy” (n = 11)	2	2	7
Score 3 “Moderate” (n = 13)	4	6	3
Score 4 “Difficult” (n = 19)	4	8	7
Score 5 “Very difficult” (n = 15)	10	2	3
Score 6 “Impossible” (n = 26)	9	7	10

* Group S, intubated in the “sniffing” position; group E, intubated in the “extension” position; and group N, intubated in the “neutral” position.

method.¹³ In the “Difficult Airway Algorithm” of the American Society of Anesthesiologists,¹⁴ the TL is also shown to be one of the instruments used as an intubation method for patients who are easy to ventilate manually but are found to have difficult laryngoscopy after induction of anesthesia. Park et al⁶ reported that the laryngoscopic view with the 9-cm pillow was significantly superior to that with other pillows and without pillow; therefore, we applied a 9-cm cushion to the sniffing position. Hung and Stewart¹⁵ reported that TL should be avoided in patients with known anatomic abnormalities of the upper airway. Therefore, postoperative patients who had anatomic abnormalities of the upper airway resulting from oral surgery were excluded from this study.

TL intubation was successful with an average time of 18.3 seconds, and it is very similar to the report (17.6 $\pm$ 6.6 seconds) of Cheng et al.¹⁶ The success rate of nasotracheal intubation by TL in this study was 91%. Cheng et al¹⁶ also reported that the success rate of nasotracheal intubation by TL was 96.7% at first attempt and 100% at second attempt, but they performed the tracheal tube cuff inflation technique¹⁷ on all patients.

One of the reasons for this low success rate is that cases in which patients could not be intubated in 3 attempts were classified as “unsuccessful” based on the report by Fox et al,¹⁸ which showed that the average number of attempts required for blind nasotracheal intubation by an experienced anesthesiologist is 3.1 times. However, we have had cases in which TL nasotracheal intubation was successful after 4 or more attempts in the clinical setting. An additional reason that could be considered is trying all 3 consecutive attempts in the same head position. Appropriate head position of nasotracheal intubation with TL may be individually different, as it is with blind nasotracheal intubation. We recommend that the nasotracheal intubation with TL should be started with the neutral position because this position is the intermediate position of the sniffing position and the extension position.

There are 2 causes of unsuccessful nasotracheal intubation by TL: when the tube is inserted into the esophagus and when the tube is lodged in the tissue surrounding the epiglottis and cannot be advanced into the trachea. In the former case, it is often possible to insert the tube by extending the neck to raise the tube tip. When it is difficult to perform the intubation in this manner, changing the tube to the Endotrol tube is contemplated.¹⁹ However, during oral and maxillofacial surgery, the use of the RAE tube with suitable flexion is encouraged because the connector of the anesthetic circuit and the tracheal tube disturb the sight of the surgeon when it is located above the patient’s nose. Iseki

et al²⁰ reported that the tip of the tracheal tube can be raised by the alteration of the TL; however, this method cannot be applied to the RAE tube. In clinical situations, we raised the tip of the tube by the tracheal tube cuff inflation technique¹⁷ and had good results. This technique is reported to increase the success rate of blind nasotracheal intubation.

Management of the case wherein the tube cannot advance into the trachea is more difficult. In such cases, we try various head positions; however, most of these cases came to be “unsuccessful” cases.

In case of intubation using a lightwand, the advantage is that the precise position of an endotracheal tube tip is known by checking the light at the sternal notch.²¹ Since the length up to a bend of the RAE tube is provided, it is necessary to set an appropriate fixed length.

It has been accepted that intubation with a direct laryngoscopy is difficult for grades 3 and 4 of the Cormack and Lehane classification. Ainsworth and Howells²² reported that the difficulty of oral intubation with lightwand has no relation to the grade of laryngoscopic view according to the Cormack and Lehane classification. Hung et al⁵ reported that the Mallampati classification correlated with the intubation time by using the laryngoscope but did not correlate with the time by using TL. Manabe et al⁹ reported that there is no correlation between the difficulty of nasotracheal intubation using TL and the Mallampati classification or the Cormack and Lehane classification.

There was no correlation between the airway parameters and the time required for intubation using the TL. In other words, nasotracheal intubation using TL was not influenced by anatomic variability in this population.

In addition, intubation using TL is a light-guided technique without visualization of the laryngeal structures; therefore, there is a potential risk of trauma to the upper airway associated with its use, but there were no adverse outcomes in this study.

Our study has several limitations. First, all intubations were performed by 1 operator; therefore, the results may be biased toward the expertise of that operator. However, patients were sampled randomly so that the head positions could be equally dispersed through the study period, and the operator had a great deal of experience in nasotracheal intubation using TL from the beginning of this study. Second, premedication or anesthesia methods were not unified. Nevertheless, all patients received 0.1 mg/kg of vecuronium bromide; therefore, we consider that the difference in anesthesia method could be ignored.

In conclusion, Trachlight is an effective alternative for patients who require nasotracheal intubation, and there was no relationship between the ease of nasotracheal intubation using the TL and the head position. We

recommend that nasotracheal intubation with TL be started in the neutral position and that the appropriate head position be determined on an individual basis.

REFERENCES

1. Hung OR, Pytko S, Morris I, Murphy M, Stewart RD. Lightwand intubation. Clinical trial of a new lightwand for tracheal intubation in patients with difficult airway. *Can J Anaesth*. 1995;42:826–830.
2. Weis FR, Hatton MN. Intubation by use of the light wand: experience in 253 patients. *J Oral Maxillofac Surg*. 1989;47:577–580.
3. Roberto F, Paolo T, Francesco DL, et al. Effective nasotracheal intubation using a modified transillumination technique. *Can J Anesth*. 2002;49:91–95.
4. Friedman PG, Rosenberg MK, Lebenbom-Mansour M. A comparison of lightwand and suspension laryngoscopic intubation techniques in outpatients. *Anesth Analg*. 1997;85:578–582.
5. Hung OR, Pytko S, Morris I. Clinical trial of a new lightwand device (Trachlight) to intubate the trachea. *Anesthesiology*. 1995;83:509–514.
6. Park SH, Park HP, Jeon YT. A comparison of direct laryngoscopic view depending pillow height. *J Anesth*. 2010;24:526–530.
7. Mallampati SR, Gatt SP, Gugino LD. A clinical sign to predict difficult tracheal intubation: a prospective study. *Can Anaesth Soc*. 1985;32:429–434.
8. Samsoon GLT, Young JRB. Difficult tracheal intubation: a retrospective study. *Anaesthesia*. 1987;42:487–490.
9. Manabe Y, Seto M, Iwamoto S. The success rate of nasotracheal intubation using lightwand does not depend on the laryngoscopic view. *J Anesth*. 2011;25:350–355.
10. Cormack RS, Lehane J. Difficult tracheal intubation in obstetrics. *Anaesthesia*. 1984;39:1105–1111.
11. Kim YC, Lee SH, Noh GJ. Thermosoftening treatment of the nasotracheal tube before intubation can reduce epistaxis and nasal damage. *Anesth Analg*. 2000;91:698–701.
12. Yamamura H, Yamamoto T, Kamiyama M. Device for blind nasal intubation. *Anesthesiology*. 1959;20:221.
13. Inoue Y, Koga K, Shigematsu A. A comparison of two tracheal intubation techniques with Trachlight in patients with cervical spine disorders. *Anesth Analg*. 2002;94:667–671.
14. American Society of Anesthesiologists Task Force on Management of the Difficult Airway. Practice guidelines for management of the difficult airway: an updated report by the American Society of Anesthesiologists Task Force on Management of the Difficult Airway. *Anesthesiology*. 2003;98:1269–1277.
15. Hung OR, Stewart RD. Lightwand intubation: I—a new lightwand device. *Can J Anaesth*. 1995;42:820–825.
16. Cheng KI, Chang MC, Lai TW, et al. Modified lightwand-guided nasotracheal intubation technique for oromaxillofacial surgical patients. *J Clin Anesth*. 2009;21:258–263.
17. Elstraete ACV, Pennant JH, Gajraj NM, Victory RA. Tracheal tube cuff inflation as an aid to blind nasotracheal intubation. *Br J Anaesth*. 1993;70:691–693.
18. Fox DJ, Castro T, Rastrelli AJ. Comparison of intubation techniques in the awake patient: the Flexi-lum surgical light (lightwand) versus blind nasal approach. *Anesthesiology*. 1987;66:69–71.
19. Asai T. Endotrol tube for blind nasotracheal intubation. *Anaesthesia*. 1996;51:507.
20. Iseki K, Murakawa M, Tase C, Otsuki M. Use of a modified lightwand for nasal intubation. *Anesthesiology*. 1999;90:635.
21. Locker GJ, Staudinger T, Knapp S, et al. Assessment of the proper depth of endotracheal tube placement with Trachlight. *J Clin Anesth*. 1998;10:389–393.
22. Ainsworth QP, Howells TH. Transilluminated tracheal intubation. *Br J Anaesth*. 1989;62:494–497.