Published in final edited form as:

Psychoneuroendocrinology. 2013 July ; 38(7): 1208–1212. doi:10.1016/j.psyneuen.2012.10.017.

Neonatal CSF Oxytocin Levels are Associated with Parent Report of Infant Soothability and Sociability

Catherine L. Clark, PhD¹, Nicholas St. John, PhD², Anca M. Pasca, MD¹, Shellie A. Hyde³, Kirsten Hornbeak³, Marina Abramova, MD¹, Heidi Feldman, MD¹, Karen J. Parker, PhD³, and Anna A. Penn, MD, PhD¹

¹Dept of Pediatrics, Div of Neonatal and Developmental Medicine, Stanford University School of Medicine, Stanford, CA 94305

²Developmental-Behavioral Pediatrics, Lucile Packard Children's Hospital at Stanford, Palo Alto, CA 94304

³Dept of Psychiatry, Stanford University School of Medicine, Stanford, CA 94305

Abstract

Oxytocin (OT) has been linked to social behavior in rodents, non-human primates, and adult humans, but almost nothing is known about brain OT activity in human newborns or its impact on social development. To better understand the role of OT biology in human social functioning, a multi-disciplinary, longitudinal study was conducted. Cerebral spinal fluid (CSF) OT levels from 18 human neonates were evaluated and examined in relationship to social-seeking behavior at term, at 3 months, and at 6 months of age. Higher neonatal CSF OT levels were consistently associated with solicitation of parental soothing and interest in social engagement with others. This is the first study to link CSF OT levels to normative human social functioning. Research is now required to test whether early OT levels serve as a biomarker for subsequent social abnormalities.

Keywords

Oxytocin; CSF; newborn; social behavior; soothability; social engagement

© 2012 Elsevier Ltd. All rights reserved.

Co-corresponding authors: Anna A. Penn, MD, PhD, 300 Pasteur Dr., S228, Dept of Pediatrics, Stanford, CA 94305-5208, apenn@stanford.edu, Phone: 650-723-5711, Fax: 650-725-7724; Karen J. Parker, PhD, 1201 Welch Rd., MSLS P-104, Department of Psychiatry, Stanford, CA 94305-5485, kjparker@stanford.edu, Phone: 650-736-9863, Fax: 650-498-7761.

Conflict of interest statement: None declared for any authors

Contributors: Karen J. Parker and Anna A. Penn conceived of and directed the study. Catherine Clark was the primary study coordinator, performing all follow up evaluations and statistical analyses. Nicholas St. John provided expert guidance on behavioral evaluation of infants in the NICU. Anca Pasca, Kirsten Hornbeak and Marina Abramova obtained consent and processed samples prior to storage. Shelly Hyde performed all OT measurements in the laboratory of Dr. Parker. Catherine Clark, Heidi Feldman, Karen Parker and Anna Penn all contributed to the writing of the manuscript.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Impairments in social functioning—observed in autism or extreme anxiety—can lead to devastating, lifelong consequences for the individual. Understanding the biology of early aberrations in infant social behavior may offer insights into, and early markers for, these developmental disorders. Oxytocin (OT) has emerged as a strong candidate in the search for the biological underpinnings of human social functioning (Meyer-Lindenberg et al., 2011).

Central OT plays a critical role in a variety of mammalian social behaviors including mother-infant attachment, affiliative contact-seeking, and social recognition (Insel, 2010). Young peer-reared rhesus monkeys display aberrant social behaviors; as a group these monkeys had lower cerebrospinal fluid (CSF) OT levels over the course of development when compared to maternally-reared controls (Winslow et al., 2003). Strikingly, individual expressions of affiliative behaviors also correlated with CSF OT levels. While CSF OT levels are positively correlated with social functioning in this non-human primate model, this relationship is not evident in plasma OT assessments (Winslow et al., 2003), highlighting the complex regulation of central and peripheral OT systems (Amico et al., 1990; Veening and Barendregt, 2010). Finally, central OT blockade, achieved pharmacologically or genetically, results in a variety of social impairments (Carter, 2007; Insel, 2010; Meyer-Lindenberg et al., 2011; Modi and Young, 2012).

There is limited information on the role of central OT biology in adult human social functioning, primarily due to the need for invasive measurement techniques (i.e., lumbar puncture), and nothing is presently known about OT biology in human infants. We therefore capitalized on the unique opportunity to examine CSF from newborns undergoing clinical sepsis evaluation, including lumbar puncture. The vast majority (>95%) of such infants are later found not to be septic, thus allowing measurement of potential biomarkers in normal newborn CSF. Here we describe the results of a pilot study designed to measure newborn CSF OT levels in the immediate postnatal period, followed by periodic behavioral assessments during the first year of life to test for the first time the relationships between human newborn CSF OT levels and later social behavior.

Methods

Participants

The study was approved by the Stanford University Institutional Review Board, (Protocol 10663). Eighteen neonates (12 males, 6 females) undergoing clinically indicated sepsis evaluation for standard risk factors (maternal fever, prolonged rupture of membranes, respiratory distress) were included. Exclusion criteria included known chromosomal anomalies and major malformations. With parental consent, 0.1-0.5 ml additional CSF was obtained at the time of lumbar puncture using standard sterile procedures within 72 hours of birth. All subjects received 48 hours of antibiotics and were found to be sepsis-negative. Gestational age ranged from 27 to 40 weeks ($M = 36.14$, $SD = 3.80$). Ethnicity and other demographics are presented in Table 1.

Fluid storage and processing procedures

CSF was transferred on ice, aliquoted, snap frozen, and stored at -80°C. Unextracted CSF samples were assayed for OT (in duplicate, 100 ul per well) by enzyme immunoassay (Enzo Life Sciences, Farmingdale, NY) as previously described (Parker et al., 2010). Per Enzo Life Sciences literature, the cross-reactivity with vasopressin is 0.6% and the limit of assay sensitivity where the curve is no longer linear is ~10pg/ml. All samples were run on a single microplate. The intra-assay coefficient of variation was <10%.

Soothability at term or term equivalent

Prior to hospital discharge, parents were asked to review a list of soothing techniques and check which ones helped to calm their baby (e.g., talking, touching, holding, providing a pacifier, feeding, not needing to be soothed). Responses were scored categorically. Parents were blind to OT levels.

Sociability follow-up at 3 months of age

At three months term-adjusted age, parents were contacted by phone to complete the Infant Behavior Questionnaire (Rothbart, 1981). Interviews were conducted in English or Spanish, as appropriate. The revised very short form of the instrument consists of 37 items assessing extroversion and positive affect, negative affect, and effortful control.

Sociability follow-up at 6 months of age

At six months term-adjusted age, parents were asked to return for evaluation and to complete two sets of questionnaires: (1) the IBQ and (2) the Relating to Others subscale from the Socialization section of the Vineland Adaptive Behavior Scales, 2nd Ed (Sparrow et al., 2005).

Statistical analyses

Study data were managed using REDCap and analyzed in SPSS v.19. Univariate statistical assessments were used to test relationships between CSF OT levels, perinatal factors, demographic factors, and behaviors; t-tests were used when the independent variable was dichotomous. Pearson correlations were used when the independent variable was continuous.

Results

CSF OT concentrations and study covariate analyses

CSF OT concentrations in newborn samples ranged from 10.25 to 34.06 pg/ml ($M = 20.55$, $SD = 7.53$). OT concentrations were not related ($p > 0.10$) to: sex, ethnicity, insurance status, gestational age, birth weight, mode of delivery (vaginal vs. c-section), Apgar scores at 1 minute or 5 minutes, maternal gravida or para status. There was no difference in OT levels in babies whose mothers were exposed versus not exposed to Pitocin ($t(16) = .17, ns$). For those who were exposed, there was no significant relationship between OT and amount of Pitocin given ($r = -.20, ns$). Therefore, data was pooled across these variables. Neither

demographic factors (Table 1) nor parity were statistically associated with parent responses regarding need for soothing at term, 3 or 6 months.

Newborn CSF OT concentrations positively correlate with need for interaction to soothe at term

Parents were given a checklist of soothing methods (holding, pacifier, feeding, does not require soothing) just prior to hospital discharge when their infant reached term adjusted age. Their responses (needed help soothing using any method versus no need for soothing) suggested that higher OT levels were associated with increased need for soothing assistance (M for babies who needed assistance being soothed = 21.37, $SD = 8.32$ vs. M for babies who did not need assistance being soothed = 16.18, $SD = .58$, $t(13) = 2.18$, $p < .05$). In particular, higher OT levels were associated with needing feeding to be soothed (M for babies who needed to be fed = 22.69, $SD = 7.45$ vs. M for babies who did not need to be fed = 12.63, $SD = 3.45$, $t(13) = 2.32$, $p < .05$).

Newborn CSF OT concentrations positively correlate with crying for attention at 3 months of age

Eleven parent phone interviews using the IBQ were completed (3 female and 8 male infants). There was no significant difference in neonatal CSF OT levels in the retained subjects versus those lost to follow-up ($t(13) = .32$). We created a separate Crying for Contact Index consisting of the mean average of two items assessing infants' need for soothing: "After sleeping, how often did the baby cry if someone doesn't come within a few minutes?" and "When you were busy with another activity, and your baby was not able to get your attention, how often did s/he cry?" Higher neonatal OT levels were significantly associated with greater crying when parents were not immediately responsive (Fig. 1A; $r = .65$, $p < .05$). Notably, IBQ measures of negative affect were not associated with higher neonatal OT levels.

Newborn CSF OT concentrations positively correlate with crying for attention and social behaviors at 6 months of age

Seven families returned to the hospital clinic for 6 month follow-up (3 female and 4 male infants). Again, despite declining subject retention, there was no significant difference in neonatal CSF OT levels in the retained subjects at 6 months versus those lost to follow up ($t(13) = .88$). The consistency between the 3-month Crying for Contact Index scores and the 6-month scores was highly significant ($r = .97$, $p < .01$). As at 3 months, higher levels of OT were associated with a greater Contact for Crying at six months (Fig. 1A; $r = .72$, $p < .10$), while negative affect was not related. On the Vineland instrument, CSF OT levels were positively related to the Relating to Others subscale (Fig. 1B; $r = .84$, $p < .05$), which includes items about watching other people, smiling, showing affection, and taking interest in other children.

Discussion

This pilot study provides the first evidence that CSF OT levels are associated with prosocial behavior at term, 3 months, and 6 months of age. These data suggest that infants with higher

CSF OT levels appear to actively seek parental social interaction for soothing, and have a greater interest in social interaction as measured at 6 months of age, in agreement with prior studies indicating that OT biology drives affiliative and attachment behaviors (Insel, 2010; Meyer-Lindenberg et al., 2011).

This study is unique in several ways. The majority of studies examining OT concentrations have been conducted in older children and adults, primarily using plasma measurements, with a focus on a limited high-risk population (Heim et al., 2009; Modahl et al., 1998). Our neonatal study population has both biological and behavioral measurements across the normal range of early social function. Moreover, we know of no prior systematic attempt to test the relationship between specific CSF biomarkers and later human social functioning.

Although few studies have examined CSF OT levels, those that have are in agreement with our findings. Two monkey studies have shown that CSF OT levels are higher in more social bonnet macaques compared to pigtail macaques, and are positively associated with affiliation in rhesus macaques (Rosenblum et al., 2002; Winslow et al., 2003). Moreover, two studies of adult humans have shown that CSF OT levels are diminished in adult survivors of childhood abuse and are negatively correlated with suicide intentionality (Heim et al., 2009; Jokinen et al., 2012). These collective data suggest that CSF OT levels are a promising means by which to assess social functioning in human and non-human primates.

As with any study, there are potential limitations to this work. Given the pilot nature of this study, we were underpowered to detect small effects of gestational age, birth weight, diurnal variation in CSF OT levels, intrapartum pitocin, and gender. In adults, there is diurnal variation in CSF OT levels (Forsling, 2000), with OT levels increasing during sleep, but the cyclicity of these levels in human newborns has not been established; therefore, a far larger cohort would be needed to assess any association with time of delivery, time of sample collection or sleep/wake cycles of the newborn. Additionally, genetic polymorphisms in CD38 gene that have been associated with changes in OT release (Higashida et al., 2012) may be relevant and should be measured in larger studies.

Attrition of participants was significant in our study, likely due to the migrant nature of the Hispanic population in our patient catchment area. The lack of differences in OT measurements between those infants who remained in the study and those lost to follow up suggests a lack of sample bias. The positive association between neonatal CSF OT levels and social interactions measured in the full cohort at term is supported by the stability of these correlations at 3 and 6 months of age, despite attrition and use of different questionnaires. Second, we were unable to concomitantly assess plasma OT levels because the plasma volume required for the OT assay exceeded the amount of blood that can safely be obtained from most newborns for research purposes. Third, parent measures were used as a proxy of infant social functioning. No well-validated tests of newborn sociability are available, making our in-hospital questionnaire results the least reliable despite their proximity to the time of OT measurement. Future studies should include more sophisticated laboratory assessments of social functioning (e.g., social initiation; eye gaze to social cues), and more thoroughly assess the roles of the variables outlined above.

In conclusion, these findings support the hypothesis that CSF OT levels may be a biomarker for later social functioning. At present, the relationships between central and peripheral OT levels are not well understood (Veening and Barendregt, 2010), but further investigations and refinement of techniques to measure OT levels should allow us to test whether peripheral measurements can be used in lieu of CSF measurements in future studies. With continued investigation this line of research could provide a critically needed early screening tool, identifying infants at high risk for developing social impairments on the basis of low early OT levels.

Acknowledgments

We are grateful to the members of the Penn, Parker, and Feldman laboratories, the nurses, doctors, and interpreters at Lucile Packard Children's Hospital, and, of course, the babies and their families. We thank Dr. Carl Feinstein, Director of the Stanford Autism Center, for his support of this line of research.

Funding sources: This work was supported by the Mosbacher Family Fund for Autism Research (to KJP); NIH Director's New Innovator Award (1DP2OD006457 to AAP); Center for Brain and Behavior at LPCH, the LPCH Autism Center and by an NIH/NCRR grant to the Stanford Center for Clinical Informatics (UL1 RR025744).

References

- Amico JA, Challinor SM, Cameron JL. Pattern of oxytocin concentrations in the plasma and cerebrospinal fluid of lactating rhesus monkeys (*Macaca mulatta*): evidence for functionally independent oxytocinergic pathways in primates. *J Clin Endocrinol Metab.* 1990; 71:1531–1535. [PubMed: 2229310]
- Carter CS. Sex differences in oxytocin and vasopressin: implications for autism spectrum disorders? *Behav Brain Res.* 2007; 176:170–186. [PubMed: 17000015]
- Forsling ML. Diurnal rhythms in neurohypophysial function. *Exp Physiol.* 2000; 85 Spec No:179S–186S. [PubMed: 10795921]
- Heim C, Young LJ, Newport DJ, Mletzko T, Miller AH, Nemeroff CB. Lower CSF oxytocin concentrations in women with a history of childhood abuse. *Mol Psychiatry.* 2009; 14:954–958. [PubMed: 18957940]
- Higashida H, Yokoyama S, Kikuchi M, Munesue T. CD38 and its role in oxytocin secretion and social behavior. *Horm Behav.* 2012; 61:351–358. [PubMed: 22227279]
- Insel TR. The challenge of translation in social neuroscience: a review of oxytocin, vasopressin, and affiliative behavior. *Neuron.* 2010; 65:768–779. [PubMed: 20346754]
- Jokinen J, Chatzittofis A, Hellstrom C, Nordstrom P, Uvnas-Moberg K, Asberg M. Low CSF oxytocin reflects high intent in suicide attempters. *Psychoneuroendocrinology.* 2012; 37:482–490. [PubMed: 21852050]
- Meyer-Lindenberg A, Domes G, Kirsch P, Heinrichs M. Oxytocin and vasopressin in the human brain: social neuropeptides for translational medicine. *Nat Rev Neurosci.* 2011; 12:524–538. [PubMed: 21852800]
- Modahl C, Green L, Fein D, Morris M, Waterhouse L, Feinstein C, Levin H. Plasma oxytocin levels in autistic children. *Biol Psychiatry.* 1998; 43:270–277. [PubMed: 9513736]
- Modi ME, Young LJ. The oxytocin system in drug discovery for autism: animal models and novel therapeutic strategies. *Horm Behav.* 2012; 61:340–350. [PubMed: 22206823]
- Parker KJ, Hoffman CL, Hyde SA, Cummings CS, Maestripieri D. Effects of age on cerebrospinal fluid oxytocin levels in free-ranging adult female and infant rhesus macaques. *Behav Neurosci.* 2010; 124:428–433. [PubMed: 20528088]
- Rosenblum LA, Smith EL, Altemus M, Scharf BA, Owens MJ, Nemeroff CB, Gorman JM, Coplan JD. Differing concentrations of corticotropin-releasing factor and oxytocin in the cerebrospinal fluid of bonnet and pigtail macaques. *Psychoneuroendocrinology.* 2002; 27:651–660. [PubMed: 12084658]

- Rothbart MK. Measurement of temperament in infancy. *Child Development*. 1981; 52:569–578.
- Sparrow, SS.; Cicchetti, DV.; Balla, DA. Vineland Adaptive Behavior Scales. 2nd. American Guidance Service; 2005.
- Veening JG, Barendregt HP. The regulation of brain states by neuroactive substances distributed via the cerebrospinal fluid; a review. *Cerebrospinal Fluid Res*. 2010; 7:1. [PubMed: 20157443]
- Winslow JT, Noble PL, Lyons CK, Sterk SM, Insel TR. Rearing effects on cerebrospinal fluid oxytocin concentration and social buffering in rhesus monkeys. *Neuropsychopharmacology*. 2003; 28:910–918. [PubMed: 12700704]

Figure 1. Neonatal CSF OT levels positively correlate with parental reports of infant behavior
(A) Neonatal CSF OT levels correlate with Infant Behavioral Questionnaire scores for “crying for contact” at 3 and 6 months **(B)** Neonatal CSF OT levels positively correlate with parent report of prosocial behaviors at 6 months of age

Table 1
Participant Demographics and Perinatal Characteristics

		Number of Participants
Sex	Females	6
	Males	12
Ethnicity	Hispanic	8
	Non-Hispanic	10
Insurance Status	Public coverage	8
	Private coverage	10
Gestation	Pre-term	9
	Term	9
Birth History	Vaginal	10
	C-section	8
Pitocin Exposure	Yes	7
	No	11